

CORRIGENDUM

DOI: 10.3892/mmr.2018.9415

Downregulation of δ opioid receptor by RNA interference enhances the sensitivity of BEL/FU drug-resistant human hepatocellular carcinoma cells to 5-FU

BO TANG, ZHIGAO HU, YANG LI, SHENGGUANG YUAN, ZHENRAN WANG, SHUIPING YU and SONGQING HE

Mol Med Rep 13: 59-66, 2016; DOI: 10.3892/mmr.2015.4511

After the publication of the article, the authors realized that they had inadvertently included the incorrect data for the 'siDOR+5-fu group' in the flow cytometric plots featured in Fig. 3A. A new version of Fig. 3 is provided, which contains the correct data for the 'siDOR+5-fu group' experiment.

The authors regret this error, and apologize to the readers for any inconvenience caused.



This work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) License.




Figure 3. Effects of DOR gene silencing (siDOR) on the apoptosis of BEL/FU cells. (A) Flow cytometric analysis using Annexin V-fluorescein isothiocyanate double staining. (B) Histogram presenting the apoptotic rates of BEL/FU cells. Data are from three independent experiments and are presented as the mean \pm standard deviation. ** $P < 0.01$, vs. siControl. DOR, δ opioid receptor; BEL/FU, BEL-7402/5-fluorouracil; PI, propidium iodide.