

Let-7a-5p may participate in the pathogenesis of diabetic nephropathy through targeting HMGA2

TAO WANG¹, HUA ZHU¹, SHUFANG YANG² and XIAOQIANG FEI²

¹Department of Clinical Laboratory, Taixing City Second People's Hospital, Taixing, Jiangsu 225400;

²Department of Endocrinology, Taizhou People's Hospital, Taizhou, Jiangsu 225300, P.R. China

Received May 17, 2018; Accepted October 26, 2018

DOI: 10.3892/mmr.2019.10057

Abstract. Diabetic nephropathy (DN) is one of the most common complications of diabetes mellitus (DM), and has been demonstrated as one of the major causes of renal failure. In a previous study, it was noted that microRNA let-7a-5p was downregulated in DN; however, the underlying mechanism requires additional investigation. The aim of the present study was to investigate the roles of let-7a-5p in the pathogenesis of DN and its associated mechanism. The renal tissues of db/db and db/m mice, and renal mesangial cells treated with high concentrations of glucose were obtained; reverse transcription-quantitative polymerase chain reaction, and western blot analysis were applied to detect the expression of let-7a-5p and high-mobility group AT-hook 2 (HMGA2) *in vivo* and *in vitro*. In addition, renal mesangial cells cultured under high-glucose conditions (20 and 30 mmol/l) were transfected with either let-7a-5p mimics or let-7a-5p inhibitors. The effects of let-7a-5p on the proliferation and apoptosis of renal mesangial cells were examined using CCK-8 and flow cytometry methods. Additionally, cells were collected and the expression of phosphoinositide 3-kinase (PI3K), phosphorylated protein kinase B (p-AKT) and HMGA2 was analyzed with western blot analysis. Finally, a dual luciferase reporter assay was performed to confirm whether HMGA2 was a direct target of let-7a-5p. Let-7a-5p was significantly downregulated and HMGA2 was markedly upregulated in the tissue samples of DN mice and renal mesangial cells cultured under high-glucose conditions. In addition, transfection of let-7a-5p mimics induced a significant decrease in the proliferation and increase in the apoptosis of renal mesangial cells cultured under high-glucose conditions; transfection of let-7a-5p inhibitors exhibited the opposite effects. Furthermore, transfection of let-7a-5p mimics also led to the inhibition of the PI3K-AKT signaling pathway; transfection

of let-7a-5p inhibitors may activate the PI3K-AKT signaling pathway through the increase in PI3K and AKT levels. Finally, a dual luciferase reporter assay confirmed that HMGA2 is a direct target of let-7a-5p. Let-7a-5p was downregulated in DN and may participate in the pathogenesis of DN via regulating HMGA2 expression and the PI3K-AKT signaling pathway.

Introduction

Diabetic nephropathy (DN) is one of the most common complications of diabetes mellitus (DM), and is known as one of the major causes of renal failure (1-3). Previously, the incidence of DN was demonstrated to be increasing yearly; according to the data from the World Health Organization, the population of patients with diabetes will reach 370 million by the year of 2025, and 30% of these patients will develop DN (4). Generally, the distinct pathological changes of DN include aberrant mesangial cell proliferation, which causes the widening of the mesangial region and abnormal accumulation of the extracellular matrix, which results in glomerular sclerosis and tubulointerstitial fibrosis. DN eventually leads to the development of chronic renal failure, which may seriously affect the quality of life of patients (5-7). At present, anti-DN therapies include the control of blood glucose and blood pressure, and the inhibition of the renin-angiotensin system; however, these methods only delay the progression of DN; such therapies are not able to reverse the progression of DN to the stages of renal function failure (7-9). Therefore, there is an urgent requirement to identify novel therapeutic targets for the management and prevention of DN.

The pathogenesis of DN remains unclear. In previous years, increasing evidence has indicated that microRNA (miRNAs) are involved in the progression of a number of diseases. miRNAs are a family of small non-coding RNAs with a length of 18-25 nucleotides. miRNAs bind to the 3'-untranslated region (3'UTR) of their target mRNAs, thereby suppressing the expression of their target genes (10-12). A variety of miRNAs have been demonstrated to participate in the pathogenesis of a number of types of disease, including cancer, autoimmune diseases, cardiovascular diseases, and acute and chronic renal disease (13-15). However, the underlying mechanism of miRNAs in renal-associated disease requires additional study.

In a previous study, it was suggested that miRNA let-7a-5p was aberrantly expressed in DN (16). The present study aimed

Correspondence to: Dr Xiaoqiang Fei, Department of Endocrinology, Taizhou People's Hospital, 399 Hailing Road, Taizhou, Jiangsu 225300, P.R. China
E-mail: feixiaoqiang0356@outlook.com

Key words: microRNA, let-7a-5p, diabetic nephropathy, proliferation, apoptosis, high-mobility group AT-hook 2

to identify the roles of let-7a-5p in the pathogenesis of DN using *in vivo* and *in vitro* models. The present study may provide novel evidence for the diagnosis and treatment of DN.

Materials and methods

Establishment of DM animal models. A total of 32 4-week-old male C57BL/KsJ-db/db mice with type II DM and an additional 32 4-week-old male db/m mice were purchased from the Animal Center of Nanjing Medical University (Nanjing, China) and included in the present study. Mice were maintained under conventional conditions with 12 h light/dark cycle at 20–22°C and were provided with standard chow and water *ad libitum*. Prior to sacrifice, mice underwent fasting for 12 h. The bilateral kidneys were then collected following laparotomy in each mouse; the connective tissues and blood vessels on the renal hilum were removed. Next, the renal specimens of one kidney from each bilateral pair were fixed in 10% neutral formalin at room temperature for 48 h, embedded in paraffin and sliced into 2- μ m sections. Then, the tissue sections underwent H&E staining and were analyzed under a light microscope (magnification, x200) to determine pathological changes within the renal tissues. Simultaneously, renal specimens of the other kidney of each bilateral pair were preserved in liquid nitrogen for RNA extraction. The present study as approved by the Ethical Committee of Taixing City Second People's Hospital.

High-glucose-stimulated renal mesangial cell model. Mouse renal mesangial cells were purchased from the Cell Bank of Type Culture Collection of Chinese Academy of Sciences (Shanghai, China; cat. no., GNM21). The renal mesangial cells were cultured at 37°C in Dulbecco's modified Eagle's medium (DMEM; Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.), and glucose at a concentration of 20 mmol/l, in an atmosphere containing 5% CO₂. After 24 h, the cells were collected for subsequent analysis.

Cell transfection. The mmu-let-7a-5p inhibitors and mmu-let-7a-5p mimics oligonucleotides were purchased from Shanghai GenePharma Co., Ltd. (Shanghai, China). Renal mesangial cells treated with 0, 20 or 30 mmol/l glucose were transfected with 50 nM mmu-let-7a-5p inhibitors or mmu-let-7a-5p mimics mixed with Lipofectamine RNAi Max (Thermo Fisher Scientific, Inc.). Cells were then cultured at 37°C in DMEM supplemented with 10% FBS in an atmosphere containing 5% CO₂ for 48 h. Cells were harvested at 48 h for the subsequent analysis. The sequences of the oligonucleotides were: mmu-let-7a-5p mimics, 5'-UGAGGUAGUAGGUUGUAUAGUU-3'; mmu-let-7a-5p mimics NC, 5'-UUCUCCGAACGUGUCACGUTT-3'; mmu-let-7a-5p inhibitors, 5'-ACUAUACAACCUACUACC UCA-3'; mmu-let-7a-5p inhibitors NC, 5'-CAGUACUUUUGU GUAGUACAA-3'.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Total RNA was isolated from cells using TRIzol® reagent (Invitrogen; Thermo Fisher Scientific, Inc.) and the reverse transcription was performed with the PrimeScript™ RT reagent kit (Takara Biotechnology Co., Ltd., Dalian, China)

with the temperature of 37°C for 15 min and 85°C for 5 sec. RT-qPCR was conducted with a SYBR ExScript RT-PCR kit (Takara Biotechnology Co., Ltd., Dalian, China) on an ABI 7500 Real-Time PCR System (Applied Biosystems; Thermo Fisher Scientific, Inc.). The thermocycling conditions were: Initial denaturation, 95°C for 30 sec; followed by 40 cycles of denaturation at 95°C for 5 sec and annealing at 60°C for 30 sec. The primers were synthesized by Sangon Biotech Co., Ltd. (Shanghai, China). The relative expression of high-mobility group AT-hook 2 (HMGA2) in each sample was normalized to that of GAPDH using the 2^{- $\Delta\Delta$ C_q} method (17). The expression of let-7a-5p was determined using the Hairpin-it™ miRNAs qPCR Quantitation kit (Shanghai GenePharma Co., Ltd.). U6 was applied for the normalization of miRNA expression. The sequences of the primers used were: Let-7a-5p, forward 5'-GCC GCTGAGGTAGTAGGTTGTA-3', reverse 5'-GTGCAGGGT CCGAGGT-3'; HMGA2, forward 5'-CAGCAGCAAGAACCA ACCG-3', reverse 5'-TGTTGTGGCCATTCCTAGGT-3'; PI3K, forward 5'-GAAATCTCCTGGGATGTGTCGT-3', reverse 5'-ATCTGGTGGCTCTCGGAGTAA-3'; AKT, forward 5'-GATGGAGGCCAGGGTACAAA-3', reverse, 5'-GCAGCG ACACCACAAAAATGA-3'; GAPDH, forward 5'-TCAACG GATTGCTCGTATTG-3', reverse, 5'-TGGGTGGAATCA TATTGGAAC-3'; U6, forward 5'-AACGCTTCACGAATT TCGCT-3', reverse 5'-AACGCTTCACGAATTTGCGT-3'.

Cell proliferation analysis. Renal mesangial cells were plated at a density of 5,000 cells/well in 96-well plates. A Cell Counting Kit-8 (CCK-8) assay was performed at 48 h after transfection to determine cell viability using a CCK-8 proliferation assay kit (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) according to the manufacturer's protocol.

Cell apoptosis analysis. For the analysis of apoptosis, 48 h after transfection or following treatment with the protein kinase B (AKT) inhibitor, MK2206 (10 μ mol/l), cells were stained using a propidium iodide/Annexin V-fluorescein isothiocyanate apoptosis detection kit (BD Biosciences, San Jose, CA, USA). Briefly, cells were collected, and 5 μ l Annexin V-FITC and propidium iodide (PI) solutions were mixed with the cells at room temperature, and incubated on ice in the dark for 30 min. Then the apoptotic rate and cell cycle in each sample were analyzed with a BD FACSVerse flow cytometer (BD Biosciences) according to the manufacturer's protocol. The data was analyzed by FlowJo version 8.7 (FlowJo LLC, Ashland, OR, USA).

Western blot analysis. Cells were lysed with radioimmunoprecipitation assay buffer (Beyotime Institute of Biotechnology, Haimen, China) and then concentration of the proteins were quantified by Enhanced BCA Protein Assay kit (Beyotime Institute of Biotechnology). Then SDS-PAGE was performed to separate the extracted proteins using 8% gel, and 30 μ g protein loaded into each lane. Then, the proteins were transferred onto polyvinylidene fluoride membranes, which were then blocked with 5% non-fat milk at room temperature for 1 h and incubated with primary antibodies [anti-phosphoinositide 3-kinase (PI3K, ab151549; 1:1,000), anti-AKT (ab8805; 1:1,000), anti-phosphorylated (p)-AKT (ab38449; 1:1,000), anti-HMGA2 (ab97276; 1:500) and anti-GAPDH (ab9485;

1:2,000) all from Abcam, Cambridge, MA, USA] overnight at 4°C. On day 2, the membranes were washed and incubated with a HRP-conjugated secondary antibody (A0208; 1:5,000; Beyotime Institute of Biotechnology) at room temperature for 45 min, and then washed and incubated with BeyoECL Plus (Beyotime Institute of Biotechnology) at room temperature for 1 min. The bands were visualized using a ChemiDoc™ XRS+ imaging system (Bio-Rad Laboratories, Inc., Hercules, CA, USA). GAPDH was applied as the internal control. Densitometric analysis was performed with Image Lab software version 3.1.1 (Bio-Rad Laboratories, Inc.).

Bioinformatics. The prediction of target gene of mmu-let-7a-5p was performed using the online bioinformatic tool Target Scan (<http://www.targetscan.org>) (18) and miRanda (<http://www.microrna.org/>) (19) as previously described (18,19). HMGA2 revealed a potential binding site for mmu-let-7a-5p in the 3'-UTR of HMGA2.

Dual luciferase reporter assay. The wild-type HMGA2 3'UTR (HMGA2-3'UTR) and mutant HMGA2 3'UTR (HMGA2-MUT) regions containing the mmu-let-7a-5p binding site were synthesized and cloned into the p-MIR-reporter plasmid (Thermo Fisher Scientific, Inc.). 293 cells (cat. no., GNHu43; Cell Bank of Type Culture Collection of Chinese Academy of Sciences, Shanghai, China) were then seeded onto 12-well plates at a density of 1×10^5 cells/well and transfected with HMGA2-3'UTR or HMGA2-MUT, in addition to either let-7a-5p mimics or NC using the Lipofectamine RNAi Max kit. Cells were harvested 48 h after transfection and the luciferase activities of the experimental groups were detected using a dual-luciferase reporter system (Dual Luciferase Reporter Gene Assay kit, Beyotime Institute of Biotechnology). Renilla luciferase was used as the internal control.

Statistical analysis. All statistical analyses were performed using SPSS 19.0 (IBM Corp., Armonk, NY, USA). Data are presented as the mean \pm standard deviation; a two-tailed independent samples t-test was performed for the comparison between two groups and one-way analysis of variance was applied for the comparison of multiple groups followed by a Dunnett's post-hoc test. Pearson correlation coefficient was used for the correlation analysis. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Downregulation of let-7a-5p and upregulation of HMGA2 in DN mice in vivo. Firstly, to explore the roles of let-7a-5p in DN, DN mice models were established. The pathological features of the kidney tissues of DN and the control mice are presented in Fig. 1A. The expression levels of let-7a-5p were examined using RT-qPCR. As demonstrated in Fig. 1B, let-7a-5p expression was significantly downregulated in the kidneys of DN mice at the mRNA level ($P < 0.05$). In addition, online computational prediction tools (TargetScan and miRanda) were applied, which predicted HMGA2 as a target gene of let-7a-5p. It has been suggested that the overexpression of HMGA2 may promote the epithelial-to-mesenchymal transition of tubular epithelial cells (20), suggesting that HMGA2 may be involved

in the pathogenesis of DN. Therefore, in the present study, the expression of HMGA2 in DN tissue and control was examined. It was observed that the expression of HMGA2 was significantly increased at the mRNA and protein levels, as detected by RT-qPCR and western blot analysis (Fig. 1B and C). Finally, the present study identified that the expression levels of let-7a-5p in DN mice were negatively correlated with that of HMGA2 (Fig. 1D; $P = 0.0088$; $r = -0.4555$).

Effects of high-glucose conditions on the expression of let-7a-5p and HMGA2 in renal mesangial cells. Renal mesangial cells were cultured under high-glucose conditions for 24 h in the present study; the effects of high-glucose conditions on the expression of let-7a-5p and HMGA2 were then examined by RT-qPCR. It was observed that high-glucose conditions induced significant decreases in the expression levels of let-7a-5p (Fig. 2A) and markedly increased the expression of HMGA2 (Fig. 2B), in a dose-dependent manner.

Let-7a-5p may affect the proliferation and apoptosis of renal mesangial cells under high-glucose conditions in vitro. To additionally explore the roles of let-7a-5p in DN, renal mesangial cells were transfected with let-7a-5p mimics or inhibitors; the effects of let-7a-5p on the proliferation and apoptosis of renal mesangial cells under high-glucose conditions were examined using CCK-8 and flow cytometry analyses. As indicated in Fig. 3A, the expression of let-7a-5p was significantly upregulated in let-7a-5p mimic-transfected cells, and markedly downregulated in let-7a-5p inhibitor-transfected cells; in addition, transfection with let-7a-5p mimics induced a significant decrease in the proliferation and an increase in the apoptosis of renal mesangial cells cultured under high-glucose conditions; transfection with let-7a-5p inhibitors exhibited opposing effects (Fig. 3B-D; $P < 0.05$).

Let-7a-5p may affect the expression of HMGA2, PI3K and p-AKT in renal mesangial cells under high-glucose conditions. To additionally explore the underlying mechanism of let-7a-5p in DN, the effects of let-7a-5p mimics or inhibitors on the expression of HMGA2, PI3K, AKT and p-AKT were examined. As demonstrated in Fig. 4, transfection with let-7a-5p mimics resulted in decreased expression levels of HMGA2 and inhibition of the PI3K/AKT signaling pathway, which was indicated by decreased expression levels of PI3K, AKT and p-AKT. Conversely, transfection with let-7a-5p inhibitors appeared to activate the PI3K-AKT signaling pathway.

HMGA2 is a direct target of let-7a-5p in vitro. As aforementioned, using the online computational prediction tools TargetScan and miRanda, HMGA2 was predicted as a target gene of let-7a-5p (Fig. 5A). Finally, a dual luciferase reporter assay was performed to investigate whether let-7a-5p may directly target HMGA2. As demonstrated in Fig. 5, transfection of let-7a-5p mimics induced significant decreases in the luciferase activities within wild-type HMGA2 3'-UTR (HMGA2-3'-UTR)-transfected 293 cells, while transfection with let-7a-5p mimics exhibited no significant effects on mutant HMGA2 3'-UTR (HMGA2-MUT)-transfected cells (Fig. 5B; $P < 0.01$). These results indicated that HMGA2 is a direct target of let-7a-5p.




Figure 1. Downregulation of let-7a-5p and upregulation of HMGA2 in DN mice *in vivo*. (A) The pathological features of kidney tissues of DN and control mice; (B) Relative expression levels of let-7a-5p and HMGA2 were examined by reverse transcription quantitative polymerase chain reaction. (C) Relative expression of HMGA2 was examined by western blot analysis. (D) Correlation between the levels of let-7a-5p and HMGA2 in DN mice. ** $P < 0.01$. DN, diabetic nephropathy; HMGA2, high-mobility group AT-hook 2.

Discussion

In previous years, increasing evidence has indicated that

miRNAs may participate in the pathogenesis of DN. For example, it has been observed that miR-29c may promote the development of DN via targeting tristetraprolin (21). In addition,




Figure 2. Downregulation of let-7a-5p and upregulation of HMGA2 in renal mesangial cell treated with high-glucose conditions. (A) Relative expression of let-7a-5p measured by RT-qPCR. (B) Relative expression of HMGA2 measured by RT-qPCR. * $P<0.05$ and ** $P<0.01$ vs. control. HMGA2, high-mobility group AT-hook 2; RT-qPCR, reverse transcription quantitative polymerase chain reaction.

miR-27a has been observed to regulate renal function through the activation of β -catenin in DN (22). Furthermore, suppression of miR-217 may ameliorate high-glucose-induced podocyte injury by inhibiting the apoptosis of cells (23). The present study focused on the roles of miRNA let-7a-5p in DN by performing *in vitro* and *in vivo* analyses. It was observed that the expression of let-7a-5p was significantly downregulated in the kidney tissues of DN mice and renal mesangial cells treated cultured under high-glucose conditions compared with the controls, which was consistent with previous data (24). Taken together, these results indicated that let-7a-5p was downregulated in DN, suggesting that let-7a-5p may serve as a biomarker for the diagnosis of DN.

At present, the majority of studies investigating let-7a are cancer-associated; let-7a-5p has been described previously in a number of studies, either as tumor suppressor or as an oncogenic miRNA. In the field of renal diseases, let-7a has been described as a tumor suppressor in renal cell carcinoma via the targeting of v-myc avian myelocytomatosis viral oncogene

homolog expression (25). In addition, it has also been observed that the detection of let-7 miRNAs in urine-derived supernatant samples may be a potential approach for the diagnosis of renal cell carcinoma. In the present study, renal mesangial cells were treated with high concentrations of glucose *in vitro* and it was observed that let-7a-5p was significantly downregulated in renal mesangial cells *in vitro*. Furthermore, transfection with let-7a-5p mimics induced significant decreases in the proliferation and promoted the apoptosis of renal mesangial cells cultured under high-glucose conditions; transfection with let-7a-5p inhibitors exhibited the opposite effects. These results conclusively indicated a pathological role of let-7a-5p in mesangial cell hypertrophy.

miRNAs exert their functions through the silencing of target genes. Using the online computational prediction tools TargetScan and miRanda, HMGA2 was predicted as a target gene of let-7a-5p in the present study, and Mayr *et al* (26) demonstrated that let-7a directly targets HMGA2. HMGA2 has been recognized as a non-histone chromosomal protein, and it binds to DNA and regulates the transcription process of its target genes (26). HMGA2 has been suggested to be involved in a number of biological processes, including cell proliferation, differentiation, embryonic development and carcinogenesis (26). It has been demonstrated previously in a numerous studies that HMGA2 is upregulated in renal diseases (27-29). However, whether HMGA2 may regulate the proliferation and apoptosis of renal mesangial cells, which leads to the development of DN, remains unclear. As it has been previously described that DN was induced by the over-proliferation of mesangial cells (30), it was hypothesized that downregulation of let-7a-5p may lead to the aberrant expression of HMGA2, which may serve critical roles in the pathogenesis.

Therefore, to explore the association between let-7a-5p and HMGA2 in DN, a series of *in vitro* and *in vivo* experiments were performed in the present study. It was observed that the expression levels of HMGA2 were markedly increased in the kidney of DN mice, and in high-glucose-treated mesangial cells. In addition, transfection with let-7a-5p mimics of high-glucose-treated mesangial cells induced a significant decrease in the expression of HMGA2; transfection with let-7a-5p inhibitors exhibited the opposite results. Finally, the results of the dual luciferase reporter assay suggested that HMGA2 is a direct target of let-7a-5p, which was consistent with the observation of Lee and Dutta (31). Taken together, these results indicated that let-7a-5p may be involved in the pathogenesis of DN by targeting HMGA2 expression.

Considerable evidence indicates that the PI3K/AKT signaling pathway is involved in the control of various cellular processes, including that of DN. For example, notoginsenoside R1 may ameliorate podocyte and renal injuries in rats with DN by activating the PI3K/AKT signaling pathway (32). Jiangtang decoction may decrease inflammation associated with DN through the regulation of the PI3K/AKT signaling pathway (33). The association between HMGA2 and the PI3K/AKT signaling pathway has been discussed previously: Tan *et al* (34) observed that HMGA2 may induce cell proliferation in acute myeloid leukemia cells via regulating the PI3K/AKT/mechanistic target of rapamycin signaling pathway; Cui *et al* (35) revealed that miR-337 may regulate




Figure 3. Let-7a-5p may affect the proliferation and apoptosis of renal mesangial cell under high-glucose conditions *in vitro*. (A) Effect of let-7a-5p mimics and inhibitors on the expression of let-7a-5p in renal mesangial cells. (B) Effect of let-7a-5p mimics and inhibitors on the proliferation of renal mesangial cell under high-glucose conditions *in vitro* measured by the Cell Counting Kit-8 method. (C) Effect of let-7a-5p mimics and inhibitors on the apoptosis of renal mesangial cell under high-glucose conditions *in vitro* measured by flow cytometry. (D) Quantification of the flow cytometry results. ** $P < 0.01$ vs. control. NC, negative control; OD, optical density; PI, propidium iodide; FITC, fluorescein isothiocyanate.




Figure 4. Let-7a-5p may affect the expressions of HMGA2 and PI3K/AKT in renal mesangial cells under high-glucose conditions. (A) Effect of let-7a-5p mimics on the mRNA expression levels of HMGA2 and PI3K/AKT in renal mesangial cells under high-glucose conditions. (B) Effect of let-7a-5p inhibitors on the mRNA expressions of HMGA2 and PI3K/AKT in renal mesangial cell under high-glucose conditions. (C) Effect of let-7a-5p inhibitors on the protein expressions of HMGA2 and PI3K/AKT in renal mesangial cell under high-glucose conditions. * $P < 0.05$ and ** $P < 0.01$ vs. control. HMGA2, high-mobility group AT-hook 2; PI3K, phosphoinositide 3-kinase; AKT, protein kinase B; p, phosphorylated; NC, negative control.




Figure 5. HMGA2 is a direct target of let-7a-5p *in vitro*. (A) HMGA2 was predicted as a target gene of mmu-let-7a-5p; (B) Relative activities of the luciferases in different groups. ** $P < 0.01$. HMGA2, high-mobility group AT-hook 2; WT, wild type; NC, negative control.

the PI3K/AKT signaling pathway to inhibit hepatocellular carcinoma progression via targeting HMGA2. The present study explored whether the PI3K/AKT signaling pathway was involved in the let-7a-5p/HMGA2-induced anti-DN effects within renal mesangial cells, and it was observed that the overexpression of let-7a-5p promoted the expression of PI3K and p-AKT in high-glucose-treated mesangial cells. Taken together, the results of the present study not only revealed the

important role of the let-7a-5p/HMGA2/PI3K/AKT signaling pathway in the development of DN, but also suggested the potential role of let-7a-5p as a potential therapeutic target for the treatment of DN.

In conclusion, the data from the present study demonstrated that let-7a-5p was downregulated in DN and that let-7a-5p may participate in the pathogenesis of DN by targeting the HMGA2/PI3K/AKT signaling pathway. The results present

let-7a-5p as a potential diagnostic marker and therapeutic targets for the diagnosis and treatment of DN.

Acknowledgements

Not applicable.

Funding

The present study was funded by grant from The Medical Clinical Science and Technology Development Fund Project of Jiangsu University (grant no. JLY20160150).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

TW performed experiments and wrote the article. HZ contributed to the design of the study and revised the manuscript. SY designed and performed the experiments and analyzed the data. XF contributed to the design of the study and revised the manuscript. All the authors have seen and approved the manuscript.

Ethics approval and consent to participate

The present animal study was approved by the Ethical Committee of Taixing City Second People's Hospital.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Barman S, Pradeep SR and Srinivasan K: Zinc supplementation alleviates the progression of diabetic nephropathy by inhibiting the overexpression of oxidative-stress-mediated molecular markers in streptozotocin-induced experimental rats. *J Nutr Biochem* 54: 113-129, 2018.
- Dissanayake AM, Wheldon MC, Ahmed J and Hood CJ: Extending metformin use in diabetic kidney disease: A pharmacokinetic study in stage 4 diabetic nephropathy. *Kidney Int Rep* 2: 705-712, 2017.
- Khitan ZJ, Tzamaloukas AH, Brodsky S and Shapiro JJ: Dihydropyridine calcium channel blockers in the elderly with diabetic nephropathy: Are they safe? *J Clin Hypertens (Greenwich)* 20: 203-204, 2018.
- Li J, Yang S, Li X, Liu D, Wang Z, Guo J, Tan N, Gao Z, Zhao X, Zhang J, *et al*: Role of endoplasmic reticulum stress in disuse osteoporosis. *Bone* 97: 2-14, 2017.
- Montero RM, Herath A, Qureshi A, Esfandiari E, Pusey CD, Frankel AH and Tam FWK: Defining phenotypes in diabetic nephropathy: A novel approach using a cross-sectional analysis of a single centre cohort. *Sci Rep* 8: 53, 2018.
- Sagoo MK and Gnudi L: Diabetic nephropathy: Is there a role for oxidative stress? *Free Radic Biol Med* 116: 50-63, 2018.
- Talsma DT, Katta K, Ettema MAB, Kel B, Kusche-Gullberg M, Daha MR, Stegeman CA, van den Born J and Wang L: Endothelial heparan sulfate deficiency reduces inflammation and fibrosis in murine diabetic nephropathy. *Lab Invest* 98: 427-438, 2018.
- Wei PZ, Kwan BC, Chow KM, Cheng PM, Luk CC, Li PK and Szeto CC: Urinary mitochondrial DNA level is an indicator of intra-renal mitochondrial depletion and renal scarring in diabetic nephropathy. *Nephrol Dial Transplant* 33: 784-788, 2018.
- Xu Z and Fan J: Islet transplantation promotes podocyte regeneration in a model of diabetic nephropathy. *Turk J Med Sci* 47: 1925-1930, 2017.
- Cai M, He H, Jia X, Chen S, Wang J, Shi Y, Liu B, Xiao W and Lai S: Genome-wide microRNA profiling of bovine milk-derived exosomes infected with *Staphylococcus aureus*. *Cell Stress Chaperones* 23: 663-672, 2018.
- Kaul V, Weinberg KI, Boyd SD, Bernstein D, Esquivel CO, Martinez OM and Krams SM: Dynamics of viral and host immune cell microRNA expression during acute infectious mononucleosis. *Front Microbiol* 8: 2666, 2018.
- Liu L, Zhang X, Nan C, Zhao Z, Ma S, Li W, Hu H and Liang Z: MicroRNA-182 targets protein phosphatase 1 regulatory inhibitor subunit 1C in glioblastoma. *Oncotarget* 8: 114677-114684, 2017.
- Tang Y, Zhang L, Tu T, Li Y, Murray D, Tu Q and Chen JJ: MicroRNA-99a is a novel regulator of KDM6B-mediated osteogenic differentiation of BMSCs. *J Cell Mol Med* 22: 2162-2176, 2018.
- Wang Y, Pang X, Wu J, Jin L, Yu Y, Gobin R and Yu J: MicroRNA hsa-let-7b suppresses the odonto/osteogenic differentiation capacity of stem cells from apical papilla by targeting MMP1. *J Cell Biochem* 119: 6545-6554, 2018.
- Wen X, Han XR, Wang YJ, Wang S, Shen M, Zhang ZF, Fan SH, Shan Q, Wang L, Li MQ, *et al*: MicroRNA-421 suppresses the apoptosis and autophagy of hippocampal neurons in epilepsy mice model by inhibition of the TLR/MyD88 pathway. *J Cell Physiol* 233: 7022-7034, 2018.
- Yan N, Wen L, Peng R, Li H, Liu H, Peng H, Sun Y, Wu T, Chen L, Duan Q, *et al*: Naringenin ameliorated kidney injury through Let-7a/TGFBR1 signaling in diabetic nephropathy. *J Diabetes Res* 2016: 8738760, 2016.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Wang LK, Xie XN, Song XH, Su T, Chang XL, Xu M, Liang B and Huang DY: Upregulation of miR-200b inhibits hepatocellular carcinoma cell proliferation and migration by targeting HMGB3 protein. *Technol Cancer Res Treat* 17: 1533033818806475, 2018.
- Yan L, Li J, Wu Q and Chen L: Specific miRNA expression profile in the blood serum of cardiac myxoma patients. *Oncol Lett* 16: 4235-4242, 2018.
- Liu H, Wang X, Liu S, Li H, Yuan X, Feng B, Bai H, Zhao B, Chu Y and Li H: Effects and mechanism of miR-23b on glucose-mediated epithelial-to-mesenchymal transition in diabetic nephropathy. *Int J Biochem Cell Biol* 70: 149-160, 2016.
- Guo J, Li J, Zhao J, Yang S, Wang L, Cheng G, Liu D, Xiao J, Liu Z and Zhao Z: MiRNA-29c regulates the expression of inflammatory cytokines in diabetic nephropathy by targeting tristetrin. *Sci Rep* 7: 2314, 2017.
- Zhou Z, Wan J, Hou X, Geng J, Li X and Bai X: MicroRNA-27a promotes podocyte injury via PPARγ-mediated β-catenin activation in diabetic nephropathy. *Cell Death Dis* 8: e2658, 2017.
- Sun J, Li ZP, Zhang RQ and Zhang HM: Repression of miR-217 protects against high glucose-induced podocyte injury and insulin resistance by restoring PTEN-mediated autophagy pathway. *Biochem Biophys Res Commun* 483: 318-324, 2017.
- Peng R, Liu H, Peng H, Zhou J, Zha H, Chen X, Zhang L, Sun Y, Yin P, Wen L, *et al*: Promoter hypermethylation of let-7a-3 is relevant to its down-expression in diabetic nephropathy by targeting UHRF1. *Gene* 570: 57-63, 2015.
- Liu Y, Yin B, Zhang C, Zhou L and Fan J: Hsa-let-7a functions as a tumor suppressor in renal cell carcinoma cell lines by targeting c-myc. *Biochem Biophys Res Commun* 417: 371-375, 2012.
- Mayr C, Hemann MT and Bartel DP: Disrupting the pairing between let-7 and Hmga2 enhances oncogenic transformation. *Science* 315: 1576-1579, 2007.
- Liu Y, Fu QZ, Pu L, Meng QG, Liu XF, Dong SF, Yang JX and Lv GY: HMGA2 expression in renal carcinoma and its clinical significance. *J Med Biochem* 34: 338-343, 2015.

28. Liu Y, Fu QZ, Pu L, Song LL, Wang YY, Liu J, Wang ZL and Wang ZM: Effect of RNA interference of the expression of HMGA2 on the proliferation and invasion ability of ACHN renal cell carcinoma cells. *Mol Med Rep* 16: 5107-5112, 2017.
29. Na N, Si T, Huang Z, Miao B, Hong L, Li H, Qiu J and Qiu J: High expression of HMGA2 predicts poor survival in patients with clear cell renal cell carcinoma. *Onco Targets Ther* 9: 7199-7205, 2016.
30. Wang M, Yao D, Wang S, Yan Q and Lu W: Long non-coding RNA ENSMUST00000147869 protects mesangial cells from proliferation and fibrosis induced by diabetic nephropathy. *Endocrine* 54: 81-92, 2016.
31. Lee YS and Dutta A: The tumor suppressor microRNA let-7 represses the HMGA2 oncogene. *Genes Dev* 21: 1025-1030, 2007.
32. Huang G, Lv J, Li T, Huai G, Li X, Xiang S, Wang L, Qin Z, Pang J, Zou B and Wang Y: Notoginsenoside R1 ameliorates podocyte injury in rats with diabetic nephropathy by activating the PI3K/Akt signaling pathway. *Int J Mol Med* 38: 1179-1189, 2016.
33. Hong JN, Li WW, Wang LL, Guo H, Jiang Y, Gao YJ, Tu PF and Wang XM: Jiangtang decoction ameliorate diabetic nephropathy through the regulation of PI3K/Akt-mediated NF- κ B pathways in KK-Ay mice. *Chin Med* 12: 13, 2017.
34. Tan L, Wei X, Zheng L, et al.: Amplified HMGA2 promotes cell growth by regulating Akt pathway in AML. *J Cancer Res Clin Oncol* 142: 389-399, 2016.
35. Cui H, Song R, Wu J, Wang W, Chen X and Yin J: MicroRNA-337 regulates the PI3K/AKT and Wnt/beta-catenin signaling pathways to inhibit hepatocellular carcinoma progression by targeting high-mobility group AT-hook 2. *Am J Cancer Res* 8: 405-421, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.