

Prediction of IER5 structure and function using a bioinformatics approach

QIANG XIONG¹, XIAOYAN JIANG^{1,2}, XIAODAN LIU³, PINGKUN ZHOU³ and KUKE DING^{1,2}

¹National Institute for Radiological Protection; ²Key Laboratory of Radiological Protection and Nuclear Emergency, Chinese Center for Disease Control and Prevention, Beijing 100088; ³Department of Radiation Toxicology and Oncology, Beijing Institute of Radiation Medicine, Beijing 100850, P.R. China

Received October 9, 2018; Accepted March 25, 2019

DOI: 10.3892/mmr.2019.10166

Abstract. Immediate-early response gene 5 (IER5) is a gene involved in the regulation of the cell cycle, and its structure and function have been investigated by bioinformatics analyses. The present study determined the sites of promoter methylation and gene ontology (GO) annotations associated with IER5. In addition, we conducted a prediction analysis to determine the physical and chemical properties, hydrophobicity/hydrophilicity, posttranslational modification, subcellular localization, transmembrane structure, signal peptide and secondary and tertiary structures of IER5. One CpG island and several methylated sites were identified close to the promoter of *IER5*. The GO analysis suggested that *IER5* could bind ions and proteins that were mainly associated with metabolic processes. IER5 comprised 327 amino acids and was reported to be an unstable hydrophilic protein with an isoelectric point of 4.91. A total of 18 O-glycosylation sites and 22 phosphorylation sites were identified within this protein. The subcellular localization of IER5 was mainly in the nucleus, and its main secondary structural element was the α -helix. Bioinformatic analyses of the features of IER5 may improve understanding of its structure and function; however, experimental verification is required.

Introduction

Immediate-early genes encode a type of polypeptides that serve a significant role in cell regulation and the response of the cell to external stimuli. The regulation of the cell cycle by these polypeptides differs among numerous types of cells, due to variations in expression. The response of some immediate-early gene family members to extracellular stimuli is characterized by slow kinetics, which delays transcription

and prolongs protein half-life (1). Immediate early response 5 (IER5) was initially reported by Williams *et al* (1), and is a member of the slow-kinetics immediate-early gene family. IER5 is a gene without introns comprising 2,350 nucleotides and is located in 1q25.3. The predicted open reading frame encodes a 327-amino-acid protein. Its amino terminus is rich in proline residues as previously noted for other homology immediate-early genes, such as *pip92/IER2/ETR101*. In contrast to *pip92/IER2/ETR101*, the transcriptional activation of *IER5* does not require induction by phosphokinase C (2).

It has been revealed that *IER5* expression was upregulated following external stimuli, inducing cell apoptosis. Therefore, it may be considered that *IER5* is involved in the regulation of the cell cycle. Savitz *et al* (3), demonstrated that the expression levels of *IER5* were increased in peripheral mononuclear cells derived from patients with depression and mood disorders compared with those of healthy subjects. Ishikawa *et al* (4,5) and Asano *et al* (6) noted that *IER5* was a positive feedback regulator of heat shock factor 1 (HSF1) dephosphorylation, following investigation of the mechanism of *HSF1* transcription. In addition, Li *et al* (7), Kawabata *et al* (8) and Nakamura *et al* (9), also observed the cell cycle regulation by *IER5* during the progression of different diseases. Our research group has demonstrated that radiation can induce upregulation of *IER5* in tumor cells (10,11) and this process can modulate the transcription of cell division cycle (*CDC*)25B by competitively binding to the *CDC25B* promoter (12). Additionally, we reported that decreased *IER5* expression could increase the population of cancer cells in the G₂/M phase of the cell cycle (13,14), and that binding of a novel transcription factor and GC binding factor (GCF) to the *IER5* promoter could act as a negative regulator of *IER5* transcriptional activity (15). Furthermore, we proposed that decreased *IER5* expression significantly lowered the efficiency of DNA double strand break repair in HeLa cells induced by ionizing radiation (16).

Although various studies have been conducted on the functional mechanism of *IER5*, only a limited number has explored the structure of the *IER5* protein. The present study aimed to determine the structure of the *IER5* protein by bioinformatics analysis, and to explain its *in vivo* function and mechanism of action, based on its structural features, so as to provide a theoretical basis for subsequent experimental determination of its structure.

Correspondence to: Professor Kuke Ding, National Institute for Radiological Protection, Chinese Center for Disease Control and Prevention, 2 Xinkang Street, Xicheng, Beijing 100088, P.R. China
E-mail: shouding@ccmu.edu.cn

Key words: immediate-early response gene 5, bioinformatics prediction, structure, function

Materials and methods

Sequence of the *IER5* gene and protein. The nucleotide sequences of 2,000 bp upstream and 1,000 bp downstream of the transcription sites of the *IER5* gene (Species, *Homo sapiens*, accession: NC_000001.11, Gene ID: 51278) were downloaded from the National Center for Biotechnology Information (NCBI; <https://www.ncbi.nlm.nih.gov/>), and the amino acid sequences of the IER5 protein (Entry: Q5VY09, Entry name: IER5_HUMAN, Length: 327-amino-acid) were downloaded from the UniProt database (<https://www.uniprot.org/>).

Prediction of the *IER5* gene sequence. The online software Promoter Scan (<https://www.bimas.cit.nih.gov/molbio/proscan/>; website decommissioned March 8, 2019) was applied for the prediction of the promoter sequence and the binding sites of the related transcription factors. The program, which recognized ~70% of primate promoter sequences, predicted promoter regions based on scoring homologies with putative eukaryotic Pol II promoter sequences. The Methprimer (<http://www.urogene.org/methprimer/>) was applied for the determination of methylation sites and CpG islands at the promoter region. The criteria for the CpG island prediction results were the following: Island size >100, GC% (percentage of G plus C) >50% and observed/expected values >0.5. Gene Ontology (GO) and annotations were investigated using the GO Enrichment Analysis using the AmiGO tool (<http://amigo.geneontology.org/amigo/landing>). GO enrichment analysis identified relevant groups of genes that functioned collectively, which reduced the thousands of molecular changes to notably fewer biological functions in order to describe a putative function corresponding to the mean number of molecular changes.

Prediction of *IER5* protein features. The physical and chemical properties of the IER5 protein were predicted by the ProtParam tool (<https://web.expasy.org/protparam/>). The protein parameters, including the molecular weight, theoretical isoelectric point, amino acid composition, atomic composition, extinction coefficient and instability index were calculated based on either compositional data or on the N-terminal amino acid residues. The hydrophobicity/hydrophilicity of the IER5 protein was predicted by the ProtScale (<https://web.expasy.org/protscale/>) which provided 57 scales defined by a numerical value assigned to each type of amino acid. The most frequently used scales were the hydrophobicity or hydrophilicity scales and the secondary structure conformational parameters scales. The O-glycosylation sites were predicted by a genetic engineering approach. The NetOGlyc 4.0 Server (<http://www.cbs.dtu.dk/services/NetOGlyc/>) was used enable a proteome-wide discovery approach of O-glycan sites by a 'bottom-up' ETD-based mass spectrometric analysis. The N-glycosylation sites of the IER5 protein were predicted by the NetNGlyc 1.0 Server (<http://www.cbs.dtu.dk/services/NetNGlyc/>) which was based on an artificial neural network in an attempt to discriminate between glycosylated and non-glycosylated sequences. The phosphorylation sites of the IER5 protein were predicted by the NetPhos 3.1 Server (<http://www.cbs.dtu.dk/services/NetPhos/>) which predicted serine, threonine and tyrosine phosphorylation sites in eukaryotic proteins using ensembles of neural networks and 17 kinases as follows:

Ataxia telangiectasia-mutated, casein kinase (CK)I, CKII, calmodulin-dependent protein kinase-II, DNA-dependent protein kinase catalytic subunit, epidermal growth factor receptor, glycogen synthase kinase (GSK)3, insulin receptor (INSR), protein kinase A (PKA), protein kinase B, protein kinase C (PKC), cGMP-dependent protein kinase, ribosomal S6 kinase, SRC, cyclin-dependent kinase 1 (cdc2), cyclin-dependent kinase 5 (cdk5) and p38 mitogen-activated kinase (MAPK). The subcellular localization of the IER5 protein was predicted by the PSORT II tool (<https://psort.hgc.jp/form2.html>) from its amino acid sequences. The location of the transmembrane, intracellular and extracellular regions was predicted by the TMHMM Server v2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>) by reading a FASTA-format protein sequence. The presence and location of signal peptide cleavage sites in the amino acid sequences were predicted by the SignalP 4.1 Server (<http://www.cbs.dtu.dk/services/SignalP/>) with a D-cutoff score of 0.5. The nuclear localization sequence (NLS) of the IER5 protein was predicted by the NLS mapper (http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi) with a cut-off score of 4.0. The secondary and tertiary structures of the IER5 protein were predicted by the PSIPRED (<http://bioinf.cs.ucl.ac.uk/psipred/>) and I-TASSER (<https://zhanglab.ccmb.med.umich.edu/I-TASSER/>) software, respectively. These two methods used a known amino acid sequence to match a template in a protein database. All-by-all TM scores were calculated for the full set of putative templates and the matrix of scores was analyzed to remove any possible outlying templates whose structure was too dissimilar with that of the full set of templates. The tertiary structures of the IER5 protein was compiled and produced by PyMOL (<https://pymol.org>).

Results

Promoter binding and methylation analysis of the *IER5* gene. The sequences located at 2,000 bp upstream and 1,000 bp downstream of the transcription sites of *IER5* were analyzed, and the promoter was located at a region between 1,722 and 1,972 bp in the plus-strand (Table I). We identified one CpG island located at a region between 1,499 and 2,944 bp and several potential methylation sites (Fig. 1). In addition, the predicted promoter sequence of *IER5* overlapped with the CpG island. We further examined the transcription factors sites close to the promoter region and identified specific transcription factors sites associated with methylation, such as the AP-2 (Table II).

Physical and chemical properties and hydrophobicity/hydrophilicity of the *IER5* protein. A total of ~12.84% (42/327) of the amino acids (Asp and Glu) were identified with negative charge, whereas 9.48% (31/327) of the amino acids (Arg and Lys) were identified with positive charge. The IER5 protein contained 11 Cys residues. Considering that all Cys residues formed cystines, the estimated molar extinction coefficient in the aqueous solution was 32,595 M⁻¹cm⁻¹, whereas for 0.1% absorbance (1 g/l) would be 0.967; however, providing all Cys residues could not form cystines, the corresponding value would be 31,970 and 0.949. The IER5 protein was predicted as an unstable protein, of which the structural formula was C₁₄₅₉H₂₂₉₄N₄₂₈O₄₆₄S₁₄ and the total molecular weight was

Table I. Promoter prediction of immediate early response 5 gene.

Promoter	Start	End	Score	Sequence
Promoter 1	1,722	1,972	72.54	AATCTGTAAC TCCAAACAGTAGCGCTCTCGAGCACCGTCCCGA ACATTCAC TCCCCACGGGCTGGTCTGCGGCCGCAAGCCGTC GCCCCCTTTAAGAGCCTGCTCCGCGGGACTAACGTTCAACG GGCCCTGGCGCCCCCTCCTCGGGCTCCGATTGGCCGTGCGCGG CGCACGAGGGCGCGCCGGCCAGCCCCGGAACGGTTGGCGGC CCTTCGTGATTGGCGGTTCGAGAAGCCTATATAAGGCGCGGG

Table II. Transcription factor sites about the promoter of immediate early response 5.

Name	Strand	Location	Weight
AP-2	+	1773	1.355
AP-2	+	1774	1.108
UCE.2	+	1793	1.278
UCE.2	-	1796	1.216
GCF	+	1856	2.361
CTF	+	1875	1.704
UCE.2	+	1879	1.278
NFI	-	1881	4.221
junB-US2	-	1882	1.51
TTR_inverted_repeat	-	1892	3.442
GCF	-	1893	2.284
UCE.2	-	1910	1.216
AP-2	-	1930	1.672
HNF1	+	1931	1.012
TFIID	+	1958	1.971
GCF	+	1968	2.361
T-Ag	+	1970	1.086

AP-2, activating protein 2; CTF, CCAAT box-binding transcription factor; GCF, GC-rich sequence DNA-binding factor; HNF1, homeobox A; junB-US2, JunB proto-oncogene, AP-1 transcription factor subunit; NFI, nuclear factor I; TFIID, transcription factor II D; UCE, ubiquitin-conjugating enzyme; T-Ag, large tumor antigen.

estimated to 33703.69. The theoretical isoelectric point was 4.91, of which the coefficient of instability was 61.1.

The aliphatic index of the IER5 protein was estimated to 60.4 and the average hydrophilic value was estimated to -0.493. Leu, the most hydrophobic amino acid, was identified at the 38th position and with an index value of 1.944. The most hydrophilic amino acid was reported at the 293th position and its index was -2.976. According to the hydrophilic/hydrophobic distribution diagram (Fig. 2), the majority of the amino acids were hydrophilic amino acids, and therefore IER5 was considered a hydrophilic protein.

Posttranslational modification of the IER5 protein. Protein modifications, such as glycosylation and phosphorylation are required to fulfill protein physiological function.




Figure 1. Methylation and CpG island prediction of immediate early response 5. The blue line denotes the nucleotide sequence, while the short vertical red lines indicate CpG sites; and green cross coarse line denotes the CpG island within the indicated region.




Figure 2. Hydrophobicity/hydrophilicity of the IER5 protein. Positive scores presented hydrophobicity and negative scores indicated hydrophilicity. The higher the absolute value, the higher the degree of hydrophobicity/hydrophilicity.

Glycosylation serves an important role in the interaction between proteins and other macromolecules (17). A total of 18 O-glycosylation sites were identified (score >0.5), in the IER5 protein; however, no N-glycosylation sites were reported. Phosphorylation is required for signal transduction (18). The results indicated 15 serine (Ser), six threonine (Thr) and one tyrosine (Tyr) protein kinase phosphorylation sites (score >0.5, Fig. 3). A total of 9 kinases associated with the phosphorylation reactions, including PKA, cdc2, INSR, PKC, cdk5, GSK3, p38MAPK, CKII and CKI were reported in addition to 'unspecified' types.

Subcellular localization, transmembrane structure and signal peptide identification of the IER5 protein. The prediction of the subcellular localization of IER5 suggested that the protein exhibited a 56.5% probability of localizing in the nucleus, whereas the possibility for cytoskeletal and mitochondrial localization was notably lower (17.4 and 13.0%, respectively). The prediction indicated a lower potential for localization in




Figure 3. Phosphorylation site of the immediate early response 5 protein. Red lines denote serine phosphorylation sites, green lines indicates threonine phosphorylation sites, the blue line denotes a tyrosine phosphorylation site and the pink line indicates the threshold.




Figure 4. Transmembrane structures of the immediate early response 5 protein. The dark purple line denotes the IER5 protein, the lower purple line indicates the outer cell membrane and the blue line represents the inner cell membrane.




Figure 5. Signal peptides of the immediate early response 5 protein. The red lines denote the C-score, green line indicates the S-score and the blue line denotes the Y-score.

the mitochondria, Golgi apparatus and vesicular secretion system (4.3%). No transmembrane structures (Fig. 4) and signal peptides (Fig. 5) were present. Further analysis indicated that the IER5 protein may possess a nuclear localization sequence (NLS) GSTPLKKPRRNLE (the position in protein sequence from the N terminal to C terminal is 235-247) with a score of 4.5 (threshold of 5.0).

Secondary structure of the IER5 protein. The secondary structure refers to a periodic structure arranged along a direction, and is the regular repeated conformation in the protein polypeptide chain. PSIPRED has been previously used to predict protein secondary structure based on a two-stage



Figure 6. Secondary structure of the immediate early response 5 protein. H denotes α -helix and C denotes disordered coil states.

neural network; the average prediction accuracy was estimated at a range of 76.5-78.3% (19). The results revealed 6 α -helices, but no β -sheet or β -turn motifs. The remaining structural parts of the proteins were determined to present as disordered coils (Fig. 6).

Tertiary structure of the IER5 protein. I-TASSER is an online integrated platform based on the ‘sequence-structure-function’ model of automatic protein structure and function prediction. Starting from the amino acid sequence, the platform produces the three-dimensional atomic-scale model through the comparison of multiple threading alignment approaches and the iterative structural assembly simulation (20,21). The platform presents five models following the completion of predicting the tertiary structure; default model 1 is considered as the best model based on comprehensive analysis of the three parameters, namely the C-score, the TM score and the RMSD. The tertiary structure was presented in a cartoon model embedded in the surface mode (Fig. 7).

GO of the IER5 gene. GO is a database established by the GO consortium, which is a unified induction, interpretation and analysis of the cytological components, molecular functions and biological approaches of genes and their products. We searched for GO terms and annotations associated with *IER5* by the AmiGO browser. The present study reported that this gene was involved in several primary biological and metabolic processes that require ion and protein binding. The main distribution of *IER5* has not been predicted in terms of cellular components (Table III).




Figure 7. Tertiary structure of the immediate early response 5 protein.

Table III. GO of immediate early response 5.

Ontology	GO ID	Term
Biological process	GO:0044238	Primary metabolic process
Cellular component	-	
Molecular function	GO:0043167	Ion binding
	GO:0042802	Identical protein binding
	GO:0005515	Protein binding
-, not predicted. GO, Gene Ontology.		

Discussion

In the present study, we investigated the structure and function of *IER5*, and its encoded protein using bioinformatics online analysis software. Hypermethylation of CpG islands at the promoter region has been reported to inhibit the transcriptional activity of the gene, whereas low promoter methylation activates gene expression (22,23). The results of the present study indicated one CpG island and several potential methylation sites. Liu *et al* (10) and Shi *et al* (11) demonstrated that radiation could upregulate the expression levels of *IER5*. Therefore, we proposed that the methylation levels of the wild type *IER5* gene may be low, but could notably increase following radiation exposure, inducing its expression (24,25). Specific transcription factors have been located at the promoter region of *IER5* and certain protein binding sites were suggested by GO analysis. Our previous study reported two GCF binding sites at the promoter region, which is in agreement with the present findings (15). Following the binding of GCF, the transcriptional activity and radiation sensitivity of *IER5* significantly decreased (15).

Glycosylation serves an important role in cellular immunity, signal transduction, protein translation regulation and protein degradation. For example, the majority of transcription factors and enzymes require glycosylation following translation (17). In addition, phosphorylation serves a key role in

protein signal transduction, gene expression and cell cycle regulation (18). The present study reported 18 O-glycosylation sites and 22 phosphorylation sites in the *IER5* protein, which reflected the complexity of *IER5* protein function.

Based on the prediction of protein subcellular localization, transmembrane region and signal peptide identification, it was speculated that the *IER5* protein was mainly localized in the nucleus. The absence of the transmembrane structure and of the signal peptide indicated that the *IER5* protein did not require entry into other membrane organelles. Following protein expression, various hydrophilic structures and lack of the transmembrane structure and of the signal peptide may facilitate the free diffusion of the *IER5* protein in the cell without its modification by the endoplasmic reticulum or the Golgi apparatus (26,27). The *IER5* protein may be channeled from the nuclear pore complex to the nucleus possibly via an NLS.

The helix-turn-helix domain (HTH) is a relatively conserved structure with various patterns that correspond to different protein families (28). HTH contains two α -helices, which are connected by one turn, and can recognize the specific base sequence of the DNA in order to regulate its transcription, replication and translation (29). Previous studies suggested that radiation increased the expression levels of the *IER5* gene and protein (10,11). Competitive binding of *IER5* to the *Cdc25B* promoter led to downregulated expression levels of *Cdc25B* (12). We demonstrated that the secondary structure of *IER5* had only 6 α -helices. Following structure prediction, it was speculated that *IER5* could possess an HTH structure based on its function. Further experiments are required to confirm this hypothesis.

Of note, the present study has certain limitations as only bioinformatics predictions were conducted. Furthermore, we did not conduct investigations using clinical samples, which may verify the results reported in the present study.

We examined the features of the *IER5* gene and protein using bioinformatics analyses, which could aid future investigation of their biological functions. Furthermore, predicting the *IER5* may provide a experimental basis for investigation into its functions in the future.

Acknowledgements

Not applicable.

Funding

The present work was supported by grants from the National Natural Science Foundation of China (grant nos. 31170806, 31770907 and 31640022) and the Beijing Natural Science Foundation (grant no. 7172146).

Availability of data and materials

All data generated or analyzed during the present study are included in this published article.

Authors' contributions

QX, XJ, XL, PZ and KD conceived and designed the study. QX wrote the paper. All authors reviewed and edited the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Williams M, Lyu MS, Yang YL, Lin EP, Dunbrack R, Birren B, Cunningham J and Hunter K: Ier5, a novel member of the slow-kinetics immediate-early genes. *Genomics* 55: 327-334, 1999.
- Takaya T, Kasatani K, Noguchi S and Nikawa J: Functional analyses of immediate early gene ETR101 expressed in yeast. *Biosci Biotechnol Biochem* 73: 1653-1660, 2009.
- Savitz J, Frank MB, Victor T, Bebak M, Marino JH, Bellgowan PS, McKinney BA, Bodurka J, Kent Teague T and Drevets WC: Inflammation and neurological disease-related genes are differentially expressed in depressed patients with mood disorders and correlate with morphometric and functional imaging abnormalities. *Brain Behav Immun* 31: 161-171, 2013.
- Ishikawa Y and Sakurai H: Heat-induced expression of the immediate-early gene IER5 and its involvement in the proliferation of heat-shocked cells. *FEBS J* 282: 332-340, 2015.
- Ishikawa Y, Kawabata S and Sakurai H: HSF1 transcriptional activity is modulated by IER5 and PP2A/B55. *FEBS Lett* 589: 1150-1155, 2015.
- Asano Y, Kawase T, Okabe A, Tsutsumi S, Ichikawa H, Tatebe S, Kitabayashi I, Tashiro F, Namiki H, Kondo T, *et al*: IER5 generates a novel hypo-phosphorylated active form of HSF1 and contributes to tumorigenesis. *Sci Rep* 6: 19174, 2016.
- Li XN, Ji C, Zhou PK and Wu YM: Establishments of IER5 silence and overexpression cervical cancer SiHa cell lines and analysis of radiosensitivity. *Int J Clin Exp Pathol* 9: 6671-6682, 2016.
- Kawabata S, Ishita Y, Ishikawa Y and Sakurai H: Immediate-early response 5 (IER5) interacts with protein phosphatase 2A and regulates the phosphorylation of ribosomal protein S6 kinase and heat shock factor 1. *FEBS Lett* 589: 3679-3685, 2015.
- Nakamura S, Nagata Y, Tan L, Takemura T, Shibata K, Fujie M, Fujisawa S, Tanaka Y, Toda M, Makita R, *et al*: Transcriptional repression of Cdc25B by IER5 inhibits the proliferation of leukemic progenitor cells through NF-YB and p300 in acute myeloid leukemia. *PLoS One* 6: e28011, 2011.
- Liu Y, Tian M, Zhao H, He Y, Li F, Li X, Yu X, Ding K, Zhou P and Wu Y: IER5 as a promising predictive marker promotes irradiation-induced apoptosis in cervical cancer tissues from patients undergoing chemoradiotherapy. *Oncotarget* 8: 36438-36448, 2017.
- Shi HM, Ding KK, Zhou PK, Guo DM, Chen D, Li YS, Zhao CL, Zhao CC and Zhang X: Radiation-induced expression of IER5 is dose-dependent and not associated with the clinical outcomes of radiotherapy in cervical cancer. *Oncol Lett* 11: 1309-1314, 2016.
- Yang C, Yang M, Feng Z, Liu X, Yin L, Zhou P and Ding K: Radiation modulated the interaction of IER5 protein and CDC25B promoter DNA in primary hepatocellular carcinoma. *Int J Clin Exp Pathol* 9: 2888-2895, 2016.
- Yang C, Wang Y, Hao C, Yuan Z, Liu X, Yang F, Jiang H, Jiang X, Zhou P and Ding K: IER5 promotes irradiation- and cisplatin-induced apoptosis in human hepatocellular carcinoma cells. *Am J Transl Res* 8: 1789-1798, 2016.
- Ding KK, Shang ZF, Hao C, Xu QZ, Shen JJ, Yang CJ, Xie YH, Qiao C, Wang Y, Xu LL and Zhou PK: Induced expression of the IER5 gene by gamma-ray irradiation and its involvement in cell cycle checkpoint control and survival. *Radiat Environ Biophys* 48: 205-213, 2009.
- Yang C, Yin L, Zhou P, Liu X, Yang M, Yang F, Jiang H and Ding K: Transcriptional regulation of IER5 in response to radiation in HepG2. *Cancer Gene Ther* 23: 61-65, 2016.
- Yu XP, Wu YM, Liu Y, Tian M, Wang JD, Ding KK, Ma T and Zhou PK: IER5 is involved in DNA double-strand breaks repair in association with PAPR1 in Hela cells. *Int J Med Sci* 14: 1292-1300, 2017.
- Ohtsubo K and Marth JD: Glycosylation in Cellular mechanisms of health and disease. *Cell* 126: 855-867, 2006.
- Singh V, Ram M, Kumar R, Prasad R, Roy BK and Singh KK: Phosphorylation: Implications in cancer. *Protein J* 36: 1-6, 2017.
- Jones DT: Protein secondary structure prediction based on position-specific scoring matrices. *J Mol Biol* 292: 195-202, 1999.
- Yang J and Zhang Y: I-TASSER server: New development for protein structure and function predictions. *Nucleic Acids Res* 43: W174-W181, 2015.
- Zhang Y: I-TASSER: Fully automated protein structure prediction in CASP8. *Proteins* 77 (Suppl 9): S100-S113, 2009.
- Hashimshony T, Zhang J, Keshet I, Bustin M and Cedar H: The role of DNA methylation in setting up chromatin structure during development. *Nat Genet* 34: 187-192, 2003.
- Balada E, Ordi-Ros J, Serrano-Acedo S, Martinez-Lostao L, Rosa-Leyva M and Vilardell-Tarrés M: Transcript levels of DNA methyltransferases DNMT1, DNMT3A and DNMT3B in CD4+ T cells from patients with systemic lupus erythematosus. *Immunology* 124: 339-347, 2008.
- Illingworth RS and Bird AP: CpG islands-'a rough guide'. *FEBS Lett* 583: 1713-1720, 2009.
- Eckhardt F, Lewin J, Cortese R, Rakyan VK, Attwood J, Burger M, Burton J, Cox TV, Davies R, Down TA, *et al*: DNA methylation profiling of human chromosomes 6, 20 and 22. *Nat Genet* 38: 1378-1385, 2006.
- Palmer KJ, Konkel JE and Stephens DJ: PCTAIRE protein kinases interact directly with the COPII complex and modulate secretory cargo transport. *J Cell Sci* 118: 3839-3847, 2005.
- Bard F, Mazelin L, Péchoux-Longin C, Malhotra V and Jurdic P: Src regulates golgi structure and KDEL receptor-dependent retrograde transport to the endoplasmic reticulum. *J Biol Chem* 278: 46601-46606, 2003.
- Wintjens R and Rooman M: Structural classification of HTH DNA-binding domains and protein-DNA interaction modes. *J Mol Biol* 262: 294-313, 1996.
- Aravind L and Koonin EV: DNA-binding proteins and evolution of transcription regulation in the archaea. *Nucleic Acids Res* 27: 4658-4670, 1999.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.