

High hydrostatic pressure induces apoptosis of retinal ganglion cells via regulation of the NGF signalling pathway

HONGJI LIU^{1*}, WEI WANG^{2*}, XIANG LI³, CHAO HUANG⁴,
ZONGDUAN ZHANG⁵, MINGYUE YUAN¹ and XIANGYU LI¹

¹College of Ophthalmology, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan 610072;

²Department of Ophthalmology, Kunming Municipal Hospital of Traditional Chinese Medicine, Kunming, Yunnan 646000;

³Department of Ophthalmology, Hospital of Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan 610072;

⁴Central Laboratory, Shenzhen Bao'an People's Hospital Affiliated to Southern Medical University, Shenzhen,

Guangdong 518100; ⁵Department of Ophthalmology, The Affiliated Eye Hospital of

Wenzhou Medical University, Wenzhou, Zhejiang 325027, P.R. China

Received July 17, 2018; Accepted April 2, 2019

DOI: 10.3892/mmr.2019.10206

Abstract. High pressure is the most important factor inducing retinal ganglion cell (RGC) apoptosis. However, the underlying mechanisms remain obscure. The present study investigated the effects of different levels of hydrostatic pressure (HP) on RGCs and the potential mechanisms involved. Primary cultured rat RGCs were exposed to five levels of HP (0, 20, 40, 60 and 80 mmHg) for 24 h. Morphological changes in RGCs were observed. The viability and apoptosis rate of RGCs were detected using a Cell Counting Kit-8 assay and Annexin V-fluorescein isothiocyanate/propidium iodide flow cytometry, respectively. Western blotting, reverse transcription-quantitative polymerase chain reaction and immunofluorescence were used to detect the expression and mRNA levels of nerve growth factor (NGF), protein kinase B (AKT), apoptosis signal-regulating kinase 1 (ASK1), forkhead box O1 (FoxO1) and cAMP response element binding protein (CREB). In the 0- and 20-mmHg groups, there were no apoptotic morphological changes. In the 40 mmHg group, parts of the cell were shrunk or disrupted. In the 60 mmHg group, neurite extension was weakened and parts of the cells were disintegrating or dying. In the 80 mmHg group, the internal structures of the cells were not visible at all. The apoptosis rates of RGCs were significantly higher and the viability rates

significantly lower under 40, 60 and 80 mmHg compared with under 0 or 20 mmHg (all $P < 0.01$). The expression and mRNA levels of NGF, AKT and CREB decreased in a dose-dependent manner in the 40-, 60- and 80-mmHg groups (all $P < 0.05$), but those of ASK1 and FoxO1 increased in a dose-dependent manner (all $P < 0.05$). Interestingly, the alterations to the expression and mRNA levels of CREB were significantly larger compared with the changes in ASK1 or FoxO1 in the 40-, 60- and 80-mmHg groups (all $P < 0.01$). The results of the present study demonstrate that elevated HP of 40, 60 or 80 mmHg reduces viability and induces apoptosis in RGCs, which may occur through effects on the NGF/ASK1/FoxO1 and NGF/AKT/CREB pathways, of which the latter is more strongly affected.

Introduction

Glaucoma, affecting ~80 million people, is the leading cause of irreversible blindness globally (1). Retinal ganglion cell (RGC) degeneration and a progressive loss of visual fields are the primary features of glaucoma (2). Normal pressure is one of the essential conditions for cellular function and cell integrity; high intraocular pressure (HIOP) is considered to be the main risk factor enhancing RGC death and visual functional damage in glaucoma (3). However, the underlying molecular mechanisms of consistent RGC loss induced by HIOP remain unclear (4). Further insight into the pathogenesis of glaucoma is critical for protecting RGCs.

Nerve growth factor (NGF) may protect RGCs against apoptosis induced by palmitic acid injury, and the effect is partly modulated by protein kinase B (AKT)/forkhead box O1 (FoxO1) (5). AKT /Activation of cAMP-responsive element binding protein 1 (CREB1) activity may play an important role to control the viability of neuronal cells during retinal development (6). FoxO1 serves a crucial function in modulating apoptosis and survival in multiple cell types (7-9). Cyclin-dependent kinase-5 can modulate of the cellular location of FoxO1 for regulating the apoptosis of neurons (10).

Correspondence to: Professor Xiang Li, Department of Ophthalmology, Hospital of Chengdu University of Traditional Chinese Medicine, 39 Shi-er-qiao Road, Jinniu, Chengdu, Sichuan 610072, P.R. China

E-mail: jeannelxiang@cudtcm.edu.cn

*Contributed equally

Key words: glaucoma, retinal ganglion cells, apoptosis, hydrostatic pressure, nerve growth factor

In addition, deletion of the apoptosis signal-regulating kinase 1 (ASK1) gene may also prevent RGC death in mouse models of glaucoma (11). CREB may also protect RGCs against death following axotomy (12). Although the above studies have suggested that NGF, AKT, ASK1, FoxO1 and CREB have crucial functions in RGC apoptosis in palmitic acid injury models or RGC death in mouse models, there is no direct evidence to demonstrate whether they function in RGCs with *in vitro* hydrostatic pressure (HP)-induced damage.

In the present study, the degree to which HP caused apoptosis in RGCs cultured for 24 h was initially determined, and subsequently the underlying mechanisms of accelerated RGC apoptosis were investigated by measuring the protein and mRNA expression levels of NGF, AKT, ASK1, FoxO1 and CREB, paving the way for the discovery of novel therapeutic strategies to protect RGCs and preserve vision in patients with glaucoma.

Materials and methods

Cell separation and primary culture. Retinas from new-born Sprague Dawley rats were dissociated by trituration and dissociated in 1.25 g/l trypsin and 2 g/l hyaluronidase (Thermo Fisher Scientific, Inc., Waltham, MA, USA). RGCs from retinas were purified by Thy-1.1 antibody panning as previously described (13). Cell suspensions were transferred to a petri dish coated with rat monoclonal anti-macrophage antibody (1:75; cat no. ab56297; Abcam, Cambridge, UK) for 45 min at room temperature, then loaded into a petri dish coated with horseradish peroxidase (HRP)-conjugated goat anti-rabbit immunoglobulin (Ig)G (1:400; cat no. ab97057; Abcam) for 30 min at room temperature to isolate microglia. Cells separated from the petri dish wall were transferred to another petri dish coated with HRP-conjugated goat anti-rabbit IgM (1:300; cat no. ab97180; Abcam) and mouse anti-rat monoclonal antibody Thy1.1 (1:100; cat no. ab6721; Abcam) for 30 min at room temperature, following which the cells that separated from the petri dish wall were removed. Next, trypsin digestion was performed at 37°C for 8 min, following which the cells were pelleted by centrifugation at 500 x g for 5 min at room temperature and seeded onto 96-well plates coated with poly-Aminutrin and laminin. Following incubation at 37°C for 24 h, purified RGCs were grown in fresh cell culture medium (97.5% Neurobasal Medium, 2% B27 and 0.5% 200 mM/l L-glutamine) at 37°C in a 5% CO₂ atmosphere. The medium was then refreshed every two days. The growth status and morphological changes were observed using an inverted phase contrast optical microscope (CKX31/41; Olympus Corporation, Tokyo, Japan) at x100 magnification. The cells were then collected for further analysis. The present study was ethically approved by the Ethics Committee for Animal Research of Chengdu University of Traditional Chinese Medicine (Chengdu, China).

Pressure system. An open pressure control system was designed to stimulate the cultured RGCs with stable, adjustable elevated HP as previously described (14). The plexiglass pressure chamber was concatenated via a low-pressure two-stage regulator (cat no. A032009; Thermo Fisher Scientific, Inc.) to a source of 5% CO₂/95% oxygen air (Thermo Fisher

Scientific, Inc.). This system provided constant HP ranging from 0-200 mmHg. Gas to the chamber was warmed and humidified by bubbling through water. The water flask and pressure chamber were maintained at 37°C via placement in an electronically controlled conventional incubator. Notably, the system had the advantages of easy and precise control of the gas flow and pressure.

Experimental group. Purified RGCs were cultured for 48 h, then subsequently divided into five groups, namely, the 0-, 20-, 40-, 60- and 80-mmHg groups. RGCs in the 0-, 20-, 40-, 60- and 80-mmHg groups were exposed to HPs of 0-, 20-, 40-, 60- and 80-mmHg, respectively, for 24 h.

Detecting viability with a Cell Counting Kit-8 (CCK-8). Following the application of different HPs for 24 h, RGC viability was measured using a CCK-8 (Dojindo Molecular Technologies, Inc., Kumamoto, Japan) assay according to the manufacturer's protocol. In brief, 0.1 ml cell suspension per well were seeded in a 96-well culture plate at a density of 5x10⁴ RGCs/ml. A total of 10 µl CCK-8 solution was added to each well of the plate. Subsequent to a 4 h incubation at 5% CO₂ and 37°C, the optical density 450 of each well of the plate was measured with a Tecan Genios reader (Varioskan™ LUX, Thermo Fisher Scientific, Inc.). There were three replicates for each group.

Measuring apoptosis rate with Annexin V fluorescein isothiocyanate (FITC)/propidium iodide (PI) flow cytometry. Subsequent to being subjected to different HPs for 24 h, RGCs were collected following trypsinization using a pancreatic enzyme (Procell Life Science and Technology Co. Ltd., Wuhan, China) without ethylenediaminetetraacetic acid and centrifuged at 200 x g for 5 min at room temperature. The collected cells were re-suspended in pre-cooled 1x phosphate buffered saline (PBS) at 200 x g for 5-10 min, then washed twice and suspended in 1x Binding Buffer of 300 µl. A total of 5 µl FITC-conjugated Annexin V (Nanjing KeyGen Biotech Co., Ltd., Nanjing, China) was added to the cell suspension. The cells were then incubated for 15 min at room temperature in the dark, then washed with binding buffer and analysed by flow cytometry using Cell Quest software (version 7.5.3; BD Biosciences, Franklin Lakes, NJ, USA). Experiments were performed at least 3 times.

Western blotting. Total protein (25 µg per sample) was extracted from the retinal tissues and the protein concentration was determined by BCA assay kit (Pierce Biotechnology; Thermo Fisher Scientific, Inc.). The proteins were subjected to 8-13% SDS-PAGE gel electrophoresis and transferred onto a polyvinylidene fluoride membrane, the membrane was immersed in 5% skim milk and incubated for 12 h at 4°C with primary antibodies, including mouse monoclonal anti-β-actin (1:200; cat no. ab6199; Abcam), rabbit anti-rat NGF polyclonal antibody (1:200; cat no. ab6199; Abcam), rabbit anti-rat AKT polyclonal antibody (1:200; cat no. ab8805; Abcam), rabbit anti-rat ASK1 polyclonal antibody (1:200 cat no. ab131506; Abcam), rabbit anti-rat FoxO1 polyclonal antibody (1:200; cat no. ab12161; Abcam) and rabbit anti-rat CREB polyclonal antibody (1:200; cat no. ab31387; Abcam). The proteins

Table I. Primers used for NGF, AKT, ASK1, FoxO1, CREB and β -actin.

mRNA	Genbank ID	Primer	Sequence (5'-3')
NGF	XM_006233053.2	Forward	CATAGCGTAATGTCCATGTTGTTCT
		Reverse	CAGTGATGTTGCGGGTCTGC
AKT	NM_033230.2	Forward	CCTGAAGCTACTGGGCAAGGG
		Reverse	ACAAAGCAGAGGCGGTCTGTG
ASK1	NM_001277694.1	Forward	GGGCCGATCTGCTTACCTCT
		Reverse	TCAGCCGCCAACTCTTTACC
FoxO1	NM_001191846.2	Forward	TCCTCGAACCAGCTCAAACGC
		Reverse	GGATACACCAGGAATGCACATC
CREB	NM_134443.1	Forward	TAGTGCCAGCAACCAAGTT
		Reverse	TGCTTCCCTGTTCTTCATTAGAC
β -actin	NM_031144.3	Forward	ACCCGCGAGTACAACCTTCTT
		Reverse	TATCGTCATCCATGGCGAACT

NGF, nerve growth factor; AKT, protein kinase B; ASK1, apoptosis signal-regulating kinase 1; FoxO1, forkhead box O1; CREB, cAMP-response element binding protein.

were then incubated with HRP-conjugated goat-anti-rabbit IgG secondary antibodies (1:500; cat no. ab97057; Abcam) for 45 min at 37°C. Following a final wash in tris-buffered saline with 0.1% Tween-20, the membrane was reacted with electrochemiluminescence reagent (Merck KGaA, Darmstadt, Germany). Densitometric analysis was performed by ImageJ software (1.48 u; National Institutes of Health, Bethesda, MD, USA) and normalized to β -actin.

Immunofluorescence staining. Cells on coverslips were fixed with 4% paraformaldehyde for 20 min at room temperature. Next, the coverslips were washed with PBS (pH 7.4) for 3 min and incubated in 5% bovine serum albumin (Sigma-Aldrich; Merck KGaA) at room temperature for 1 h. Immunoreaction was then performed overnight with monoclonal anti-mouse Thy1.1 antibody (1:100; cat no. ab6721; Abcam). All coverslips were subsequently treated with anti-mouse immunoglobulin G H&L secondary antibodies (1:200; cat no. ab6720; Abcam) for 60 min at 37°C. The cell nuclei were counterstained with DAPI for 10 min at room temperature, then covered with an anti-fading mounting medium prior to optical microscopy (CKX31/41; Olympus Corporation) at x100 and x200 magnification. Fluorescent intensities were quantified using Image-Pro® Plus 6.0 software (Media Cybernetics, Inc., Rockville, MD, USA).

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Total RNA was extracted using TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol. A reverse transcriptase cDNA synthesis kit (Toyobo Life Science, Osaka, Japan) was used for obtaining cDNA. qPCR was performed using the SYBR Green PCR Master Mix (Applied Biosystems; Thermo Fisher Scientific, Inc.) with the Applied Biosystems 7500 RT-PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc.). The RT-qPCR thermocycling conditions were as follows: A total of 2 min initial denaturation at 95°C, then 45 cycles of

denaturation at 95°C for 10 sec and annealing at 60°C for 30 sec. The relative expression levels of the target mRNAs were normalized to that of β -actin. The $2^{-\Delta\Delta C_q}$ method (15) was used to calculate the relative fold changes in mRNA expression. The sequences of the primer pairs are presented in Table I.

Statistical analysis. All data are presented as the mean \pm standard deviation. Statistical analysis was performed in SPSS 21.0 (IBM Corp., Armonk, NY, USA). Comparisons were performed using one-way analysis of variance, and the comparison between specific groups was performed using the least significant difference test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Identification and purity testing of RGCs. Immunofluorescence was used to identify RGCs and detect their purity with an anti-Thy1.1 antibody. Areas of positive fluorescent green signal observed by fluorescent microscopy mainly appeared in the cytomembrane. Certain cell neurites also exhibited green fluorescence. RGC purity was calculated as the proportion of the number of cells marked with green fluorescence out of the total number of cells in a given field of view by inverted phase contrast microscopy. The results from 12 fields of view demonstrated that the purity of the RGCs was 96.97% (Fig. 1).

Elevated HP induces apoptosis-associated morphological changes in RGCs in vitro. Ohashi *et al* (16) reported that elevated HP influences the morphological changes of cells. In the present study, morphological changes in RGCs were observed by inverted phase contrast microscopy. In the 0-mmHg group, the cells tended to congregate and present long neurites (Fig. 2A). In the 20-mmHg group, the RGCs were similar to those in the 0-mmHg group (Fig. 2B).




Figure 1. Identification and purity testing of retinal ganglion cells. Green is Thy1.1 immunofluorescence staining and blue is DAPI staining (magnification, x100 and x200).




Figure 2. Elevated HP induced apoptotic morphological changes in RGCs *in vitro*. The effect was observed by inverted phase contrast microscopy. (A) Micrograph of RGCs cultured at HP values of 0 mmHg (magnification, x100). (B) Micrograph of RGCs cultured at HP values of 20 mmHg (magnification, x100). (C) Micrograph of RGCs cultured at HP values of 40 mmHg (magnification, x100). (D) Micrograph of RGCs cultured at HP values of 60 mmHg (magnification, x100). (E) Micrograph of RGCs cultured at HP values of 80 mmHg (magnification, x100). HP, hydrostatic pressure; RGC, retinal ganglion cell.




Figure 3. Elevated HP reduced viability and induced apoptosis in RGCs *in vitro*. These effects were observed using a Cell Counting K-8 assay and by Annexin V FITC/PI flow cytometry, respectively. (A) Annexin V FITC/PI flow cytometry results at different HP values (0, 20, 40, 60 and 80 mmHg). (B) Quantification of cell viability rates. (C) Quantification of cell apoptosis rates. Data are presented as the mean \pm standard deviation (n=3). **P<0.01 vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; PI, propidium iodide; FITC, fluorescein isothiocyanate; UL, upper left; UR, upper right; LL, lower left; LR, lower right.

However, in the 40-mmHg group, sections of the cell bodies were shrunken, and certain cell membranes were disrupted, although a number of cells remained intact (Fig. 2C). In the 60-mmHg group, there were numerous floating cells with shrunken cell bodies and incomplete cell membranes. In addition, the extension of the neurites was weakened, parts of the cells were disintegrating or dying and the internal structures of the cells were only dimly visible (Fig. 2D). In the 80-mmHg group, the apoptotic morphological changes in the RGCs were similar compared with those of the 60-mmHg group, but the internal structures of the cells were not visible at all (Fig. 2E). These results indicate that HPs of 40, 60 and 80 mmHg may induce apoptotic morphological changes in RGCs *in vitro*.

Elevated HP reduces viability and induces apoptosis in RGCs in vitro. Once cells were subjected to different HPs for 24 h, Annexin V FITC/PI flow cytometry results revealed that the apoptosis rates of the 40-, 60- and 80-mmHg groups were significantly higher (P<0.01; Fig. 3A and C) and the viability rates were significantly lower (P<0.01; Fig. 3B) compared with those of the 0-mmHg group. These results are consistent with those of Li *et al* (17) and Zhang *et al* (18), whose studies demonstrated that elevated HP may cause apoptosis in RGCs *in vitro*. In the present study, the apoptosis rates gradually increased and the viability rates gradually decreased from the 40 mmHg to the 80 mmHg group (P<0.05; Fig. 3B and C). However, there was no significant difference in apoptosis rates or viability rates between the 0-mmHg group and 20-mmHg groups (P>0.05;




Figure 4. Influence of elevated HP on mRNA and protein expression levels in pressure-treated RGCs. (A) NGF, AKT, ASK1, FoxO1 and CREB protein expression levels of RGCs were detected by western blotting. (B) Quantified protein expression of ASK1, FoxO1 and CREB at 20, 40, 60 and 80 mmHg compared with 0 mmHg. (C) Quantified mRNA expression levels of ASK1, FoxO1 and CREB at 20, 40, 60 and 80 mmHg compared with 0 mmHg. To compare the NGF/ASK1/FoxO1 and NGF/AKT/CREB pathways, the expression levels at 0 mmHg were defined as the baseline; the data were calculated by subtracting the protein and mRNA expression levels at 0 mmHg from the protein and mRNA expression levels at 20, 40, 60 and 80 mmHg. (D) Quantified protein expression levels of NGF, AKT, ASK1, FoxO1 and CREB. (E) Quantified mRNA expression levels of NGF, AKT, ASK1, FoxO1 and CREB. Data are presented as the mean \pm standard deviation ($n=3$). * $P<0.01$ vs. CREB at the same level of HP. ** $P<0.05$ and *** $P<0.01$ vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; NGF, nerve growth factor; AKT, protein kinase B; ASK1, apoptosis signal-regulating kinase 1; FoxO1, forkhead box O1; CREB, cAMP response element binding protein.

Fig. 3B and C). These results indicate that elevated HP of 40, 60, or 80 mmHg notably reduces the viability rate and increases the apoptosis rate of RGCs cultured for 24 h.

Elevated HP promotes RGC apoptosis by altering the expression of NGF, AKT, ASK1, FoxO1 and CREB. Protein expression levels were analysed by western blotting and immunofluorescence, and mRNA expression levels were assessed using RT-qPCR (Figs. 4-9). Compared with the 0-mmHg group, the protein and mRNA levels of NGF, AKT and CREB in the 40-, 60- and 80- mmHg groups significantly decreased in a dose-dependent manner ($P<0.05$; Fig. 4A, D and E), while those of ASK1 and FoxO1 significantly increased in a dose-dependent manner ($P<0.05$; Fig. 4A, D and E). Similar results were confirmed by immunofluorescence ($P<0.05$; Figs. 5-9). To evaluate the differences in the NGF/ASK1/FoxO1 and NGF/AKT/CREB pathways, the expression levels in the 0-mmHg group were defined as the baseline; the data in Fig. 4B and C were calculated by subtracting the protein and mRNA expression levels at 0 mmHg from the protein and mRNA expression levels at 20, 40, 60 and 80 mmHg. Interestingly, the changes in the protein and mRNA levels of CREB were significantly greater compared those of ASK1 or FoxO1 from an HP of 40 to 80 mmHg ($P<0.01$; Fig. 4B and C). These results reveal that HPs of 40, 60 and 80 mmHg may restrain the expression of NGF, AKT and CREB and enhance the expression of ASK1 and FoxO1 to induce apoptosis in

RGCs *in vitro*. In addition, the expression of CREB changes more substantially compared with that of ASK1 or FoxO1.

Discussion

It is generally recognized that HIOP is a chief risk factor for expedited RGC apoptosis in glaucoma (19). A growing number of studies have demonstrated that increased apoptosis, proliferation and migration exist in cells treated with elevated HP (20,21). Thus, cells exposed to elevated HP may generally function as an *in vitro* model of glaucomatous RGC apoptosis, as performed in the present study (22). Although Osborne *et al* (23) revealed that elevation of HP did not cause apoptosis in human RGCs cultured at 60 mmHg for 24 h, this difference may be associated with the speed of pressure fluctuations (24). In the present study, an open pressure control system was successfully established. In this system, HP values were first set as required. Next, five pressure values (0, 20, 40, 60 and 80 mmHg) were selected to simulate HIOP injury. Finally, the present study was able to identify HP levels that resulted in the apoptosis of cultured RGCs. It was observed that treatment with 40, 60 and 80 mmHg over a period of 24 h reduced the viability rates and increased the apoptosis rates of RGCs. However, these patterns were not observed in the 0- or 20-mmHg group. Thus, the data suggests that a HP of 40, 60 or 80 mmHg may reduce viability and enhance apoptosis in RGCs *in vitro*, while 20 mmHg HP




Figure 5. Influence of elevated HP on NGF protein expression in each group of RGCs. (A) Micrographs of immunofluorescence exhibiting NGF expression in RGCs. Green is FITC immunofluorescence staining, and blue is DAPI staining (magnification, x100). (B) Quantification and statistical analysis of fluorescence intensity. Data are presented as the mean ± standard deviation (n=3). *P<0.05 vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; NGF, nerve growth factor; FITC, fluorescein isothiocyanate.

does not cause RGC apoptosis, meaning that the viability and apoptosis rates of RGCs will not noticeably change until pressure reaches 40 mmHg.

Lee *et al* (25) reported that when RGC were exposed to pressure of 100 mmHg for 20 h, 20 out of 31 examined cells exhibited apoptotic morphological changes, presenting cell




Figure 6. Influence of elevated HP on AKT protein expression in each group of RGCs. (A) Micrographs of immunofluorescence exhibiting AKT expression in RGCs. Green is FITC immunofluorescence staining, and blue is DAPI staining (magnification, x100). (B) Quantification and statistical analysis of fluorescence intensity. Data are presented as the mean \pm standard deviation (n=3). *P<0.05 vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; AKT, protein kinase B; FITC, fluorescein isothiocyanate.

body shrinkage and neurite retraction. However, only 3 out of 20 examined cells exhibited the spontaneous morphological changes of cell body shrinkage and neurite retraction under

exposure to 15 mmHg pressure (26). A report by Miao *et al* (20) had demonstrated that an elevated pressure of 60 mmHg hindered cell migration. In the present study, an elevated HP




Figure 7. Influence of elevated HP on ASK1 protein expression in each group of RGCs. (A) Micrographs of immunofluorescence exhibiting ASK1 expression in RGCs. Green is FITC immunofluorescence staining, and blue is DAPI staining (magnification, x100). (B) Quantification and statistical analysis of fluorescence intensity. Data are presented as the mean \pm standard deviation (n=3). *P<0.05 vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; ASK1, apoptosis signal-regulating kinase 1; FITC, fluorescein isothiocyanate.

of 40, 60 and 80 mmHg was revealed to induce apoptotic morphological changes including cell body shrinkage and neurite retraction of RGCs, in addition to the disintegration of

cells. These changes gradually increased with rising HP, which indicates that exposure to elevated HP may engender apoptotic morphological changes in RGCs. Lee *et al* (25) reported that




Figure 8. Influence of elevated HP on FoxO1 protein expression in each group of RGCs. (A) Micrographs of immunofluorescence exhibiting FoxO1 expression in RGCs. Green is FITC immunofluorescence staining, and blue DAPI staining (magnification, x100). (B) Quantification and statistical analysis of fluorescence intensity. Data are presented as the mean \pm standard deviation (n=3). *P<0.05 vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; FoxO1, forkhead box O1; FITC, fluorescein isothiocyanate.

caspace-3 activation was associated with the dismantling of the cytoskeleton and apoptotic morphological changes in RGCs exposed to high pressure. Cell apoptosis occurs by reduced

activity of the NGF signalling pathway in connection with active caspace-3 (27). In the present study, there was reason to hypothesise that the underlying mechanisms of apoptotic




Figure 9. Influence of elevated HP on CREB protein expression in each group of RGCs. (A) Micrographs of immunofluorescence exhibiting CREB expression in RGCs. Green is FITC immunofluorescence staining, and blue is DAPI staining (magnification, x100). (B) Quantification and statistical analysis of fluorescence intensity. Data are presented as the mean \pm standard deviation (n=3). *P<0.05 vs. the 0 mmHg group. HP, hydrostatic pressure; RGC, retinal ganglion cell; CREB, cAMP response element binding protein; FITC, fluorescein isothiocyanate.

morphological changes may be associated with the dysregulation of the NGF pathway. Thus, further research needs to be performed.

RGC apoptosis is the predominant change observed in glaucoma (28). Preliminary studies (29,30) have reported that RGC loss is associated with the downregulation of NGF and




Figure 10. Schematic illustrating the mechanism in which elevated HP induces apoptosis in RGCs through the regulation of the NGF signalling pathway. The protein and mRNA expression level of NGF were decreased in RGCs exposed to HP levels of 40, 60 and 80 mmHg HP; the binding of NGF to TRK receptors activates the PI3K/AKT signalling pathway. When AKT is activated by NGF-stimulated phosphorylation, it may influence RGC apoptosis in three ways. First, it may phosphorylate the target of FoxO1 to mediate cell fate decisions by regulating apoptosis. Second, AKT may phosphorylate ASK1 and reduce its activity; the kinase JNK1/2, downstream of ASK1, phosphorylates and activates FoxO1 protein, resulting in cell apoptosis. Third, the decrease in NGF may trigger reduced phosphorylation of AKT and CREB to induce RGC apoptosis. HP, hydrostatic pressure; RGC, retinal ganglion cell; NGF, nerve growth factor; AKT, protein kinase B; ASK1, apoptosis signal-regulating kinase 1; FoxO1, forkhead box O1; CREB, cAMP response element binding protein; JNK1/2, c-Jun N-terminal kinase 1/2; PI3K, phosphoinositide-3 kinase; TRK, tropomyosin receptor kinase.

NGF receptor expression in the retina, in contrast, treatment with NGF may substantially alleviate the damage induced by glaucoma. The binding of NGF to tropomyosin receptor kinase receptors activates the phosphoinositide-3 kinase/AKT signalling pathway, which is indispensable for maintaining the balance between neuronal life and death (31,32). When AKT is activated by NGF-stimulated phosphorylation, it phosphorylates multiple targets to promote cell survival (33). Although the above experiments revealed the association of NGF and AKT to the loss of RGCs in a rat model of glaucoma, there is no direct evidence to indicate whether they exert the same effects in RGCs with HIOP-induced damage. In the present study, it was revealed that NGF and AKT were closely associated with *in vitro* RGC apoptosis induced by HP of 40, 60 and 80 mmHg. In addition, downstream of NGF/AKT, ASK1/FoxO1 and CREB have important functions in cell apoptosis. Forkhead box proteins, including FoxO1, are among the targets of AKT and may mediate cell fate decisions by regulating proliferation, differentiation and migration, in addition to apoptosis and survival (7,34). Dysregulation and mutation of forkhead box proteins, including FoxO1, frequently cause human genetic diseases, including glaucoma

and tumorigenesis (35,36). Similarly, ASK1 has been reported to be phosphorylated by AKT at serine 83, which reduces its activity (37,38). Studies by Harada *et al* (39,40) and Katome *et al* (11) have revealed that the deletion of the ASK1 gene prevented RGC death in a variety of mouse models of glaucoma. The kinase c-Jun N-terminal kinase 1/2, downstream of ASK1, may phosphorylate and activate FoxO1 protein, resulting in cell apoptosis (41). On the other hand, Phan *et al* (42) had reported that netleatin together with NGF enhanced the phosphorylation of AKT and CREB, helping to protect neurons from cell death and promoting neurogenesis in serum-deprived PC12 cells. In the present study, the increased expression levels of ASK1 and FoxO1 were observed in RGCs exposed to 40, 60 and 80 mmHg HP. CREB, another downstream target of NGF/AKT, had reduced protein and mRNA expression levels in RGCs exposed to HP levels of 40, 60 and 80 mmHg, and there were also trends toward decreased protein and mRNA expression levels of NGF and AKT. Interesting, taking 0 mmHg as the baseline, it was revealed that the changes in the protein and mRNA expression levels of CREB, part of the NGF/AKT/CREB pathway, were substantially larger compared with those in ASK1 or FoxO1

of the NGF/ASK1/FoxO1 pathway from HP values of 40 to 80 mmHg. This implies that the NGF/AKT/CREB pathway may serve a notable role in the apoptosis of RGCs exposed to HP values of 40 to 80 mmHg, as presented in Fig. 10.

In conclusion, the observations of the present study suggest that HPs of 40, 60 and 80 mmHg may induce apoptotic morphological changes and increase the apoptosis rate of RGCs *in vitro*. Regulation of the NGF/ASK1/FoxO1 and NGF/AKT/CREB pathways may be notable mechanisms resulting in RGC apoptosis upon exposure to elevated HP. In addition, the NGF/AKT/CREB pathway may be the primary mechanism of apoptosis in RGCs exposed to elevated HP *in vitro*. Therefore, it would be worthwhile further investigating the underlying mechanisms in order to extend present understanding of RGC apoptosis under HIOP. The results of the present study provide novel evidence for the molecular mechanisms of RGC apoptosis under elevated HP and pave the way for discovering novel therapeutic methods for patients with glaucoma.

Acknowledgements

The authors would like to thank the Ophthalmic Laboratory of the Hospital of Chengdu University of Traditional Chinese Medicine for the use of its laboratory and equipment and to Professor Junguo Duan and Professor Xuejing Lu of the Chengdu University of Traditional Chinese Medicine for their help in experimental research.

Funding

The present study was supported by the National Natural Science Foundation of China (grant no. 81373695).

Availability of data and materials

The datasets used and/or analysed during the present study are available from the corresponding author on reasonable request.

Authors' contributions

XL designed the present study and approved this submission. HL and WW performed the experiments, and wrote the manuscript. CH, ZZ, MY and XYL collected and analyzed the data, and revised the manuscript. All authors read and approved the final version of the manuscript.

Ethics approval and consent to participate

The present study was ethically approved by the Ethics Committee for Animal Research of Chengdu University of Traditional Chinese Medicine (Chengdu, China).

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Tham YC, Li X, Wong TY, Quigley HA, Aung T and Cheng CY: Global prevalence of glaucoma and projections of glaucoma burden through 2040: A systematic review and meta-analysis. *Ophthalmology* 121: 2081-2090, 2014.
2. Zhu Y, Zhang L, Sasaki Y, Milbrandt J and Gidday JM: Protection of mouse retinal ganglion cell axons and soma from glaucomatous and ischemic injury by cytoplasmic overexpression of Nmnat1. *Invest Ophthalmol Vis Sci* 54: 25-36, 2013.
3. Caprioli J: Glaucoma: A disease of early cellular senescence. *Invest Ophthalmol Vis Sci* 54: ORSF60-67, 2013.
4. Cordeiro MF and Levin LA: Clinical evidence for neuroprotection in glaucoma. *Am J Ophthalmol* 152: 715-716, 2011.
5. Yan PS, Tang S, Zhang HF, Guo YY, Zeng ZW and Wen Q: Nerve growth factor protects against palmitic acid-induced injury in retinal ganglion cells. *Neural Regen Res* 11: 1851-1856, 2016.
6. Socodato R, Brito R, Portugal CC, de Oliveira NA, Calaza KC and Paes-de-Carvalho R: The nitric oxide-cGKII system relays death and survival signals during embryonic retinal development via AKT-induced CREB1 activation. *Cell Death Differ* 21: 915-928, 2014.
7. Zhang X, Tang N, Hadden TJ and Rishi AK: Akt, FoxO and regulation of apoptosis. *Biochim Biophys Acta* 1813: 1978-1986, 2011.
8. Cheng C, Jiao JT, Qian Y, Guo XY, Huang J, Dai MC, Zhang L, Ding XP, Zong D and Shao JF: Curcumin induces G2/M arrest and triggers apoptosis via FoxO1 signaling in U87 human glioma cells. *Mol Med Rep* 13: 3763-3770, 2016.
9. Ma Y, Wang X, Peng Y and Ding X: Forkhead box O1 promotes INSI cell apoptosis by reducing the expression of CD24. *Mol Med Rep* 13: 2991-2998, 2016.
10. Zhou J, Li H, Li X, Zhang G, Niu Y, Yuan Z, Herrup K, Zhang YW, Bu G, Xu H and Zhang J: The roles of Cdk5-mediated subcellular localization of FOXO1 in neuronal death. *J Neurosci* 35: 2624-2635, 2015.
11. Katome T, Namekata K, Guo X, Semba K, Kittaka D, Kawamura K, Kimura A, Harada C, Ichijo H, Mitamura Y and Harada T: Inhibition of ASK1-p38 pathway prevents neural cell death following optic nerve injury. *Cell Death Differ* 20: 270-280, 2013.
12. Sun MM, Wang YC, Li Y, Guo XD, Chen YM and Zhang ZZ: Effect of ATF3-deletion on apoptosis of cultured retinal ganglion cells. *Int J Ophthalmol* 10: 691-695, 2017.
13. He S, Park YH, Yorito T and Krishnamoorthy RR: Endothelin-mediated changes in gene expression in isolated purified rat retinal ganglion cells. *Invest Ophthalmol Vis Sci* 56: 6144-6161, 2015.
14. Tok L, Naziroglu M, Uguz AC and Tok O: Elevated hydrostatic pressures induce apoptosis and oxidative stress through mitochondrial membrane depolarization in PC12 neuronal cells: A cell culture model of glaucoma. *J Recept Signal Transduct Res* 34: 410-416, 2014.
15. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
16. Ohashi T, Sugaya Y, Sakamoto N and Sato M: Hydrostatic pressure influences morphology and expression of VE-cadherin of vascular endothelial cells. *J Biomech* 40: 2399-2405, 2007.
17. Li Y, Chen YM, Sun MM, Guo XD, Wang YC and Zhang ZZ: Inhibition on apoptosis induced by elevated hydrostatic pressure in retinal ganglion cell-5 via laminin upregulating beta1-integrin/Focal adhesion kinase/protein kinase B signaling pathway. *Chin Med J (Engl)* 129: 976-983, 2016.
18. Zhang SH, Gao FJ, Sun ZM, Xu P, Chen JY, Sun XH and Wu JH: High pressure-induced mtDNA alterations in retinal ganglion cells and subsequent apoptosis. *Front Cell Neurosci* 10: 254, 2016.
19. Guo L, Moss SE, Alexander RA, Ali RR, Fitzke FW and Cordeiro MF: Retinal ganglion cell apoptosis in glaucoma is related to intraocular pressure and IOP-induced effects on extracellular matrix. *Invest Ophthalmol Vis Sci* 46: 175-182, 2005.
20. Miao H, Crabb AW, Hernandez MR and Lukas TJ: Modulation of factors affecting optic nerve head astrocyte migration. *Invest Ophthalmol Vis Sci* 51: 4096-4103, 2010.
21. Agar A, Li S, Agarwal N, Coroneo MT and Hill MA: Retinal ganglion cell line apoptosis induced by hydrostatic pressure. *Brain Res* 1086: 191-200, 2006.
22. Wang S, Hu T, Wang Z, Li N, Zhou L, Liao L, Wang M, Liao L, Wang H, Zeng L, *et al*: Macroglia-derived thrombospondin 2 regulates alterations of presynaptic proteins of retinal neurons following elevated hydrostatic pressure. *PLoS One* 12: e0185388, 2017.

23. Osborne A, Aldarwesh A, Rhodes JD, Broadway DC, Everitt C and Sanderson J: Hydrostatic pressure does not cause detectable changes in survival of human retinal ganglion cells. *PLoS One* 10: e0115591, 2015.
24. Resta V, Novelli E, Vozzi G, Scarpa C, Caleo M, Ahluwalia A, Solini A, Santini E, Parisi V, Di Virgilio F and Galli-Resta L: Acute retinal ganglion cell injury caused by intraocular pressure spikes is mediated by endogenous extracellular ATP. *Eur J Neurosci* 25: 2741-2754, 2007.
25. Lee JK, Lu S and Madhukar A: Real-Time dynamics of Ca²⁺, caspase-3/7, and morphological changes in retinal ganglion cell apoptosis under elevated pressure. *PLoS One* 5: e13437, 2010.
26. Maeno E, Ishizaki Y, Kanaseki T, Hazama A and Okada Y: Normotonic cell shrinkage because of disordered volume regulation is an early prerequisite to apoptosis. *Proc Natl Acad Sci USA* 97: 9487-9492, 2000.
27. Mnich K, Carleton LA, Kavanagh ET, Doyle KM, Samali A and Gorman AM: Nerve growth factor-mediated inhibition of apoptosis post-caspase activation is due to removal of active caspase-3 in a lysosome-dependent manner. *Cell Death Dis* 5: e1202, 2014.
28. Rokicki W, Dorecka M and Romaniuk W: Retinal ganglion cells death in glaucoma-mechanism and potential treatment. Part I. *Klin Oczna* 109: 349-352, 2007 (In Polish).
29. Colafrancesco V, Parisi V, Sposato V, Rossi S, Russo MA, Coassin M, Lambiase A and Aloe L: Ocular application of nerve growth factor protects degenerating retinal ganglion cells in a rat model of glaucoma. *J Glaucoma* 20: 100-108, 2011.
30. Wang H, Wang R, Thrimawithana T, Little PJ, Xu J, Feng ZP and Zheng W: The nerve growth factor signaling and its potential as therapeutic target for glaucoma. *Biomed Res Int* 2014: 759473, 2014.
31. Andjelkovic M, Suidan HS, Meier R, Frech M, Alessi DR and Hemmings BA: Nerve growth factor promotes activation of the alpha, beta and gamma isoforms of protein kinase B in PC12 pheochromocytoma cells. *Eur J Biochem* 251: 195-200, 1998.
32. Crowder RJ and Freeman RS: Phosphatidylinositol 3-kinase and Akt protein kinase are necessary and sufficient for the survival of nerve growth factor-dependent sympathetic neurons. *J Neurosci* 18: 2933-2943, 1998.
33. Franke TF, Hornik CP, Segev L, Shostak GA and Sugimoto C: PI3K/Akt and apoptosis: Size matters. *Oncogene* 22: 8983-8998, 2003.
34. Tzivion G, Dobson M and Ramakrishnan G: FoxO transcription factors; Regulation by AKT and 14-3-3 proteins. *Biochim Biophys Acta* 1813: 1938-1945, 2011.
35. Wierstra I: FOXM1 (Forkhead box M1) in tumorigenesis: Overexpression in human cancer, implication in tumorigenesis, oncogenic functions, tumor-suppressive properties, and target of anticancer therapy. *Adv Cancer Res* 119: 191-419, 2013.
36. Hannehalli S and Kaestner KH: The evolution of Fox genes and their role in development and disease. *Nat Rev Genet* 10: 233-240, 2009.
37. Kim AH, Khursigara G, Sun X, Franke TF and Chao MV: Akt phosphorylates and negatively regulates apoptosis signal-regulating kinase 1. *Mol Cell Biol* 21: 893-901, 2001.
38. Yuan ZQ, Feldman RI, Sussman GE, Coppola D, Nicosia SV and Cheng JQ: AKT2 inhibition of cisplatin-induced JNK/p38 and Bax activation by phosphorylation of ASK1. Implication of AKT2 in chemoresistance. *J Biol Chem* 278: 22832-23840, 2003.
39. Harada C, Namekata K, Guo X, Yoshida H, Mitamura Y, Matsumoto Y, Tanaka K, Ichijo H and Harada T: ASK1 deficiency attenuates neural cell death in GLAST-deficient mice, a model of normal tension glaucoma. *Cell Death Differ* 17: 1751-1759, 2010.
40. Harada C, Nakamura K, Namekata K, Okumura A, Mitamura Y, Iizuka Y, Kashiwagi K, Yoshida K, Ohno S, Matsuzawa A, *et al*: Role of apoptosis signal-regulating kinase 1 in stress-induced neural cell apoptosis in vivo. *Am J Pathol* 168: 261-269, 2006.
41. Essers MA, Weijzen S, de Vries-Smits AM, Saarloos I, de Ruiter ND, Bos JL and Burgering BM: FOXO transcription factor activation by oxidative stress mediated by the small GTPase Ral and JNK. *EMBO J* 23: 4802-4812, 2004.
42. Phan CW, Sabaratnam V, Bovicelli P, Righi G and Saso L: Neglectin as a neuroprotectant enhances the action of nerve growth factor and induces neurite outgrowth in PC12 cells. *Biofactors* 42: 591-599, 2016.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.