

# Characteristics of olfactory ensheathing cells and microarray analysis in *Tupaia belangeri* (Wagner, 1841)

NA LIN<sup>1,2\*</sup>, XIU-JUAN DONG<sup>3\*</sup>, TING-YONG WANG<sup>1</sup>, WEN-JI HE<sup>4</sup>, JING WEI<sup>5</sup>,  
HAI-YING WU<sup>6</sup> and TING-HUA WANG<sup>1,7</sup>

<sup>1</sup>Institute of Neuroscience, Kunming Medical University, Kunming, Yunnan 650500;

<sup>2</sup>Kunming Lin Ming Biological Technology Co., Ltd., Kunming, Yunnan 650000; <sup>3</sup>Department of Physical Education, Hainan Normal University, Haikou, Hainan 571100; <sup>4</sup>Department of Ultrasonic Cardiogram, Kunming Children's Hospital, Kunming Medical University, Kunming, Yunnan 650228; Departments of <sup>5</sup>Pharmacy and <sup>6</sup>Emergency, First Affiliated Hospital of Kunming Medical University, Kunming, Yunnan 650000; <sup>7</sup>Institute of Neuroscience, and Animal Zoology Department, Kunming Medical University, Kunming, Yunnan 650500, P.R. China

Received April 15, 2016; Accepted May 25, 2017

DOI: 10.3892/mmr.2019.10422

**Abstract.** Tree shrews are most closely related to the primates and so possess a number of advantages in experimental studies; they have been used as an animal model in bacterial and virus infection, cancer, endocrine system disease, and certain nervous system diseases. Their olfactory ensheathing cells (OECs) are able to release several cytokines to promote neuronal survival, regeneration and remyelination. The present study used western blot analysis to identify antibody specificity in protein extracts from whole tree shrew brains to identify the specificity of p75 nerve growth factor receptor (NGFR) derived from rabbits (75 kDa). OECs were cultured and isolated, then stained and identified using the antibodies for p75NGFR. To investigate the capacity of OECs to express cytokines and growth factors, microarray technology was used, and the analysis revealed that OECs were able to express 9,821 genes. Of these genes, 44 genes were from the neurotrophic factor family, which may indicate their potential in transplantation *in vivo*. The present study considered the function of OECs as revealed by other studies, and may contribute to future research.

## Introduction

Tree shrews (*Tupaia belangeri*, Wagner, 1841) are small animals originally regarded as either primates or insectivores and later classified as a separate order, Scandentia (1,2). They are most closely related to the primates, with the advantages of being readily available, easy to tame and possessing a strong reproductive capacity (3). Due to these advantages they have been successfully employed as an animal model for bacterial (4) or viral infection (5,6), tumors (7), endocrine system disease (3,8), and certain nervous system diseases, including myopia (9-11), psychosocial stress (12,13), cerebral ischemia (14) and aging (15).

Olfactory ensheathing cells (OECs) are the glial cells that derive from the olfactory placode and envelop olfactory axons in the course of migration from the olfactory epithelium to the bulb (16). A number of studies have identified that OECs can not only release various cytokines (17,18) that can promote neuronal survival (19), regeneration (20,21) and remyelination (22,23), *in vitro* and *in vivo*, but can also migrate into lesion regions (24) and facilitate functional repair (25,26) following injuries. In the present study, OECs of tree shrews were isolated and purified successfully for the first time, to the best of the authors' knowledge. Cytokine or the receptor mRNA expression was also investigated using a microarray. The results of the present study may contribute to the use of OECs in transplantation and the evaluation of their curative effect in fundamental research and future clinical trials.

## Materials and methods

**Animals.** Tree shrews (2 newborn for cell culture and 1 adult male, weight 140±10 g at the start of the experiment) and 1 adult male Sprague-Dawley rats (for western blot analysis, weight 140±10 g at the start of the experiment) were provided by Kunming Medical University Animal Center (Kunming, China) and employed in this study. The animals were raised at 20±5°C, 40-60% humidity and a 12-h light/dark cycle with

**Correspondence to:** Professor Ting-Hua Wang, Institute of Neuroscience, Kunming Medical University, 191 Western Renmin Road, Kunming, Yunnan 650500, P.R. China  
E-mail: 395160081@qq.com

Professor Hai-Ying Wu, Department of Emergency, First Affiliated Hospital of Kunming Medical University, 295 Xichang Road, Kunming, Yunnan 650000, P.R. China  
E-mail: 874705340@qq.com

\*Contributed equally

**Key words:** tree shrew, olfactory ensheathing cells, microarray analysis

food and water *ad libitum*. All animal work was conducted according to the guidelines for the care and use of experimental animals established by the Ministry of Science and Technology of the People's Republic of China (approval no. 2006-398), and was approved by the ethics committee of Institute of Neuroscience, Kunming Medical University (Kunming, China). All animals were bred in separated cages.

**Primary cell cultures.** Primary cultures of OECs were set up from newborn tree shrews and the animals were sacrificed. Olfactory bulbs were aseptically removed. The meningeal layer was stripped off with fine forceps and the tissue was enzymatically (trypsin 0.25%; 1:30 tissues to trypsin ratio) and mechanically dissociated, then incubated in 5% CO<sub>2</sub> at 37°C for 15 min. The cells were recovered by centrifugation (100 x g for 5 min at 37°C) in culture medium [Dulbecco's modified Eagle's medium/F12 (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (Thermo Fisher Scientific, Inc.)] and seeded at 1x10<sup>5</sup> cells/ml into culture plates and incubated in 5% CO<sub>2</sub> at 37°C. Cells were cultured for 10 h and then the supernatant was removed to another culture plate.

**Antibodies.** Antibodies against p75 neurotrophin receptor (p75NGFR) and BrdU were used to study the purity and proliferation of OECs. For each antibody, a minimum of two whole series of different animals were studied. Western blot analysis with protein extracts from whole tree shrew and rat brains ensured specificity of the antibodies used. The tissues were homogenized in RIPA buffer (Beyotime Institute of Biotechnology, Haimen, China) that contained a mixture of proteinase inhibitors and the supernatant was collected by centrifuging at 4°C at 22,500 x g for 15 min. The protein concentration was determined by the DC protein assay from Bio-Rad Laboratories, Inc. (Hercules, CA, USA), and protein samples (100 µg) of each cell lysate in SDS loading buffer (Biosharp) were electrophoresed on 10% SDS-PAGE gel. Electrophoresis was performed and the proteins were transferred onto polyvinylidene difluoride membranes (Pall Life Sciences, Port Washington, NY, USA) using electroblotting apparatus (Bio-Rad Laboratories, Inc.). The membranes were blocked at 37°C for 1 h in TBS containing 0.1% Tween-20 and 5% dried milk, and were then incubated at 4°C overnight with NGFRp75 antibody (1:1,000; cat. no. AB1554; EMD Millipore, Billerica, MA, USA). They were incubated with the IRDye 800-conjugated affinity purified goat anti-rabbit immunoglobulin (Ig)G (1:5,000; cat. no. 611-1302; Rockland Immunochemicals, Inc., Pottstown, PA, USA) for 1 h at room temperature and visualization using an Odyssey laser scanning system (LI-COR Biosciences, Lincoln, NE, USA). Blots were reprobbed with the monoclonal mouse anti-β actin antibody (1:5,000; cat. no. A2228; Sigma-Aldrich; Merck KGaA, Darmstadt, Germany), followed by reaction with IRDye 800-conjugated affinity purified goat anti-mouse IgG (1:5,000; cat. no. 610-1302; Rockland Immunochemicals, Inc.) at 37°C for 2 h to confirm equal protein loading.

**Immunofluorescence.** Immunofluorescence was performed on OECs. The cells were fixed with 4% formalin solution at room temperature for 5 min then washed with PBS and





Figure 1. (A) Specificity of the antibody. The bands of p75NGFR with protein extracts from whole rat and tree shrew brain. The molecular marker was a restrained marker and did not react with ECL. (B) OEC biological characteristics *in vitro*. (B1) Morphology of OECs cultured for 24 h (magnification, x200). (B2) Morphology of OECs cultured for 6 days (magnification, x200). (B3) Immunocytochemical staining of BrdU (magnification, x200). (B4) p75NGFR positive cells (magnification, x400). (B5) DAPI positive cells (magnification, x400). (B6) Merged view of the p75NGFR and DAPI double positive cells (magnification, x400). OECs, olfactory ensheathing cells; NGFR, nerve growth factor receptor; R, rat; TS, tree shrew.

permeabilized with 0.1% Triton X-100. The cells were then blocked with 1% bovine serum albumin (10082-139; Gibco; Thermo Fisher Scientific, Inc.) at 37°C for 30 min and incubated for 18 h at 4°C with anti-p75NGFR (1:500; AB1554; EMD Millipore) and anti-BrdU (1:200; mAb #5292; Cell Signaling Technology, Inc., Danvers, MA, USA) antibodies. Following washing with PBS, the cells were incubated at 37°C for 2 h with respective anti-Mouse IgG H&L (Alexa Fluor 647; 1:500; ab150115; Abcam, Cambridge, MA, USA). Following incubation, the slides were washed with PBS, mounted and examined under a fluorescence microscope (Leica TCS SP2; Leica Microsystems GmbH, Wetzlar, Germany). Fluoroshield mounting medium with DAPI (Sigma-Aldrich; Merck KGaA) was used for DAPI staining and the staining was performed according to the manufacturer's protocol.

**Microarray analysis.** Total RNA from OECs was quantified using a NanoDrop ND-1000 (Thermo Fisher Scientific, Inc.) and RNA integrity was assessed using standard denaturing agarose gel electrophoresis (27) (performed by KangChen Biotech Co., Ltd., Shanghai, China). Total RNA from each sample was used for labeling and array hybridization as



Gene Ontology (<http://www.geneontology.org>) to show distinguishable gene expression profiling among samples.

**Specificity of the antibody.** Western blotting with protein extracts from whole tree shrew and rat brains ensured specificity of the antibodies used in the present study. Western blotting demonstrated that the antibodies produced bands that were located at the same level in the gel for rat and tree shrew brain protein extracts. Notably, the western blots also demonstrated that these bands had the proper molecular weight for the p75NGFR antibody examined (Fig. 1A).

**OEC biological characteristics in vitro.** To investigate the method of culturing OECs from tree shrews, the bulbous olfactorius of newborn tree shrews was used. Cultured for 24 h, certain cells began to demonstrate adherence and assume a round or star-like shape (Fig. 1B1). Then, 6 days following

Table I. Genes with significant expression difference.

Sequence identification	Normalized tree shrew OECs	Gene name	Synonyms
NM_001002915	1914.4823	IGFL2	VPRI645
M37484	1709.5454	IGF1	IGFI
NM_004970	1441.7056	IGFALS	ALS
NM_005117	1399.2097	FGF19	n/a
NM_201284	849.4621	EGFR	ERBB, ERBB1, mENA
NM_000614	879.9232	CNTF	HCNTF
BC105131	818.3038	FGF17	FGF-13
BC000125	658.60596	TGFB1	CED, DPD1, TGFB
M19154	643.18677	TGFB2	MGC116892, TGF- $\beta$ 2
BC040499	623.899	TGFB2	AAT3, HNPCC6, MFS2, RIIC, TGFR-2, TGF $\beta$ -RII
NM_001007225	620.82745	IGF2BP2	IMP-2, IMP2, VICKZ2, p62
M29644	592.314	IGF1	IGFI
AY358869	576.01324	FGF17	FGF-13
NM_001004356	551.01306	FGFRL1	FGFR5, FHFR
BC037785	484.15298	FGFR1OP	FOP
AK024388	476.04077	FGFR1	BFGFR, C-FGR, CD331, CEK, FLG, FLT2, H2, H3, H4, H5, KAL2, N-SAM
NM_015927	474.31787	TGFB1I1	ARA55, HIC-5, HIC5, TSC-5
NM_001024847	474.17993	TGFB2	AAT3, HNPCC6, MFS2, RIIC, TGFR-2, TGF $\beta$ -RII
NM_006119	463.30542	FGF8	AIGF, HBGF-8
NM_001012978	436.89304	NGFRAP1L1	BEX5, MGC104434, MGC126446
NM_005247	398.42584	FGF3	HBGF-3, INT2
NM_006206	383.68063	PDGFRA	CD140A, MGC74795, PDGFR2
NM_002607	299.87314	PDGFA	PDGF-A, PDGF1
M63888	280.45587	FGFR1	BFGFR, C-FGR, CD331, CEK, FLG, FLT2, H2, H3, H4, H5, KAL2, N-SAM
NM_020638	267.7028	FGF23	ADHR, HPDR2, HYPF, PHPTC
BC004972	260.64957	TGFB1	BIGH3, CDB1, CDG2, CDGG1, CSD, CSD1, CSD2, CSD3, LCD1
NM_198283	246.34807	EGFL11	KIAA0663, dJ1018A4.2
BC069692	240.3946	FGF9	GAF, HBFG-9, MGC119914, MGC119915
NM_023028	225.70663	FGFR2	BEK, BFR-1, CD332, CEK3, CFD1, ECT1, JWS, K-SAM, KGFR, TK14, TK25
BC032502	210.7425	FGF11	FHF3, FLJ16061, MGC102953, MGC45269,
S85192	207.95761	VEGF	MGC70609, VEGFA, VPF
BC093731	185.4517	EGF	URG
NM_003867	183.04973	FGF17	FGF-13
NM_002643	168.185	PIGF	MGC32646, MGC33136
BC011453	165.53284	IGFBP5	IBP5
BC098252	157.41748	FGF23	ADHR/HPDR2/HYPF/PHPTC
BC026352	151.22711	TGFB1	BIGH3, CDB1, CDG2, CDGG1, CSD, CSD1, CSD2, CSD3, LCD1
NM_000142	151.22061	FGFR3	ACH, CD333, CEK2, HSFGFR3EX, JTK4
NM_001007563	149.79308	IGFBPL1	IGFBP-RP4, bA113O24.1
NM_020637	140.37721	FGF22	n/a
AK172829	139.57779	FGFRL1	FGFR5, FHFR
NM_022976	125.33181	FGFR2	BEK, BFR-1, CD332, CEK3, CFD1, ECT1, JWS, K-SAM, KGFR, TK14, TK25
M27281	120.924126	VEGF	MGC70609, VEGFA, VPF
NM_004114	102.02009	FGF13	FGF2, FHF2
NM_015633	100.07872	FGFR1OP2	DKFZp564O1863, HSPC123-like

OECs, olfactory ensheathing cells; n/a, not applicable.



adherent culture, three classic morphologies of OECs were observed: Bipolar or fusiform and oblate (Fig. 1B2). Among these shapes, fusiform and oblate were more common, and an evident three-dimensional appearance. After 14 days, the number of cells did not significantly increase, although the cells were still growing rapidly (data not shown). Cells were stained against BrdU (Fig. 1B3) to confirm their capacity for proliferation, which demonstrated that all stages of OECs nuclei were positively stained, demonstrating that they possessed proliferative ability. Immunocytochemical staining against p75NGFR (Fig. 1B4) demonstrated that the fusiform cells or their nuclei were positively stained, demonstrating that they were OECs. All cell nuclei were positively stained with DAPI (Fig. 1B5) and the merged image is demonstrated in Fig. 1B6.

**Gene chip assay.** In order to investigate RNA integrity and genomic DNA contamination, the denaturing agarose gel electrophoresis test was used. The result demonstrated that optical density (OD) A260/A280 ratio was 2.00 and the OD A260/A230 ratio was 2.35. These results confirmed that the RNA was pure [for spectrophotometer, the O.D. A260/A280 ratio should be close to 2.0 for pure RNA (ratios between 1.8 and 2.1 are acceptable) The OD A260/A230 ratio should be >1.8]. The 28S and 18S ribosomal RNA bands were fairly sharp, intense bands and there was a diffuse smear of ethidium bromide-stained material migrating between the 18S and 28S ribosomal bands, possibly comprising mRNA and other heterogeneous RNA species (Fig. 2A). Prior to analysis a boxplot was performed (Fig. 2B). Then, gene chip assay suggested that OECs expressed 9,821 genes (Fig. 2C) and 45 important genes are listed in Table I and Fig. 2D and E.

## Discussion

**Specificity of the antibodies.** Tree shrews possess a number of advantages as animal models, thus, studies on them are popular. However, there are no specific reagents for tree shrews, particularly antibodies. The present study used western blot analysis on protein extracts from whole tree shrew and rat brains to ensure specificity of the antibodies used. Protein extracts from rat brain were used as a positive control, as the specificity of the antibodies has already been verified in this species. The results demonstrated that the antibodies produced bands that were located at the same level. Notably, the molecular weight for p75NGFR was ~75 kDa (28). These data suggested that the antibodies used were specific and the results reliable.

**OECs biological characteristics in vitro.** The present study was the first to successfully isolate and purify tree shrew OECs, to the best of the authors' knowledge. As in a previous study, the cultured cells exhibited the three classical morphologies of OECs: Bipolar or fusiform, multi-ecphyma shape and oblate (26). Previous studies have demonstrated that p75NGFR was the specific marker of OEC (29,30) and that the antibody to BrdU is useful in detecting S-phase cells, as BrdU-positive cells are proliferating (31,32). The present study demonstrated that the majority of cells were p75NGFR positive and BrdU positive. These data further proved that the

majority of the cultured cells were OECs and the majority of them proliferated well.

**Gene chip assay.** The microarray technique allows the monitoring of the expression of tens of thousands of genes simultaneously in one hybridization experiment. DNA segments are closely arranged on a slide. cDNA was reverse transcribed from mRNA derived from tree shrew OECs, labelled with a Cyanine3 (Cy3) fluorophore and hybridized on the microarray slide. Through this technique, the detection of differentially expressed genes is greatly facilitated. In the present study, prior to the microarray analysis, RNA quantification and quality assurance was performed to ensure reliability of the results (33-35). Prior to the statistical analysis, a boxplot test was performed. The boxplot is a traditional method for visualizing the distribution of a dataset (36) and is useful for comparing the distributions of several datasets (37). In the present study, a boxplot was used to observe the distributions of expression values for the samples in the experiment following normalization. The data demonstrated that tree shrew OECs can express 9,821 genes, containing 44 cytokines, including the fibroblast growth factors family (38), the insulin-like growth factor family (39-42), the transforming growth factor family (38), epidermal growth factor (43,44), ciliary neurotrophic factor (40), vascular endothelial growth factor (45) and platelet derived growth factor receptor  $\alpha$  (46,47), which were hypothesized as possessing the capacity to promote neuronal survival, regeneration, remyelination or function improvement.

In conclusion, tree shrew OECs were isolated and purified successfully for the first time, to the best of the authors' knowledge, and the cells were able to secrete a number of cytokines that may contribute to nervous system diseases. Although there was no specific kit, tests were performed to ensure that the data were reliable. These results may lead to future fundamental research or clinical trials.

## Acknowledgements

Not applicable.

## Funding

The authors of the present study would like to thank the Chinese Ministry of the National Science-Technology Support Plan Project for financial support (grant no. 2014BAI01B01).

## Availability of data and materials

All the materials included in the manuscript, including all relevant raw data, will be made freely available to any researchers who wish to use them for non-commercial purposes, while preserving any necessary confidentiality and anonymity.

## Authors' contributions

T-HW, NL and H-YW designed the study. T-YW performed cell cultures, X-JD, W-JH, and T-YW performed immunofluorescence, NL and JW performed microarray analysis,

western blotting and wrote the manuscript. All authors read and approved the final manuscript.

### Ethics approval and consent to participate

All animal work was conducted according to the guidelines for the care and use of experimental animals established by the Ministry of Science and Technology of the People's Republic of China (approval no. 2006-398), and was approved by the ethics committee of Institute of Neuroscience, Kunming Medical University.

### Patient consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### References

- Diogo R: The head and neck muscles of the Philippine colugo (Dermoptera: *Cynocephalus volans*), with a comparison to tree-shrews, primates, and other mammals. *J Morphol* 270: 14-51, 2009.
- Xu L, Chen SY, Nie WH, Jiang XL and Yao YG: Evaluating the phylogenetic position of Chinese tree shrew (*Tupaia belangeri chinensis*) based on complete mitochondrial genome: Implication for using tree shrew as an alternative experimental animal to primates in biomedical research. *J Genet Genomics* 39: 131-137, 2012.
- Collins PM, Tsang WN and Urbanski HF: Endocrine correlates of reproductive development in the male tree-shrew (*Tupaia belangeri*) and the effects of infantile exposure to exogenous androgens. *Gen Comp Endocrinol* 154: 22-30, 2007.
- Li SA, Lee WH and Zhang Y: Two bacterial infection models in tree shrew for evaluating the efficacy of antimicrobial agents. *Dongwuxue Yanjiu* 33: 1-6, 2012.
- Yang ZF, Zhao J, Zhu YT, Wang YT, Liu R, Zhao SS, Li RF, Yang CG, Li JQ and Zhong NS: The tree shrew provides a useful alternative model for the study of influenza H1N1 virus. *Virology* 448: 110-111, 2013.
- Xu G, Gao Z, He W, Ma Y, Feng X, Cai T, Lu F, Liu L and Li W: microRNA expression in hepatitis B virus infected primary tree-shrew hepatocytes and the independence of intracellular miR-122 level for de novo HBV infection in culture. *Virology* 448: 247-254, 2014.
- Xu Y, Köck J, Lu Y, Yang D, Lu M and Zhao X: Suppression of hepatitis B virus replication in *Tupaia* hepatocytes by tumor necrosis factor alpha of *Tupaia belangeri*. *Comp Immunol Microbiol Infect Dis* 34: 361-368, 2011.
- van der Hart MG, de Biurrun G, Czech B, Rupniak NM, den Boer JA and Fuchs E: Chronic psychosocial stress in tree shrews: Effect of the substance P (NK1 receptor) antagonist L-760735 and clomipramine on endocrine and behavioral parameters. *Psychopharmacology (Berl)* 181: 207-216, 2005.
- Arumugam B and McBrien NA: Muscarinic antagonist control of myopia: Evidence for M4 and M1 receptor-based pathways in the inhibition of experimentally-induced axial myopia in the tree shrew. *Invest Ophthalmol Vis Sci* 53: 5827-5837, 2012.
- Guo L, Frost MR, He L, Siegwart JT Jr and Norton TT: Gene expression signatures in tree shrew sclera in response to three myopiagenic conditions. *Invest Ophthalmol Vis Sci* 54: 6806-6819, 2013.
- Zhu X, McBrien NA, Smith EL III, Troilo D and Wallman J: Eyes in various species can shorten to compensate for myopic defocus. *Invest Ophthalmol Vis Sci* 54: 2634-2644, 2013.
- Czech B, Simon M, Schmelting B, Hiemke C and Fuchs E: Astroglial plasticity in the hippocampus is affected by chronic psychosocial stress and concomitant fluoxetine treatment. *Neuropsychopharmacology* 31: 1616-1626, 2006.
- Zambello E, Fuchs E, Abumaria N, Rygula R, Domenici E and Caberlotto L: Chronic psychosocial stress alters NPY system: Different effects in rat and tree shrew. *Prog Neuropsychopharmacol Biol Psychiatry* 34: 122-130, 2010.
- Li S, Meng Q and Zhang L: Experimental therapy of a platelet-activating factor antagonist (ginkgolide B) on photochemically induced thrombotic cerebral ischaemia in tree shrews. *Clin Exp Pharmacol Physiol* 26: 824-825, 1999.
- Keuker JJ, de Biurrun G, Luiten PG and Fuchs E: Preservation of hippocampal neuron numbers and hippocampal subfield volumes in behaviorally characterized aged tree shrews. *J Comp Neurol* 468: 509-517, 2004.
- Yang H, He BR and Hao DJ: Biological roles of olfactory ensheathing cells in facilitating neural regeneration: A systematic review. *Mol Neurobiol* 51: 168-179, 2014.
- Lang BC, Zhang Z, Lv LY, Liu J, Wang TY, Yang LH, Liao DQ, Zhang WS and Wang TH: OECs transplantation results in neuropathic pain associated with BDNF regulating ERK activity in rats following cord hemisection. *BMC Neurosci* 14: 80, 2013.
- Pastrana E, Moreno-Flores MT, Avila J, Wandosell F, Minichiello L and Diaz-Nido J: BDNF production by olfactory ensheathing cells contributes to axonal regeneration of cultured adult CNS neurons. *Neurochem Int* 50: 491-498, 2007.
- He BR, Xie ST, Wu MM, Hao DJ and Yang H: Phagocytic removal of neuronal debris by olfactory ensheathing cells enhances neuronal survival and neurite outgrowth via p38MAPK Activity. *Mol Neurobiol* 49: 1501-1512, 2014.
- Huo SJ, Li Y, Raisman G and Yin ZQ: Transplanted olfactory ensheathing cells reduce the gliotic injury response of Muller cells in a rat model of retinitis pigmentosa. *Brain Res* 1382: 238-244, 2011.
- Sandvig I, Hoang L, Sardella TC, Barnett SC, Brekken C, Tvedt K, Berry M, Haraldseth O, Sandvig A and Thuen M: Labelling of olfactory ensheathing cells with micron-sized particles of iron oxide and detection by MRI. *Contrast Media Mol Imaging* 7: 403-410, 2012.
- Sasaki M, Lankford KL, Radtke C, Honmou O and Kocsis JD: Remyelination after olfactory ensheathing cell transplantation into diverse demyelinating environments. *Exp Neurol* 229: 88-98, 2011.
- Radtke C and Kocsis JD: Peripheral nerve injuries and transplantation of olfactory ensheathing cells for axonal regeneration and remyelination: Fact or fiction? *Int J Mol Sci* 13: 12911-12924, 2012.
- Andrews MR and Stelzner DJ: Evaluation of olfactory ensheathing and schwann cells after implantation into a dorsal injury of adult rat spinal cord. *J Neurotrauma* 24: 1773-1792, 2007.
- Sun T, Ye C, Zhang Z, Wu J and Huang H: Cotransplantation of olfactory ensheathing cells and Schwann cells combined with treadmill training promotes functional recovery in rats with contused spinal cords. *Cell Transplant* 22 (Suppl 1): S27-S38, 2013.
- Wang YC, Xia QJ, Ba YC, Wang TY, Lin N, Zou Y, Shang FF, Zhou XF, Wang TH, Fu XM and Qi JG: Transplantation of olfactory ensheathing cells promotes the recovery of neurological functions in rats with traumatic brain injury associated with downregulation of Bad. *Cytotherapy* 16: 1000-1010, 2014.
- Aranda PS, LaJoie DM and Jorczyk CL: Bleach gel: A simple agarose gel for analyzing RNA quality. *Electrophoresis* 33: 366-369, 2012.
- Vanella L, Barbagallo I, Acquaviva R, Di Giacomo C, Cardile V, Abraham NG and Sorrenti V: Ellagic acid: Cytodifferentiating and antiproliferative effects in human prostatic cancer cell lines. *Curr Pharm Des* 19: 2728-2736, 2013.
- Woodhouse A, Vincent AJ, Kozel MA, Chung RS, Waite PM, Vickers JC, West AK and Chuah MI: Spinal cord tissue affects ensheathing cell proliferation and apoptosis. *Neuroreport* 16: 737-740, 2005.
- Raucci F, Tiong JD and Wray S: P75 nerve growth factor receptors modulate development of GnRH neurons and olfactory ensheathing cells. *Front Neurosci* 7: 262, 2013.
- Dowling MR, Kan A, Heinzel S, Zhou JH, Marchingo JM, Wellard CJ, Markham JF and Hodgkin PD: Stretched cell cycle model for proliferating lymphocytes. *Proc Natl Acad Sci USA* 111: 6377-6382, 2014.
- Gallagher ZR, Johnston ST and Czaja K: Neural proliferation in the dorsal root ganglia of the adult rat following capsaicin-induced neuronal death. *J Comp Neurol* 522: 3295-3307, 2014.

33. Lubelchek RJ, Max B, Sandusky CJ, Hota B and Barker DE: Reliability at the lower limits of HIV-1 RNA quantification in clinical samples: A comparison of RT-PCR versus bDNA assays. *PLoS One* 4: e6008, 2009.
34. Senechal B and James VL: Ten years of external quality assessment of human immunodeficiency virus type 1 RNA quantification. *J Clin Microbiol* 50: 3614-3619, 2012.
35. Rodriguez RA, Polston PM, Wu MJ, Wu J and Sobsey MD: An improved infectivity assay combining cell culture with real-time PCR for rapid quantification of human adenoviruses 41 and semi-quantification of human adenovirus in sewage. *Water Res* 47: 3183-3191, 2013.
36. Hong Y, Davis B, Marron JS, Kwitt R, Singh N, Kimbell JS, Pitkin E, Superfine R, Davis SD, Zdanski CJ and Niethammer M: Statistical atlas construction via weighted functional boxplots. *Med Image Anal* 18: 684-698, 2014.
37. Gleiss A, Sanchez-Cabo F, Perco P, Tong D and Heinze G: Adaptive trimmed t-statistics for identifying predominantly high expression in a microarray experiment. *Stat Med* 30: 52-61, 2011.
38. Diemel LT, Jackson SJ and Cuzner ML: Role for TGF-beta1, FGF-2 and PDGF-AA in a myelination of CNS aggregate cultures enriched with macrophages. *J Neurosci Res* 74: 858-867, 2003.
39. Cannella B, Pitt D, Capello E and Raine CS: Insulin-like growth factor-1 fails to enhance central nervous system myelin repair during autoimmune demyelination. *Am J Pathol* 157: 933-943, 2000.
40. Copelman CA, Cuzner ML, Groome N and Diemel LT: Temporal analysis of growth factor mRNA expression in myelinating rat brain aggregate cultures: Increments in CNTF, FGF-2, IGF-I, and PDGF-AA mRNA are induced by antibody-mediated demyelination. *Glia* 30: 342-351, 2000.
41. Saito F, Kanagawa M, Ikeda M, Hagiwara H, Masaki T, Ohkuma H, Shimizu T, Sonoo M, Toda T and Matsumura K: Overexpression of LARGE suppresses muscle regeneration via down-regulation of insulin-like growth factor 1 and aggravates muscular dystrophy in mice. *Hum Mol Genet* 23: 4543-4558, 2014.
42. Sheng, MH, Lau KH and Baylink DJ: Role of osteocyte-derived insulin-like growth factor I in developmental growth, modeling, remodeling, and regeneration of the bone. *J Bone Metab* 21: 41-54, 2014.
43. Gezginci-Oktayoglu S, Karatug A and Bolkent S: The relation among NGF, EGF and insulin is important for triggering pancreatic beta cell apoptosis. *Diabetes Metab Res Rev* 28: 654-662, 2012.
44. Munemasa Y, Chang CS, Kwong JM, Kyung H, Kitaoka Y, Caprioli J and Piri N: The neuronal EGF-related gene Nell2 interacts with Macf1 and supports survival of retinal ganglion cells after optic nerve injury. *PLoS One* 7: e34810, 2012.
45. Nowacka M and Obuchowicz E: BDNF and VEGF in the pathogenesis of stress-induced affective diseases: An insight from experimental studies. *Pharmacol Rep* 65: 535-546, 2013.
46. Awuah PK, Nejak-Bowen KN and Monga SP: Role and regulation of PDGFRalpha signaling in liver development and regeneration. *Am J Pathol* 182: 1648-1658, 2013.
47. McGowan SE and McCoy DM: Platelet-derived growth factor-A and sonic hedgehog signaling direct lung fibroblast precursors during alveolar septal formation. *Am J Physiol Lung Cell Mol Physiol* 305: L229-L239, 2013.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.