

# Identification of *CRB1* mutations in two Chinese consanguineous families exhibiting autosomal recessive retinitis pigmentosa

XIAOXIN GUO<sup>1\*</sup>, JIE LI<sup>2\*</sup>, QINGWEI WANG<sup>1,3\*</sup>, YI SHU<sup>1</sup>, JIN WANG<sup>1</sup>, LIJIA CHEN<sup>4</sup>, HOUBIN ZHANG<sup>1</sup>, YI SHI<sup>1</sup>, JIYUN YANG<sup>1</sup>, FANG LU<sup>1</sup>, LI JIANG<sup>1</sup>, CHAO QU<sup>2</sup> and BO GONG<sup>1,4</sup>

<sup>1</sup>Key Laboratory for Human Disease Gene Study of Sichuan Province and Department of Laboratory Medicine,

<sup>2</sup>Department of Ophthalmology, Sichuan Academy of Medical Sciences and Sichuan Provincial People's Hospital, School of Medicine, University of Electronic Science and Technology of China, Chengdu, Sichuan 610072;

<sup>3</sup>Institute of Chengdu Biology, Sichuan Translational Medicine Hospital, Chinese Academy of Sciences, Chengdu, Sichuan 610000; <sup>4</sup>Department of Ophthalmology and Visual Sciences,

The Chinese University of Hong Kong, Hong Kong, SAR, P.R. China

Received January 24, 2019; Accepted June 12, 2019

DOI: 10.3892/mmr.2019.10495

**Abstract.** Retinitis pigmentosa (RP) is a leading cause of inherited blindness characterized by progressive loss of retinal photoreceptor cells. The present study aimed to identify the causative gene mutations in two Chinese families with autosomal recessive retinitis pigmentosa (arRP). Two Chinese consanguineous arRP families (RP-2284 and RP-2360) were recruited in this study, involving totally three affected and 25 unaffected members. All the affected members underwent a complete ophthalmic examination, including fundus photography, multifocal electroretinography (ERG) and full field ERG. Exome sequencing was performed on the three RP patients in the two families, followed by direct Sanger sequencing in all the family members and in 1,260 unrelated controls for validation of the mutations identified. Two homozygous missense mutations in the crumbs homolog 1 (*CRB1*) gene, which is known to cause severe retinal dystrophies, were found

to be related to the phenotype of the two arRP families. The homozygous missense mutation c.1997 T>A in *CRB1* was detected in two patients in the RP-2284 family. The proband in the RP-2360 family was the only RP patient and was found to carry the novel homozygous missense mutation c.2426 A>C in *CRB1*. The two mutations were heterozygous or absent in the other healthy family members, and they were absent in the 1,260 controls. The amino acid changes in the CRB1 protein affected by the two mutations were predicted to be damaging by Polyphen-2. Our study reported two *CRB1* mutations causing arRP in two Chinese families, which expands the *CRB1* mutation spectrum of RP in the Chinese population and emphasizes the causative role of *CRB1* in RP.

## Introduction

Retinitis pigmentosa (RP) is caused by a progressive degeneration of photoreceptor cells that initially primarily affect the rods whereas the function of the cones is compromised as the disease progresses (1). Due to its heterogeneity and diversity of inheritance patterns, the clinical presentation of RP is variable. Some RP patients lose vision in their childhood, whereas other patients retain normal vision until mid-adulthood. The clinical symptoms of RP include night blindness and gradual loss of the peripheral visual field caused by rod photoreceptor lost. Later with the loss of cone photoreceptors, the central vision is impaired and patients may ultimately suffer from legal blindness ultimately (1). The worldwide prevalence of RP is approximately 1 in 4,000, or more than 1 million individuals (2).

RP is clinically and genetically heterogeneous. It can be inherited as an autosomal dominant (30-40%), autosomal recessive (50-60%), or X-linked (5-15%) trait (3). Some patients who have no known family history are classified as simplex RP. Autosomal recessive RP (arRP) is the most common form of RP worldwide. To date, more than 80 genes (<https://sph.uth.edu/retnet/>) have been implicated in syndromic and non-syndromic forms of RP. However, these identified

*Correspondence to:* Dr Bo Gong, Key Laboratory for Human Disease Gene Study of Sichuan Province and Department of Laboratory Medicine, Sichuan Academy of Medical Sciences and Sichuan Provincial People's Hospital, School of Medicine, University of Electronic Science and Technology of China, 32 Road West 2, The First Ring, Chengdu, Sichuan 610072, P.R. China  
E-mail: gongbo2007@hotmail.com

Dr Chao Qu, Department of Ophthalmology, Sichuan Academy of Medical Sciences and Sichuan Provincial People's Hospital, School of Medicine, University of Electronic Science and Technology of China, 32 Road West 2, The First Ring, Chengdu, Sichuan 610072, P.R. China  
E-mail: lucyjeffersonqu@hotmail.com

\*Contributed equally

**Key words:** autosomal recessive retinitis pigmentosa, crumbs homolog 1 gene, *CRB1*, homozygous mutation, exome sequencing

genes account for no more than 60% of RP cases, and there has been limited success in screening known genes for RP using conventional Sanger sequencing. Recently, exome sequencing has been successfully used to identify genes associated with Mendelian disorders (4,5). Coupled with DNA capture technology, the next-generation sequencing (NGS) platform enables rapid and cost-effective parallel sequencing of a large panel of disease genes. In some recent studies, NGS has provided a promising alternative for the molecular diagnosis and gene identification of RP (6-10).

In the present study, the identification of disease-causing mutations of RP was presented by using exome sequencing in two Chinese families with arRP. This report concerns the clinical features and genetic findings of the two arRP families.

## Patients and methods

**Study subjects.** Two arRP pedigrees with Chinese origin were recruited from Sichuan Provincial People's Hospital, including a four-generation family with two consanguineous arRP patients and 16 unaffected members (RP-2284; Fig. 1A), and a four-generation family with one consanguineous RP patient and nine unaffected members (RP-2360; Fig. 1B). In addition, 1,260 unrelated healthy individuals of Chinese Han origin were also recruited. This study was approved by the Institutional Review Board of the Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China. Informed written consent was obtained from each individual prior to participation in this study. All procedures were carried out in accordance with the tenets of the Declaration of Helsinki.

**Clinical diagnosis.** The clinical information of the two families is listed in Table I. Each family member underwent a complete ophthalmic examination, including best-corrected visual acuity (BCVA), slit-lamp biomicroscopy, fundus photography if possible, visual field testing (Octopus; Interzeag AG) and electroretinography (ERG). ERGs were performed using a multifocal ERG recorder (GT-2008V-IV; Chongqing Kanghua Ruiming S&T Co., Ltd) and corneal contact lens electrodes. The ERG protocol complied with the standards of the International Society for Clinical Electrophysiology of Vision. Diagnosis of RP in the three patients was based on poor night vision at an early age, followed by progressive loss of peripheral visual fields, and extinguished ERG responses, along with abnormal fundus changes (2). In contrast, all the unaffected family members, and the unrelated control individuals had no signs of retinal diseases.

**DNA extraction.** Genomic DNA were extracted from the peripheral blood of all family members (RP-2284 and RP-2360) and 1,260 controls, using a TIANamp Blood DNA kit (cat. no. DP348) according to the manufacturer's protocol (Tiangen Biotech Co., Ltd.).

**Exome sequencing and variant detection.** Exome sequencing was performed on patients IV:2 and IV:4 in RP-2284 and patient IV:3 in RP-2360 by Genergy Biotechnology, Co., Ltd. The sequenced samples were prepared according to the Illumina protocols of the SureSelect Target Enrichment

System Capture Process (Illumina, Inc.) and exome sequencing analysis was performed as described previously (11). Briefly, the reads were mapped against UCSC hg19 (<http://genome.ucsc.edu/>) by BWA (<http://bio-bwa.sourceforge.net/>). The SNPs and Indels were detected by SAMTOOLS (<http://samtools.sourceforge.net/>). The detected variants were annotated and filtered based on public and in-house databases, under the following conditions: i) variants within intergenic, intronic, and UTR regions and synonymous mutations were excluded from downstream analysis; ii) variants in dbSNP138 (<http://www.ncbi.nlm.nih.gov/projects/SNP/>), 1000 Genome project (<ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp>), Exome Aggregation Consortium, YH Database (<http://yh.genomics.org.cn/>), the HapMap Project (<ftp://ftp.ncbi.nlm.nih.gov/hapmap>) and our in-house database generated from 2,010 samples of exome sequencing, were excluded; iii) Polyphen2 (<http://genetics.bwh.harvard.edu/pph2/>) was used to predict the possible damaging impacts of each mutation.

**Mutation validation.** The *CRB1* mutations identified by exome sequencing were further confirmed in all members of the two families and in 1,260 controls by direct Sanger sequencing, using the PCR primers (*CRB1*-c.1997T>A-F: 5'-AGTGGCATT TCGTGGAGGTA-3' and *CRB1*-c.1997T>A-R: 5'-GCTGTT TCTGCTCTGCTCTG-3'; *CRB1*-c.2426A>C-F: 5'-TTTGTC CGAACGCTTCAACC-3' and *CRB1*-c.2426A>C-R: 5'-TCT TGCTTGTCAGGTAGGCC-3') designed by Primer 3.0 (<http://primer3.ut.ee/>) and synthesized by Invitrogen (Thermo Fisher Scientific, Inc.). Direct sequencing was performed in an ABI 3130XL genetic analyzer according to ABI BigDye sequencing protocols (Thermo Fisher Scientific, Inc.).

## Results

**Patients and clinical information.** Two Chinese consanguineous arRP families (RP-2284 and RP-2360, Fig. 1) were recruited in this study. Two subjects (IV:2 and IV:4) were diagnosed with RP in family RP-2284, which exhibited a pattern of recessive inheritance. They presented typical clinical symptoms, including night blindness rapidly progressing to severe visual disability. The fundus examination of the proband (IV:2) showed pallor in the optic nerve head, retinal artery attenuation, and mid-peripheral diffuse and dense bone-spicule pigmentation (Fig. 2A and B). Another affected member (IV:4) had similar clinical manifestations to the proband and presented with macular degeneration (Fig. 2C). Family RP-2360 included one RP patient (IV:3, the proband), who had experienced night blindness since 23 years of age and progressive bilateral visual loss (Table I). She presented with early-onset and markedly decreased visual acuity (OD: 20/400, OS: 20/400) in both eyes (Table I). The fundus image in the right eye was blurred due to dense posterior subcapsular cataract. Fundus examination of her left eye showed similar features to those of patient IV:2 in RP-2284 (data not shown). None of the three patients exhibited glaucoma features, and their ERG examination showed no recordable response under either scotopic or photopic condition, indicating significant loss of function in the both rods and cones (Fig. 2D and E).

Table I. Genotypes and phenotypes of the two RP family members.

## A, Family RP-2284

Family member	Age (years)/sex	Phenotype	Age at onset (years)	Visual acuity (OD/OS)	Fundus appearance	Mutation	Mutation type
V:1	6/F	Normal	-	-	-	C.1997 T>A	Het
IV:1	30/F	Normal	-	1.0/1.0	Normal	No mutation	-
IV:2	31/M	RP	24	Light perception	RVA and DBSP	C.1997 T>A	Hom
IV:3	28/M	Normal	-	1.0/1.0	Normal	C.1997 T>A	Het
IV:4	23/M	RP	22	Counting fingers	RVA, DBSP and MD	C.1997 T>A	Hom
IV:5	26/M	Normal	-	1.0/1.0	Normal	No mutation	-
III:1	50/M	Normal	-	-	-	C.1997 T>A	Het
III:2	52/F	Normal	-	1.0/0.9	Normal	No mutation	-
III:3	54/F	Normal	-	1.0/1.0	Normal	C.1997 T>A	Het
III:4	55/M	Normal	-	1.0/0.8	Normal	C.1997 T>A	Het
III:5	53/F	Normal	-	-	-	C.1997 T>A	Het
III:6	50/F	Normal	-	0.8/0.8	Normal	No mutation	-
III:7	48/F	Normal	-	-	-	C.1997 T>A	Het
III:8	49/F	Normal	-	0.8/0.8	Normal	No mutation	-
III:9	51/M	Normal	-	0.9/0.8	Normal	C.1997 T>A	Het
III:10	48/F	Normal	-	0.8/1.0	Normal	C.1997 T>A	Het
III:11	46/F	Normal	-	-	-	No mutation	-
II:1	78/M	Cataract	-	-	-	C.1997 T>A	Het

## B, Family RP-2360

Family member	Age (years)/sex	Phenotype	Age at onset (years)	Visual acuity (OD/OS)	Fundus appearance	Mutation	Mutation type
V:1	18/F	Normal	-	-	Normal	C.2426 A>C	Het
IV:1	41/F	Normal	-	1.0/1.0	Normal	C.2426 A>C	Het
IV:2	45/M	Normal	-	1.0/0.9	Normal	No mutation	-
IV:3	43/F	RP	23	Counting fingers	OS: RVA and DBSP	C.2426 A>C	Hom
IV:4	46/F	Normal	-	-	-	No mutation	-
III:1	70/M	Normal	-	0.8/0.7	Normal	C.2426 A>C	Het
III:2	69/F	Normal	-	0.7/0.8	Normal	C.2426 A>C	Het
III:3	67/F	Normal	-	-	-	C.2426 A>C	Het
III:4	66/F	Normal	-	0.8/0.8	Normal	No mutation	-
II:1	92/M	Cataract	-	-	-	No mutation	-

RP, retinitis pigmentosa; Hom, homozygous; Het, heterozygous; F, female; M, male; RVA, retinal vascular attenuation; DBSP, dense bone-spicule pigmentation; MD, macular degeneration; OD, right eye; OS, left eye; -, not available.

**Exome sequencing and data analysis.** Exome sequencing was performed on two patients (IV:2 and IV:4) in the RP-2284 family and on one patient (IV:3) in the RP-2360 family with a mean read depth of target regions of 52.3x. There were 18,793 SNPs in the coding regions (8,763 non-synonymous SNPs, 9,836 synonymous SNPs, and 873 other types of SNPs) and 387 coding indels identified. To identify the disease-causing mutations, the researchers paid close attention

to the functional SNPs/indels in homozygous or compound heterozygous status including frameshift coding-region insertion or deletions and non-synonymous variants which were more likely to be pathogenic. Detected variants were compared with HapMap Project, dbSNP138, 1000 Genomes Project, Exome Aggregation Consortium, YH Database, and our in-house database. The rare homozygous variants shared by the two affected patients in RP-2284 and RP-2360 families




Figure 1. Pedigrees of two Chinese consanguineous families with arRP. (A and B) Two arRP families were recruited in this study. The arrow indicates the proband in each family. Affected individuals are indicated by filled symbols and open symbols indicated unaffected individuals. RP-2284, a four-generation family with two consanguineous arRP patients and 16 unaffected members; RP-2360, a four-generation family with one consanguineous RP patient and nine unaffected members are shown. arRP, autosomal recessive retinitis pigmentosa.




Figure 2. Representative clinical features of arRP patients. (A-C) Compared to the normal individual (IV:3), peripheral pigmentation and retinal vascular attenuation were noted in the proband's fundus photograph, as well as peripheral diffuse pigmentation complicated with macular degeneration in another patient (IV:4) in family RP-2284. (D and E) ERG records showed no detectable rod and cone responses in either eye of the proband in family RP-2284. OD, right eye; OS, left eye. arRP, autosomal recessive retinitis pigmentosa.

are listed in Table S1. Using the autosomal recessive mode, two homozygous mutations, c.1997 T>A and c.2426 A>C, were identified in the *CRB1* gene (NM\_201253.2) in the RP-2284 and RP-2360 families, respectively.

**Mutation validation and analysis.** In the RP-2284 family, the presence of homozygous mutation c.1997 T>A in two patients (IV:2 and IV:4) was confirmed by Sanger sequencing (Fig. 3B). The parents and other unaffected siblings of these two patients were either unaffected heterozygous carriers of c.1997 T>A, or had a normal *CRB1* sequence without any mutation (Table I). In the RP-2360 family, direct Sanger sequencing confirmed the presence of homozygous mutation c.2426 A>C (Fig. 3A) in the

proband (IV:3), whereas all other unaffected family members were heterozygotes or normal. The two homozygous mutations described above were absent in the 1,260 Han Chinese healthy controls. These results reveal a complete co-segregation of the two *CRB1* mutations with RP within the respective families. The genotypes of the members of families RP-2284 and RP-2360 are shown in Table I.

The c.1997 T>A (p.V666D) homozygous mutation is expected to change valine (hydrophobic non-polar) to aspartic acid (hydrophilic positively charged) at codon 666 in exon 6. The substituted amino acid is predicted to alter the hydrophobicity and chargeability of the CRB1 protein. The c.2426 A>C (p.Q809T) mutation introduced a substitution




Figure 3. *CRB1* mutations were identified in two Chinese consanguineous arRP families. (A) A normal sequence of *CRB1* from an unaffected member, a heterozygous A to C substitution at codon 2,426 from the proband's parents, and a homozygous mutation from the patients were identified in family RP-2284. (B) A normal sequence of *CRB1* from the proband's sister, a heterozygous T to A substitution at codon 1,997 from the proband's parents, and a homozygous mutation from the proband were identified in family RP-2236. arRP, autosomal recessive retinitis pigmentosa; *CRB1*, crumbs homolog 1.




Figure 4. Orthologous protein sequence alignment and structure diagram of *CRB1*. (A) Orthologous protein sequence alignment showed the seven mutation sites occurring in a highly conserved region of *CRB1* among different species. (B) Fifteen mutations that were identified in Chinese RP patients occurred mainly at exon 6, 7 and 9 within a laminin A globular (AG)-like domain. The boxed mutations in red were newly identified in the present study. RP, retinitis pigmentosa; *CRB1*, crumbs homolog 1.

of glutamine for threonine at codon 809 in exon 7 (Fig. 4B). Comparative amino acid sequence alignment of the *CRB1*

proteins across human and other species revealed that the two mutations occurred at highly conserved positions (Fig. 4A).

These changes were predicted to be damaging to *CRB1* using Polyphen-2.

## Discussion

A previous study demonstrated that the *CRB1* gene was responsible for retinitis pigmentosa (RP) at the RP12 locus (12) and more than 150 disease-associated variants in *CRB1* have been reported (13). Most of the mutations occur in exon 7 (27%) and exon 9 (41%), which encode the second and the third laminin AG-like domains, respectively (13). In the present study, two homozygous mutations c.1997 T>A and c.2426 A>C in *CRB1* causing autosomal recessive retinitis pigmentosa (arRP) in two Chinese families were identified using exome sequencing. This result expanded the *CRB1* mutation spectrum causing RP, which may contribute to improvement of the molecular prognosis of RP.

Studies have reported that approximately 194 mutations in *CRB1* have been found to be related to the pathogenesis of arRP in diverse populations, including 189 mutations in the extracellular region (14–17). *CRB1* is a human homolog of *Drosophila* crumbs, a well conserved gene with homologue across multiple phyla (18). It is located at 1q31.3 and consists of 12 exons spanning 210 kb of genomic DNA (12). This gene encodes two different proteins of 1,376 and 1,406 amino acids, and both protein are composed of three parts: 19 epidermal growth factor (EGF)-like domains, three laminin A globular (AG)-like domains and a signal peptide sequence. Beyond that, the longer isoform also contains additional transmembrane and cytoplasmic domains (19). *CRB1* is a transmembrane protein and is mainly expressed in the inner segments of mammalian photoreceptors as well as in the brain (12,18). It plays an essential role in the development of the retina (20,21). *CRB1* is involved in photoreceptor morphogenesis in the retina (22). Mutations in *CRB1* may restrain retinal development and result in a loss of photoreceptor signaling (23). In the mouse and human retina, the protein *Crb1* or *CRB1* is concentrated in the vicinity of the outer limiting membrane and is expressed in photoreceptor inner segments, in Muller cell and the epithelial cells (24). The *Crb1* mutant in mouse showed a developmental defect of retina, including a disrupted outer limiting membrane and folded retina (25).

The present study involved two Han Chinese arRP families. An analysis of the exome sequencing data revealed that only the two homozygous mutations c.1997 T>A (p.V666D) and c.2426 A>C (p.Q809T) in *CRB1* were found to be co-segregated with arRP in the families. The two mutations have never been reported in the ExAC database (<http://exac.broadinstitute.org/>) or the human gene mutation database (<http://www.hgmd.org/>). The clinical findings associated with retinal dystrophies caused by *CRB1* mutations include congenital blindness, LCA, to early-onset rod-cone dystrophy and arRP. Mutations in *CRB1* account for 10–15% of all patients with LCA and as many as 6.5% of all patients with arRP. In this study, the age of onset of patients in the two RP families was approximately 22–24 years of age. The homozygous mutation c.2426 A>C found in family RP-2360 is located at exon 7 of the *CRB1* gene, encoding the second laminin AG-like domain. Apart from the proband (IV:3) who carried the homozygous mutation c.2426 A>C, other family members carried

either heterozygous mutation or no mutation. The proband's parents had a consanguineous marriage, and they were each heterozygous for c.2426 A>C, which was also found in members V:1, IV:1 and III:3 (Table I). As previously reported, laminin AG-like domain is predicted to affect interactions among proteins, calcium binding, and protein folding (26). This mutation introduces a substitution of glutamine to threonine, which may dramatically affect the function of the second laminin AG-like domain of the *CRB1* protein (PolyPhen2 scores ~0.824). There were many disease causative mutations reported in the laminin AG-like domain, which indicates that this domain plays an important role in *CRB1* function (24). The mutation c.1997 T>A identified in family RP-2284, was previously reported by Wang *et al* as present in RP patients in compound heterozygote (27), which further confirmed its pathogenic role in RP. In this pedigree with a consanguineous marriage, the proband's parents (III:3 and III:4), daughter (V:1), aunts (III:5 and III:7), uncle (III:1) and brother (IV:3), as well as another patient's parents (III:9 and III:10), were found to carry the heterozygous mutation (Table I). The change could convert valine to aspartic acid at codon 666 (p.V666 D). This substitution was predicted to be damaging by Polyphen-2 and to alter the hydrophobicity and chargeability of the *CRB1* protein. Nevertheless, how these mutations exactly affect the function of *CRB1* protein remains unclear. Further functional research is essential to elucidate the underlying mechanism governing how the two homozygous mutations result in RP.

In conclusion, the present study revealed two homozygous mutations c.1997 T>A (p.V 666D) and c.2426 A>C (p.Q809T) in the *CRB1* gene in two consanguineous Chinese families with arRP via exome sequencing. These data expand the *CRB1* mutation spectrum causing RP and may contribute to an improvement in the molecular diagnosis for retinal dystrophies.

## Acknowledgements

Not applicable.

## Funding

The present study was supported by grants from the Natural Science Foundation of China (81670853, 81700841, 81670893, 81570882 and 81670877), CAS 'Light of West China' Program (to BG), Science and Technology Department of Sichuan Province (2019JDJQ0031 and 2014FZ0124), Foundation for Technology & Science & Technology Bureau of Chengdu (2015-HM02-00094-SF and 2018-YF05-00348-SN).

## Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Authors' contributions

BG, CQ, JL, QW and JY conceived the experiment. XG, JW and YS performed the experiments. YS, FL, JY, LJ, CQ and HZ recruited the patients. BG, FL, LC, QW, LJ, CQ and HZ

analyzed the data. XG and JL drafted the manuscript. JY and HZ critically revised the manuscript for important intellectual content. All authors read and approved the manuscript and agreed to be accountable for all aspects of the research in ensuring that the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### Ethics approval and consent to participate

This study was approved by the Institutional Review Board of the Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China. Informed written consent was obtained from each individual prior to participation in this study. All procedures were carried out in accordance with the tenets of the Declaration of Helsinki.

### Patient consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### References

- Bird AC: Retinal photoreceptor dystrophies LI. Edward Jackson Memorial Lecture. Am J Ophthalmol 119: 543-562, 1995.
- Hartong DT, Berson EL and Dryja TP: Retinitis pigmentosa. Lancet 368: 1795-1809, 2006.
- Rivolta C, Sharon D, DeAngelis MM and Dryja TP: Retinitis pigmentosa and allied diseases: Numerous diseases, genes, and inheritance patterns. Hum Mol Genet 11: 1219-1227, 2002.
- Ng SB, Turner EH, Robertson PD, Flygare SD, Bigham AW, Lee C, Shaffer T, Wong M, Bhattacharjee A, Eichler EE, et al: Targeted capture and massively parallel sequencing of 12 human exomes. Nature 461: 272-276, 2009.
- Ng SB, Buckingham KJ, Lee C, Bigham AW, Tabor HK, Dent KM, Huff CD, Shannon PT, Jabs EW, Nickerson DA, et al: Exome sequencing identifies the cause of a mendelian disorder. Nat Genet 42: 30-35, 2010.
- Neveling K, Collin RW, Gilissen C, van Huet RA, Visser L, Klint MP, Gijsen SJ, Zonneveld MN, Wieskamp N, de Ligt J, et al: Next-generation genetic testing for retinitis pigmentosa. Hum Mutat 33: 963-972, 2012.
- O'Sullivan J, Mullaney BG, Bhaskar SS, Dickerson JE, Hall G, O'Grady A, Webster A, Ramsden SC and Black GC: A paradigm shift in the delivery of services for diagnosis of inherited retinal disease. J Med Genet 49: 322-326, 2012.
- Katagiri S, Akahori M, Sergeev Y, Yoshitake K, Ikeo K, Furuno M, Hayashi T, Kondo M, Ueno S, Tsunoda K, et al: Whole exome analysis identifies frequent CNGA1 mutations in Japanese population with autosomal recessive retinitis pigmentosa. PLoS One 9: e108721, 2014.
- Wu J, Chen L, Tam OS, Huang XF, Pang CP and Jin ZB: Whole exome sequencing reveals genetic predisposition in a large family with retinitis pigmentosa. Biomed Res Int 2014: 302487, 2014.
- Villanueva A, Willer JR, Bryois J, Dermitzakis ET, Katsanis N and Davis EE: Whole exome sequencing of a dominant retinitis pigmentosa family identifies a novel deletion in PRPF31. Invest Ophthalmol Vis Sci 55: 2121-2129, 2014.
- Zhou Y, Shuai P, Li X, Wang J, Yang Y, Hao F, Lin H, Zhang D and Gong B: Association of SOD2 polymorphisms with primary open angle glaucoma in a Chinese population. Ophthalmic Genet 36: 43-49, 2015.
- den Hollander AI, ten Brink JB, de Kok YJ, van Soest S, van den Born LI, van Driel MA, van de Pol DJ, Payne AM, Bhattacharya SS, Kellner U, et al: Mutations in a human homologue of *Drosophila* crumbs cause retinitis pigmentosa (RP12). Nat Genet 23: 217-221, 1999.
- Slavotinek AM: The family of crumbs genes and human disease. Mol Syndromol 7: 274-281, 2016.
- Cordovez JA, Traboulsi EI, Capasso JE, Sadagopan KA, Ganesh A, Rychwalski PJ, Neely KA, Brodie SE and Levin AV: Retinal dystrophy with intraretinal cystoid spaces associated with mutations in the crumbs homologue (CRB1) gene. Ophthalmic Genet 36: 257-264, 2015.
- Booij JC, Florijn RJ, ten Brink JB, Loves W, Meire F, van Schooneveld MJ, de Jong PT and Bergen AA: Identification of mutations in the AIPL1, CRB1, GUCY2D, RPE65, and RPGRIP1 genes in patients with juvenile retinitis pigmentosa. J Med Genet 42: e67, 2005.
- Beryozkin A, Zelinger L, Bandah-Rozenfeld D, Harel A, Strom TA, Merin S, Chowers I, Banin E and Sharon D: Mutations in CRB1 are a relatively common cause of autosomal recessive early-onset retinal degeneration in the Israeli and Palestinian populations. Invest Ophthalmol Vis Sci 54: 2068-2075, 2013.
- Chen Y, Lin Y, Vithana EN, Jia L, Zuo X, Wong TY, Chen LJ, Zhu X, Tam PO, Gong B, et al: Common variants near ABCA1 and in PMM2 are associated with primary open-angle glaucoma. Nat Genet 46: 1115-1119, 2014.
- Richard M, Roepman R, Aartsen WM, van Rossum AG, den Hollander AI, Knust E, Wijnholds J and Cremers FP: Towards understanding CRUMBS function in retinal dystrophies. Hum Mol Genet 15: R235-R243, 2006.
- den Hollander AI, Johnson K, de Kok YJ, Klebes A, Brunner HG, Knust E and Cremers FP: CRB1 has a cytoplasmic domain that is functionally conserved between human and *Drosophila*. Hum Mol Genet 10: 2767-2773, 2001.
- Tepass U: Crumbs, a component of the apical membrane, is required for zonula adherens formation in primary epithelia of *Drosophila*. Dev Biol 177: 217-225, 1996.
- Pellikka M, Tanentzapf G, Pinto M, Smith C, McGlade CJ, Ready DF and Tepass U: Crumbs, the *Drosophila* homologue of human CRB1/RP12, is essential for photoreceptor morphogenesis. Nature 416: 143-149, 2002.
- Beheshtian M, Saeed Rad S, Babanejad M, Mohseni M, Hashemi H, Eshghabadi A, Hajizadeh F, Akbari MR, Kahrizi K, Razi Esfahani M and Najmabadi H: Impact of whole exome sequencing among Iranian patients with autosomal recessive retinitis pigmentosa. Arch Iran Med 18: 776-785, 2015.
- Thompson DA, Jancke AR, Lange J, Feathers KL, Hübner CA, McHenry CL, Stockton DW, Rammesmayr G, Lupski JR, Antinolo G, et al: Retinal degeneration associated with RDH12 mutations results from decreased 11-cis retinal synthesis due to disruption of the visual cycle. Hum Mol Genet 14: 3865-3875, 2005.
- Bujakowska K, Audo I, Mohand-Saïd S, Lancelot ME, Antonio A, Germain A, Léveillard T, Letexier M, Saraiva JP, Lonjou C, et al: CRB1 mutations in inherited retinal dystrophies. Hum Mutat 33: 306-315, 2012.
- Mehalow AK, Kameya S, Smith RS, Hawes NL, Denegre JM, Young JA, Bechtold L, Haider NB, Tepass U, Heckenlively JR, et al: CRB1 is essential for external limiting membrane integrity and photoreceptor morphogenesis in the mammalian retina. Hum Mol Genet 12: 2179-2189, 2003.
- Zhou Y, Tao S, Chen H, Huang L, Zhu X, Li Y, Wang Z, Lin H, Hao F, Yang Z, et al: Exome sequencing analysis identifies compound heterozygous mutation in ABCA4 in a Chinese family with Stargardt disease. PLoS One 9: e91962, 2014.
- Wang H, Wang X, Zou X, Xu S, Li H, Soens ZT, Wang K, Li Y, Dong F, Chen R and Sui R: Comprehensive molecular diagnosis of a large Chinese Leber congenital Amaurosis cohort. Invest Ophthalmol Vis Sci 56: 3642-3655, 2015.