

COL12A1, a novel potential prognostic factor and therapeutic target in gastric cancer

XIAOXIA JIANG^{1,2}, MENGJIE WU^{1,2}, XIN XU^{1,2}, LIWEI ZHANG³, YINGYING HUANG^{1,2}, ZHENZHEN XU^{1,2}, KUIFENG HE¹, HAIYONG WANG^{1,2}, HAOHAO WANG^{1,2} and LISONG TENG^{1,2}

¹Cancer Center, The First Affiliated Hospital, College of Medicine, Zhejiang University;

²Key Laboratory of Precision Diagnosis and Treatment for Hepatobiliary and Pancreatic Tumor of Zhejiang Province, Hangzhou, Zhejiang 310003; ³Department of Obstetrics and Gynecology, Renji Hospital, School of Medicine, Shanghai Jiaotong University, Shanghai 200127, P.R. China

Received January 30, 2019; Accepted July 5, 2019

DOI: 10.3892/mmr.2019.10548

Abstract. Dysregulation of collagen type XII $\alpha 1$ chain (*COL12A1*) has been found in several cancer types and could be involved in tumor progression. However, its clinical significance in gastric cancer (GC) remains under exploration. Online databases (Gene Expression Omnibus and UALCAN), reverse transcription-quantitative PCR and immunohistochemistry were utilized in the present study to evaluate the expression of *COL12A1* in GC tissues and cell lines. It was found that *COL12A1* expression was notably upregulated in GC. Clinicopathological analysis showed that elevated *COL12A1* expression was positively correlated with tumor invasiveness, metastasis and advanced clinical stage. The prognostic analysis suggested that high *COL12A1* expression contributed to poor overall survival. Multivariate Cox analysis indicated that *COL12A1* overexpression was a powerful independent prognostic indicator in patients with GC (hazard ratio, 1.896; 95% CI, 1.267-2.837; $P=0.002$). The results highlighted the importance of *COL12A1* in GC and suggested its potential role as a candidate for clinical outcome prediction and targeted therapy in patients with GC.

Introduction

Gastric cancer (GC), as one of the most common malignancies, is frequently accompanied by an unfavorable prognosis and low survival rate (1). Although progress has been made in therapeutic strategies, and early detection and clinical management have improved its prognosis, GC remains a complex clinical challenge, with the majority of patients

being diagnosed during the later stages of the disease (2,3). Previously, it was reported that the aberrant expression of specific biomarkers was related to cancer development and a worse prognosis, and that some biomarkers affected the sensitivity of cancer to chemotherapeutic drugs (4). Therefore, it is of great benefit to identify novel biomarkers characterized by high sensitivity as well as high specificity that can be used as predictors for GC diagnosis and precise treatment.

Collagen type XII $\alpha 1$ chain (*COL12A1*), encoded by a gene whose chromosomal location is 6q12-q13, is a member of the fibril-associated collagen family harboring interrupted triple-helical collagen domains (5). *COL12A1* acts as a bridge between fibrils and its mutations are reported to be related to myopathy (6). *COL12A1* has received increasing attention due to its essential roles in human cancer, given that its overexpression has been identified in several different cancer types. A comparison of differentially expressed genes between 279 metastatic samples from multiple adenocarcinoma types and unmatched primary adenocarcinomas revealed elevated expression of collagen genes, including *COL12A1*, closely related to poor prognosis in several cancer types (7-9). Notably, *COL12A1* was identified as a desmoplastic marker during myofibroblast differentiation in colorectal cancer (CRC), suggesting that *COL12A1* upregulation is related to the malignant potential of the tumor (10). Besides, differential gene expression from microarray analyses showed that *COL12A1* was highly expressed in renal cancer (8). Moreover, in ovarian cancer, *COL12A1* expression exhibited a significantly increased level of expression in cisplatin- and doxorubicin-resistant cell lines relative to parental cell lines, and was associated with poor overall survival (OS) (7).

Although these results suggest a relationship between *COL12A1* and cancer development and progression, its significance in GC remains an underexplored issue. Therefore, the present study evaluated *COL12A1* expression using online databases, followed by quantification of *COL12A1* mRNA and protein levels in clinical samples via reverse transcription-quantitative PCR (RT-qPCR) and immunohistochemistry (IHC), respectively. Furthermore, associations between *COL12A1* expression and clinical features, as well as the clinical outcomes of patients with GC, were assessed.

Correspondence to: Professor Lisong Teng, Cancer Center, The First Affiliated Hospital, College of Medicine, Zhejiang University, 79 Qingchun Road, Hangzhou, Zhejiang 310003, P.R. China
E-mail: lsteng@zju.edu.cn

Key words: bioinformatics analysis, immunohistochemistry, collagen type XII $\alpha 1$ chain, extracellular matrix, prognosis

Materials and methods

Patients and specimens. To measure *COL12A1* expression, 64 paired fresh GC and matched control tissues were obtained between October 2011 and April 2014 at the First Affiliated Hospital of Zhejiang University. Table I shows the clinical features of the 64 GC patients. Paired formalin fixed paraffin-embedded (FFPE) samples (107 GC and 107 matched normal gastric tissues) were collected between January 2012 and December 2014. All 171 samples were obtained from 130 male and 41 female patients with a mean age of 64 years (range from 28-82 years) diagnosed with GC based on histopathology. Additionally, a tissue microarray containing 93 GC and 77 matched normal gastric tissues was purchased from Shanghai Outdo Biotech Co., Ltd. Patient clinicopathological information was collected; this included age, gender, tumor size, differentiation and the tumor-node-metastasis (TNM) stage pathological classification, according to the 8th edition of the American Joint Committee on Cancer guidelines (11). The study protocol was approved by the ethics committee of the First Affiliated Hospital of Zhejiang University.

Cell lines and culture conditions. Human GC cell lines (AGS, HGC-27, MKN45 and MGC-803) were obtained from the Institute of Cellular Biology (Chinese Academy of Science, Shanghai, China) and cultured in RPMI 1640 medium (Gibco; Thermo Fisher Scientific, Inc.) supplemented with 10% FBS (Gibco; Thermo Fisher Scientific, Inc.). All cells were maintained at 37°C in a 5% CO₂ atmosphere.

RNA extraction and RT-qPCR. TRIzol® reagent (Invitrogen; Thermo Fisher Scientific, Inc.) was used for total RNA extraction from fresh tissue samples and GC cell lines. Total RNA (1 µg) was reverse transcribed using the PrimeScript RT-PCR kit (Takara Bio, Inc.) with the reaction conditions: 15 min at 37°C and stored at 4°C. Then, RT-qPCR was conducted to evaluate *COL12A1* expression on a StepOnePlus system (Applied Biosystems; Thermo Fisher Scientific, Inc.) and SYBR Green (Takara Bio, Inc.) with a two-step reaction condition: 30 sec at 95°C followed by 5 sec at 95°C and 60 sec at 60°C repeated for 40 cycles. The primers were as follows: *COL12A1*, forward, 5'-CCACAGGTTCAAGAGGTCCC-3' and reverse, 5'-TGT GTTAGCCGGAACCTGGA-3'; and *GAPDH* forward, 5'-TGG GTGTGAACCATGAGAAGT-3' and reverse, 5'-TGAGTC CTTCCACGATACCAA-3'. *COL12A1* expression levels were quantified relative to those of *GAPDH*, which was used as the endogenous control, using the 2^{-ΔΔCq} method (12).

IHC analysis. FFPE tissues were cut into 4-µm sections and the IHC staining against *COL12A1* (cat. no. ab123104; monoclonal; raised in rabbit; 1:40 dilution; Abcam) was performed as previously described (13).

IHC staining evaluation. Each stained tissue section was reviewed and scored by two independent observers. Each section was examined in five fields at high magnification (x200) under a light microscope. The percentage of positive staining and the intensity of staining were scored based on previously published standards (14). The final immunoreactivity score (0-12) was equivalent to the score of the percentage

Table I. Clinicopathological parameters in 64 patients with gastric cancer.

Variable	No. cases
Age (<60 years/≥60 years)	22/42
Sex (male/female)	45/19
Tumor size (≤5 cm/>5 cm)	38/26
Lauren's classification (intestinal type/diffuse type)	26/38
Differentiation (well to moderate/poor)	28/36
Tumor invasion (T1/T2/T3/T4)	8/16/0/40
Lymph node metastasis (N0/N1/N2/N3)	16/10/11/27

of positive staining multiplied by the staining intensity score. Based on the analysis, the final score were defined as 'high' if the score was >4, and 'low' if the score was ≤4.

Bioinformatics analysis. A total of three gastric cancer gene expression profile datasets [GSE2701 (15) GSE63089 (16) and GSE79973 (17)] containing 95 GC tissues and 55 normal gastric tissues were downloaded from the Gene Expression Omnibus (GEO) database (18). The data were analyzed using the online tool GEO2R (<https://www.ncbi.nlm.nih.gov/geo/geo2r>) to analyze the mRNA expression of *COL12A1* expression in GC and normal gastric tissues. UALCAN (19) was used to identify the up-or downregulation of *COL12A1* in The Cancer Genome Atlas (TCGA) gastric cancer portal. The Human Protein Atlas (HPA) website (<http://www.proteinatlas.org/>) was used to explore the basic expression of *COL12A1* in different organs. Three datasets (15,20,21) from Oncomine database (22) were used to analysis the relationship between *COL12A1* mRNA expression and Lauren classification. The prognostic significance of *COL12A1* was evaluated in an Affymetrix array (225,664_at; Thermo Fisher Scientific, Inc.) using Kaplan Meier-plotter (KM-plotter) (<http://kmplot.com/analysis/index.php?p=service&cancer=gastric>). The autoselected best threshold value was used to identify the high and low groups.

Statistical analysis. SPSS 19.0 (IBM Corp.) and GraphPad Prism 5.01 (GraphPad Software, Inc.) software were used to perform all statistical analyses. *COL12A1* mRNA expression in GC relative to normal gastric tissues was analyzed by paired t-test. The χ^2 test was used to analyze the relationships between *COL12A1* expression and clinicopathological variables, and survival curves between high and low *COL12A1* expression were plotted using the Kaplan-Meier method. Univariate and further multivariate analyses were conducted using the Cox proportional hazards model in order to identify prognostic indicators. P<0.05 was considered to indicate a statistically significant difference.

Results

mRNA expression of *COL12A1* is significantly elevated in GC tissues and cell lines. To evaluate *COL12A1* mRNA expression in GC, three microarray datasets from GEO were analyzed and




Figure 1. *COL12A1* mRNA expression is upregulated in GC tissues and cell lines. (A) The expression of *COL12A1* in three datasets. GSE7901 contains 45 GC and 5 normal gastric tissues. GSE63089 and GSE79973 contain 45 and 10 paired GC and normal gastric tissues, respectively. (B) The data-mining analysis showed that *COL12A1* was upregulated in the TCGA gastric cancer portal (34 normal gastric samples and 415 primary tumor samples). (C) The data-mining analysis showed the expression status of *COL12A1* in the TCGA gastric cancer portal based on different TNM stages (normal gastric samples and tumor samples classified to stage1, stage2, stage3 and stage4). (D) Basic protein expression levels of *COL12A1* in normal organs. (E) Basic mRNA expression levels of *COL12A1* in normal organs. (F) *COL12A1* expression levels in 64 pairs of fresh normal gastric and GC tissues (*COL12A1*/*GAPDH*). (G) *COL12A1* expression levels in 64 patients with GC (normalized to normal samples). (H) *COL12A1* mRNA expression was examined by reverse transcription-quantitative PCR and normalized to *GAPDH* expression in four GC cell lines. *** $P < 0.001$. *COL12A1*, collagen type XII $\alpha 1$ chain; TCGA, The Cancer Genome Atlas; GC, gastric cancer.

the results revealed that *COL12A1* was highly expressed in GC (Fig. 1A; $P < 0.001$). The results from the TCGA dataset also indicated that *COL12A1* mRNA was markedly increased in GC compared with normal controls, and this was more apparent at advanced clinical stages (Fig. 1B and C). Additionally, the basic expression of *COL12A1* in different organs was explored using HPA. As shown in Fig. 1D and E, the basic expression level of *COL12A1* in the stomach was moderate compared to that in other organs. Validation of *COL12A1* expression in 64 paired frozen GC and matched controls using RT-qPCR showed a significant elevation of *COL12A1* mRNA levels in cancer samples ($P < 0.0001$; Fig. 1F and G) relative to those in adjacent normal tissues. This was consistent with expectations. In addition, as shown in Fig. 1H, *COL12A1* mRNA was highly expressed in three (HGC-27, MKN45 and MGC-803) of the four GC cell lines. Furthermore, the relationship between *COL12A1* mRNA expression and Lauren classification was evaluated using the Oncomine database. As shown in Fig. 2, *COL12A1* was markedly elevated in all the tumor tissues, regardless of Lauren classification.

***COL12A1* protein expression status in GC tissues.** The elevated *COL12A1* mRNA expression in GC suggested its carcinogenic role in this disease. To investigate its protein expression profiles in GC, IHC staining of *COL12A1* protein

was performed in a total of 200 GC and 184 corresponding normal tissues. IHC staining showed that *COL12A1* protein expression was not observed in normal control tissues in the present cohort, while it was present in the stroma surrounding the GC cells. Representative *COL12A1* protein expression images in normal or GC tissues are shown in Fig. 3A. These results suggested that *COL12A1* is exclusively expressed in GC and not in matched normal tissues, and thus it may be a highly specific biomarker of GC.

Correlation between COL12A1 protein expression and clinicopathological features. Subsequently, the association between *COL12A1* expression and clinical parameters was analyzed. Based on the median value of 4 for final immunoreactivity scoring for *COL12A1* staining, elevated *COL12A1* protein expression was present in 86 of 200 (43%) GC samples, with the other 114 (67%) samples exhibiting low expression profiles relative to normal controls. Additionally, in GC samples, the IHC scores of *COL12A1* were increased in patients with more advanced tumor stages (stage III and IV; $P < 0.01$; Fig. 3B) and in patients > 60 years old ($P < 0.01$; Fig. 3C). Furthermore, as shown in Table II, *COL12A1* protein overexpression was positively correlated with sex ($P = 0.0027$), tumor invasion ($P = 0.0001$), lymph node metastasis ($P = 0.0121$), distant metastasis ($P = 0.0366$) and advanced TNM stage ($P = 0.0048$);




Figure 2. COL12A1 protein expression is upregulated in all Lauren types. Three datasets (15,20,21) from Oncomine were used to evaluate the association between COL12A1 expression and Lauren classification. COL12A1, collagen type XII α 1 chain.




Figure 3. COL12A1 protein expression is upregulated in GC. (A) Representative images of COL12A1 IHC staining in GC samples. The IHC score is equal to the percentage of positive staining multiples staining intensity. Magnification, x200. The IHC scores were significantly higher in (B) patients of advanced stage (III and IV) and (C) in patients >60 years old. $^{**}P<0.01$. COL12A1, collagen type XII α 1 chain; IHC, immunohistochemistry; GC, gastric cancer; TNM, tumor-node-metastasis.

however, no significant associations with other clinical factors (such as age, tumor size and differentiation degree) were observed.

Survival analysis of GC patients. As described, elevated COL12A1 expression was positively related to advanced TNM stage. Therefore, the question of whether high COL12A1

Table II. Association between COL12A1 protein expression and clinicopathological characteristics in 200 patients with gastric cancer.

Characteristics	No. patients	COL12A1 expression		P-value
		High	Low	
All cases	200	86	114	
Age, years				0.764
<60	68	28	40	
≥60	132	58	74	
Sex				0.0027
Male	143	71	72	
Female	57	15	42	
Tumor size, cm				
≤5	98	41	57	0.7764
>5	102	45	57	
Tumor invasion (T)				0.0001
T1+T2	31	3	25	
T3+T4	169	83	89	
Lymph node metastasis (N)				0.0121
N0	48	13	35	
N1+N2+N3	152	73	79	
Distant metastasis (M)				0.0366
M0	184	75	109	
M1	16	11	5	
Differentiation				0.7751
Well to moderate	86	38	48	
Low	114	48	66	
TNM stage				0.0048
I+II	74	22	52	
III+IV	126	64	62	
Sample type				<0.0001
Tumor	200	86	115	
Normal	184	0	184	

TNM, tumor-node-metastasis; COL12A1, collagen type XII α 1 chain.

expression influenced GC prognosis was investigated. Of the 200 patients included in the study, 22 lacked survival information. Therefore, 178 patients were used to explore the correlations between COL12A1 expression level and survival status. The results revealed that patients with low expression levels of COL12A1 protein exhibited more extended periods of OS than those with high COL12A1 protein levels (Fig. 4A; $P<0.0001$). Similar results were acquired in patients at intermediate TNM stage II (Fig. 4B; $P=0.0004$), although not in those at stage III (Fig. 4C; $P>0.05$). When stratified by age, sex or lymph node metastasis, the results indicated that high COL12A1 expression was positively associated with worse OS in patients >60 years old ($P<0.0001$), in male patients ($P<0.0001$), and in patients with lymph node metastasis ($P=0.0013$) (Fig. 4D-F).

To further verify the prognostic role of COL12A1, the KM-plotter database, which contains a total of 875 GC

samples, was used. For COL12A1, 631 GC samples from one Affymetrix array (225664_at; Thermo Fisher Scientific, Inc.) were analyzed in the present study. It was predicted that in this cohort, COL12A1 overexpression was positively related to worse OS [Fig. 5A; hazard ratio (HR), 1.4; 95% CI, 1.13-1.72; log-rank $P=0.002$] and progression free survival (PFS) (Fig. 5D; HR, 1.43; 95% CI, 1.13-1.82; log-rank $P=0.0028$). Further analysis in male patients and patients with lymph node metastasis revealed similar results in OS (Fig. 5B and C) and PFS (Fig. 5E and F). Stratification analysis showed that COL12A1 was an indicator of inferior OS for intestinal-and mixed-type GC and an indicator of inferior PFS for intestinal-and diffuse-type GC (Fig. 6). Moreover, the univariate analysis suggested tumor invasiveness, lymph node invasion, distant metastasis, advanced clinical stage and COL12A1 high expression were significantly associated with OS, and further multivariate Cox analysis confirmed




Figure 4. Kaplan-Meier survival analysis. (A) COL12A1 overexpression was positively associated with worse OS. The association between COL12A1 expression and OS in patients of (B) stage II and (C) stage III is presented. Stratified analysis indicated that COL12A1 was associated with a worse OS in (D) patients >60 years old, (E) male patients and (F) patients with lymph node metastasis. OS, overall survival; COL12A1, collagen type XII α 1 chain.




Figure 5. KM plotter predicts the prognostic role of COL12A1 in patients with different Lauren classifications. OS in (A) intestinal-, (B) diffuse- and (C) mixed-type GC patients; PFS in (D) intestinal-, (E) diffuse- and (F) mixed-type GC patients. OS, overall survival; PFS, progression-free survival; COL12A1, collagen type XII α 1 chain; GC, gastric cancer.

that distant metastasis and COL12A1 overexpression could be regarded as powerful independent prognostic factors for OS

in GC (P<0.05; Table III); large-scale studies are warranted to verify these results further.




Figure 6. KM plotter predicts a poorer prognosis for gastric cancer patients with high expression of COL12A1. (A) OS in all patients (n=631); (B) OS in male patients; and (C) OS in patients with lymph node metastasis. (D) PFS (n=522) in all patients; (E) PFS in male patients; and (F) PFS in patients with lymph node metastasis. OS, overall survival; PFS, progression-free survival; COL12A1, collagen type XII α 1 chain.

Discussion

Early diagnosis of GC is a long-standing problem; however, early diagnosis can significantly benefit the clinical outcomes of GC patients by facilitating the implementation of prompt and more effective treatment. Therefore, extensive efforts have been expended to better understand the potential mechanisms associated with cancer development. Among these mechanisms, dysregulation of the extracellular matrix (ECM), a network of macromolecules between cells that is mainly composed of collagens, has attracted substantial attention (23). The steady state of ECM remodeling, in which damaged or old molecules are degraded and simultaneously replaced by new versions, is a key process involved in maintaining tissue homeostasis (24). Recently, dysregulation of ECM proteins was revealed in numerous cancer types (23,25). Alterations in ECM composition and quantity strongly influence tumor and stromal cell properties, which activate downstream signaling that modulates cell behavior and contributes to tumor progression (26). A review noted that the ECM is involved in neoplastic progression by modulating cancer hallmarks, and that its influence might vary in different cancer types (27). Together, these data indicate that ECM proteins are critical in the development of human malignancies and may represent promising candidates for early diagnosis and the personalization of clinical treatments.

COL12A1 is an ECM protein and, to date, its known disorders are mainly restricted to connective tissue pathology. Specifically, *COL12A1* mutations are involved in blended muscle and connective tissue phenotypes, often resulting in poorer prognoses compared with patients without *COL12A1* mutations (28). Additionally, elevated *COL12A1* expression is related to anterior cruciate ligament (ACL) tears, with higher expression levels in acute compared with chronic ACL damage (29).

The protein encoded by *COL12A1*, collagen type XII, interacts specifically with fibers, especially those containing collagen I (30). Recently, dysregulation of the ECM has been observed in a large spectrum of human cancer types, including CRC, and ovarian, breast and renal cell carcinomas. In CRC, *COL12A1* expression is increased and *COL12A1* levels are upregulated in tumors with microsatellite stability relative to tumors exhibiting high degrees of microsatellite instability (31). Notably, proteomics analyses have been conducted to search for markers of CRC-invasion fronts, with the results suggesting that type XII collagen is a potent marker of myofibroblast differentiation, implying that the COL12A1 protein is closely related to tumor metastasis in CRC (32). In ovarian cancer cell lines, the *COL12A1* expression level is significantly increased in cells resistant to cisplatin and/or doxorubicin (33), and high *COL12A1* expression levels are positively correlated with worse OS in patients with ovarian cancer or renal cell carcinoma (8,34).

Table III. Univariate and multivariate analyses of COL12A1 expression and patient survival.

Characteristics	HR	Overall survival		P-value
		(95% CI)		
Univariate analysis (n=178)				
Age (≥60 years vs. <60 years)	1.436	0.943	2.18	0.091
Sex (male vs. female)	0.835	0.56	1.24	0.377
Tumor size (>5 cm vs. ≤5 cm)	1.6	0.15	1.097	2.33
Tumor invasion (T3+T4 vs. T1+T2)	2.018	1.12	3.606	0.018
Lymph node metastasis (N1-3 vs. N0)	1.85	1.148	2.981	0.011
Distant metastasis (M1 vs. M0)	4.232	2.277	7.864	<0.001
Differentiation (poor vs. well to moderate)	1.352	0.922	1.983	0.123
TNM stage (III+IV vs. I+II)	2.348	1.546	3.566	<0.001
COL12A1 expression (high vs. low)	2.361	1.614	3.453	<0.001
Multivariate analysis				
Tumor invasion (T3+T4 vs. T1+T2)	1.516	0.833	2.76	0.174
Lymph node metastasis (N1-3 vs. N0)	1.474	0.903	2.406	0.12
Distant metastasis (M1 vs. M0)	2.919	1.547	5.51	0.001
COL12A1 expression (high vs. low)	1.896	1.267	2.837	0.002

Patients with $P < 0.05$ in univariate analysis were selected for multivariate analysis of survival. HR, hazard ratio; COL12A1, collagen type XII $\alpha 1$ chain; TNM, tumor-node-metastasis.

Although increased attention has been focused on elucidating the significance of COL12A1 in human cancer, the detailed mechanism of its involvement remains unknown. The present study, to the best of our knowledge, is the first to analyze COL12A1 expression in GC at both the mRNA and protein levels. Specifically, the present data revealed that COL12A1 expression was markedly increased in GC tissues compared with corresponding normal controls, both at the mRNA and protein levels. Additionally, COL12A1 mRNA was highly expressed in several GC cell lines except AGS, which may be attributed to the fact that different cell lines have different backgrounds, and thus it is reasonable for a certain gene to be expressed differentially in different cell lines. The IHC assay of COL12A1 showed that COL12A1 was mainly expressed in the stromal cells, especially in fibroblasts as showed in Fig. 3. Fibroblasts are essential components of the ECM and are responsible for the production of ECM. Cancer-associated fibroblasts (CAFs) are a cluster of cells characterized by their morphology and the expression of protein markers, including α -smooth muscle actin, fibroblast activated protein, neuron-glial antigen-2 and fibroblast specific protein (35-37). CAFs mainly originate from local fibroblasts or fibroblast precursors. Studies have linked CAFs to tumor progression, metastasis and poor clinical outcomes in various cancer types (36-38). The present results suggested that COL12A1 may be used as a potential marker of CAFs.

Additionally, Lauren classification is widely used for the histological classification of gastric adenocarcinoma (39). Based on these criteria, the majority of GC cases can be divided into either intestinal or diffuse types, with the remaining cases classified as mixed or unclassified. The two GC types differ from each other in a number of ways (40-43). For example, compared to

intestinal-type GC, diffuse-type GC is characterized by a lower response rate to chemotherapy, a worse prognosis, and higher peritoneal or distant recurrence rates (44,45). Therefore, in-depth elucidation of the underlying mechanisms of diffuse-type GC is required. In the present study, it was found that COL12A1 was highly expressed in intestinal-and diffuse-type GC, and was associated with worse clinical outcomes (OS and PFS).

Further analysis suggested that high COL12A1 protein expression was associated with tumor invasiveness, clinical metastasis and a worse TNM stage. Prognostic analysis demonstrated a positive association between high expression of COL12A1 and worse prognosis, and the results were in accordance with the prognosis prediction of KM-plotter. Therefore, the present findings revealed a tumor-promoting role of COL12A1 in GC. However, there are several limitations to the present study. For example, the study was retrospectively designed and the Lauren classification status of the samples was lacking. Large-scale prospective studies with more comprehensive information are required. *In vitro* and *in vivo* studies validating the tumor-promoting role of COL12A1 in the aggressiveness of GC and fibroblast cells by up- or downregulating its expression are required. Furthermore, explorations of the underlying molecular biological mechanisms of COL12A1 in GC are also warranted.

In conclusion, the present study revealed that COL12A1 expression was notably increased in GC tissues, and elevated COL12A1 protein levels were positively associated with aggressive clinical features. Importantly, COL12A1 protein expression levels represent a potential prognostic indicator for OS in patients with GC. These results indicated that COL12A1 plays a tumor-promoting role and might be a potential candidate for therapeutic targeting, based on its putative role in tumor progression and prognosis prediction in patients with GC.

Acknowledgements

Not applicable.

Funding

The present study was supported by grants from the National Natural Science Foundation of China (grant nos. 81201089 and 81272676) and the Natural Science Foundation of Zhejiang Province, China (grant nos. LY15H160026 and LY15H160012).

Availability of data and materials

The data used and/or analyzed in the present study are available from the corresponding author on reasonable request.

Authors' contributions

XJ, MW, XX and LT designed the study. XJ, MW, YH and LZ wrote the manuscript and conducted the immunohistochemical analyses. XJ, MW, XX, KH, YH and ZX collected the clinical samples and managed the clinical data. XJ, KH, HYW and HHW contributed to bioinformatics analysis. KH, HYW and ZX were involved in project management, and contributed to preparing and making figures and tables. HYW, HHW and LT supervised the study. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The study protocol was approved by the ethics committee of the First Affiliated Hospital of Zhejiang University and written informed consent was obtained from all participants in the study.

Patient consent for publication

Written informed consent was obtained from all participants.

Competing interests

The authors declare that they have no competing interests.

References

- Siegel RL, Miller KD and Jemal A: Cancer Statistics, 2017. *CA Cancer J Clin* 67: 7-30, 2017.
- Yang SY, Roh KH, Kim YN, Cho M, Lim SH, Son T, Hyung WJ and Kim HI: Surgical outcomes after open, laparoscopic and robotic gastrectomy for gastric cancer. *Ann Surg Oncol* 24: 1770-1777, 2017.
- Miller KD, Siegel RL, Lin CC, Mariotto AB, Kramer JL, Rowland JH, Stein KD, Alteri R and Jemal A: Cancer treatment and survivorship statistics, 2016. *CA Cancer J Clin* 66: 271-289, 2016.
- Pu X, Ye Y and Wu X: Development and validation of risk models and molecular diagnostics to permit personalized management of cancer. *Cancer* 120: 11-19, 2014.
- Gerecke DR, Olson PF, Koch M, Knoll JH, Taylor R, Hudson DL, Champlaud MF, Olsen BR and Burgeson RE: Complete primary structure of two splice variants of collagen XII and assignment of alpha 1 (XII) collagen (COL12A1), alpha 1 (IX) collagen (COL9A1) and alpha 1 (XIX) collagen (COL19A1) to human chromosome 6q12-q13. *Genomics* 41: 236-242, 1997.
- Hicks D, Farsani GT, Laval S, Collins J, Sarkozy A, Martoni E, Shah A, Zou Y, Koch M, Bönnemann CG, *et al*: Mutations in the collagen XII gene define a new form of extracellular matrix-related myopathy. *Hum Mol Genet* 23: 2353-2363, 2014.
- Sun F, Ding W, He JH, Wang XJ, Ma ZB and Li YF: Stomatin-like protein 2 is overexpressed in epithelial ovarian cancer and predicts poor patient survival. *BMC Cancer* 15: 746, 2015.
- Zhao H, Jungberg B, Grankvist K, Rasmuson T, Tibshirani R and Brooks JD: Gene expression profiling predicts survival in conventional renal cell carcinoma. *PLoS Med* 3: e13, 2006.
- Ramaswamy S, Ross KN, Lander ES and Golub TR: A molecular signature of metastasis in primary solid tumors. *Nat Genet* 33: 49-54, 2003.
- Torres S, Bartolomé RA, Mendes M, Barderas R, Fernandez-Aceñero MJ, Peláez-García A, Peña C, Lopez-Lucendo M, Villar-Vázquez R, de Herreros AG, *et al*: Proteome profiling of cancer-associated fibroblasts identifies novel proinflammatory signatures and prognostic markers for colorectal cancer. *Clin Cancer Res* 19: 6006-6019, 2013.
- Ji X, Bu ZD, Yan Y, Li ZY, Wu AW, Zhang LH, Zhang J, Wu XJ, Zong XL, Li SX, *et al*: The 8th edition of the american joint committee on cancer tumor-node-metastasis staging system for gastric cancer is superior to the 7th edition: Results from a Chinese mono-institutional study of 1663 patients. *Gastric Cancer* 21: 643-652, 2018.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Gao B, Lin J, Jiang Z, Yang Z, Yu H, Ding L, Yu M, Cui Q, Dunavin N, Zhang M and Li M: Upregulation of chemokine CXCL10 enhances chronic pulmonary inflammation in tree shrew collagen-induced arthritis. *Sci Rep* 8: 9993, 2018.
- Baptista MZ, Sarian LO, Vassallo J, Pinto GA, Soares FA and de Souza GA: Prognostic significance of GRP78 expression patterns in breast cancer patients receiving adjuvant chemotherapy. *Int J Biol Markers* 26: 188-196, 2011.
- Chen X, Leung SY, Yuen ST, Chu KM, Ji J, Li R, Chan AS, Law S, Troyanskaya OG, Wong J, *et al*: Variation in gene expression patterns in human gastric cancers. *Mol Biol Cell* 14: 3208-3215, 2003.
- Zhang X, Ni Z, Duan Z, Xin Z, Wang H, Tan J, Wang G and Li F: Overexpression of E2F mRNAs associated with gastric cancer progression identified by the transcription factor and miRNA co-regulatory network analysis. *PLoS One* 10: e0116979, 2015.
- He J, Jin Y, Chen Y, Yao HB, Xia YJ, Ma YY, Wang W and Shao QS: Downregulation of ALDOB is associated with poor prognosis of patients with gastric cancer. *Oncotargets Ther* 9: 6099-6109, 2016.
- Clough E and Barrett T: The gene expression omnibus database. *Methods Mol Biol* 1418: 93-110, 2016.
- Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Ponce-Rodriguez I, Chakravarthi BVSK and Varambally S: UALCAN: A portal for facilitating tumor subgroup gene expression and survival analyses. *Neoplasia* 19: 649-658, 2017.
- Cho JY, Lim JY, Cheong JH, Park YY, Yoon SL, Kim SM, Kim SB, Kim H, Hong SW, Park YN, *et al*: Gene expression signature-based prognostic risk score in gastric cancer. *Clin Cancer Res* 17: 1850-1857, 2011.
- D'Errico M, de Rinaldis E, Blasi MF, Viti V, Falchetti M, Calcagnile A, Sera F, Saieva C, Ottini L, Palli D, *et al*: Genome-wide expression profile of sporadic gastric cancers with microsatellite instability. *Eur J Cancer* 45: 461-469, 2009.
- Rhodes DR, Yu J, Shanker K, Deshpande N, Varambally R, Ghosh D, Barrette T, Pandey A and Chinnaiyan AM: ONCOMINE: A cancer microarray database and integrated data-mining platform. *Neoplasia* 6: 1-6, 2004.
- Karsdal MA, Nielsen SH, Leeming DJ, Langholm LL, Nielsen MJ, Manon-Jensen T, Siebuhr A, Gudmann NS, Rønnov S, Sand JM, *et al*: The good and the bad collagens of fibrosis: their role in signaling and organ function. *Adv Drug Deliv Rev* 121: 43-56, 2017.
- Hansen NU, Genovese F, Leeming DJ and Karsdal MA: The importance of extracellular matrix for cell function and in vivo likeness. *Exp Mol Pathol* 98: 286-294, 2015.
- Gilkes DM, Semenza GL and Wirtz D: Hypoxia and the extracellular matrix: Drivers of tumour metastasis. *Nat Rev Cancer* 14: 430-439, 2014.
- Millar MW, Corson N and Xu L: The adhesion g-protein-coupled receptor, GPR56/ADGRG1, inhibits cell-extracellular matrix signaling to prevent metastatic melanoma growth. *Front Oncol* 8: 8, 2018.

27. Pickup MW, Mouw JK and Weaver VM: The extracellular matrix modulates the hallmarks of cancer. *EMBO Rep* 15: 1243-1253, 2014.
28. Punetha J, Kesari A, Hoffman EP, Gos M, Kamińska A, Kostera-Pruszczyk A, Hausmanowa-Petrusewicz I, Hu Y, Zou Y, Bönnemann CG and Jędrzejowska M: Novel Col12A1 variant expands the clinical picture of congenital myopathies with extracellular matrix defects. *Muscle Nerve* 55: 277-281, 2017.
29. Brophy RH, Tycksen ED, Sandell LJ and Rai MF: Changes in transcriptome-wide gene expression of anterior cruciate ligament tears based on time from injury. *Am J Sports Med* 44: 2064-2075, 2016.
30. Bader HL, Keene DR, Charvet B, Veit G, Driever W, Koch M and Ruggiero F: Zebrafish collagen XII is present in embryonic connective tissue sheaths (fascia) and basement membranes. *Matrix Biol* 28: 32-43, 2009.
31. Ortega P, Moran A, Fernandez-Marcelo T, De Juan C, Frias C, Lopez-Asenjo JA, Sanchez-Pernaute A, Torres A, Diaz-Rubio E, Iniesta P and Benito M: MMP-7 and SGCE as distinctive molecular factors in sporadic colorectal cancers from the mutator phenotype pathway. *Int J Oncol* 36: 1209-1215, 2010.
32. Karagiannis GS, Petraki C, Prassas I, Saraon P, Musrap N, Dimitromanolakis A and Diamandis EP: Proteomic signatures of the desmoplastic invasion front reveal collagen type XII as a marker of myofibroblastic differentiation during colorectal cancer metastasis. *Oncotarget* 3: 267-285, 2012.
33. Januchowski R, Świerczewska M, Sterzynska K, Wojtowicz K, Nowicki M and Zabel M: Increased expression of several collagen genes is associated with drug resistance in ovarian cancer cell lines. *J Cancer* 7: 1295-1310, 2016.
34. Chudasama D, Bo V, Hall M, Anikin V, Jeyaneethi J, Gregory J, Pados G, Tucker A, Harvey A, Pink R and Karteris E: Identification of novel cancer biomarkers of prognostic value using specific gene regulatory networks (GRN): A novel role of RAD51AP1 for ovarian and lung cancers. *Carcinogenesis* 39: 407-417, 2018.
35. Huang L, Xu AM, Liu S, Liu W and Li TJ: Cancer-associated fibroblasts in digestive tumors. *World J Gastroenterol* 20: 17804-17818, 2014.
36. Ostman A and Augsten M: Cancer-associated fibroblasts and tumor growth-bystanders turning into key players. *Curr Opin Genet Dev* 19: 67-73, 2009.
37. Kalluri R and Zeisberg M: Fibroblasts in cancer. *Nat Rev Cancer* 6: 392-401, 2006.
38. Luo H, Tu G, Liu Z and Liu M: Cancer-associated fibroblasts: A multifaceted driver of breast cancer progression. *Cancer Lett* 361: 155-163, 2015.
39. Lauren P: The two histological main types of gastric carcinoma: Diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Pathol Microbiol Scand* 64: 31-49, 1965.
40. Sunakawa Y and Lenz HJ: Molecular classification of gastric adenocarcinoma: Translating new insights from the cancer genome atlas research network. *Curr Treat Options Oncol* 16: 17, 2015.
41. Riquelme I, Saavedra K, Espinoza JA, Weber H, García P, Nervi B, Garrido M, Corvalán AH, Roa JC and Bizama C: Molecular classification of gastric cancer: Towards a pathway-driven targeted therapy. *Oncotarget* 6: 24750-24779, 2015.
42. Vauhkonen M, Vauhkonen H, Sajantila A and Sipponen P: Differences in genomic instability between intestinal-and diffuse-type gastric cancer. *Gastric Cancer* 8: 238-244, 2005.
43. Ma J, Shen H, Kapesa L and Zeng S: Lauren classification and individualized chemotherapy in gastric cancer. *Oncol Lett* 11: 2959-2964, 2016.
44. Marrelli D, Polom K, Neri A and Roviello F: Clinical impact of molecular classifications in gastric cancer. *Updates Surg* 70: 225-232, 2018.
45. Lee JH, Chang KK, Yoon C, Tang LH, Strong VE and Yoon SS: Lauren histologic type is the most important factor associated with pattern of recurrence following resection of gastric adenocarcinoma. *Ann Surg* 267: 105-113, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.