

Calycosin induces apoptosis in osteosarcoma cell line via ER β -mediated PI3K/Akt signaling pathways

WEI TIAN, ZHI-WEI WANG, BAO-MING YUAN and YONG-GE BAO

Department of Orthopedics, Affiliated Hospital of Inner Mongolia University for Nationalities,
Tongliao, Inner Mongolia Autonomous Region 028007, P.R. China

Received September 12, 2019; Accepted February 24, 2020

DOI: 10.3892/mmr.2020.11039

Abstract. Previous studies have shown that calycosin, a natural phytoestrogen which is structurally similar to estrogen, inhibits proliferation and induces apoptosis in estrogen-dependent cancer types via the estrogen receptor (ER) β -induced inhibition of PI3K/Akt. Therefore, the aims of the present study were to investigate the effects of calycosin on human osteosarcoma (OS), and to examine the molecular mechanisms associated with ER β . Human OS MG-63 cells were treated with various concentrations of calycosin, and MTT and flow cytometry assays were used to assess the effects of calycosin on cellular proliferation and apoptosis. In addition, protein expression levels of ER β , phosphorylated (p)-PI3K, p-Akt, cleaved poly (ADP-ribose) polymerase 1 (PARP) and cleaved caspase-3 were evaluated by western blot analysis. The present results suggested that calycosin inhibited proliferation and induced apoptosis in MG-63 cells. Furthermore, increased ER β expression was detected in OS MG-63 cells treated with calycosin, and an ER β inhibitor (PHTPP) reversed calycosin-induced cytotoxicity and apoptosis. Moreover, phosphorylation levels of PI3K and Akt were significantly downregulated after calycosin treatment, whereas PHTPP reversed their phosphorylation. ER β -mediated PI3K/Akt downstream signaling pathways were found to influence the activity of poly (ADP-ribose) polymerase 1 and caspase-3. Thus, the present results indicated that calycosin inhibited proliferation and induced apoptosis in OS MG-63 cells, and that these effects were mediated by ER β -dependent inhibition of the PI3K/Akt pathways.

Introduction

Osteosarcoma (OS) is the most common primary malignant bone tumor in pediatrics and young adolescents (1).

Furthermore, the standard care for patients with OS, which includes surgical resection in combination with systemic chemotherapy, has a 5-year overall survival rate of 70% (2). Treatment of OS often fails due to the development of chemotherapy resistance and metastasis (3). It has been previously reported that sex hormones are involved in the occurrence and development of human OS, thus suggesting that novel endocrine therapy may be development for OS using clinically available estrogen inhibitors (4).

Plant-derived phytoestrogens and mammalian estrogens have similar structures and functions, and can cause anti-estrogen or estrogen-like effects (5). For this reason, plant-derived phytoestrogens are a current topic in research (6). Phytoestrogenic compounds are widely found in nature and can be divided into four categories: i) Isoflavones; ii), stilbene; iii), coumarins; and iv) lignans (7). Calycosin, a bioactive phytoestrogen isoflavone that is extracted from *Trifolium pratense* (red clover), has been shown to inhibit proliferation and induce apoptosis in cancer types (8,9). In addition, calycosin has been shown to induce apoptosis in human estrogen receptor (ER)-positive OS cells, but has no effect on ER-negative OS cells, suggesting that the inhibition of calycosin on ER-positive OS cells may be achieved by increasing ER expression (10).

ER belongs to the steroid hormone receptor family and consists of two subtypes, ER α and ER β (11). A high ER α :ER β ratio leads to increased cell proliferation, whereas a higher level of ER β than ER α leads to decreased proliferation (12,13). Since the expression of ER β has been shown to decrease during tumor progression, ER β has been considered a potential tumor suppressor and therapeutic target in various types of cancer, including breast cancer and renal cell carcinoma (14,15). Moreover, ER β agonists may be novel potential therapeutic candidates for OS endocrine therapy (16). Therefore, it was hypothesized that upregulation of ER β may inhibit tumor development and progression.

The PI3K/Akt signaling pathway, an important regulator of cellular functions, has been found to be frequently hyperactivated in OS and contributes to tumorigenesis, proliferation, invasion, cell cycle progression and inhibition of apoptosis (17). Thus, suppressing this signaling pathway could inhibit disease initiation and development (18). Moreover, ER β has an anti-tumor effect on OS cells by regulating the PI3K/Akt signaling pathway (16). In addition, ER β can

Correspondence to: Professor Yong-Ge Bao, Department of Orthopedics, Affiliated Hospital of Inner Mongolia University for Nationalities, 1742 Huolin He Street, Tongliao, Inner Mongolia Autonomous Region 028007, P.R. China
E-mail: baoyongge001@163.com

Key words: calycosin, osteosarcoma, estrogen receptor β , apoptosis

mediate inhibition of proliferation and activation of apoptosis in various types of cancer, including breast cancer and colorectal cancer following treatment with calycosin, which is shown to regulate the PI3K/Akt pathway (8,9). Therefore, the present study investigated whether calycosin had anti-tumor effects on OS MG-63 cells by mediating the ER β -dependent PI3K/Akt signaling pathway.

Collectively, along with the anti-proliferative effect of calycosin on OS MG-63 cells, the present study examined the role of the ER β -mediated PI3K/Akt pathway in OS MG-63 cells to help facilitate the current understanding of the molecular mechanism underlying calycosin functions.

Materials and methods

Calycosin. Calycosin (purity 98%; Tianjin JAHE Science and Technology Co. Ltd.) solution was diluted into a 250 μ g/ml stock solution with DMSO (Sigma-Aldrich; Merck KGaA).

Cell culture. Human OS cells (MG-63) and human fetal osteoblast cells (hFOB1.19; Shanghai Institute of Biochemistry and Cell Biology) were incubated in DMEM (Gibco; Thermo Fisher Scientific, Inc.) supplemented with 10% FBS (Gibco; Thermo Fisher Scientific, Inc.) and antibiotics (100 U/ml penicillin and 100 μ g/ml streptomycin) in a humidified incubator containing 5% CO₂ at 37°C. The medium was changed every 48 h.

MTT assay. Cell viability was determined by MTT (Sigma-Aldrich; Merck KGaA) assay. Cells were harvested using 0.25% trypsin and seeded into 96-well plates at a density of 3x10⁴ cells/well at 37°C for 24 h. Then, cells were treated with calycosin (at concentrations of 0, 25, 50 or 100 μ M) at 37°C for 24, 48 and 72 h, or with 100 μ M calycosin in the presence or absence of the ER β inhibitor PHTPP (50 μ M; MedChemExpress) at 37°C for 48 h. In total, 20 μ l MTT (5 mg/ml) was added to cells for 4 h at 37°C. Following incubation, DMSO (100 μ l) was added to dissolve the formazan crystals and shaken at room temperature for 10 min. Subsequently, cell viability was assessed by measuring the absorbance at 570 nm using a microplate reader (Thermo Fisher Scientific, Inc.). Proliferation rate (%) was calculated as follows: Optical density (OD) treatment group/OD control x100%.

Flow cytometry assay. Flow cytometry was used to study the effects of calycosin treatment on apoptosis of the OS cells using the FITC Annexin V Apoptosis Detection Kit I (BD Biosciences), according to the manufacturer's protocol. MG-63 cells were treated with calycosin (0, 25, 50 or 100 μ M), or calycosin (100 μ M) in the presence or absence of PHTPP at 37°C for 48 h. Cells were harvested and washed with PBS. Apoptotic cells were identified by double staining with 5 μ l FITC-conjugated Annexin V and 5 μ l PI. Data were obtained and analyzed using a FACS-Canto flow cytometer (Beckman Coulter, Inc.) with Cell Quest software (version 5.1; BD Biosciences). Cells stained positive for Annexin V-FITC and negative for PI were considered early apoptotic, and cells stained positive for Annexin V-FITC and positive for PI were considered in late apoptosis.

Western blot analysis. After being treated with calycosin (0, 25, 50 and 100 μ M), or calycosin (100 μ M) in the presence or absence of PHTPP for 48 h, MG-63 cells were harvested with ice-cold PBS and lysed on ice in lysis buffer (Beyotime Institute of Biotechnology) for 30 min. The lysates were centrifuged at 4°C at 700 x g for 10 min and collected. Then, the total protein was measured with a bicinchoninic protein assay kit (Tiangen Biotech Co., Ltd.). The samples (20 μ g) were separated via 12% SDS-PAGE and then transferred to PVDF membranes (EMD Millipore). The membranes were blocked with 5% non-fat dried milk in TBST (0.2% Tween-20) buffer for 1 h at room temperature, and then incubated with the following primary antibodies overnight at 4°C: ER β (cat. no. sc8974; 1:500), PI3K (cat. no. 4255; 1:1,000), phosphorylated (p)-PI3K (cat. no. 17366; 1:1,000), Akt (cat. no. 9271; 1:1,000), p-Akt (cat. no. 9611; 1:2,000), cleaved caspase-3 (cat. no. 9661; 1:1,000), cleaved poly (ADP-ribose) polymerase 1 (cleaved PARP-1; cat. no. 9185; 1:1,000) and β -actin (cat. no. 7077; 1:1,000). After three washes with TBST, the membranes were subsequently incubated with a horseradish peroxidase-conjugated secondary antibody (cat. no. 7074; 1:5,000) for 1 h at room temperature. The protein signal was detected via electrochemiluminescence with an ECL-Plus kit (Beyotime Institute of Biotechnology) and analyzed using ImageJ software (National Institutes of Health). Anti-ER β was purchased from Santa Cruz Biotechnology, Inc. The other antibodies were purchased from Cell Signaling Technology, Inc.

Statistical analysis. Data were obtained from ≥ 3 independent experiments and are presented as the mean \pm SD relative to the control value. Statistical analysis for multiple comparisons was performed using one-way ANOVA followed by Tukey's post hoc test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Inhibition of proliferation and induction of apoptosis in OS MG-63 cells by calycosin. To evaluate the anti-proliferative effect of calycosin, MG-63 and hFOB1.19 cells were incubated with different concentrations of calycosin for 24, 48 and 72 h. It was found that calycosin caused a time- and concentration-dependent inhibition on the proliferation of MG-63 cells (Fig. 1A). However, the inhibitory effect of calycosin on the proliferation of hFOB1.19 cells was not significant (Fig. 1B), thus suggesting that the effect of calycosin was negligible on healthy osteoblasts.

Consistent with the aforementioned results, flow cytometry assay results demonstrated that calycosin induced MG-63 cell apoptosis in a concentration-dependent manner, and induced a significant increase in the percentage of early and late apoptotic cells (Fig. 1C and D).

Upregulation of ER β in OS MG-63 cells by calycosin. ER β is a traditional estrogen receptor, whose activity is inversely related to the occurrence and development of tumors (19). Therefore, the expression of ER β was examined in MG-63 cells following treatment with calycosin. It was demonstrated that ER β protein expression was increased significantly in a dose-dependent manner after calycosin treatment (Fig. 2A and B). Moreover,




Figure 1. Inhibition of proliferation and apoptosis of osteosarcoma MG-63 cells by calycosin. Cells were incubated with varying concentrations of calycosin for 24, 48 and 72 h. Cell viability of (A) MG-63 cells and (B) hFOB1.19 cells were detected by MTT. Apoptosis of MG-63 cells was determined by (C) flow cytometry and (D) the results were quantified. These independent experiments were performed three times. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. 0 μM calycosin. PI, propidium iodide.

MG-63 cells were treated with 100 μM calycosin in the presence or absence of PHTPP, and then the expression of ER β was assessed. It was found that calycosin in combination with PHTPP significantly decreased ER β expression ($P < 0.01$; Fig. 2C and D). Therefore, the present results suggested that the inhibition of calycosin on MG-63 cells occurred via the regulation of ER β .

ER β -mediated inhibition of proliferation and activity of apoptosis in MG-63 cells by calycosin. Previous studies have shown that ER β is a key negative mediator of cell proliferation, and positive regulator of apoptosis in cancer (8,9,16). Therefore, to study the effect of ER β on proliferation and apoptosis, MG-63 cells were pretreated with 100 μM calycosin in the presence or absence of PHTPP for 48 h. It was identified that calycosin + PHTPP reversed the calycosin-mediated inhibition of cell proliferation ($P < 0.001$; Fig. 3A). In addition,

calycosin + PHTPP abolished calycosin-induced apoptosis ($P < 0.001$, Fig. 3B and C). Consistent with the findings of previous studies, calycosin-induced cytotoxicity and apoptosis were found to be mediated by the ER β -signaling pathway.

Regulation of ER β -mediated PI3K/Akt signaling in OS MG-63 cells by calycosin. It has been previously shown that inhibiting the PI3K/Akt signaling pathway may be a novel treatment strategy for OS (17). Therefore, the PI3K/Akt signaling pathway was examined in the present study. After a 48 h exposure to calycosin (0, 25, 50 or 100 μM) in MG-63 cells, PI3K phosphorylation and Akt expression were found to be downregulated in a concentration-dependent manner (Fig. 4A and B). Thus, the present results indicated that calycosin inactivated the PI3K/Akt signaling pathway in MG-63 cells. In addition, to assess the relationship between




Figure 2. Calycosin upregulates ER β expression in osteosarcoma MG-63 cells. (A) Western blotting results of (B) protein expression of ER β after treatment with different concentrations of calycosin. (C) Western blotting results of (D) protein expression of ER β after 100 μ M calycosin treatment in the presence or absence of PHTPP for 48 h. These independent experiments were performed three times. * $P < 0.05$, ** $P < 0.01$ vs. 0 μ M calycosin. ## $P < 0.01$ vs. calycosin alone. & $P < 0.05$ vs. PHTPP alone. ER β , estrogen receptor β .




Figure 3. Estrogen receptor β -mediated inhibition of proliferation and apoptosis osteosarcoma MG-63 cells by calycosin. Cells were incubated with 100 μ M calycosin in the presence or absence of PHTPP for 48 h. (A) Cell viability was detected by MTT. (B) Apoptosis was determined by flow cytometry and (C) the results were quantified. These independent experiments were performed three times. * $P < 0.05$, *** $P < 0.001$ vs. 0 μ M calycosin. ### $P < 0.001$ vs. calycosin. & $P < 0.05$, && $P < 0.01$, &&& $P < 0.001$ vs. PHTPP. PI, propidium iodide.

ER β and the PI3K/Akt signaling pathway, MG-63 cells were treated with calycosin in the presence or absence of PHTPP. It was demonstrated that the combination of calycosin + PHTPP

reversed the decrease in the phosphorylation levels of PI3K and Akt ($P < 0.01$; Fig. 4C and D), while no changes were observed in the expression levels of total Akt and total PI3K. Therefore,




Figure 4. Regulation of estrogen receptor β -mediated PI3K/Akt signaling in osteosarcoma MG-63 cells by calycosin. (A) Western blotting results of (B) protein expression levels of p-PI3K/PIK3 and p-Akt/Akt after treatment with different concentrations of calycosin. (C) Western blotting results of (D) protein expression levels of p-PI3K/PIK3 and p-Akt/Akt after treatment with 100 μ M calycosin, in the presence or absence of PHTPP for 48 h. These independent experiments were performed three times. * $P < 0.05$, ** $P < 0.01$ vs. 0 μ M calycosin. ## $P < 0.01$ vs. calycosin. & $P < 0.05$ vs. PHTPP. p, phosphorylated.

the present results suggested that the ER β -mediated decrease in PI3K/Akt activity was involved in calycosin-induced cell death.

ER β -mediated increase in the expression levels of apoptotic-associated protein in OS MG-63 cells treated with calycosin. Apoptosis is often associated with the cleavage of specific substrates, such as cleaved PARP-1 and caspase-3, whose excessive activation can promote apoptosis (20). When MG-63 cells were treated with 0, 25, 50 and 100 μ M calycosin for 48 h, it was found that the protein expression levels of PARP-1 and cleaved caspase-3 increased in a time- and dose-dependent manner (Fig. 5A and B). This suggested that activation of PARP-1 and cleavage of caspase-3 may be involved in calycosin-mediated cell apoptosis. In addition, calycosin + PHTPP treatment reduced PARP-1 and caspase-3 cleavage protein expression levels ($P < 0.05$; Fig. 5C and D). Collectively, the present results indicated that PARP-1 and caspase-3 may be downstream targets of the ER β -mediated PI3K/Akt pathway.

Discussion

OS usually occurs in adolescence when the synthesis of sex hormones, such as estrogen or androgen, peaks, thus indicating that sex steroids and their receptors may be involved in the development of OS (21). Phytoestrogens have recently received increased attention due to their ability to bind to ERs. Previous studies have suggested that certain phytoestrogens

exhibit antiestrogenic activity via ER-mediated signaling pathways (22,23). Moreover, calycosin is a phytoestrogen isoflavone that exhibits estrogenic activity and anti-tumor effects on several cancer types, by inducing apoptosis of tumor cells *in vitro* and *in vivo* (8,9). Furthermore, *in vitro* and *in vivo* studies have also shown that calycosin has antiapoptotic and antimetastatic activities against OS (10,24). Consistent with these previous studies, the present results suggested that calycosin effectively inhibited cell proliferation and induced apoptosis in a time- and dose-dependent manner in OS MG-63 cells. In addition, calycosin had a low cytotoxicity in osteoblast hFOB1.19 cells.

Several previous studies have shown that calycosin inhibits tumorigenesis and tumor progression by regulating ER β expression (8,9,25). Moreover, the downregulation of ER β has been observed in various types of cancer (8,9). In the present study, ER β protein expression in OS cells was found to be decreased, while its expression increased significantly in a dose-dependent manner after calycosin treatment. A previous study showed that estrogen extracted from *Astragalus membranaceus* could inhibit etoposide-induced apoptosis of human OS cells by activating ER β (26). Furthermore, despite the reduction in ER β expression, which may be related to different subtypes of OS cells, the estrogen inhibitor fulvestrant exhibits significant anti-OS activity in OS 143B cells (27). The present results suggested that the ER β inhibitor PHTPP reversed the increase in the protein expression of ER β , and significantly reversed the cytotoxicity and apoptosis detected following calycosin treatment in MG-63 cells. Therefore, the present




Figure 5. Induction of ER β -mediated expression of apoptotic-associated proteins in ER-positive cells by calycosin. (A) Western blotting results of (B) protein expression levels of cleaved PARP and cleaved caspase-3 after treatment with different concentrations of calycosin. (C) Western blotting results of (D) protein expression levels cleaved PARP and cleaved caspase-3 after treatment with 100 μ M calycosin, in the presence or absence of PHTPP for 48 h. These independent experiments were performed three times. * $P < 0.05$, ** $P < 0.01$ vs. calycosin (0 μ M or 0 h). # $P < 0.05$ vs. calycosin. * $P < 0.05$ vs. PHTPP. ER, estrogen receptor; PARP, poly (ADP-ribose) polymerase 1.

results indicated that the anti-tumor effects of calycosin were mediated by the ER β signaling pathway.

In addition, the mechanism underlying the anti-tumor effect of ER β was examined in the present study. The association between calycosin and ER β was investigated, and calycosin was found to reduce OS cell proliferation by inhibiting the ER β signaling pathway, in particular a potential downstream effector. The PI3K/Akt signaling pathway is frequently hyperactivated in OS and has been shown to be involved in tumorigenesis, proliferation, invasion, cell cycle progression and inhibition of apoptosis (17). Furthermore, ER β can independently predict the prognosis of triple-negative breast cancer by interacting with the PI3K/Akt pathway (19). Estrogen can activate the PI3K/Akt pathway via ER β in breast cancer (28). Moreover, ER β has a significant anti-tumor effect on OS U2-OS cells by regulating the PI3K/Akt signal pathway (16). It has been shown that the inhibitory effects of calycosin on OS MG-63 cells are mediated by the PI3K/Akt pathway (10). In the present study, it was demonstrated that calycosin inactivated the PI3K/Akt signaling pathway in OS MG-63 cells, whereas the ER β inhibitor PHTPP enhanced the phosphorylation of PI3K and Akt. Thus, the present results suggested that the anti-tumor effects of ER β were associated with the PI3K/Akt signaling pathway.

Defects in apoptosis play an important role in tumor pathogenesis, and the cytotoxic effects of many antineoplastic drugs are usually accompanied by an increase in apoptosis (29).

Experimental studies of OS cells and 143B-harbored nude mice have shown that calycosin possesses an anti-osteosarcoma effect, and the underlying mechanism is associated with the activation of apoptosis (30). In the present study, calycosin was found to induce apoptosis in MG-63 cells, as indicated by morphological changes, and the activation of caspase-3 and PARP-1. Caspases are known for their role as initiators and executors of apoptosis (31). Activation of different caspase cascades plays an important role in apoptosis by cleaving key factors involved in cellular function and viability (32). Moreover, the apoptotic executor factor caspase-3 can cleave the caspase substrate PARP-1 into two specific fragments, thus contributing to cell death (33). Therefore, cleaved PARP-1 and caspase-3 are considered as apoptotic markers. The present results indicated that there were increased expression levels of cleaved PARP-1 and caspase-3 following treatment with calycosin in a concentration-dependent manner, thus indicating the involvement of PARP-1 and caspase-3 in the effects of calycosin. Moreover, the PHTPP inactivated PARP-1 and caspase-3 cleavage, indicating that ER β mediated PARP-1 and caspase-3 activity in calycosin-induced apoptosis.

Collectively, the present study provided further evidence for the interaction between calycosin and ER β . In addition, the antiproliferative effects of calycosin were found to be mediated by the ER β -dependent regulation of the PI3K/Akt pathways. Therefore, the present study provides a theoretical basis for the potential use of calycosin as a therapeutic to treat OS.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

WT performed the majority of the experiments and drafted the manuscript. ZWW helped perform the experiments. BMV analyzed the data and drafted the manuscript. YGB conceived the study, supervised the experiments and edited the manuscript. designed the study. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Pingping B, Yuhong Z, Weiqi L, Chunxiao W, Chunfang W, Yuanjue S, Chenping Z, Jianru X, Jiade L, Lin K, *et al*: Incidence and mortality of sarcomas in Shanghai, China, during 2002-2014. *Front Oncol* 9: 662, 2019.
- Harrison DJ, Geller DS, Gill JD, Lewis VO and Gorlick R: Current and future therapeutic approaches for osteosarcoma. *Expert Rev Anticancer Ther* 18: 39-50, 2018.
- Lin YH, Jewell BE, Gingold J, Lu L, Zhao R, Wang LL and Lee DF: Osteosarcoma: Molecular pathogenesis and iPSC modeling. *Trends Mol Med* 23: 737-755, 2017.
- Hu Q, Li S, Chen C, Zhu M, Chen Y and Zhao Z: 17 β -estradiol treatment drives Sp1 to upregulate MALAT-1 expression and epigenetically affects physiological processes in U2OS cells. *Mol Med Rep* 15: 1335-1342, 2017.
- Li M, Zeng M, Zhang Z, Zhang J, Zhang B, Zhao X, Zheng X and Feng W: Uridine derivatives from the seeds of *Lepidium apetalum* Willd. And their estrogenic effects. *Phytochemistry* 155: 45-52, 2018.
- van Duursen MB, Smeets EE, Rijk JC, Nijmeijer SM and van den Berg M: Phytoestrogens in menopausal supplements induce ER-dependent cell proliferation and overcome breast cancer treatment in an in vitro breast cancer model. *Toxicol Appl Pharmacol* 269: 132-140, 2013.
- Lin AH, Li RW, Ho EY, Leung GP, Leung SW, Vanhoutte PM and Man RY: Differential ligand binding affinities of human estrogen receptor- α isoforms. *PLoS One* 8: e63199, 2013.
- Zhao X, Li X, Ren Q, Tian J and Chen J: Calycosin induces apoptosis in colorectal cancer cells, through modulating the ER β /MiR-95 and IGF-1R, PI3K/Akt signaling pathways. *Gene* 591: 123-128, 2016.
- Chen J, Hou R, Zhang X, Ye Y, Wang Y and Tian J: Calycosin suppresses breast cancer cell growth via ER β -dependent regulation of IGF-1R, p38 MAPK and PI3K/Akt pathways. *PLoS One* 9: e91245, 2014.
- Sun H, Yin M, Qian W and Yin H: Calycosin, a Phytoestrogen isoflavone, induces apoptosis of estrogen receptor-positive MG-63 osteosarcoma cells via the phosphatidylinositol 3-kinase (PI3K)/AKT/mammalian target of rapamycin (mTOR) pathway. *Med Sci Monit* 24: 6178-6186, 2018.
- Renoir JM: Estradiol receptors in breast cancer cells: Associated co-factors as targets for new therapeutic approaches. *Steroids* 77: 1249-1261, 2012.
- Zhang Y, Wei F, Zhang J, Hao L, Jiang J, Dang L, Mei D, Fan S, Yu Y and Jiang L: Bisphenol A and estrogen induce proliferation of human thyroid tumor cells via an estrogen-receptor-dependent pathway. *Arch Biochem Biophys* 633: 29-39, 2017.
- Treeck O, Diepolder E, Skrzypczak M, Schüler-Toprak S and Ortmann O: Knockdown of estrogen receptor β increases proliferation and affects the transcriptome of endometrial adenocarcinoma cells. *BMC Cancer* 19: 745, 2019.
- Li H, Tu Z, An L, Qian Z, Achilefu S and Gu Y: Inhibitory effects of ER β on proliferation, invasion, and tumor formation of MCF-7 breast cancer cells-prognostication for the use of ER β -selective therapy. *Pharm Biol* 50: 839-849, 2012.
- Yu CP, Ho JY, Huang YT, Cha TL, Sun GH, Yu DS, Chang FW, Chen SP and Hsu RJ: Estrogen inhibits renal cell carcinoma cell progression through estrogen receptor- β activation. *PLoS One* 8: e56667, 2013.
- Yang M, Liu B, Jin L, Tao H and Yang Z: Estrogen receptor β exhibited anti-tumor effects on osteosarcoma cells by regulating integrin, IAP, NF-kB/BCL-2 and PI3K/Akt signal pathway. *J Bone Oncol* 9: 15-20, 2017.
- Zhang J, Yu XH, Yan YG, Wang C and Wang WJ: PI3K/Akt signaling in osteosarcoma. *Clin Chim Acta* 444: 182-192, 2015.
- Xu Y, Li N, Xiang R and Sun P: Emerging roles of the p38 MAPK and PI3K/AKT/mTOR pathways in oncogene-induced senescence. *Trends Biochem Sci* 39: 268-76, 2014.
- Wang J, Zhang C, Chen K, Tang H, Tang J, Song C and Xie X: ER β 1 inversely correlates with PTEN/PI3K/AKT pathway and predicts a favorable prognosis in triple-negative breast cancer. *Breast Cancer Res Treat* 152: 255-269, 2015.
- Gupta S, Kass GE, Szegezdi E and Joseph B: The mitochondrial death pathway: A promising therapeutic target in diseases. *J Cell Mol Med* 13: 1004-1033, 2009.
- Botter SM, Neri D and Fuchs B: Recent advances in osteosarcoma. *Curr Opin Pharmacol* 16: 15-23, 2014.
- Hwang KA, Park MA, Kang NH, Yi BR, Hyun SH, Jeung EB and Choi KC: Anticancer effect of genistein on BG-1 ovarian cancer growth induced by 17 β -estradiol or bisphenol A via the suppression of the crosstalk between estrogen receptor α and insulin-like growth factor-1 receptor signaling pathways. *Toxicol Appl Pharmacol* 272: 637-646, 2013.
- Thelen P, Wuttke W and Seidlová-Wuttke D: Phytoestrogens selective for the estrogen receptor beta exert anti-androgenic effects in castration resistant prostate cancer. *J Steroid Biochem Mol Biol* 139: 290-293, 2014.
- Qiu R, Li X, Qin K, Chen X, Wang R, Dai Y, Deng L and Ye Y: Antimetastatic effects of calycosin on osteosarcoma and the underlying mechanism. *Biofactors* 45: 975-982, 2019.
- Chen J, Zhao X, Ye Y, Wang Y and Tian J: Estrogen receptor beta-mediated proliferative inhibition and apoptosis in human breast cancer by calycosin and formononetin. *Cell Physiol Bioch* 32: 1790-1797, 2013.
- Zhou R, Chen H, Chen J, Chen X, Wen Y and Xu L: Extract from *Astragalus membranaceus* inhibit breast cancer cells proliferation via PI3K/AKT/mTOR signaling pathway. *BMC Complement Altern Med* 18: 83, 2018.
- Gorska M, Wyszowska RM, Kuban-Jankowska A and Wozniak M: Impact of apparent antagonism of estrogen receptor β by fulvestrant on anticancer activity of 2-methoxyestradiol. *Anticancer Res* 36: 2217-2226, 2016.
- Kanaizumi H, Higashi C, Tanaka Y, Hamada M, Shinzaki W, Azumi T, Hashimoto Y, Inui H, Houjou T and Komoike Y: PI3K/Akt/mTOR signalling pathway activation in patients with ER-positive, metachronous, contralateral breast cancer treated with hormone therapy. *Oncol Lett* 17: 1962-1968, 2019.

29. Hassan M, Watari H, AbuAlmaaty A, Ohba Y and Sakuragi N: Apoptosis and molecular targeting therapy in cancer. *Biomed Res Int* 2014: 150845, 2014.
30. Qiu R, Ma G, Zheng C, Qiu X, Li X, Li X, Mo J, Li Z, Liu Y, Mo L, *et al*: Antineoplastic effect of calycosin on osteosarcoma through inducing apoptosis showing in vitro and in vivo investigations. *Exp Mol Pathol* 97: 17-22, 2014.
31. Kopeina GS, Prokhorova EA, Lavrik IN and Zhivotovsky B: Alterations in the nucleocytoplasmic transport in apoptosis: Caspases lead the way. *Cell Prolif* 51: e12467, 2018.
32. Akanda MR, Kim MJ, Kim IS, Ahn D, Tae HJ, Rahman MM, Park YG, Seol JW, Nam HH, Choo BK and Park BY: Neuroprotective effects of sigesbeckia pubescens extract on glutamate-induced oxidative stress in HT22 cells via downregulation of MAPK/caspase-3 pathways. *Cell Mol Neurobiol* 38: 497-505, 2018.
33. Rogalska A, Bukowska B and Marczak A: Metformin and epothilone A treatment up regulate pro-apoptotic PARP-1, Casp-3 and H2AX genes and decrease of AKT kinase level to control cell death of human hepatocellular carcinoma and ovary adenocarcinoma cells. *Toxicol In Vitro* 47: 48-62, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.