

Hypoxia-inducible factor-1 α regulates Lipin1 differently in pre-adipocytes and mature adipocytes

YOSHITAKA KIHIRA^{1,2}, YOSHINO FUJIMURA¹, SHUHEI TOMITA³,
TOSHIAKI TAMAKI^{2,4} and EIJI SATO¹

¹Department of Clinical Pharmacy, Faculty of Pharmacy and Pharmaceutical Sciences, Fukuyama University, Fukuyama, Hiroshima 729-0292; ²Department of Pharmacology, Institute of Biomedical Sciences, Tokushima University Graduate School, Kuramoto-cho, Tokushima 770-8503; ³Department of Pharmacology, Osaka City University Graduate School of Medicine, Osaka-shi 558-8585; ⁴Anan Medical Center, Anan, Tokushima 774-0045, Japan

Received November 26, 2019; Accepted January 29, 2020

DOI: 10.3892/mmr.2020.11076

Abstract. Hypoxia-inducible factor (HIF)-1 α is a transcription factor that is activated in low oxygen conditions. Adipose tissues are poorly oxygenated in patients with obesity. The low oxygen conditions in obese adipose tissues induce HIF-1 α in adipocytes. Previous studies using genetically modified mice suggest that HIF-1 α contributes to dysfunction in adipocytes. Lipin1 is a bifunctional protein that works as a phosphatidate phosphatase and transcriptional coactivator, which regulates lipid metabolism and adipogenesis, respectively. HIF-1 α directly regulates Lipin1 in hepatocytes. However, the regulation of Lipin1 by HIF-1 α in adipocytes is not well determined. Therefore, the present study investigated the regulation of Lipin1 by HIF-1 α in adipocytes. Expression levels of Lipin1 were reduced in epididymal adipose tissues of adipocyte-specific HIF-1 α knockout mice, indicating that HIF-1 α regulates Lipin1 in adipocytes. In differentiated mature adipocytes, a HIF-1 α activator, dimethylxallyl glycine (DMOG), was demonstrated to increase Lipin1, and a HIF-1 α inhibitor, 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole (YC-1), reversed this increase, indicating that HIF-1 α regulates Lipin1 in differentiated adipocytes. However, during differentiation of pre-adipocytes into adipocytes, YC-1 increased Lipin1 even though HIF-1 α was decreased. The differentiation efficiency increased with YC-1 treatment. In addition, DMOG reduced Lipin1 expression levels during differentiation despite increased HIF-1 α . Under these conditions, differentiation efficiency was reduced. These results suggest that Lipin1 is

negatively regulated by HIF-1 α in pre-adipocytes. Our results show that regulation of Lipin1 by HIF-1 α is different in adipocytes and pre-adipocytes.

Introduction

Obesity is one of the causes of type 2 diabetes. In obesity, adipocytes are hypertrophied, and their functions are aberrant, leading to glucose intolerance or type 2 diabetes. It is widely suggested that adipose tissues are poorly oxygenated in obesity (1-5). Hypoxia activates many genes for cellular adaptation to hypoxic environments. Hypoxia-inducible factor (HIF)-1 α is a transcription factor that responds to low oxygen conditions (6). Prolyl hydroxylase enzymes (PHDs) sense cellular oxygen, leading to degradation of HIF-1 α under normoxic conditions. However, HIF-1 α is stabilized under hypoxic conditions and is transferred to the nucleus where it activates its target genes (7,8). HIF-1 α is expressed in adipocytes from obese adipose tissue under hypoxic conditions (4,9). Because HIF-1 α regulates glucose metabolism, cell survival, and inflammation (6), it is expected that HIF-1 α expression in adipocytes is dysregulated causing inflammation in adipose tissue, thereby inducing the onset of type 2 diabetes. Indeed, previous reports show expression of HIF-1 α in adipocytes leads to inflammation and fibrosis in adipose tissue, secretion defects of hormones and cytokines from adipocytes, increased lipid storage, and whole-body glucose intolerance (5,10-12). In addition, adipocyte-specific knockout of HIF-1 α shows reduced adipocyte sizes in obese adipose tissues (11,12). Transgenic HIF-1 α mice show increased adipocyte size in subcutaneous white adipose tissues (5). These results indicate that HIF-1 α participates in adipogenesis.

Lipin1 plays important roles in lipid homeostasis and metabolism as an enzyme for lipid synthesis and as a nuclear receptor coactivator (13). The enzymatic function of Lipin1 is as a phosphatidate phosphatase (PAP), which catalyzes phosphatidic acid to diacylglycerol, contributing to lipid storage in adipocytes (14,15). As a transcriptional coactivator, Lipin1 forms a complex with peroxisome proliferator-activated receptor (PPAR) α and PPAR γ coactivator 1 (PGC-1) regu-

Correspondence to: Dr Yoshitaka Kihira, Department of Clinical Pharmacy, Faculty of Pharmacy and Pharmaceutical Sciences, Fukuyama University, 1 Sanzo, Gakuen-cho, Fukuyama, Hiroshima 729-0292, Japan
E-mail: kihira@fukuyama-u.ac.jp

Key words: hypoxia-inducible factor, lipin1, adipocytes, obesity, differentiation

lating gene expression for fatty acid oxidation (16). In addition, Lipin1 is expressed in differentiating pre-adipocytes. Lipin1 activation during differentiation of adipocyte requires the adipogenic transcription factors PPAR γ and CCAAT/enhancer binding protein α (C/EBP α) (17). Knockdown of Lipin1 in pre-adipocytes inhibits differentiation into adipocytes, whereas Lipin1 overexpression enhances adipocyte differentiation (18,19). Thus, Lipin1 is important for adipogenesis. Lipin1 is regulated by HIF-1 α in cells of non-adipocyte origin (20). However, the regulation of Lipin1 by HIF-1 α in adipocytes is unknown.

In the present study, we investigated the HIF-1 α regulation of Lipin1 in adipocytes. Lipin1 expression levels in epididymal adipose tissue of adipocyte-specific HIF-1 α knockout mice were significantly decreased relative to wild type mice, indicating that HIF-1 α regulates Lipin1. In differentiated 3T3-L1 adipocytes, HIF-1 α activation induced Lipin1 and HIF-1 α inhibition reduced Lipin1. However, during differentiation, HIF-1 α activation reduced Lipin1 and HIF-1 α inhibition induced Lipin1. Lipin1 expression levels correlated with adipocyte differentiation efficiency. Together, our results indicate that regulation of Lipin1 by HIF-1 α is different in pre-adipocytes and mature adipocytes.

Materials and methods

Chemicals and antibodies. 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole (YC-1) and dimethylxallyl glycine (DMOG) were purchased from Cayman Chemical. Anti-HIF-1 α antibody (Cayman Chemical), anti-Lipin1 antibody (Cell Signaling), and anti- β -actin antibody (Cell Signaling) were used.

Adipocyte-specific HIF-1 α knockout mice. All the experimental procedures were performed in accordance with the guidelines of the Animal Research Committee, Tokushima University. The protocol was approved by the Animal Research Committee, Tokushima University (approval no: 14129). HIF-1 α -floxed mice containing loxP sites flanking exons 13–15 of the HIF-1 α gene (21) were crossed with mice harboring the Cre recombinase under the control of the aP2 promoter (aP2-Cre mice; a gift from Ronald M. Evans, Salk Institute for Biological Studies), generating adipocyte-specific HIF-1 α knockout (ahKO) mice. All mice were C57BL/6 and only male mice were used for experiments. The mice were maintained under temperature- and light-controlled environmental settings with free access to water. Six-week-old mice were fed a high fat diet (HFD) (57% kcal consisting of fat; high fat diet 32 (CLEA Japan)) for 15 weeks. The epididymal fat pads were resected from the wild type and ahKO mice.

Cell culture. 3T3-L1 cells were cultured until confluence, allowed to grow for 2 days postconfluency, and then differentiated with the addition of 500 μ M 3-isobutyl-1-methylxanthine, 1 μ M dexamethasone, and 10 μ g/ml insulin for 2 days. The medium was changed to growth medium supplemented with 10 μ g/ml insulin (differentiation medium) every 2 days. For chemical treatment of differentiated 3T3-L1 adipocytes, the adipocytes were cultured in serum-free media for 24 h and then treated with 500 μ M DMOG or 50 μ M YC-1. For measurement of differentiation efficiency, the area of differentiated adipo-




Figure 1. Lipin1 expression in epididymal adipose tissues in wild type and adipocyte-specific HIF-1 α knockout (ahKO) mice. WT and ahKO mice were fed a high fat diet for 15 weeks. Then, epididymal fat pads were resected. The expression levels of Lipin1 were observed using western blot. (A) Representative western blot and (B) means expression levels (mean \pm S.E.M; n =3) of Lipin1. * P <0.05. HIF, Hypoxia-inducible factor; WT, wild-type.

cytes was divided by the total area in a microscopic field taken at magnification, $\times 100$.

Western blot analysis. Epididymal fat pads and cells were lysed in lysis buffer (20 mM Tris-HCl, pH 8.0), 0.15 M NaCl, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, protease inhibitor mixture (2 g/ml aprotinin, 1 μ g/ml leupeptin, 2 μ g/ml antipain, and 10 μ g/ml benzamidine), and phosphatase inhibitor mixture (10 mM NaF, 60 mM β -glycerophosphate, 10 mM sodium pyrophosphate, and 2 mM sodium orthovanadate). Proteins were separated on SDS polyacrylamide gels and electrophoretically transferred to polyvinylidene fluoride membranes. The membranes were incubated with a primary antibody overnight at 4°C and probed with an HRP-conjugated secondary antibody (KPL). Immunoreactive bands were detected with ECL (GE Healthcare) and visualized by exposing the membranes to X-ray films (GE Healthcare). The proteins were quantified by densitometric analysis using ImageJ analysis software.

Statistical analysis. Data are presented as mean values \pm standard error of the mean (S.E.M.). Statistical significance was assessed using the Student's *t*-test or two-way analysis of variance (ANOVA) with Sidak's multiple comparisons test, where values of P <0.05 were considered to indicate a statistically significant difference. Prism version 6.0h (GraphPad Software) was used for data analysis.

Results

Lipin1 is decreased in ahKO mice. Lipin1 is regulated by HIF-1 α in cells of non-adipocyte origin (20). However, the regulation of Lipin1 by HIF-1 α in adipocytes is unknown.




Figure 2. Effects of activation and inhibition of HIF-1 α on Lipin1 expression in differentiated adipocytes. Differentiated 3T3-L1 cells were treated with 500 μ M DMOG and 50 μ M YC-1 for 4 h. (A) Representative western blot analysis. Expression levels (mean \pm S.E.M; n=3) of (B) HIF-1 α and (C) Lipin1, respectively. *P<0.05. HIF, Hypoxia-inducible factor; DMOG, dimethyloxallyl glycine; YC-1, 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole.

Therefore, we investigated the expression of Lipin1 in adipose tissue from ahKO mice. Wild type (WT) and ahKO mice were fed an HFD for 15 weeks and then the epididymal adipose tissues were resected. Western blots showed that the expression levels of Lipin1 in the epididymal adipose tissues of ahKO mice significantly decreased compared with WT mice (Fig. 1). This result indicates that HIF-1 α regulates Lipin1 in adipocytes.

Lipin1 regulation by HIF-1 α in differentiated 3T3-L1 adipocytes. To assess whether HIF-1 α regulates Lipin1 in adipocytes, 3T3-L1 cells were differentiated to adipocytes and treated with a HIF-1 α activator, DMOG, for 4 h. DMOG treatment significantly increased HIF-1 α expression levels in the adipocytes (Fig. 2A and B). In addition, DMOG also elevated Lipin1 expression levels in the differentiated adipocytes under the same conditions (Fig. 2A and C). To confirm the regulation of Lipin1 by HIF-1 α , the effect of YC-1, an inhibitor of HIF-1 α , on the elevation of Lipin1 was studied. The DMOG-induced increases of HIF-1 α and Lipin1 were decreased by YC-1 (Fig. 2A-C), suggesting that HIF-1 α regulates Lipin1 in the differentiated adipocytes.

DMOG suppresses the differentiation of 3T3-L1 pre-adipocytes into adipocytes. Lipin1 has two different functions: one is as an enzyme regulating lipid metabolism (14,15) and the other is as a mediator of differentiation into adipocytes (17). Therefore, we studied the relationship between HIF-1 α and Lipin1 along with the effect of DMOG on the differentiation

of 3T3-L1 pre-adipocytes into adipocytes. we studied the effect of DMOG on differentiation of 3T3-L1 cells. Treatment with DMOG significantly increased HIF-1 α expression levels in 3T3-L1 cells on days 2, 4, and 8 during differentiation (Fig. 3A and B). In contrast, the expression levels of Lipin1 significantly decreased on day 6 and 8. The differentiation efficiency of 3T3-L1 cells was significantly reduced with DMOG (Fig. 3D and E). These results show that DMOG reduces Lipin1 expression levels and differentiation efficiency, whereas HIF-1 α expression levels increase.

YC-1 accelerates the differentiation of 3T3-L1 pre-adipocytes into adipocytes. The effect of the HIF-1 α inhibitor, YC-1, during differentiation of 3T3-L1 cells, was investigated. HIF-1 α was induced in the initial 24 h and then gradually decreased under the normal conditions for differentiation (Fig. 4A and B). In contrast, Lipin1 increased at day 4 after differentiation initiation. Next, the effect of YC-1 on differentiation of 3T3-L1 pre-adipocytes into adipocytes was investigated. 3T3-L1 pre-adipocytes were differentiated in differentiation medium containing YC-1. Addition of YC-1 during differentiation of 3T3-L1 cells reduced HIF-1 α expression levels in the initial 24 h of the differentiation. Under the same conditions, Lipin1 increased at day 6 and day 8 after differentiation initiation. YC-1 treatment significantly increased the differentiation efficiency of 3T3-L1 cells (Fig. 4D and E). The expression of PPAR γ (an adipogenic transcription factor) and adiponectin (an adipose-secreted protein) was significantly increased in YC-1 treated adipocytes at day 8 (Fig. 5).




Figure 3. Effect of activation of HIF-1 α on expression of Lipin1 and differentiation efficiency of pre-adipocytes. 3T3-L1 pre-adipocytes were differentiated in differentiation medium with or without 500 μ M DMOG. (A) Representative western blot. The day after differentiation initiation is indicated on the top of the blot. (B and C) Expression levels of HIF-1 α and Lipin1, respectively. Values are mean \pm S.E.M.; (n=3). (D) Cells 8 days after differentiation initiation. Magnification, $\times 100$. (E) Differentiation efficiency. * $P < 0.05$. HIF, Hypoxia-inducible factor; DMOG, dimethylxallyl glycine; S.E.M., standard error of the mean.

Discussion

Lipin1 is regulated by HIF-1 α in cells of non-adipocyte origin (20). However, the regulation of Lipin1 by HIF-1 α in adipocytes is unknown. Therefore, in the present study, we focused on the regulation of Lipin1 by HIF-1 α . We found that Lipin1 was significantly decreased in epididymal adipose tissues of *ahKO* mice (Fig. 1). This result indicates that HIF-1 α regulates Lipin1 in adipocytes in the adipose tissue of mice. In addition, HIF-1 α upregulates Lipin1 in mature adipocytes but downregulates Lipin1 in pre-adipocytes.

We found that DMOG, a HIF-1 α activator, induced Lipin1 in differentiated adipocytes (Fig. 2). In addition, YC-1, a HIF-1 α inhibitor, canceled the induction of Lipin1 with DMOG (Fig. 2). These results indicate that HIF-1 α regulates Lipin1 in differentiated adipocytes. Previously, it was shown that Lipin1 is regulated by HIF-1 α in cells of non-adipocyte origin (20,22,23). Our results show that Lipin1 is regulated by HIF-1 α in differentiated adipocytes as well. Previously, it has been shown that HIF-1 α -induced Lipin1 causes the accumulation of lipids in hepatocytes (20). Therefore, HIF-1 α activation in adipocytes can also cause excess lipid accumulation, leading to metabolic disorders.

During differentiation of pre-adipocytes into adipocytes, HIF-1 α levels gradually decreased (Figs. 3 and 4). However, although HIF-1 α expression had gradually decreased by day 6 in the DMOG-treated cells its expression increased at day 8 (Fig. 3). HIF-1 α is constantly synthesized and degraded (6). Therefore, the balance between synthesis and degradation of HIF-1 α changes to synthesis dominant around day 8 after increased differentiation, suggesting that the effects of DMOG is increased. In addition, because HIF-1 α mRNA levels increase in the initial 3 to 6 h after differentiation (24), mRNA regulation partly contributed to the initial increased expression of HIF-1 α protein after differentiation.

During differentiation, YC-1, an inhibitor of HIF-1 α , reduced HIF-1 α expression levels in the pre-adipocytes. However, regulation of Lipin1 by HIF-1 α was opposite in differentiating pre-adipocytes compared with differentiated adipocytes. YC-1 reduced HIF-1 α expression levels during the initial 24 h, whereas it increased Lipin1 expression levels 6 to 8 days after differentiation initiation (Fig. 4). This result indicates that HIF-1 α inhibition indirectly participated in the upregulation of Lipin1. To activate Lipin1 during differentiation of adipocytes, the adipogenic transcription factors PPAR γ and C/EBP α are required (17). In the study, PPAR γ was increased




Figure 4. Effect of HIF-1 α inhibition on expression of Lipin1 and differentiation efficiency of pre-adipocytes. 3T3-L1 pre-adipocytes were differentiated in differentiation medium with or without 50 μ M YC-1. (A) Representative western blot analysis. The day after initiating differentiation is indicated on the top of the blot. Expression levels of (B) HIF-1 α and (C) Lipin1, respectively. Values are mean \pm S.E.M. (n=3). (D) Cells 8 days after differentiation initiation. Magnification, $\times 100$. (E) Differentiation efficiency. *P<0.05, **P<0.01. HIF, Hypoxia-inducible factor; YC-1, 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole, S.E.M., standard error of the mean.




Figure 5. Effect of HIF-1 α inhibition on the expression of PPAR γ and adiponectin. 3T3-L1 pre-adipocytes were differentiated in a differentiation medium with or without 50 μ M YC-1. (A) Representative western blot. The day after initiating differentiation is indicated on the top of the blot. (B and C) Expression levels of PPAR γ and adiponectin, respectively. Values are mean \pm S.E.M. (n=3). *P<0.05. YC-1, 3-(5'-hydroxymethyl-2'-furyl)-1-benzylindazole; HIF, Hypoxia-inducible factor; PPAR γ , peroxisome proliferator-activated receptor; S.E.M., standard error of the mean.

in YC-1 treated adipocytes (Fig. 5). Further, it was reported that HIF-1 α indirectly downregulates PPAR γ (25). Therefore, the inhibition of HIF-1 α probably contributes to upregulation of PPAR γ , leading to Lipin1 induction during YC-1-affected differentiation of adipocytes. Adiponectin is an adipokine derived from the adipocytes and has anti-inflammatory and insulin-sensitizing effects (26). In obese adipose tissues, the secretion of adiponectin is decreased (27-30). YC-1-affected differentiation increased adiponectin expression, indicating that the differentiated adipocytes are healthy. In addition, DMOG increased HIF-1 α expression levels 2, 4, and 8 days after starting differentiation. On the contrary, Lipin1 was decreased 6 and 8 days after differentiation initiation. In this condition, differentiation efficiency was quite low (Fig. 3). Previous reports showed that hypoxia inhibits adipogenesis of 3T3-L1 cells through HIF-1 α (25,31,32). Our results suggest that the hypoxic inhibition of adipogenesis may participate in Lipin1 downregulation.

In *in vivo* studies, adipocyte-specific knockout of HIF-1 α protects obese mice from insulin resistance and inflammation (11,12), whereas transgenic mice expressing a constitutively active form of HIF-1 α have insulin resistance and tissue fibrosis (5). In the studies, the size of adipocytes were smaller in the adipocyte-specific HIF-1 α knockout mice than in WT mice (11,12). In the overexpression study of a constitutively active form of HIF-1 α , the transgenic mice showed increased adipocyte size in subcutaneous white adipose tissues (5). In addition, HIF overexpression with PHD2 deletion in mice reduces lipolysis and increases lipid storage (33). Our results suggest that lipid accumulation in the knockout mice might be decreased by reduced Lipin1 expression in differentiated adipocytes.

Some limitations exist in the present study. Our study showed that HIF-1 α upregulates Lipin1 in mature adipocytes but downregulates it in pre-adipocytes. However, the regulation of Lipin1 by HIF-1 α in ahKO mice was not clear. To further study this, the effects of YC-1 on the adipose tissue of ahKO mice, or HIF-1 α knockdown as reported in the previous study (34), should be observed.

It is possible that regulation mechanisms of Lipin1 by HIF-1 α are different between pre-adipocytes and adipocytes. HIF-1 α reduces Lipin1 during differentiation of pre-adipocytes and reduces differentiation efficiency. HIF-1 α directly increases Lipin1 in differentiated adipocytes and regulates lipid metabolism. Regulation of the HIF-1 α - Lipin1 system may be a potential therapeutic target for the treatment of obesity and type 2 diabetes.

Acknowledgements

Not applicable.

Funding

The present study was partially supported by KAKENHI (grant no. 16K08272).

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

YK and ST contributed to the conception and design of the study, acquired and analyzed the data and drafted the manuscript. YF contributed to acquiring and analyzing the data. TT and ES contributed to the design of the study, revised the manuscript and approved the final manuscript.

Ethics approval and consent to participate

All the experimental procedures were performed in accordance with the guidelines of the Animal Research Committee, Tokushima University. The protocol was approved by the Animal Research Committee, Tokushima University (approval no. 14129).

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Hosogai N, Fukuhara A, Oshima K, Miyata Y, Tanaka S, Segawa K, Furukawa S, Tochino Y, Komuro R, Matsuda M, *et al*: Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes* 56: 901-911, 2007.
- Rausch ME, Weisberg S, Vardhana P and Tortorello DV: Obesity in C57BL/6J mice is characterized by adipose tissue hypoxia and cytotoxic T-cell infiltration. *Int J Obes* 32: 451-463, 2008.
- Wang B, Wood IS and Trayhurn P: Dysregulation of the expression and secretion of inflammation-related adipokines by hypoxia in human adipocytes. *Pflugers Arch* 455: 479-492, 2007.
- Ye J, Gao Z, Yin J and He Q: Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice. *Am J Physiol Endocrinol Metab* 293: E1118-E1128, 2007.
- Halberg N, Khan T, Trujillo ME, Wernstedt-Asterholm I, Attie AD, Sherwani S, Wang ZV, Landskroner-Eiger S, Dineen S, Magalang UJ, *et al*: HIF 1 α induces fibrosis and insulin resistance in white adipose tissue. *Mol Cell Biol* 29: 4467-4483, 2009.
- Semenza GL: Targeting HIF-1 for cancer therapy. *Nat Rev Cancer* 3: 721-732, 2003.
- Brahimi-Horn MC and Pouyssegur J: HIF at a glance. *J Cell Sci* 122: 1055-1057, 2009.
- Huang LE, Arany Z, Livingston DM and Bunn HF: Activation of hypoxia-inducible transcription factor depends primarily upon redox-sensitive stabilization of its α subunit. *J Biol Chem* 271: 32253-32259, 1996.
- Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, Salic A, Asara JM, Lane WS and Kaelin WG Jr: HIF α targeted for VHL-mediated destruction by proline hydroxylation: Implications for O₂ sensing. *Science* 292: 464-468, 2001.
- Kihira Y, Miyake M, Hirata M, Hoshina Y, Kato K, Shirakawa H, Sakaue H, Yamano N, Izawa-Ishizawa Y, Ishizawa K, *et al*: Deletion of hypoxia-inducible factor-1 α in adipocytes enhances glucagon-like peptide-1 secretion and reduces adipose tissue inflammation. *PLoS One* 9: e93856, 2014.
- Jiang C, Kim JH, Li F, Qu A, Gavrilova O, Shah YM and Gonzalez FJ: Hypoxia-inducible factor 1 α regulates a SOCS3-STAT3-adiponectin signal transduction pathway in adipocytes. *J Biol Chem* 288: 3844-3857, 2013.
- Jiang C, Qu A, Matsubara T, Chanturiya T, Jou W, Gavrilova O, Shah YM and Gonzalez FJ: Disruption of hypoxia-inducible factor 1 in adipocytes improves insulin sensitivity and decreases adiposity in high-fat diet-fed mice. *Diabetes* 60: 2484-2495, 2011.
- Reue K and Dwyer JR: Lipin proteins and metabolic homeostasis. *J Lipid Res* 50 (Suppl): S109-S114, 2009.

14. Phan J and Reue K: Lipin, a lipodystrophy and obesity gene. *Cell Metab* 1: 73-83, 2005.
15. Langner CA, Birkenmeier EH, Ben-Zeev O, Schotz MC, Sweet HO, Davisson MT and Gordon JI: The fatty liver dystrophy (fld) mutation. A new mutant mouse with a developmental abnormality in triglyceride metabolism and associated tissue-specific defects in lipoprotein lipase and hepatic lipase activities. *J Biol Chem* 264: 7994-8003, 1989.
16. Finck BN, Gropler MC, Chen Z, Leone TC, Croce MA, Harris TE, Lawrence JC Jr and Kelly DP: Lipin 1 is an inducible amplifier of the hepatic PGC-1alpha/PPARalpha regulatory pathway. *Cell Metab* 4: 199-210, 2006.
17. Phan J, Péterfy M and Reue K: Lipin expression preceding peroxisome proliferator-activated receptor-gamma is critical for adipogenesis in vivo and in vitro. *J Biol Chem* 279: 29558-29564, 2004.
18. Péterfy M, Phan J, Xu P and Reue K: Lipodystrophy in the fld mouse results from mutation of a new gene encoding a nuclear protein, lipin. *Nat Genet* 27: 121-124, 2001.
19. Koh YK, Lee MY, Kim JW, Kim M, Moon JS, Lee YJ, Ahn YH and Kim KS: Lipin1 is a key factor for the maturation and maintenance of adipocytes in the regulatory network with CCAAT/enhancer-binding protein alpha and peroxisome proliferator-activated receptor gamma 2. *J Biol Chem* 283: 34896-34906, 2008.
20. Mylonis I, Sembongi H, Befani C, Liakos P, Siniossoglou S and Simos G: Hypoxia causes triglyceride accumulation by HIF-1-mediated stimulation of lipin 1 expression. *J Cell Sci* 125: 3485-3493, 2012.
21. Tomita S, Ueno M, Sakamoto M, Kitahama Y, Ueki M, Maekawa N, Sakamoto H, Gassmann M, Kageyama R, Ueda N, *et al*: Defective brain development in mice lacking the Hif-1alpha gene in neural cells. *Mol Cell Biol* 23: 6739-6749, 2003.
22. Arai T, Tanaka M and Goda N: HIF-1-dependent lipin1 induction prevents excessive lipid accumulation in choline-deficient diet-induced fatty liver. *Sci Rep* 8: 14230, 2018.
23. Kourti M, Ikonomou G, Giakoumakis NN, Rapsomaniki MA, Landegren U, Siniossoglou S, Lygerou Z, Simos G and Mylonis I: CK1δ restrains lipin-1 induction, lipid droplet formation and cell proliferation under hypoxia by reducing HIF-1α/ARNT complex formation. *Cell Signal* 27: 1129-1140, 2015.
24. Imagawa M, Tsuchiya T and Nishihara T: Identification of inducible genes at the early stage of adipocyte differentiation of 3T3-L1 cells. *Biochem Biophys Res Commun* 254: 299-305, 1999.
25. Lin Q, Lee YJ and Yun Z: Differentiation arrest by hypoxia. *J Biol Chem* 281: 30678-30683, 2006.
26. Yun Z, Maecker HL, Johnson RS and Giaccia AJ: Inhibition of PPAR gamma 2 gene expression by the HIF-1-regulated gene DEC1/Stral3: A mechanism for regulation of adipogenesis by hypoxia. *Dev Cell* 2: 331-341, 2002.
27. Zhou S, Lechpammer S, Greenberger JS and Glowacki J: Hypoxia inhibition of adipocytogenesis in human bone marrow stromal cells requires transforming growth factor-beta/Smad3 signaling. *J Biol Chem* 280: 22688-22696, 2005.
28. Ouchi N and Walsh K: Adiponectin as an anti-inflammatory factor. *Clin Chim Acta* 380: 24-30, 2007.
29. Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, *et al*: The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med* 7: 941-946, 2001.
30. Fruebis J, Tsao TS, Javorschi S, Ebbets-Reed D, Erickson MR, Yen FT, Bihain BE and Lodish HF: Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci USA* 98: 2005-2010, 2001.
31. Berg AH, Combs TP, Du X, Brownlee M and Scherer PE: The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 7: 947-953, 2001.
32. Kern PA, Di Gregorio GB, Lu T, Rassouli N and Ranganathan G: Adiponectin expression from human adipose tissue: Relation to obesity, insulin resistance, and tumor necrosis factor-α expression. *Diabetes* 52: 1779-1785, 2003.
33. Michailidou Z, Morton NM, Moreno Navarrete JM, West CC, Stewart KJ, Fernández-Real JM, Schofield CJ, Seckl JR and Ratcliffe PJ: Adipocyte pseudohypoxia suppresses lipolysis and facilitates benign adipose tissue expansion. *Diabetes* 64: 733-745, 2015.
34. Wang F, Zhang G, Xing T, Lu Z, Li J, Peng C, Liu G and Wang N: Renalase contributes to the renal protection of delayed ischaemic preconditioning via the regulation of hypoxia-inducible factor-1α. *J Cell Mol Med* 19: 1400-1409, 2015.