Anti-inflammatory effects of the NF- κ B inhibitor dehydroxymethylepoxyquinomicin on ARPE-19 cells

YOSHIMASA ANDO¹, YASUHIKO SATO², AKIHIKO KUDO³, TAKAYO WATANABE¹,
AKITO HIRAKATA¹, ANNABELLE A. OKADA¹, KAZUO UMEZAWA⁴ and HIROSHI KEINO¹

¹Department of Ophthalmology; ²Division of Radioisotope Research; ³Department of Anatomy,
Kyorin University School of Medicine, Tokyo 181-8611; ⁴Department of Molecular Target
Medicine Screening, Aichi Medical University, Nagakute, Aichi 480-1195, Japan

Received August 27, 2019; Accepted March 26, 2020

DOI: 10.3892/mmr.2020.11115

Abstract. The retinal pigment epithelium (RPE) is a polarized, monolayer of pigmented cells that forms the outer retinal layer. A key function of the RPE is to maintain the integrity of the photoreceptors mainly via phagocytosis and recycling of the digested photoreceptor outer segments. Moreover, RPE cells are a major source of inflammatory cytokines and chemokines, which play important roles in the activation of other immune cells under inflammatory conditions in the posterior segment of the eye. Dehydroxymethylepoxyquinomicin (DHMEQ) is a NF- κ B inhibitor and its structure is related to that of epoxyquinomicin C, which is an antibiotic. The present study evaluated the anti-inflammatory effects of DHMEQ on a human retinal pigment epithelial cell line (ARPE-19). It was revealed that high concentrations of DHMEQ (100 μ g/ml) induced apoptosis and necrosis of tumor necrosis factor (TNF)- α -stimulated ARPE-19 cells. Furthermore, the percentage of intercellular adhesion molecule 1 (ICAM-1)-positive TNF- α -stimulated cells was significantly reduced in the presence of DHMEQ (10 μ g/ml), as determined by flow cytometry. It was also demonstrated that DHMEQ exposure significantly decreased the levels of interleukin (IL)-8 and monocyte chemoattractant protein-1 (MCP-1) in the supernatant of cultured ARPE-19 cells as determined by ELISA. Moreover, the protein expression levels of IL-8 and MCP-1 were significantly reduced in ARPE-19 cells exposed to DHMEQ compared with cells exposed to dexamethasone. PCR array analysis revealed that DHMEQ reduced the expression levels of MCP-1, ICAM-1, IL-6, Toll-like receptor (TLR)2, TLR3 and TLR4. Therefore, the present results indicated that DHMEQ has anti-inflammatory effects on TNF- α -stimulated ARPE-19 cells. Thus,

DHMEQ may have therapeutic potential for TNF- α -mediated inflammatory disorders of the eye.

Introduction

The retinal pigment epithelium (RPE) is a polarized, monolayer of pigmented cells that forms the outer retinal layer, and maintains the integrity of the photoreceptors, primarily by phagocytosing and recycling the retinal photoreceptor outer segments (1). RPE cells are a major source of proinflammatory cytokines, including interleukin (IL)-6, and chemokines, such as monocyte chemotactic protein (MCP)-1 and IL-8 (1). RPE cells also secrete regulated on the activation of normal T-cell expressed and secreted (RANTES) and interferon (IFN)- γ induced protein (IP)-10 kDa (IP-10) (1-3). Furthermore, these cytokines and chemokines secreted by RPE cells play important roles in the activation of other immune cells under inflammatory conditions of the posterior segment of the eye (1).

Tumor necrosis factor- α (TNF- α) is an inflammatory cytokine that contributes to the progression of non-infectious uveitis, and it has been shown that blocking TNF- α is effective for treating refractory uveitis (4-8). Furthermore, TNF- α receptors activate the NF- κ B signaling pathway (9). NF- κ B, a member of a family of ubiquitously expressed proteins, is usually found in an inactive state in the cytoplasm, except during immune and inflammatory responses (9,10). The NF- κ B family includes REL proto-oncogene (Rel)A, RelB, c-Rel, p50/p105 and p52/p100, all of which form homo- or heterodimers with each other (11). Inhibitors of NF- κ B (I κ B) proteins are phosphorylated and are degraded by proteasomes (12). Moreover, released NF- κ B dimers translocate into the nucleus and bind to κ B sites in the promoter and enhancer regions of targeted genes of various inflammatory cytokines and chemokines, including IL-1, IL-2, IL-6, TNF and macrophage inflammatory protein-1/2, and adhesion molecules, such as intercellular adhesion molecule-1 (ICAM-1) (11,13).

Dehydroxymethylepoxyquinomicin (DHMEQ) is a low molecular weight inhibitor of the NF- κ B signaling pathway, and its structure is related to that of epoxyquinomicin C, which is an antibiotic (14,15). DHMEQ suppresses the TNF- α -induced nuclear translocation of NF- κ B, but it does not prevent the phosphorylation and degradation of I κ B (16).

Correspondence to: Dr Hiroshi Keino, Department of Ophthalmology, Kyorin University School of Medicine, 6-20-2 Shinkawa, Mitaka, Tokyo 181-8611, Japan
E-mail: keino@eye-center.org

Key words: NF- κ B, dehydroxymethylepoxyquinomicin, ARPE-19, dexamethasone, tumor necrosis factor- α

A previous study also revealed that DHMEQ binds directly to the Rel-family proteins to prevent their DNA-binding activity (17). Furthermore, it has been revealed that DHMEQ is able to suppress inflammation and the progression of cancer in animal models without obvious adverse effects (18).

The cells of the human RPE cell line, ARPE-19, are frequently used in *in vitro* studies to investigate the mechanisms involved in posterior segment inflammatory disorders, including uveitis and age-related macular degeneration (19-21). Therefore, the aim of the present study was to determine whether DHMEQ has inhibitory effects on the expression of ICAM-1 in TNF- α -stimulated ARPE-19 cells. In addition, the present study examined whether DHMEQ can affect the production of TNF- α -stimulated ARPE-19 cells and the expression of NF- κ B related-genes in TNF- α -stimulated ARPE-19 cells treated with DHMEQ.

Materials and methods

Materials. DHMEQ was synthesized by Umezawa and Chaicharoenpong (16), and for the present study it was dissolved in 100% DMSO at a concentration of 10 mg/ml and stored at -30°C (14,16). Before use in cell cultures, DHMEQ was diluted with the culture medium (DMEM/F-12; Invitrogen; Thermo Fisher Scientific, Inc.) to a final concentration of $\leq 0.1\%$. Dexamethasone was purchased from Sigma-Aldrich (Merck KGaA).

Cell cultures. ARPE-19 cells were purchased from the American Type Culture Collection and maintained in DMEM/F-12 supplemented with 10% FBS, 100 U/ml penicillin and 0.1 mg/ml streptomycin (all from Invitrogen; Thermo Fisher Scientific, Inc.) at 37°C in 5% CO₂ in air. After reaching confluency, ARPE-19 cells were detached with a trypsin-EDTA solution (0.05%) (Thermo Fisher Scientific, Inc.) and plated for subcultures. Cells were passaged every 4-6 days, and those used in each experiment were confluent and exhibited no visible pigmentation. Moreover, cells were maintained for 3 weeks before the experimental procedures and were used at passages 4-6.

Cell viability determined by MTT assay. The effects of various concentrations of DHMEQ on ARPE-19 cells were evaluated by MTT. Cells were grown at 37°C in 96-well plates at a density of 2×10^4 cells/well for 24 h. Upon confluency, the medium was replaced with a serum-free medium, and ARPE-19 cells were cultured at 37°C for 24 h with 0.1, 1.0, 5.0, 10.0, 50.0 or 100.0 μ g/ml DHMEQ or without DHMEQ at 37°C for 24 h. After 24 h, the assay was performed by adding 10 μ l MTT solution to the wells (Biotium, Inc.). After incubation for 4 h at 37°C, 200 μ l DMSO was added to the cells. After incubation at room temperature for 5 min, the optical density (OD) at 570 nm (signal absorbance) and 630 nm (background absorbance) was measured using a microplate reader, and the normalized absorbance values (OD at 570 nm and OD at 630) were determined.

Flow cytometric analyses. ARPE-19 cells were seeded in 6-well plates at 2×10^5 cells/well and cultured for 24 h. Upon confluency, the medium was replaced with serum-free

medium, and cells were or were not exposed to 20 ng/ml TNF- α (R&D Systems, Inc.) with DHMEQ (1.0, 10.0 and 100.0 μ g/ml) or without DHMEQ at 37°C for 24 h. After exposure, cells were washed with phosphate-buffered saline (PBS, pH 7.4), detached by trypsin-EDTA (0.05%) and suspended in PBS. For staining of prepared cells, Annexin V-FITC solution (5 μ l; Nacalai Tesque, Inc.) was added to 100 μ l of cell suspension (1×10^6 cells), then $<1\%$ propidium iodide (PI) solution (5 μ l; Nacalai Tesque, Inc.) was added to the cell suspension, according to the manufacturer's instructions (cat. no. 15342; Nacalai Tesque, Inc.). The cells were incubated at room temperature for 15 min and analyzed by flow cytometry (FACSCalibur) using CellQuest Pro software version 6.0 (both BD Biosciences).

ARPE-19 cells were seeded in 6-well plates at 2×10^5 cells/well and cultured for 24 h. Upon confluency, the medium was replaced with serum-free medium, and cells were or were not exposed to 20 ng/ml TNF- α (R&D Systems, Inc.) with DMSO (0.1%) or DHMEQ (1.0 or 10.0 μ g/ml) at 37°C for 24 h. After exposure, cells were washed with PBS (pH 7.4), detached by trypsin-EDTA (0.05%) and suspended in PBS. The prepared ARPE-19 cells were incubated with phycoerythrin-conjugated monoclonal antibody to ICAM-1 (1:100; cat. no. 555511; BD Biosciences) at 4°C for 20 min and analyzed by flow cytometry (FACSCalibur) using CellQuest Pro software version 6.0 (both BD Biosciences).

Chemokine assay in culture supernatants. ARPE-19 cells were seeded in 6-well plates at a density of 2×10^5 cells/well and cultured for 24 h. Upon confluency (80-90% confluency), the medium was replaced with serum-free medium and cells were exposed to 20 ng/ml TNF- α and DMSO (0.1%), DHMEQ (1.0 μ g/ml=4.0 μ M, 10 μ g/ml=40 μ M) or dexamethasone (40 μ M) at 37°C for 24 h. After exposure, the supernatant was collected from each well, and the levels of IL-8 and MCP-1 in the supernatant were determined by Quantikine® Colorimetric Sandwich ELISA kits (R&D Systems, Inc.; IL-8, cat. no. D8000C; MCP-1, cat. no. DCP00).

In another experiment, ARPE-19 cells were exposed to 20 ng/ml TNF- α and DMSO (0.1%) or DHMEQ (10 μ g/ml) at 37°C for 6, 12 and 24 h. After exposure, the supernatant was collected from each well and the levels of IL-8 and MCP-1 in the supernatant was determined by the Quantikine® ELISA kits (R&D Systems, Inc.) as described above.

NF- κ B-associated gene expression level assay. ARPE-19 cells were seeded in 6-well plates at a density of 2×10^5 cells/well and cultured for 24 h. Upon confluency, the medium was replaced with serum-free medium and cells were exposed to 20 ng/ml TNF- α in the absence or presence of DHMEQ (10 μ g/ml) at 37°C for 24 h. After exposure, cells were washed with PBS, detached by trypsin-EDTA (0.05%) and suspended in PBS. Total RNA from the prepared ARPE-19 cells was extracted with ISOGEN (Nippon Gene Co., Ltd.) according to the manufacturer's instructions. Briefly, 10 μ g total RNA from each sample was reverse-transcribed using the High Capacity RNA-to-cDNA kit (Applied Biosystems; Thermo Fisher Scientific, Inc.; reverse transcription conditions were as follows: Initial incubation at 37°C for 60 min and 95°C for 5 min), then loaded onto Human NF κ B Pathway TaqMan®

Array plates (cat. no. 4414095; Applied Biosystems; Thermo Fisher Scientific, Inc.) for profiling of NF- κ B-associated 92 genes expression levels. PCR was performed on a QuantStudio™ 12K Flex Real-Time PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc.), according to the manufacturer's instructions. Raw cycle threshold (Cq) values were calculated with SDS software 1.2.3. (Applied Biosystems; Thermo Fisher Scientific, Inc.). The sequences of the forward and reverse primers are not commercially available. Data were analyzed according to the comparative Cq method, and the global median normalization method was used (22). The $2^{-\Delta\Delta Cq}$ method was performed to calculate the expression level of the fold change (23). The fold change in ARPE-19 cells exposed to 20 ng/ml TNF- α in the absence or presence of DHMEQ was calculated for each gene; genes with a 2-fold increase in this ratio were defined arbitrarily as upregulated in cells exposed to DHMEQ, whereas those with a 2-fold decrease were defined as downregulated genes.

Quantitative PCR analysis. In order to validate the expression levels of genes [lymphotoxin β receptor (LTBR), MCP-1, and Toll-like receptor 4 (TLR4)], which were either upregulated or downregulated in ARPE-19 cells treated with DHMEQ, quantitative PCR was carried out in duplicate using the TaqMan® Universal PCR Master Mix (Applied Biosystems; Thermo Fisher Scientific, Inc.) on a QuantStudio™ 12K Flex Real-Time PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc.), following the manufacturer's protocol. ARPE-19 cells were seeded in 6-well plates at a density of 2×10^5 cells/well and cultured for 24 h. Upon confluency, the medium was replaced with serum-free medium and cells were exposed to 20 ng/ml TNF- α exposed to DMSO (0.1%) or DHMEQ (10 μ g/ml) at 37°C for 24 h. After exposure, cells were washed with PBS, detached by trypsin-EDTA (0.05%) and suspended in PBS. Total RNA from the prepared ARPE-19 cells was extracted with ISOGEN (Nippon Gene Co., Ltd.) as described above and reverse-transcribed using the High Capacity RNA-to-cDNA kit (Applied Biosystems; Thermo Fisher Scientific, Inc.) PCR conditions were as follows: Initial incubation at 50°C for 2 min and 95°C for 10 min, followed by 40 cycles two-step cycling (denaturing at 95°C for 15 sec, annealing/extension at 60°C for 60 sec). The TaqMan primers/probes pairs were obtained from Applied Biosystems (Thermo Fisher Scientific, Inc.) using inventoried TaqMan gene expression assays [LTBR assay ID, Hs01101194_m1; MCP-1 assay ID, Hs00234140_m1; TLR4 assay ID, Hs00152939_m1]. For an endogenous control mRNA, the β -actin (assay ID, Hs99999903_m1, Thermo Fisher Scientific, Inc.) was used for data normalization of the mRNA expression levels.

Statistical analyses. Data are presented as the mean $\pm$ SD. Statistical significance was evaluated with unpaired t-tests. To compare data among ≥ 3 groups, one-way ANOVA using the Bonferroni's multiple comparison tests was performed. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Effect of DHMEQ on the viability of ARPE-19 cells. Confluent ARPE-19 cells were exposed to various concentrations of

Figure 1. Viability of ARPE-19 cells incubated with DHMEQ. Cells were incubated with or without DHMEQ (0-100 μ g/ml). After 24 h, cell viability was evaluated by MTT assay. Data are presented as the mean $\pm$ SD, n=4. * $P < 0.05$. DHMEQ, dehydroxymethylepoxyquinomicin; OD, optical density.

DHMEQ, and it was revealed that DHMEQ at concentrations ≤ 10 μ g/ml did not have toxic effects on cells. However, DHMEQ at concentrations of 50 and 100 μ g/ml significantly inhibited the viability of ARPE-19 cells compared with the viability of cells cultured without DHMEQ (Fig. 1). Therefore, these results indicated that higher concentrations of DHMEQ, such as 50 and 100 μ g/ml, reduce the viability of ARPE-19 cells.

Effect of DHMEQ on the induction of apoptosis and necrosis of ARPE-19 cells. To determine whether DHMEQ induces apoptosis or necrosis in ARPE-19 cells, Annexin-V and/or PI-positive cells were analyzed by flow cytometry. It was demonstrated that concentrations of 1.0 and 10 μ g/ml DHMEQ did not alter the percentage of apoptotic cells (Annexin-V-positive and PI-negative; Fig. 2A and B). However, a dose of 100 μ g/ml DHMEQ significantly increased the number of apoptotic cells compared with cells treated with TNF- α (20 ng/ml) without DHMEQ (Fig. 2A and B). In addition, concentrations of 100 μ g/ml DHMEQ significantly increased the number of necrotic cells compared with cells treated with TNF- α (20 ng/ml) without DHMEQ (Annexin-V-positive and PI-positive; Fig. 2A and B). Collectively, the results indicated that a high concentration of DHMEQ, such as 100 μ g/ml, promoted the induction of apoptosis and necrosis of ARPE-19 cells. Based on these findings, a concentration of DHMEQ < 100 μ g/ml was used in subsequent experiments.

Suppression of ICAM-1 in ARPE-19 cells by DHMEQ. Previous studies have revealed that TNF- α can enhance the protein expression of ICAM-1 synthesized by ARPE-19 cells (24). To assess these findings, the present study stimulated ARPE-19 cells with TNF- α (20 ng/ml) for 24 h and evaluated the protein expression of ICAM-1 by flow cytometry. The results revealed that the expression of ICAM-1 was increased 7-fold in cells stimulated by TNF- α compared with cells cultured without TNF- α (Fig. 3A and B).

The effects of DHMEQ on the protein expression level of ICAM-1 were also examined in ARPE-19 cells stimulated with

Figure 2. Effect of DHMEQ on apoptosis and necrosis of ARPE-19 cells. (A) Cells were cultured without TNF- α (non-treated) or with TNF- α (20 ng/ml) in the absence or presence of DHMEQ (1, 10 or 100 μ g/ml) for 24 h. The percentage of apoptotic cells (PI-negative and Annexin-V FITC-positive) and necrotic cells (PI-positive and Annexin-V FITC-positive) was measured by flow cytometry. (B) Percentages of apoptotic and necrotic cells. Data are presented as the mean $\pm$ SD, n=3. *P<0.05. PI, propidium iodide; DHMEQ, dehydroxymethylapoxyquinomicin; TNF- α , tumor necrosis factor- α ; NS, not significant.

TNF- α . It was revealed that DHMEQ (10 μ g/ml) significantly reduced the expression of ICAM-1 in cells by ~50% compared with cells treated with DMSO (Fig. 3C-E). Thus, it was speculated that DHMEQ may be able to decrease TNF- α -induced ICAM-1 expression in ARPE-19 cells.

Suppressive effect of DHMEQ on chemokine production in TNF- α -stimulated ARPE-19 cells. MCP-1 and IL-8 have

been revealed to be the major chemokines produced by TNF- α -stimulated ARPE-19 cells (25), and dexamethasone has been reported to have anti-inflammatory effects on ARPE-19 cells (20). Therefore, the present study investigated whether DHMEQ is able to decrease the protein expression levels of MCP-1 and IL-8 in ARPE-19 cells stimulated with TNF- α . Moreover, the anti-inflammatory effect of DHMEQ was compared with that of dexamethasone in cells. It was demonstrated that the

Figure 3. DHMEQ-mediated reduction of ICAM-1 expression in TNF- α -stimulated ARPE-19 cells. Cells were cultured (A) without TNF- α or (B) with TNF- α (20 ng/ml) in the presence of DMSO (0.1%) or (C) 1.0 or (D) 10 $\mu\text{g/ml}$ DHMEQ for 24 h, and PE-labeled ICAM-1-positive cells were measured by flow cytometry. (E) Percentages of PE-labeled ICAM-1-positive cells. Data are presented as the mean $\pm$ SD, $n=3$. * $P<0.05$. NS, not significant; PE, phycoerythrin; ICAM-1, intercellular adhesion molecule 1; DHMEQ, dehydroxymethyldeoxyquinomicin.

production of IL-8 and MCP-1 from cells treated with DHMEQ (40 μM =10 $\mu\text{g/ml}$) was significantly decreased compared with ARPE-19 cells treated with DMSO. In addition, there was a significant difference in the production of IL-8 and MCP-1 between cells treated with DHMEQ (40 μM) and those treated with dexamethasone (40 μM ; Fig. 4A and B). These findings revealed that DHMEQ has strong suppressive effects on the production of MCP-1 and IL-8 by ARPE-19 cells compared with dexamethasone. Furthermore, the results indicated that DHMEQ significantly reduced the protein expression levels of MCP-1 and IL-8 at 6, 12 and 24 h (Fig. 4C and D). Therefore, DHMEQ may be able to decrease the TNF- α -induced chemokine production in ARPE-19 cells at several time-points after co-culturing.

Suppression of NF- κ B-related inflammatory gene expression levels of ARPE-19 cells by DHMEQ. To determine the alterations of the expression levels of NF- κ B-associated inflammatory genes in ARPE-19 cells exposed to DHMEQ, the present study compared RNA isolated from TNF- α -stimulated cells in the absence or presence of DHMEQ, using the Human NF- κ B Pathway TaqMan[®] Array Plates that analyze 92 NF- κ B-associated inflammatory genes. Moreover, summaries of the differentially expressed genes between the two cell populations are presented in Table I. A total of 19 genes were revealed to be upregulated and 25 genes were revealed to be downregulated in cells exposed to DHMEQ compared with those in the absence of DHMEQ. The differentially expressed genes are presented in Tables I and II. The gene

expression levels of cytokines and chemokines, including MCP-1, ICAM-1, IL-6 and IL-8, and TLR2, TLR3 and TLR4, were downregulated in ARPE-19 cells treated with DHMEQ (Table I). In addition, DHMEQ suppressed TNF superfamily member 15 (TNFSF15) and TNF- α -induced protein 3 (TNFAIP3; Table I). However, it was revealed that DHMEQ increased the expression levels of numerous genes associated with the NF- κ B signaling pathway, including prostaglandin E synthase (PTGES), mitogen-activated protein kinase 14 (MAP3K14), LTBR and TNFRSF1A associated via death domain (TRADD).

In addition, representative genes, LTBR, MCP-1 and TLR4, which were identified by NF- κ B Pathway Array, were assessed by quantitative PCR analysis. Although the fold changes were not exactly the same between the two methods, the gene expression level of LTBR was significantly increased in the presence of DHMEQ compared to that in the presence of DMSO, whereas the gene expression levels of MCP-1 and TLR4 were significantly decreased in the presence of DHMEQ compared to that in the presence of DMSO (Fig. 5).

Discussion

The present results indicated that DHMEQ significantly decreased the protein expression of TNF- α -induced ICAM-1 in ARPE-19 cells, and also decreased the production of IL-8 and MCP-1 by cells stimulated with TNF- α . In addition, it was determined that DHMEQ at higher concentrations

Figure 4. Effect of DHMEQ and DEXA on the levels of MCP-1 and IL-8 in ARPE-19 cells stimulated by TNF- α . Cells were cultured with TNF- α (20 ng/ml) in the presence of DMSO (0.1%), DHMEQ (1 μ g/ml=4 μ M, 10 μ g/ml=40 μ M) or DEXA (40 μ M) for 24 h. The effect of DHMEQ on levels of (A) MCP-1 and (B) IL-8 in the culture supernatants of cells stimulated with TNF- α . ARPE-19 cells were cultured with TNF- α (20 ng/ml) in the presence of DMSO (0.1%) or DHMEQ (10 μ g/ml) for 6, 12 and 24 h. The effect of DHMEQ on levels of (C) MCP-1 and (D) IL-8 in the culture supernatants of cells stimulated with TNF- α , as measured by ELISA. Data are presented as the mean $\pm$ SD, n=3. *P<0.05. ND, not detected; DEXA, dexamethasone; NS, not significant; TNF- α , tumor necrosis factor- α ; MCP-1, monocyte chemoattractant protein-1; IL, interleukin; DHMEQ, dehydroxymethylepoxyquinomicin.

Figure 5. Validation of array data by quantitative PCR. mRNA expression levels of LTBR, MCP-1 and TLR4 of TNF- α -stimulated ARPE-19 cells treated with DMSO (0.1%) or DHMEQ (10 μ g/ml) for 24 h were determined by quantitative PCR. Duplicate assays were performed for each gene. Gene expression levels were normalized to β -actin as endogenous control. Fold changes in mRNA expression from qPCR are presented as the mean $\pm$ SDs, n=4. *P<0.05. DHMEQ, dehydroxymethylepoxyquinomicin; TNF- α , tumor necrosis factor- α ; TLR4, Toll-like receptor 4; LTBR, lymphotoxin β receptor.

Table I. Summary of downregulated genes in ARPE-19 cells stimulated with TNF- α in the presence of DHMEQ.

Target gene	Probe ID	Fold change
BIRC5	Hs00977611_g1	0.040
TLR2	Hs00152932_m1	0.053
TRAF5	Hs00182979_m1	0.082
TNFSF15	Hs00353710_s1	0.088
BCL10	Hs00184839_m1	0.127
CHUK	Hs00989507_m1	0.130
CSF2	Hs00171266_m1	0.164
BCL2	Hs00608023_m1	0.228
HPRT1	Hs99999909_m1	0.236
MCP-1	Hs00234140_m1	0.238
FADD	Hs00538709_m1	0.249
TLR4	Hs00152939_m1	0.274
MALT1	Hs00198984_m1	0.275
EDARADD	Hs00369830_m1	0.281
TNFAIP3	Hs00234713_m1	0.388
ICAM1	Hs00164932_m1	0.389
IRAK1BP1	Hs00418138_m1	0.396
CD83	Hs00188486_m1	0.414
TLR3	Hs00152933_m1	0.416
REL	Hs00968436_m1	0.437
CSF1	Hs00174164_m1	0.456
CXCL1	Hs00236937_m1	0.456
ZNF675	Hs00603247_m1	0.456
IL6	Hs00174131_m1	0.490
RIPK1	Hs00169407_m1	0.490

DHMEQ, dehydroxymethylepoxyquinomicin; TNF- α , tumor necrosis factor- α ; TLR, Toll-like receptor; TNFSF15 TNF superfamily member 15; TNFAIP3, TNF- α -induced protein 3.

had increased anti-inflammatory effects on ARPE-19 cells compared with dexamethasone. The results also indicated that exposure to DHMEQ decreased the expression level of ICAM-1 in ARPE-19 cells. Moreover, the present results are consistent with those from a previous study, which reported that DHMEQ decreases the expression level of ICAM-1 in the retina of diabetic mice and that it reduces the number of retinal-adherent leukocytes (26). Furthermore, the expression of ICAM-1 in RPE cells has been revealed to be elevated under inflammatory conditions, leading to the enhancement of leukocyte-RPE cell interactions (27,28). Previous studies have also identified increased ICAM-1 expression levels in ocular tissues of patients with uveitis and revealed that antibody-based blockage of ICAM-1 led to a suppression of experimental autoimmune uveoretinitis (29-31). Collectively, both the present findings and previous results indicated that DHMEQ may be a potential anti-inflammatory compound for RPE cells due to its ability to reduce the expression of ICAM-1.

Elnor *et al* (25) revealed that RPE cells produce several chemokines, including IL-8 and MCP-1. The present results revealed that DHMEQ inhibited the production of IL-8

Table II. Summary of upregulated genes in ARPE-19 cells stimulated with TNF- α in the presence of DHMEQ.

Target gene	Probe ID	Fold change
PTGES	Hs00610420_m1	18.850
MAP3K14	Hs00177695_m1	6.446
LTBR	Hs00158922_m1	5.488
TRADD	Hs00182558_m1	5.024
BCL3	Hs00180403_m1	4.300
NKIRAS2	Hs00383387_m1	3.855
TNFRSF10A	Hs00269492_m1	3.773
IKBKG	Hs00415849_m1	3.609
MAP3K7IP1	Hs00196143_m1	3.600
MYC	Hs00153408_m1	3.194
ZFP36	Hs00185658_m1	3.091
IRAK1	Hs00155570_m1	3.065
ENPP2	Hs00196470_m1	2.653
TRAF1	Hs00194638_m1	2.614
CARD10	Hs00367225_m1	2.574
NFKBIB	Hs00182115_m1	2.570
IRAK2	Hs00176394_m1	2.569
RELA	Hs00153294_m1	2.446
NFKB2	Hs00174517_m1	2.341

DHMEQ, dehydroxymethylepoxyquinomicin; TNF- α , tumor necrosis factor- α ; PTGES, prostaglandin E synthase; MAP3K14, mitogen-activated protein kinase 14; LTBR, lymphotoxin β receptor; TRADD, TNFRSF1A associated via death domain.

and MCP-1 in TNF- α -stimulated ARPE-19 cells, although the effect of treatment with IL-8 on ARPE-19 cells in the presence or absence of DHMEQ was not examined in the present study. It has been demonstrated that MCP-1, IL-8 and RANTES are elevated in the ocular tissues of experimental autoimmune uveitis, which suggests that these upregulated chemokines are potent chemoattractants in the pathogenesis of uveitis (32-34). Wakamatsu *et al* (35) revealed that DHMEQ had a positive therapeutic effect on established murine arthritis, and Iwata *et al* (36) revealed that DHMEQ was able to ameliorate experimental autoimmune uveoretinitis. Our previous study revealed that DHMEQ has anti-inflammatory effects on ocular inflammation induced by lipopolysaccharide via the inhibition of TNF- α and IL-6 expression levels in the aqueous humor, which indicated that DHMEQ may be a potential candidate to treat intraocular inflammatory diseases (37).

Local and systemic corticosteroids have been used to control ocular inflammation in patients with uveitis; however, long-term corticosteroid treatment can lead to adverse local and systemic side effects (38). The present results indicated that TNF- α -stimulated ARPE-19 cells were resistant to dexamethasone in relation to the protein expression levels of IL-8 and MCP-1, but DHMEQ significantly decreased the production of IL-8 and MCP-1 of TNF- α -stimulated cells treated with DHMEQ. Furthermore, previous studies have revealed that TNF signaling can suppress the action of the glucocorticoid

receptor by interfering with the transactivation function of glucocorticoid (GC) (39), which contributes to tissue resistance to GCs in several pathologic inflammatory states (39). These findings indicate the possibility that DHMEQ may have anti-inflammatory properties, even in inflammatory conditions with glucocorticoid resistance.

In the present study, NF- κ B-associated gene array analysis identified that the gene expression levels of cytokines and chemokines, including MCP-1, ICAM-1, IL-6, TNFSF15 and TNFAIP3, and TLR2, TLR3 and TLR4 were downregulated in ARPE-19 cells treated with DHMEQ, which was further demonstrated by quantitative PCR analysis. Moreover, it was revealed that DHMEQ increased the expression levels of several genes related to the NF- κ B signaling pathway, including PTGES, MAP3K14, LTBR and TRADD. However, the present study did not examine the protein expression levels of the gene products either upregulated or downregulated in ARPE-19 cells by DHMEQ. In addition, the translocation of p65-NF- κ B into the nucleus in the presence of DHMEQ in TNF- α -stimulated cells was not investigated. Therefore, future studies are required to assess post-transcriptional regulation by DHMEQ with western blotting and to examine the translocation of p65-NF- κ B in the presence of DHMEQ with electrophoretic mobility shift assay.

The present results demonstrated that 50 and 100 μ g/ml DHMEQ had severe cytotoxic effects on cultured ARPE-19 cells, and that high concentrations of DHMEQ (100 μ g/ml) induced apoptosis and necrosis in TNF- α -stimulated cells. NF- κ B is known to play important roles in protecting cells from apoptosis (40,41). Furthermore, previous studies have revealed that DHMEQ is able to induce apoptosis of cancer cells (42,43). Although the present study used an RPE cell line, it remains to be determined whether high concentrations of DHMEQ have cytotoxic or apoptotic effects on primary cultured RPE cells, and healthy and inflamed RPE cells *in vivo*. Thus, the relationship between the anti-inflammatory potential of DHMEQ and the induction of apoptosis by DHMEQ in healthy or inflamed RPE cells requires further examination.

In conclusion, the present results indicated that DHMEQ may have an anti-inflammatory effect on TNF- α -stimulated ARPE-19 cells. However, it is not fully understood whether DHMEQ has a suppressive effect on the expression of ICAM-1 and chemokine production in primary cultured human RPE cells and *in vivo* RPE monolayers, thus further studies are required to assess the anti-inflammatory effects and safety of DHMEQ on human RPE cells.

Acknowledgements

The authors would like to thank Ms Mirai Kano (Department of Ophthalmology, Kyorin University, School of Medicine) for technical assistance and Professor Emeritus Duco Hamasaki (Bascom Palmer Eye Institute, University of Miami, Miami, Florida, USA) for editing the manuscript.

Funding

The present study was supported by Grant-in-Aid for Scientific Research (grant. no. 15K10901) from the Ministry

of Education, Culture, Sports, Science and Technology, Japan and Research Grant from Kyorin University, Tokyo, Japan.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YA, HK and YS performed the experiments. YA and HK designed the experiments. AK contributed to the design of the methodology. YA, HK, TW, AH and AAO analyzed the results. YA and HK wrote the paper. KU prepared DHMEQ and analyzed the results. All authors read and approved the manuscript and agree to be accountable for all aspects of the research in ensuring that the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Holtkamp GM, Kijlstra A, Peek R and de Vos AF: Retinal pigment epithelium-immune system interactions: Cytokine production and cytokine-induced changes. *Prog Retin Eye Res* 20: 29-48, 2001.
- Momma Y, Nagineni CN, Chin MS, Srinivasan K, Detrick B and Hooks JJ: Differential expression of chemokines by human retinal pigment epithelial cells infected with cytomegalovirus. *Invest Ophthalmol Vis Sci* 44: 2026-2033, 2003.
- Elner SG, Delmonte D, Bian ZM, Lukacs NW and Elner VM: Differential expression of retinal pigment epithelium (RPE) IP-10 and interleukin-8. *Exp Eye Res* 83: 374-379, 2006.
- Dick AD: Doyne lecture 2016: Intraocular health and the many faces of inflammation. *Eye (Lond)* 31: 87-96, 2017.
- Sugita S, Kawazoe Y, Imai A, Yamada Y, Horie S and Mochizuki M: Inhibition of Th17 differentiation by anti-TNF-alpha therapy in uveitis patients with Behcet's disease. *Arthritis Res Ther* 14: R99, 2012.
- Okada AA, Goto H, Ohno S and Mochizuki M; Ocular Behcet's Disease Research Group Of Japan: Multicenter study of infliximab for refractory uveoretinitis in Behcet disease. *Arch Ophthalmol* 130: 592-598, 2012.
- Takeuchi M, Kezuka T, Sugita S, Keino H, Namba K, Kaburaki T, Maruyama K, Nakai K, Hijioka K, Shibuya E, *et al*: Evaluation of the long-term efficacy and safety of infliximab treatment for uveitis in Behcet's disease: A multicenter study. *Ophthalmology* 121: 1877-1884, 2014.
- Jaffe GJ, Dick AD, Brezin AP, Nguyen QD, Thorne JE, Kestelyn P, Barisani-Asenbauer T, Franco P, Heiligenhaus A, Scales D, *et al*: Adalimumab in patients with active noninfectious uveitis. *N Engl J Med* 375: 932-943, 2016.
- Brenner D, Blaser H and Mak TW: Regulation of tumour necrosis factor signalling: Live or let die. *Nat Rev Immunol* 15: 362-374, 2015.

10. Hayden MS and Ghosh S: NF- κ B in immunobiology. *Cell Res* 21: 223-244, 2011.
11. Okamoto H, Cujec TP, Yamanaka H and Kamatani N: Molecular aspects of rheumatoid arthritis: Role of transcription factors. *FEBS J* 275: 4463-4470, 2008.
12. Kawai T and Akira S: Signaling to NF- κ B by Toll-like receptors. *Trends Mol Med* 13: 460-469, 2007.
13. Wu J, Ding J, Yang J, Guo X and Zheng Y: MicroRNA roles in the nuclear factor kappa B signaling pathway in cancer. *Front Immunol* 9: 546, 2018.
14. Ariga A, Namekawa J, Matsumoto N, Inoue J and Umezawa K: Inhibition of tumor necrosis factor- α -induced nuclear translocation and activation of NF- κ B by dehydroxymethylepoxyquinomicin. *J Biol Chem* 277: 24625-24630, 2002.
15. Umezawa K: Inhibition of tumor growth by NF- κ B inhibitors. *Cancer Sci* 97: 990-995, 2006.
16. Umezawa K and Chaichareonpong C: Molecular design and biological activities of NF- κ B inhibitors. *Mol Cells* 14: 163-167, 2002.
17. Yamamoto M, Horie R, Takeiri M, Kozawa I and Umezawa K: Inactivation of NF- κ B components by covalent binding of (-)-dehydroxymethylepoxyquinomicin to specific cysteine residues. *J Med Chem* 51: 5780-5788, 2008.
18. Lin Y, Ukaji T, Koide N and Umezawa K: Inhibition of late and early phases of cancer metastasis by the NF- κ B inhibitor DHMEQ derived from microbial bioactive metabolite epoxyquinomicin: A review. *Int J Mol Sci* 19: E729, 2018.
19. Juel HB, Faber C, Udsen MS, Folkersen L and Nissen MH: Chemokine expression in retinal pigment epithelial ARPE-19 cells in response to coculture with activated T cells. *Invest Ophthalmol Vis Sci* 53: 8472-8480, 2012.
20. Zanon Cde F, Sonehara NM, Girol AP, Gil CD and Oliani SM: Protective effects of the galectin-1 protein on in vivo and in vitro models of ocular inflammation. *Mol Vis* 21: 1036-1050, 2015.
21. Yang PM, Wu ZZ, Zhang YQ and Wung BS: Lycopene inhibits ICAM-1 expression and NF- κ B activation by Nrf2-regulated cell redox state in human retinal pigment epithelial cells. *Life Sci* 155: 94-101, 2016.
22. Mestdagh P, Van Vlierberghe P, De Weer A, Muth D, Westermann F, Speleman F and Vandesompele J: A novel and universal method for microRNA RT-qPCR data normalization. *Genome Biol* 10: R64, 2009.
23. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
24. Chen YH, Chen CL, Liang CM, Liang JB, Tai MC, Chang YH, Lu DW and Chen JT: Silibinin inhibits ICAM-1 expression via regulation of N-linked and O-linked glycosylation in ARPE-19 cells. *Biomed Res Int* 2014: 701395, 2014.
25. Elner VM, Burnstine MA, Strieter RM, Kunkel SL and Elner SG: Cell-associated human retinal pigment epithelium interleukin-8 and monocyte chemotactic protein-1: Immunochemical and in-situ hybridization analyses. *Exp Eye Res* 65: 781-789, 1997.
26. Nagai N, Izumi-Nagai K, Oike Y, Koto T, Satofuka S, Ozawa Y, Yamashiro K, Inoue M, Tsubota K, Umezawa K and Ishida S: Suppression of diabetes-induced retinal inflammation by blocking the angiotensin II type 1 receptor or its downstream nuclear factor- κ B pathway. *Invest Ophthalmol Vis Sci* 48: 4342-4350, 2007.
27. Chen JT, Liang JB, Chou CL, Chien MW, Shyu RC, Chou PI and Lu DW: Glucosamine sulfate inhibits TNF- α and IFN- γ -induced production of ICAM-1 in human retinal pigment epithelial cells in vitro. *Invest Ophthalmol Vis Sci* 47: 664-672, 2006.
28. Chen JT, Chen PL, Chang YH, Chien MW, Chen YH and Lu DW: Glucosamine sulfate inhibits leukocyte adhesion in response to cytokine stimulation of retinal pigment epithelial cells in vitro. *Exp Eye Res* 83: 1052-1062, 2006.
29. Whitcup SM, Chan CC, Li Q and Nussenblatt RB: Expression of cell adhesion molecules in posterior uveitis. *Arch Ophthalmol* 110: 662-666, 1992.
30. Whitcup SM, DeBarge LR, Caspi RR, Harning R, Nussenblatt RB and Chan CC: Monoclonal antibodies against ICAM-1 (CD54) and LFA-1 (CD11a/CD18) inhibit experimental autoimmune uveitis. *Clin Immunol Immunopathol* 67: 143-150, 1993.
31. Uchio E, Kijima M, Tanaka S and Ohno S: Suppression of experimental uveitis with monoclonal antibodies to ICAM-1 and LFA-1. *Invest Ophthalmol Vis Sci* 35: 2626-2631, 1994.
32. Crane JJ, McKillop-Smith S, Wallace CA, Lamont GR and Forrester JV: Expression of the chemokines MIP-1 α , MCP-1, and RANTES in experimental autoimmune uveitis. *Invest Ophthalmol Vis Sci* 42: 1547-1552, 2001.
33. Foxman EF, Zhang M, Hurst SD, Muchamuel T, Shen D, Wawrousek EF, Chan CC and Gery I: Inflammatory mediators in uveitis: Differential induction of cytokines and chemokines in Th1- versus Th2-mediated ocular inflammation. *J Immunol* 168: 2483-2492, 2002.
34. Keino H, Takeuchi M, Kezuka T, Yamakawa N, Tsukahara R and Usui M: Chemokine and chemokine receptor expression during experimental autoimmune uveoretinitis in mice. *Graefes Arch Clin Exp Ophthalmol* 241: 111-115, 2003.
35. Wakamatsu K, Nanki T, Miyasaka N, Umezawa K and Kubota T: Effect of a small molecule inhibitor of nuclear factor- κ B nuclear translocation in a murine model of arthritis and cultured human synovial cells. *Arthritis Res Ther* 7: R1348-R1359, 2005.
36. Iwata D, Kitaichi N, Miyazaki A, Iwabuchi K, Yoshida K, Namba K, Ozaki M, Ohno S, Umezawa K, Yamashita K, *et al*: Amelioration of experimental autoimmune uveoretinitis with nuclear factor- κ B inhibitor dehydroxymethyl epoxyquinomicin in mice. *Invest Ophthalmol Vis Sci* 51: 2077-2084, 2010.
37. Ando Y, Keino H, Kudo A, Hirakata A, Okada AA and Umezawa K: Anti-inflammatory effect of dehydroxymethylepoxyquinomicin, a nuclear factor- κ B inhibitor, on endotoxin-induced uveitis in rats in vivo and in vitro. *Ocul Immunol Inflamm* 28: 240-248, 2020.
38. Gaudio PA: A review of evidence guiding the use of corticosteroids in the treatment of intraocular inflammation. *Ocul Immunol Inflamm* 12: 169-192, 2004.
39. Van Bogaert T, De Bosscher K and Libert C: Crosstalk between TNF and glucocorticoid receptor signaling pathways. *Cytokine Growth Factor Rev* 21: 275-286, 2010.
40. Ukaji T and Umezawa K: Novel approaches to target NF- κ B and other signaling pathways in cancer stem cells. *Adv Biol Regul* 56: 108-115, 2014.
41. de Castro Barbosa ML, da Conceicao RA, Fraga AGM, Camarinha BD, de Carvalho Silva GC, Lima AGF, Cardoso EA and de Oliveira Freitas Lione V: NF- κ B signaling pathway inhibitors as anticancer drug candidates. *Anticancer Agents Med Chem* 17: 483-490, 2017.
42. Miyake A, Dewan MZ, Ishida T, Watanabe M, Honda M, Sata T, Yamamoto N, Umezawa K, Watanabe T and Horie R: Induction of apoptosis in Epstein-Barr virus-infected B-lymphocytes by the NF- κ B inhibitor DHMEQ. *Microbes Infect* 10: 748-756, 2008.
43. Fukushima T, Kawaguchi M, Yorita K, Tanaka H, Takeshima H, Umezawa K and Kataoka H: Antitumor effect of dehydroxymethylepoxyquinomicin, a small molecule inhibitor of nuclear factor- κ B, on glioblastoma. *Neuro Oncol* 14: 19-28, 2012.