

# ***Panax notoginseng* Saponins protect auditory cells against cisplatin-induced ototoxicity by inducing the AKT/Nrf2 signaling-mediated redox pathway**

BING FEI<sup>1</sup>, ZHIBIAO LIU<sup>1</sup>, LISHENG XIE<sup>1</sup>, LINGYUN LV<sup>1</sup>, WENYAN ZHU<sup>1</sup>,  
JING LIU<sup>2,3</sup>, YANHONG DAI<sup>2,3</sup> and WANDONG SHE<sup>1,2</sup>

<sup>1</sup>Department of Otolaryngology-Head and Neck Surgery, Nanjing Drum Tower Hospital Clinical College of Nanjing Medical University; <sup>2</sup>Department of Otolaryngology-Head and Neck Surgery, Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School, Jiangsu Provincial Key Medical Discipline; <sup>3</sup>Hearing Center, Otorhinolaryngology Research Institute of Nanjing Drum Tower Hospital, Nanjing, Jiangsu 210008, P.R. China

Received October 16, 2019; Accepted June 15, 2020

DOI: 10.3892/mmr.2020.11390

**Abstract.** Cisplatin-induced cytotoxicity, such as nephrotoxicity, neurotoxicity and ototoxicity, restricts the clinical application of this compound. *Panax notoginseng* Saponins (PNS) exhibit potent free radical scavenging and antioxidant activity. PNS have been demonstrated to reduce cisplatin-induced nephrotoxicity and neurotoxicity. The present study investigated the ability of PNS to protect the auditory HEI-OC1 cell line against ototoxicity induced by cisplatin. PNS induced activation of the AKT/nuclear factor erythroid 2-related factor 2 (Nrf2) signaling pathway. Following pretreatment with PNS, HEI-OC1 cells were treated with cisplatin and cultured for 24 h. The viability of HEI-OC1 cells was examined using a Cell Counting Kit-8 assay. Double staining analysis was used to measure cell apoptosis. The ability of PNS to reduce reactive oxygen species (ROS) levels was assessed by flow cytometry. The levels of phosphorylated (p)-AKT, heme oxygenase 1 (HO-1), NAD(P)H quinone dehydrogenase 1 (NQO1), glutamate-cysteine ligase catalytic (GCLC) and Nrf2 were measured by western blotting. HEI-OC1 cells that were pretreated with PNS exhibited significantly increased cell viability compared with that noted in cells treated only with cisplatin. In addition, PNS suppressed the induction of apoptosis and ROS production following cisplatin treatment. The upregulation of NQO1, HO-1 and GCLC expression in PNS-pretreated cells was associated with p-AKT

levels and the activation of Nrf2. These findings suggested that PNS protected auditory cells against ototoxicity induced by cisplatin by activating AKT/Nrf2 signaling. PNS may serve as a potential candidate in regulating cisplatin-induced cytotoxicity.

## **Introduction**

Cisplatin is a typical chemotherapeutic drug that is widely used in the treatment of a variety of solid tumors (1,2). However, the dose-limiting side effects of high concentrations of cisplatin, such as renal damage, neurotoxicity and ototoxicity can restrict its clinical application (3-5). Among patients that receive cisplatin chemotherapy, >90% may end up with ear-associated adverse effects (3,6). However, currently no effective treatment exists to avoid or reduce ototoxicity induced by cisplatin (5,7-9). Therefore, the prevention of ototoxicity induced by cisplatin has become a major focus in tumor therapy.

To the best of our knowledge, the mechanism of the aforementioned process has not yet been elucidated. However, additional evidence has demonstrated that overproduction of reactive oxygen species (ROS) serves a key role in this process (10,11). High accumulation of cisplatin in cochlear tissues may reduce the expression levels of nuclear factor erythroid 2-related factor 2 (Nrf2), a well-known transcription factor responsible for the antioxidant-pro-oxidant balance (12,13). This is caused by modulation of the expression levels of various antioxidant enzymes, including heme oxygenase 1 (HO-1), NAD(P)H quinone dehydrogenase 1 (NQO1) and glutamate-cysteine ligase catalytic (GCLC) (14-16). This pathological process disrupts the balance between ROS production, and antioxidant enzyme activity and expression, leading to auditory cell injury and apoptosis (17). Therefore, upregulation of Nrf2 expression and the increase in antioxidant enzyme expression has become an important target for treating cisplatin-induced ototoxicity.

*Panax notoginseng* Saponins (PNS) are compounds that are extracted from *Panax notoginseng* (Sanqi).

**Correspondence to:** Dr Wandong She, Department of Otolaryngology-Head and Neck Surgery, Nanjing Drum Tower Hospital Clinical College of Nanjing Medical University, 321 Zhongshan Road, Nanjing, Jiangsu 210008, P.R. China  
E-mail: shewandong@163.com

**Key words:** cisplatin, ototoxicity, oxidative stress, nuclear factor erythroid 2-related factor 2, apoptosis

*Panax notoginseng* is routinely used to treat acute cerebral infarction and acute myocardial infarction (18,19). A number of previous reports have demonstrated that PNS can inhibit cisplatin-induced nephrotoxicity and neurotoxicity (20-23). However, whether PNS can inhibit cisplatin-induced ototoxicity remains unknown. A previous study indicated that PNS could regulate Nrf2 antioxidant signaling via the AKT signaling pathway (24). Therefore, the present study examined whether PNS could protect against ototoxicity induced by cisplatin via activation of the AKT/Nrf2 signaling pathway. HEI-OC1 cells were incubated with cisplatin with or without pretreatment of PNS and subsequently cell viability and apoptotic rate were determined. Additionally, the associated signaling mechanisms were examined.

## Materials and methods

**Chemicals and materials.** PNS was obtained from Chengdu Manst Biotechnology Co., Ltd. (www.cdmust.com). and dissolved in DMSO (cat. no. A1222; purity  $\geq 98\%$ ). House Ear Institute-Organ of Corti 1 (HEI-OC1) cell lines derived from the auditory organ of a transgenic mouse were obtained from House Ear Institute. The antibodies for active caspase 3, Nrf2, NQO1 and Bcl-2 were obtained from Abcam. The antibodies for HO-1, GCLC, AKT, and phosphorylated-AKT (p-AKT) were obtained from Abcam. The Hoechst 33258 staining kit was obtained from Biyuntian Biotechnology Co., Ltd. Cisplatin and the propidium iodide (PI) staining kit were obtained from Sigma-Aldrich; Merck KGaA. The oxygen species assay kit was purchased from Nanjing KeyGen Biotech Co., Ltd. The p-AKT inhibitor LY294002 was purchased from Selleck Chemicals. Cells were treated with 25  $\mu\text{mol/l}$  LY294002 for 24 h at room temperature before transfection. The culture media were obtained from Gibco; Thermo Fisher Scientific, Inc.

**Cell culture and cell viability assay.** The HEI-OC1 cell line is a frequently used model of ototoxicity assessment and expresses several molecular characteristics of Corti sensory cells (25,26). This cell line was selected for use in the present study. The cells were cultivated in complete medium containing high-glucose DMEM (Beijing Solarbio Science & Technology Co., Ltd.) supplemented with 10% FBS (Beijing Solarbio Science & Technology Co., Ltd.) and 50 U/ml interferon- $\gamma$  without antibiotics. Cells were cultured at 33°C in the presence of 10% CO<sub>2</sub> in a humidified incubator (permissive conditions). Cell toxicity was assessed as follows: A total of 5x10<sup>3</sup> HEI-OC1 cells/well in 100  $\mu\text{l}$  complete medium were seeded in a 96-well plate and cultured at 33°C in the presence of 10% CO<sub>2</sub>. The cells were pretreated with different concentrations of PNS (0, 25, 50, 100, 200, 400 and 800  $\mu\text{g/ml}$ ) for 2 h at room temperature. Subsequently, 20  $\mu\text{M}$  cisplatin was added to each well of the plate. Following a 24-h cultivation, 10  $\mu\text{l}$  Cell Counting Kit-8 (CCK-8; cat. no. CK-4; Dojindo Molecular Technologies, Inc.) solution was added to each well and incubated for an additional 30 min, according to the manufacturer's protocol. A microplate reader (Lab Systems Multiskan Ascent 354; MTX Lab Systems, Inc.) was used to detect cell viability at 450 nm. Finally, the appropriate doses of PNS were selected for further *in vitro* studies depending on the initial toxicity data.

**Hoechst 33258 and PI double staining.** A total of 1x10<sup>5</sup> HEI-OC1 cells were diluted in 4 ml complete medium and incubated in 6-well plates with 100  $\mu\text{g/ml}$  PNS for 2 h prior to treatment with cisplatin (20  $\mu\text{M}$ ) at 33°C in a humidified 10% CO<sub>2</sub> environment for 24 h. Following two washes with PBS, the cells were fixed with 4% paraformaldehyde for 30 min at 4°C, and stained with 2  $\mu\text{g/ml}$  Hoechst 33258 and 1  $\mu\text{g/ml}$  PI for 20 min in the dark at 4°C. Finally, following two washes with PBS, the cells were visualized using a Leica DMi8 fluorescence microscope (magnification, x200; Leica Microsystems, Inc.). The highest emission was recorded at  $\sim 460$  nm for PI staining and at 485 nm for Hoechst dyes. The amount of dead cells was quantified using ImageJ software version 1.8.0 (National Institutes of Health) and calculated as follows: Amount of Hoechst-positive cells/amount of PI-positive cells.

**TUNEL staining.** To assess the DNA fragmentation using TUNEL, cells were fixed with 1% formaldehyde for 10 min at room temperature. Following incubated with 100  $\mu\text{g/ml}$  PNS for 2 h prior to treatment with cisplatin (20  $\mu\text{M}$ ) at 33°C in a humidified 10% CO<sub>2</sub> environment for 24 h, a total of 1x10<sup>5</sup> HEI-OC1 cells were incubated with 20  $\mu\text{g/ml}$  proteinase K for 60 min at room temperature. The slides were then rinsed with PBS for 3 min, dried and incubated in 20  $\mu\text{l}$  TUNEL reaction mixture for 1 h at room temperature. Subsequently, cells were treated with 2% H<sub>2</sub>O<sub>2</sub> for 30 min at room temperature. After washing with PBS, sections were incubated with 150 U/ml anti-digoxigenin peroxidase conjugate for another 30 min at room temperature in a humidified atmosphere. Finally, sections were stained with 10  $\mu\text{g/ml}$  diaminobenzene in the dark for 10 min at room temperature and counterstained with 20  $\mu\text{g/ml}$  hematoxylin at room temperature for 10 min. Cell nuclei were stained with 0.1  $\mu\text{g/ml}$  DAPI at 37°C for another 10 min. After being washed with PBS for 3 min, 1% ammonia (cat. no. SJ00815F17008; Sigma-Aldrich; Merck KGaA), and rehydrated using different concentration of alcohol, the sections were sealed and detected with a fluorescent microscope (magnification, x200) with VECTASHIELD mounting media (Vector Laboratories, inc.) for fluorescence microscope. TUNEL positive (apoptotic) cells were quantified by counting amber-colored cells in 10 fields (6,000  $\mu\text{m}^2/\text{field}$ ). The specimens were visualized using a Leica confocal laser scanning microscope and processed with Photoshop CS6 (Adobe Systems, Inc.).

**Determination of intracellular ROS.** A total of 1x10<sup>5</sup> HEI-OC1 cells were diluted in 4 ml complete medium and cultured in 6-well plates. The cells were pre-incubated with 100  $\mu\text{g/ml}$  PNS for 2 h prior to incubation with cisplatin (20  $\mu\text{M}$ ) at 33°C in a humidified 10% CO<sub>2</sub> environment for 24 h. Subsequently, the cells were washed with cold PBS twice and incubated with 2',7'-dichlorofluorescein diacetate (DCFH-DA; Invitrogen; Thermo Fisher Scientific, Inc.) for 20 min at room temperature. The cells were harvested into 1.5 ml tubes using trypsin-EDTA and washed with PBS twice.

**Flow cytometry analysis.** The quantification of cell death was determined via flow cytometry using the Annexin V-FITC apoptosis detection kit according to the manufacturer's protocol (BD Pharmingen; BD Bioscience) on a BD




Figure 1. PNS prevents CP-induced cytotoxicity. (A) HEI-OC1 cell viability was determined following administration of various concentrations of PNS with or without 20  $\mu$ M CP to the cells. (B) HEI-OC1 cell viability following treatment with 100  $\mu$ g/ml PNS and 20  $\mu$ M CP (n=3 per group). All data are presented as the mean  $\pm$  SEM. \*\*P<0.01 vs. CP group. CP, cisplatin; PNS, *Panax notoginseng* Saponins.

FACSCalibur flow cytometer (BD Biosciences) at an excitation and emission wavelength of 488 and 525 nm, respectively. Briefly, (10,000 events per sample) DCFH-DA-treated cells were seeded into each Petri dish (size, 30 mm) and after a 24 h incubation, various concentrations of the test compound (20  $\mu$ M CP, 100  $\mu$ g/ml PNS or 20  $\mu$ M CP + 100  $\mu$ g/ml PNS) were added and incubated for 24, 48 and 72 h at room temperature. The cells were washed with PBS, suspended in Annexin V binding buffer and then added to an Annexin V-FITC solution and propidium iodide (PI) for 10 min at room temperature. The percentage of apoptotic cells was calculated using CellQuest software version 5.1 (Becton, Dickinson and Company). Data are presented as the percentage of Annexin V-stained cells, and all experiments were performed in triplicate. Morphological changes were imaged under a phase contrast microscope (magnification, x200; Olympus Corporation).

**Western blot analysis.** Total protein was extracted from cells using the Cell Total Protein Extraction kit (Sigma-Aldrich; Merck KGaA). The concentration of the proteins was measured using the bicinchoninic acid method. The 20  $\mu$ g/lane protein samples were separated by 10% SDS-PAGE and analyzed by electrophoresis. The proteins were transferred to PVDF membranes and blocked in 3% BSA (Thermo Fisher Scientific, Inc.) for 1 h at room temperature. Subsequently, the membranes were incubated at 4°C overnight with the following primary antibodies: Active caspase-3 (dilution, 1:2,000; rabbit; cat. no. ab214430; Abcam), Bcl-2 (dilution, 1:1,000; rabbit; cat. no. ab182858; Abcam), GCLC (dilution, 1:2,000; rabbit; cat. no. ab190685; Abcam), NQO1 (dilution, 1:2,000; rabbit; cat. no. ab80588; Abcam), HO-1 (dilution, 1:10,000; rabbit; cat. no. ab68477; Abcam), Nrf2 (dilution, 1:2,000; rabbit; cat. no. ab62352; Abcam), AKT (dilution, 1:2,000; rabbit; cat. no. ab179463; Abcam), p-AKT (dilution, 1:2,000; rabbit; cat. no. ab192623; Abcam) or GAPDH (dilution, 1:3,000; rabbit; cat. no. ab181602; Abcam). The next morning, the membranes were incubated with secondary antibodies conjugated to horseradish peroxidase IgG (dilution, 1:2,000; cat. no. ab181658; Abcam) for 1 h. Finally, the proteins were visualized using the WesternBright™ ECL kit (cat. no. E-IR-R301; Elabscience, Inc.) and analyzed using ImageJ software version 1.8.0 (National Institutes of Health).

**Statistical analysis.** SPSS v22.0 software (IBM Corp.) was used for the statistical analysis and GraphPad Prism 6 software (GraphPad Software, Inc.) was used for the generation of statistical charts. All data are presented as the mean  $\pm$  standard error of the mean. All the experiments in this study were conducted in triplicate. Statistical analysis was performed by one-way ANOVA followed by Tukey's post hoc test. P<0.05 was considered to indicate a statistically significant difference.

## Results

**PNS prevents cisplatin-induced cytotoxicity.** As shown in Fig. 1A, at concentrations up to 100  $\mu$ g/ml PNS, the cell viability of HEI-OC1 was not significantly affected. However, there was a significant variation between 100 and 200  $\mu$ g/ml PNS. The results of the cell viability assay indicated that in the presence or absence of cisplatin, the viability of HEI-OC1 cells was not affected by pretreatment with PNS concentrations <100  $\mu$ g/ml. Therefore, 100  $\mu$ g/ml PNS could be used in the subsequent experiments in order to avoid toxicity. Furthermore, after treated with 20  $\mu$ M CP, it was found that 100  $\mu$ g/ml PNS was able to prevent cisplatin-induced cytotoxicity in HEI-OC1 cells (Fig. 1B;  $0.42 \pm 0.11$  vs.  $0.71 \pm 0.19$ ; P<0.01). Therefore, this concentration (100  $\mu$ g/ml PNS and 20  $\mu$ M/l CP) was selected for use in subsequent experiments.

**PNS reduces cisplatin-mediated apoptosis in HEI-OC1 cells.** The ImageJ software version 1.8.0 (National Institutes of Health) was used to measure the rate of cell apoptosis. The induction of HEI-OC1 cell apoptosis was not evident in the control group. Following exposure to cisplatin, the number of HEI-OC1 cells that had undergone apoptosis was significantly increased compared with control or PNS group (Fig. 2A-D). This effect was significantly relieved following pretreatment of the cells with PNS (Fig. 2A-D). To further verify the anti-apoptotic effect of PNS, western blotting was used to assess the expression levels of apoptosis-associated proteins and the results indicated that HEI-OC1 cells treated with cisplatin exhibited increased expression levels of active caspase-3 and reduced Bcl-2 expression levels compared with the corresponding levels noted in the control group (Fig. 2E).




Figure 2. PNS attenuates CP-mediated apoptosis in HEI-OC1 cells. (A) Hoechst 33258 staining and PI staining of different groups. (B) TUNEL staining results in different groups. (C) Percentage of Hoechst 33258 positive staining and PI positive staining (n=3 per group). \*\*P<0.01, CP vs. control; PNS + CP vs. CP. (D) Quantified results of TUNEL staining in different groups. \*\*P<0.01, PNS + CP vs. CP; PNS vs. CP. (E) Western blot analysis of the protein levels of active caspase-3 and Bcl-2. GAPDH was used as an internal control for equal loading. All data are presented as the mean  $\pm$  SEM. CP, cisplatin; PNS, *Panax notoginseng* Saponins; PI, propidium iodide.

*PNS attenuates cisplatin-mediated ROS production and increases antioxidant enzyme expression.* To further

investigate the mechanism by which PNS attenuated apoptosis induction by cisplatin in HEI-OC1 cells, changes






Figure 3. PNS attenuates cisplatin-mediated ROS production and increases antioxidant enzyme expression levels. (A) Flow cytometry results and statistical analysis of the data from the different groups. (B) Western blot analysis of the protein expression levels of GCLC, NQO1 and HO-1. GAPDH was used as an internal control for equal loading. All data are presented as the mean  $\pm$  SEM. \*\* $P < 0.01$  vs. CP group. CP, cisplatin; PNS, *Panax notoginseng* Saponins; ROS, reactive oxygen species; GCLC, glutamate-cysteine ligase catalytic; NQO1, NAD(P)H quinone dehydrogenase 1; HO-1, heme oxygenase 1.

in the production of ROS and in the expression levels of antioxidant enzymes were examined. The ROS levels in the CP group was higher compared with the PNS + CP, PNS or control groups (Fig. 3). This suggested that PNS could significantly reduce the increase in ROS levels caused by cisplatin. Western blot analysis indicated that cisplatin could reduce the expression levels of GCLC, NQO1 and HO-1 (Fig. 3B). These results indicated that PNS could alleviate cisplatin-induced cell damage and apoptosis by restoring the balance between oxidative stress and the expression levels of antioxidant enzymes.

**PNS increases Nrf2 expression by upregulating AKT phosphorylation in HEI-OC1 cells.** A recent study has demonstrated that the AKT-Nrf2 axis regulates the expression of antioxidant enzymes (27). Initially, the effects of PNS pretreatment on Nrf2 expression were examined. Following cisplatin administration, Nrf2 expression in HEI-OC1 cells has no significant variation at 6 h, whereas it was markedly decreased at 12 and 24 h (Fig. 4A and C). However, HEI-OC1 cells pretreated with PNS expressed lower levels of Nrf2 at 6 and 12 h compared with the cisplatin group. In addition, the phosphorylation levels of AKT were examined after 6 h of PNS treatment of cells (Fig. 4B and D). PNS treatment increased AKT phosphorylation levels induced by cisplatin at

6 h. These results indicated that the protective effect of PNS on cisplatin-induced cytotoxicity may be achieved via the AKT/Nrf2 axis.

**Protective effects of PNS on HEI-OC1 cells are blocked by AKT inhibition.** To demonstrate that the inhibition of cisplatin-induced cytotoxicity by PNS was mediated via the AKT/Nrf2 axis, the AKT inhibitor LY294002 was used. Treatment of the cells with MK2002 blocked the protective effect of PNS on cisplatin-induced cytotoxicity ( $0.43 \pm 0.16$  vs.  $0.76 \pm 0.17$ ,  $P < 0.01$ ; Fig. 5).

## Discussion

Cisplatin is a complex of heavy metal platinum and was officially approved for clinical chemotherapy in the United States in 1972 (28,29). As one of the most widely used broad-spectrum antitumor drugs in clinical settings, cisplatin exerts potent antitumor activity and is widely used to treat malignant tumors of the head and neck (1,30,31). However, cisplatin causes ototoxicity that mainly manifests as deafness (2,3,32). This adverse effect limits its clinical application (2,3,32). Deafness is usually the most apparent cause of bilateral hearing impairment (33). As the dose of cisplatin increases, hearing loss can gradually develop from the low frequency




Figure 4. PNS increases Nrf2 expression by upregulating AKT phosphorylation in HEI-OC1 cells. (A) Western blot analysis of the protein expression levels of Nrf2 at different time points. (B) Western blot analysis of the protein levels of AKT and p-AKT at 6 h. (C) Quantified results of A are presented. (D) Quantified results of B are presented. GAPDH was used as an internal control for equal loading. \* $P < 0.05$  vs. control. \*\* $P < 0.01$ , PNS + CP vs. CP; CP vs. control. PNS, *Panax notoginseng* Saponins; Nrf2, nuclear factor erythroid 2-related factor 2; p-, phosphorylated.




Figure 5. Protective effect of PNS on HEI-OC1 cells is blocked by AKT inhibitor treatment. HEI-OC1 cell viability was determined following treatment of the cells with 100 mg/ml PNS, 20  $\mu$ M CP and 20 nM MK. All the data are presented as the mean  $\pm$  SEM. \*\* $P < 0.01$ . PNS, *Panax notoginseng* Saponins; CP, cisplatin; MK, MK202.

region and affect the language frequency, eventually leading to permanent hearing loss (31). Accumulating evidence has demonstrated that cisplatin primarily destroys hearing by inducing auditory cell apoptosis (31-33). In the present study, the auditory HEI-OC1 cell line was used. Following culture with cisplatin, the viability of HEI-OC1 cells was reduced and the apoptotic rate was increased, which was consistent with observations from other studies (34,35).

PNS is an active ingredient extracted from *Panax notoginseng* (Sanqi). It contains >20 types of dammarane-type saponins and is prepared as a Xueshuantong capsule injection (36,37). PNS is commonly used to treat acute cerebral infarction and acute myocardial infarction (18,19). Previous studies have demonstrated that PNS can inhibit cisplatin-induced nephrotoxicity and neurotoxicity (20-23). However, whether PNS can inhibit ototoxicity induced by cisplatin has not yet been elucidated. The present study

demonstrated that PNS attenuated apoptosis induced by cisplatin in HEI-OC1 cells and increased their viability, which was consistent with the changes in the apoptosis-associated proteins Bcl-2 and caspase-3. The results indicated that PNS exhibited a therapeutic effect on ototoxicity induced by cisplatin.

At present, the underlying mechanisms of ototoxicity induced by cisplatin have not been fully demonstrated, whereas the excessive induction of oxidative stress and auditory cell apoptosis caused by increased free radical production has been widely accepted (3,10,38). Previous studies have reported that cisplatin can produce an excess of free radicals and deplete the activity of antioxidant enzyme systems in the cochlea, resulting in DNA, lipid and protein damage, as well as hair cell apoptosis, via the mitochondrial signaling pathway (38-40). Therefore, the restoration of the balance between oxidative stress and the expression of antioxidant enzymes is considered an important way to treat cisplatin-induced ototoxicity (4). Previous studies have demonstrated that PNS is dependent on potent free radical scavenging and antioxidant action in order to exert its therapeutic effects on cardiovascular and cerebrovascular conditions (36,41,42). In the present study, PNS administration could inhibit ROS production in HEI-OC1 cells exposed to cisplatin. Further experiments revealed that the protein expression levels of the antioxidant enzymes, including HO-1, NQO1 and GCLC, were markedly increased following PNS pretreatment, which indicated that remission of HEI-OC1 cell apoptosis may be associated with PNS-induced restoration of ROS balance.

Nrf2 is a transcription factor that exerts its function by regulating the redox homeostatic gene network (43). A previous study demonstrated that the activated Nrf2 protein binds to the antioxidant response element and increases the transcription of various cytoprotective genes that encode antioxidant proteins, including NQO1, HO-1 and GCLC (27). The increase in the expression levels of antioxidant enzymes, including GCLC, NQO1 and HO-1, could counteract the induction of oxidative stress (11-14). These enzymes reduce the intracellular ROS levels and, therefore, protect cells from oxidative damage (44,45). Additionally, the Nrf2 protein can upregulate the expression levels of Bcl-2 in order to prevent cell apoptosis (31). In the present study, PNS increased the levels of AKT phosphorylation and Nrf2 expression. These effects were reversed by treatment of the cells with an AKT inhibitor. These results suggested that PNS protected against ototoxicity induced by cisplatin via the activation of the AKT/Nrf2/ signaling-mediated redox pathway.

However, *in vivo* experiments should be conducted in the future in order to further discover the underlying effects of PNS on regulation of cisplatin-induced ototoxicity in auditory cells, which is considered to be a limitation of the present study.

In conclusion, the present study indicated the potential mechanism of the anti-cytotoxic and anti-apoptotic effects of PNS against cisplatin-induced ototoxicity. This mechanism was mediated by activation of the AKT/Nrf2 signaling pathway. The results indicated that PNS may function as a promising candidate in the prevention and treatment of cisplatin-induced ototoxicity.

## Acknowledgements

Not applicable.

## Funding

The present study was supported by a grant from the National Natural Science Foundation of China (grant no. 81670931), a grant from the Six Talent Peaks Project of Jiangsu Province (grant no. WSW-075), the Medical Science and Technology Development Key Foundation Nanjing Department of Health (grant no. ZKX17019) and the Jiangsu Provincial Key Medical Discipline (grant no. ZDXKB2016015), China.

## Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Authors' contributions

BF, ZBL, LSX, LYL, WYZ, JL, YHD and WDS made substantial contributions to conception, design and acquisition of data. BF, LYL, WYZ and JL contributed to manuscript drafting, analysis and interpretation of data. YHD and WDS were involved in manuscript revising, general supervision. All authors read and approved the final manuscript, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy of integrity of any part of the work are appropriately investigated and resolved.

## Ethics approval and consent to participate

The present study was approved by the Ethics Committee of Nanjing Drum Tower Hospital, The Affiliated Hospital of Nanjing University Medical School and in accordance with the Declaration of Helsinki. Written consent was obtained from each participant.

## Patient consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

1. Kelland L: The resurgence of platinum-based cancer chemotherapy. *Nat Rev Cancer* 7: 573-584, 2007.
2. Chovanec M, Abu Zaid M, Hanna N, El-Kouri N, Einhorn LH and Albany C: Long-term toxicity of cisplatin in germ-cell tumor survivors. *Ann Oncol* 28: 2670-2679, 2017.
3. Rybak LP, Mukherjee D, Jajoo S and Ramkumar V: Cisplatin ototoxicity and protection: Clinical and experimental studies. *Tohoku J Exp Med* 219: 177-186, 2009.
4. Sheth S, Mukherjee D, Rybak LP and Ramkumar V: Mechanisms of cisplatin-induced ototoxicity and otoprotection. *Front Cell Neurosci* 11: 338, 2017.
5. Kursunluoglu G, Taskiran D and Kayali HA: The investigation of the antitumor agent toxicity and capsaicin effect on the electron transport chain enzymes, catalase activities and lipid peroxidation levels in lung, heart and brain tissues of rats. *Molecules* 23: 3267, 2018.

6. Breglio AM, Rusheen AE, Shide ED, Fernandez KA, Spielbauer KK, McLachlin MM, Hall MD, Amable L and Cunningham LL: Cisplatin is retained in the cochlea indefinitely following chemotherapy. *Nat Commun* 8: 1654, 2017.
7. Moroso MJ and Blair RL: A review of cis-platinum ototoxicity. *J Otolaryngol* 12: 365-379, 1983.
8. Sakamoto M, Kaga K and Kamio T: Extended high-frequency ototoxicity induced by the first administration of cisplatin. *Otolaryngol Head Neck Surg* 122: 828-833, 2000.
9. Waissbluth S and Daniel SJ: Cisplatin-induced ototoxicity: Transporters playing a role in cisplatin toxicity. *Hear Res* 299: 37-45, 2013.
10. Ravi R, Somani SM and Rybak LP: Mechanism of cisplatin ototoxicity: Antioxidant system. *Pharmacol Toxicol* 76: 386-394, 1995.
11. Rybak LP, Ravi R and Somani SM: Mechanism of protection by diethyldithiocarbamate against cisplatin ototoxicity: Antioxidant system. *Fundam Appl Toxicol* 26: 293-300, 1995.
12. So H, Kim H, Kim Y, Kim E, Pae HO, Chung HT, Kim HJ, Kwon KB, Lee KM, Lee HY, *et al*: Evidence that cisplatin-induced auditory damage is attenuated by downregulation of pro-inflammatory cytokines via Nrf2/HO-1. *J Assoc Res Otolaryngol* 9: 290-306, 2008.
13. Vomund S, Schäfer A, Parnham MJ, Brüne B and Knethen A: Nrf2, the master regulator of anti-oxidative responses. *Int J Mol Sci* 18: 2772, 2017.
14. Kopke RD, Liu W, Gabaizadeh R, Jacono A, Feghali J, Spray D, Garcia P, Steinman H, Malgrange B, Ruben RJ, *et al*: Use of organotypic cultures of Corti's organ to study the protective effects of antioxidant molecules on cisplatin-induced damage of auditory hair cells. *Am J Otol* 18: 559-571, 1997.
15. Kim SJ, Park C, Han AL, Youn MJ, Lee JH, Kim Y, Kim ES, Kim HJ, Kim JK, Lee HK, *et al*: Ebselen attenuates cisplatin-induced ROS generation through Nrf2 activation in auditory cells. *Hear Res* 251: 70-82, 2009.
16. Na HK and Surh YJ: Oncogenic potential of Nrf2 and its principal target protein heme oxygenase-1. *Free Radic Biol Med* 67: 353-365, 2014.
17. Vriend J and Reiter RJ: The Keap1-Nrf2-antioxidant response element pathway: A review of its regulation by melatonin and the proteasome. *Mol Cell Endocrinol* 401: 213-220, 2015.
18. Li C, Xu T, Zhou P, Zhang J, Guan G, Zhang H, Ling X, Li W, Meng F, Liu G, *et al*: Post-marketing safety surveillance and re-evaluation of Xueshuantong injection. *BMC Complement Altern Med* 18: 277, 2018.
19. Huang Y, Guo B, Shi B, Gao Q and Zhou Q: Chinese herbal medicine xueshuantong enhances cerebral blood flow and improves neural functions in Alzheimer's disease mice. *J Alzheimers Dis* 63: 1089-1107, 2018.
20. Liang X, Yang Y, Huang Z, Zhou J, Li Y and Zhong X: *Panax notoginseng* saponins mitigate cisplatin induced nephrotoxicity by inducing mitophagy via HIF-1 $\alpha$ . *Oncotarget* 8: 102989-103003, 2017.
21. Liu X, Huang Z, Zou X, Yang Y, Qiu Y and Wen Y: *Panax notoginseng* saponins attenuates cisplatin-induced nephrotoxicity via inhibiting the mitochondrial pathway of apoptosis. *Int J Clin Exp Pathol* 7: 8391-8400, 2014.
22. Liu X, Huang Z, Zou X, Yang Y, Qiu Y and Wen Y: Possible mechanism of PNS protection against cisplatin-induced nephrotoxicity in rat models. *Toxicol Mech Methods* 25: 347-354, 2015.
23. Liu SJ and Zhou SW: *Panax notoginseng* saponins attenuated cisplatin-induced nephrotoxicity. *Acta Pharmacol Sin* 21: 257-260, 2000.
24. Hu S, Wu Y, Zhao B, Hu H, Zhu B, Sun Z, Li P and Du S: *Panax notoginseng* saponins protect cerebral microvascular endothelial cells against oxygen-glucose deprivation/reperfusion-induced barrier dysfunction via activation of PI3K/Akt/Nrf2 antioxidant signaling pathway. *Molecules* 23: 2781, 2018.
25. Kalinec G, Thein P, Park C and Kainec F: HEI-OC1 cells as a model for investigating drug cytotoxicity. *Hear Res* 335: 105-117, 2016.
26. Kalinec GM, Webster P, Lim DJ and Kalinec F: A cochlear cell line as an in vitro system for drug ototoxicity screening. *Audiol Neurotol* 8: 177-189, 2003.
27. Unoki T, Akiyama M, Kumagai Y, Gonçalves FM, Farina M, da Rocha JBT and Aschner M: Molecular pathways associated with methylmercury-induced Nrf2 modulation. *Front Genet* 9: 373, 2018.
28. Hill JM, Loeb E, MacLellan A, Hill NO, Khan A and King JJ: Clinical studies of platinum coordination compounds in the treatment of various malignant diseases. *Cancer Chemother Rep* 59: 647-659, 1975.
29. Hill JM and Speer RJ: Organo-platinum complexes as antitumor agents (review). *Anticancer Res* 2: 173-186, 1982.
30. Sun CY, Zhang QY, Zheng GJ and Feng B: Phytochemicals: Current strategy to sensitize cancer cells to cisplatin. *Biomed Pharmacother* 110: 518-527, 2019.
31. Cepeda V, Fuertes MA, Castilla J, Alonso C, Quevedo C and Pérez JM: Biochemical mechanisms of cisplatin cytotoxicity. *Anticancer Agents Med Chem* 7: 3-18, 2007.
32. Trendowski MR, El Charif O, Dinh PC Jr, Travis LB and Dolan ME: Genetic and modifiable risk factors contributing to cisplatin-induced toxicities. *Clin Cancer Res* 25: 1147-1155, 2019.
33. Travis LB, Fossa SD, Sessa HD, Frisina RD, Herrmann DN, Beard CJ, Feldman DR, Pagliaro LC, Miller RC, Vaughn DJ, *et al*: Chemotherapy-induced peripheral neurotoxicity and ototoxicity: New paradigms for translational genomics. *J Natl Cancer Inst* 106: dju044, 2014.
34. Im GJ, Chang J, Lee S, Choi J, Jung HH, Lee HM, Ryu SH, Park SK, Kim JH and Kim HJ: Protective role of edaravone against cisplatin-induced ototoxicity in an auditory cell line. *Hear Res* 330: 113-118, 2015.
35. Kim SJ, Park C, Lee JN, Lim H, Hong GY, Moon SK, Lim DJ, Choe SK and Park R: Erdosteine protects HEI-OC1 auditory cells from cisplatin toxicity through suppression of inflammatory cytokines and induction of Nrf2 target proteins. *Toxicol Appl Pharmacol* 288: 192-202, 2015.
36. Wang T, Guo R, Zhou G, Zhou X, Kou Z, Sui F, Li C, Tang L and Wang Z: Traditional uses, botany, phytochemistry, pharmacology and toxicology of *Panax notoginseng* (Burk.) F.H. Chen: A review. *J Ethnopharmacol* 188: 234-258, 2016.
37. Zhao H, Han Z, Li G, Zhang S and Luo Y: Therapeutic potential and cellular mechanisms of *Panax notoginseng* on prevention of aging and cell senescence-associated diseases. *Aging Dis* 8: 721-739, 2017.
38. Rybak LP, Mukherjee D and Ramkumar V: Mechanisms of cisplatin-induced ototoxicity and prevention. *Semin Hear* 40: 197-204, 2019.
39. Callejo A, Sedó-Cabezón L, Juan ID and Llorens J: Cisplatin-induced ototoxicity: Effects, mechanisms and protection strategies. *Toxics* 3: 268-293, 2015.
40. Op de Beeck K, Schacht J and Van Camp G: Apoptosis in acquired and genetic hearing impairment: The programmed death of the hair cell. *Hear Res* 281: 18-27, 2011.
41. Yang PF, Song XY and Chen NH: Advances in pharmacological studies of *Panax notoginseng* saponins on brain ischemia-reperfusion injury. *Yao Xue Xue Bao* 51: 1039-1046, 2016 (In Chinese).
42. Su P, Wang L, Du SJ, Xin WF and Zhang WS: Advance in studies of *Panax notoginseng* saponins on pharmacological mechanism of nervous system disease. *Zhongguo Zhong Yao Za Zhi* 39: 4516-4521, 2014 (In Chinese).
43. Ooi BK, Goh BH and Yap WH: Oxidative stress in cardiovascular diseases: Involvement of Nrf2 antioxidant redox signaling in macrophage foam cells formation. *Int J Mol Sci* 18: 2336, 2017.
44. Lourenço Dos Santos S, Petropoulos I and Friguet B: The oxidized protein repair enzymes methionine sulfoxide reductases and their roles in protecting against oxidative stress, in ageing and in regulating protein function. *Antioxidants (Basel)* 7: 191, 2018.
45. Machado-Silva A, Cerqueira PG, Grazielle-Silva V, Gadelha FR, Peloso Ede F, Teixeira SM and Machado CR: How trypanosoma cruzi deals with oxidative stress: Antioxidant defence and DNA repair pathways. *Mutat Res Rev Mutat Res* 767: 8-22, 2016.