

Low-calorie sweetener D-psicose promotes hydrogen peroxide-mediated apoptosis in C2C12 myogenic cells favoring skeletal muscle cell injury

ZHEN-JIE WEI^{1,2*}, LU SUN^{3*}, YU-LIN LI^{2*}, JIBRAN SUALEH MUHAMMAD⁴, YING WANG¹, QIAN-WEN FENG², YAN-ZHUO ZHANG¹, HIDEKUNI INADERA², ZHENG-GUO CUI^{2,5} and CHENG-AI WU¹

¹Department of Molecular Orthopedics, Beijing Research Institute of Traumatology and Orthopedics, Beijing Jishuitan Hospital, Beijing 100035, P.R. China; ²Department of Public Health, Graduate School of Medicine and Pharmaceutical Sciences, University of Toyama, Toyama 930-0194, Japan; ³Department of Pediatric Cardiology, Heart Center, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, Guangdong 510623, P.R. China; ⁴Department of Basic Medical Sciences, College of Medicine, University of Sharjah, Sharjah 27272, United Arab Emirates; ⁵Department of Environmental Health, School of Medical Science, University of Fukui, Fukui 910-1193, Japan

Received October 7, 2020; Accepted April 13, 2021

DOI: 10.3892/mmr.2021.12175

Abstract. Diet and exercise are the most effective approaches used to induce weight loss. D-psicose is a low-calorie sweetener that has been shown to reduce weight in obese individuals. However, the effect of D-psicose on muscle cells under oxidative stress, which is produced during exercise, requires further investigation. The present study aimed to determine the effects of D-psicose on C2C12 myogenic cells *in vitro*. Hydrogen peroxide (H₂O₂) was used to stimulate the generation of intracellular reactive oxygen species (ROS) in muscle cells to mimic exercise conditions. Cell viability was analyzed using a MTT assay and flow cytometry was used to analyze the levels of apoptosis, mitochondrial membrane potential (MMP), the generation of ROS and the cell cycle distribution following treatment. Furthermore, protein expression levels were analyzed using western blotting and cell proliferation was determined using a colony formation assay. The results

of the present study revealed that D-psicose alone exerted no toxicity on C2C12 mouse myogenic cells. However, in the presence of low-dose (100 μM) H₂O₂-induced ROS, D-psicose induced C2C12 cell injury and significantly decreased C2C12 cell viability in a dose-dependent manner. In addition, the levels of apoptosis and the generation of ROS increased, while the MMP decreased. MAPK family molecules were also activated in a dose-dependent manner following treatment. Notably, the combined treatment induced G₂/M phase arrest and reduced the proliferation of C2C12 cells. In conclusion, the findings of the present study suggested that D-psicose may induce toxic effects on muscle cells in a simulated exercise situation by increasing ROS levels, activating the MAPK signaling pathway and disrupting the MMP.

Introduction

Over the past four decades, the prevalence of obesity has tripled worldwide (1). As the number of individuals that are overweight and obese increases, the consumption of low-calorie sweeteners has also gradually increased each year (2). Although low-calorie synthetic sweeteners, such as aspartame and saccharine, may have a beneficial effect on weight loss, several studies have reported adverse effects (3). Therefore, nature-derived sugar substitutes with low caloric contents, such as D-psicose, have been developed.

D-psicose, also known as D-allulose, is produced in small quantities in various fruits. It is an epimer of fructose that can be enzymatically produced using D-psicose 3-epimerase (4). In addition to its anti-obesity effects, D-psicose has also been found to inhibit cancer cell proliferation by inducing apoptosis and cell cycle arrest (5-7). Since the purpose of using low-calorie sweeteners is to help lose weight, it is likely that individuals will also be regularly exercising while consuming D-psicose. However, the effect of the chronic consumption

Correspondence to: Dr Cheng-Ai Wu, Department of Molecular Orthopedics, Beijing Research Institute of Traumatology and Orthopedics, Beijing Jishuitan Hospital, 31 Xijiekou East Street, Xicheng, Beijing 100035, P.R. China
E-mail: wuchengai@jst-hosp.com.cn

Dr Zheng-Guo Cui, Department of Environmental Health, School of Medical Science, University of Fukui, 23-3 Matsuoka Shimoaizuki Eiheiji, Fukui 910-1193, Japan
E-mail: sai@u-fukui.ac.jp

*Contributed equally

Key words: D-psicose, muscle cell, reactive oxygen species, apoptosis, MAPK signaling pathway

of D-psicose on exercising muscle tissue remains unclear. A previous *in vivo* study reported that the oral administration of D-psicose was well absorbed and distributed throughout the body; the highest accumulation was found in the liver and kidneys, followed by the lungs, spleen and skeletal muscle tissue (8). Skeletal muscles comprise >40% of body organ systems. The muscles in those who exercise regularly are highly metabolically active and use the majority of the simple sugars available. Thus, the use of D-psicose by an obese individual who regularly exercises may lead to muscle cells being highly exposed to this natural sugar. Therefore, the effect of D-psicose on muscle cells under exercise-induced oxidative stress requires further investigation.

During exercise, the rate of muscle contractile activity exponentially increases, which leads to overactive mitochondria in muscle cells and the production of reactive oxygen species (ROS), such as superoxide free radicals (9). Certain types of free radicals have been detected in muscle tissue and increased free-radical activity has been shown to lead to extensive muscle damage. The produced reactive species can positively or negatively modulate muscle cells (10). Previous studies that have used hydrogen peroxide (H_2O_2) treatment for C2C12 myogenic cells to mimic exercise-induced alterations in skeletal muscles have revealed that oxidative stress could damage proliferating myoblasts (11,12).

Apoptosis is an evolutionarily conserved process that serves an important role in the musculoskeletal system during development, homeostasis and disease pathology (13). H_2O_2 and its reactive by-products act as potential mediators of apoptosis induced by diverse stimuli (14). Numerous signaling pathways have been found to be regulated by exercise-induced ROS generation, leading to muscle cell remodeling or death by apoptosis (10,11,13). Among these, MAPK was demonstrated to have a crucial role in exercise physiology. Apoptosis is a type of programmed cell death that occurs in response to unnecessary cell proliferation and facilitates the elimination of injured cells. The ratio of endogenous pro- to anti-apoptotic proteins is the major determining factor of cell fate. The overactivation of proapoptotic proteins in cells is known to inhibit cell cycle progression and cell proliferation (15). Notably, several previous studies have reported that D-psicose upregulated the expression levels of CDKs, leading to cell cycle arrest and apoptosis in several cell types (5-7).

The present study aimed to determine the effects of D-psicose on C2C12 myogenic cells. Low dose H_2O_2 was used to mimic the generation of ROS during exercise and to determine how D-psicose affected skeletal muscle cells under oxidative stress-induced conditions, such as exercise. The results revealed that D-psicose, in the presence of physiological concentrations of H_2O_2 , induced the generation of ROS, triggered cell cycle arrest and initiated apoptosis in C2C12 myogenic cells. To the best of our knowledge, this was the first study to demonstrate the toxic effects of D-psicose on muscle cells treated with H_2O_2 , which were suggested to occur via the MAPK signaling pathway.

Materials and methods

Chemicals. D-psicose, H_2O_2 , N-acetylcysteine (NAC), JNK inhibitor VIII and SB203580 were all purchased from Sigma-Aldrich (Merck KGaA).

Cell culture and treatment. C2C12 cells were obtained from the Human Sciences Research Resources Bank (Japan Human Sciences Foundation, Tokyo, Japan). Cells were cultured in low glucose DMEM (Thermo Fisher Scientific, Inc.), supplemented with 10% heat-inactivated FBS (Gibco; Thermo Fisher Scientific, Inc.), and maintained at 37°C in a humidified atmosphere with 5% CO_2 . For each experiment, cells were seeded in a 6-well plate and pretreated with 1, 2 or 5 mM D-psicose for 3 h and then with 100 μM H_2O_2 for 2 h, after which the medium was changed, followed by incubation for 19 h.

MTT assay. An MTT assay was used to evaluate cell viability. Briefly, cells were seeded into 96-well plates at a density of 3×10^3 cells/well and incubated overnight. Following the incubation, 20 μl sterile MTT dye (5 mg/ml) was added to each well and incubated for another 3 h at 37°C. After removal of the medium, 100 μl DMSO was added to each well and incubated for a further 10 min. The absorbance was measured at a wavelength of 570 nm using a microplate reader (16). Eight replicate wells were used for each concentration.

Flow cytometric analysis of apoptosis. Cells were stained with Annexin V-FITC and propidium iodide (PI; cat. no. BMS500FI-300; Thermo Fisher Scientific, Inc.) as previously described (17). Apoptotic cells were subsequently analyzed using flow cytometry (BD FACSCanto II flow cytometer and FACSDiva ver. 6.1; BD Biosciences) according to the manufacturer's protocol.

Colony formation assay. Cells (1×10^3 /well) were seeded into 6-well plates and cultured in the indicated media for 10-15 days. Subsequently, the media was removed, cells were washed twice in PBS, fixed with 4% paraformaldehyde for 1 h at room temperature and stained with crystal violet for 40 min at room temperature. Plates were then thoroughly washed with water and air-dried.

Cell cycle distribution analysis. Cells were harvested after 24 h of treatment and centrifuged at 200 x g for 5 min at room temperature. Cell pellets were suspended in 100 μl PBS prior to being fixed with 75% (v/v) cold ethanol for 2 h at room temperature and then stained with a PI solution containing DNase-free RNase A for 30 min at room temperature in the dark. Analysis was performed using a flow cytometer according to the manufacturer's protocol.

Measurement of intracellular ROS levels. Intracellular ROS levels were detected using dihydroethidium (DHE; Molecular Probes; Thermo Fisher Scientific, Inc.) using flow cytometry. DHE produces blue fluorescence in the cytosol until it is oxidized by superoxide to 2-hydroxyethidium, which then intercalates within cellular DNA and stains the nucleus a bright fluorescent red. Following incubation with 4 μM DHE for 15 min at 37°C, the cells were washed twice with PBS and the intracellular ROS levels (%) were analyzed using flow cytometry.

Measurement of the mitochondrial membrane potential (MMP). Following treatment, the cells were collected and treated with 10 nM tetramethylrhodamine methyl ester perchlorate (TMRM;

Molecular Probes; Thermo Fisher Scientific, Inc.) in 1 ml FBS (1%) diluted in PBS for 15 min at 37°C. TMRM is a cationic fluorophore widely used for staining cellular mitochondria and the mitochondrial matrix. The percentage of cells with potential MMP loss was analyzed by flow cytometry gated for red TMRM fluorescence, using an excitation wavelength of 488 nm and an emission wavelength of 575 nm (18).

Western blotting. Cells were harvested, washed with PBS and total protein was extracted using RIPA lysis buffer [150 mM NaCl, 1% Triton X-100 (v/v), 1% sodium deoxycholate, 0.1% SDS, 1 µg/ml each of aprotinin, pepstatin and leupeptin, 1 mM EGTA, 50 mM Tris-HCl; pH 7.5] for 20 min on ice. Following a brief sonication, lysates were centrifuged at 13,000 x g for 10 min at 4°C, the supernatant was collected and the protein contents in the supernatant were measured using a Coomassie (Bradford) Protein Assay kit (Thermo Fisher Scientific, Inc.). An equal amount of protein (10 µg) was loaded per lane and separated via SDS-PAGE. The separated proteins were subsequently transferred onto PVDF membranes and blocked with 5% skimmed milk in TBS-Tween 20 (150 mM NaCl, 50 mM Tris, 0.1% Tween-20; pH 7.5) for 1 h at room temperature. The membrane was then incubated overnight at 4°C with the following primary antibodies: Anti-MCL1 apoptosis regulator BCL2 family member (Mcl-1; cat. no. CST68542; 1:1,000), anti-cleaved caspase-3 (cat. no. CST9661; 1:1,000), anti-JNK (cat. no. CST9258; 1:1,000), anti-phosphorylated (p)-JNK (cat. no. CST4668; 1:1,000), anti-ERK1/2 (cat. no. CST4695; 1:1,000), anti-p-ERK1/2 (cat. no. CST4370; 1:2,000), anti-p38 (cat. no. CST8690; 1:1,000), anti-p-p38 (cat. no. CST4511; 1:1,000), anti-p-checkpoint kinase 1 (Chk1; cat. no. CST2348; 1:1,000), anti-p-DNA polymerase δ 1, catalytic subunit (CDC2; cat. no. CST4539; 1:1,000), anti-sirtuin 3 (SIRT3; cat. no. CST5490; 1:1,000) (all Cell Signaling Technology, Inc.), anti-Bax (cat. no. ab32503; 1:5,000), anti-cleaved-poly(ADP-ribose) polymerase 1 (PARP1; cat. no. ab32064; 1:1,000; both Abcam), anti-cell division cycle 25 C (CDC25C; cat. no. sc327; 1:1,000), anti-superoxide dismutase 2 (SOD2; cat. no. sc30080; 1:1,000), anti-Bcl-2 (cat. no. sc7382; 1:400) and anti-β-actin (cat. no. sc8432; 1:1,000; all Santa Cruz Biotechnology, Inc.). Following the primary antibody incubation, the membranes were incubated for 2 h at room temperature with the following secondary antibodies: Anti-rabbit IgG, HRP-linked Antibody (cat. no. CST7074; 1:3,000) and anti-mouse IgG, HRP-linked Antibody (cat. no. CST7076; 1:2,000; both Cell Signaling Technology, Inc.). Protein bands were visualized using ECL (Amersham; Cytiva) according to the manufacturer's protocol (19). Densitometric analysis was performed using ImageJ software 1.50i (National Institutes of Health).

Statistical analysis. All experiments are representative of three independent biological replicates. Data are presented as the mean ± SD. Multiple comparisons were performed using a one-way or two-way ANOVA with a Tukey's post hoc test. P<0.05 was considered to indicate a statistically significant difference.

Results

D-psicose treatment exerts no cytotoxic effects on resting muscle cells. D-psicose is a ketohexose monosaccharide with




Figure 1. Safety of D-psicose in C2C12 myoblasts. (A) Two-dimensional and (B) three-dimensional chemical structure of D-psicose. (C) C2C12 cells were treated with 0, 0.1, 0.2, 1, 2 or 5 mM D-psicose for 24 or 48 h and cell viability was subsequently analyzed using a MTT assay. The results revealed that D-psicose exerted no cytotoxic effects on C2C12 myoblasts.

the following molecular formula: C₆H₁₂O₆. It is a C3 epimer of fructose that was first derived from the antibiotic psicofurarine, which contains one ketone group that acts as a reducing agent (4). The three-dimensional structure produces a white, odorless powder that is soluble in water (Fig. 1A and B). The present study first investigated the effect of D-psicose on C2C12 myogenic cells in resting conditions (without oxidative stress). The results demonstrated that increasing concentrations of D-psicose (0-5 mM) exerted no cytotoxic effects on resting C2C12 cells at 24 or 48 h post-treatment (Fig. 1C), which suggested the potential safe application of D-psicose as a sugar replacement in daily food products.

D-psicose exacerbates H₂O₂-induced cell damage in C2C12 cells. To mimic exercise-induced oxidative stress in muscle cells, C2C12 cells were treated with H₂O₂. The results revealed that as the concentration of H₂O₂ increased, the viability of C2C12 cells decreased; ~10% of the cells had decreased viability following 100 µM H₂O₂ treatment and only 60% of cells remained viable following 500 µM H₂O₂ treatment (Fig. 2A). Thus, 100 µM H₂O₂ was selected as the optimal concentration to act as an exercise mimetic in further experiments.

C2C12 cells were pre-incubated with D-psicose for 3 h and then treated with H₂O₂ for 2 h, after which the cell medium was replaced with fresh cell medium, and cells were incubated for a further 19 h. Cell viability analysis revealed a dose-dependent decrease in cell viability in cells treated with D-psicose in the presence of H₂O₂. Notably, viability was lost in >50% of cells following 5 mM D-psicose treatment (Fig. 2B).

Subsequently, whether H₂O₂ induced apoptosis in D-psicose pretreated cells was determined (Fig. 2C and D).




Figure 2. D-psicose enhances H₂O₂-induced C2C12 cell damage. (A) C2C12 cells were pretreated with 0–500 μM H₂O₂ for 2 h, after which the medium was changed. Following incubation for 24 h, a MTT assay was used to analyze cell viability. The concentration of 100 μM H₂O₂ was determined to be a moderate inducer of oxidative stress. C2C12 cells were pretreated with 1, 2 or 5 mM D-psicose for 3 h and then with 100 μM H₂O₂ for 2 h, after which the medium was changed. Following incubation for 19 h, (B) an MTT assay, (C) Annexin V/propidium iodide staining, (D) flow cytometry, (J and K) tetramethylrhodamine methyl ester perchlorate staining and (E–I) western blotting were used to analyze cell viability, cell apoptosis, MMP loss and the expression levels of apoptosis-related proteins. Data are presented as the mean ± SEM of independent experiments. *P<0.05, **P<0.01 and ***P<0.001 vs. NT; †P<0.05, ††P<0.01 and †††P<0.001 vs. H₂O₂ (n=3). H₂O₂, hydrogen peroxide; MMP, mitochondrial membrane potential; Mcl-1, MCL1 apoptosis regulator BCL2 family member; PARP-1, anti-cleaved-poly(ADP-ribose) polymerase 1; NT, No treatment.




Figure 3. D-psicose enhances H₂O₂-induced apoptosis of C2C12 cells through the MAPK signaling pathway. C2C12 cells were pretreated with 1, 2 or 5 mM D-psicose for 3 h and then 100 μM H₂O₂ for 2 h, after which the medium was changed. Following incubation for 3 h, protein expression levels of (A) p-JNK and JNK, (B) p-P38 and P38, and (C) p-ERK and ERK were analyzed using western blotting. (D) Representative western blotting image of p-JNK, JNK, p-P38, P38, p-ERK and ERK. (E) C2C12 cells were pretreated with 5 mM D-psicose + 100 μM H₂O₂ or 5 mM D-psicose + 10 μM JNK inhibitor VIII + 100 μM H₂O₂ for 1 h. The expression levels of JNK, p-JNK, cleaved caspase-3, Bcl-2 and Bax were analyzed using western blotting. (F) C2C12 cells were pretreated with 5 mM D-psicose + 100 μM H₂O₂ or 5 mM D-psicose + 20 μM SB203580 + 100 μM H₂O₂ for 1 h. The expression levels of P38, p-P38, cleaved caspase-3, Bcl-2 and Bax were analyzed using western blotting. β-actin was used as the loading control. Data are presented as the mean ± SEM of independent experiments. **P<0.01 and ***P<0.001 vs. NT; *P<0.05 and ***P<0.001 vs. H₂O₂; §P<0.05 vs. D-psicose + H₂O₂ (n=3). H₂O₂, hydrogen peroxide; p-, phosphorylated.

To investigate the effect of D-psicose on the apoptotic pathway, expression levels of proteins associated with apoptosis were analyzed using western blotting. D-psicose was found to potentiate H₂O₂-induced apoptosis by downregulating the expression levels of the anti-apoptotic proteins, Bcl-2 and Mcl-1, and upregulating the expression levels of the proapoptotic protein Bax (Fig. 2E-I). A concomitant increase in the cleavage of caspase-3 and PARP1 was also observed, suggesting that D-psicose initiated caspase-mediated

apoptosis. When cells are under immense oxidative stress, mitochondria lose their membrane integrity and exercising muscles rely on mitochondria for the majority of their energy production. Thus, the effects of D-psicose on mitochondrial function were investigated. The treatment with H₂O₂ or D-psicose alone did not affect the mitochondria of muscle cells; however, in the presence of D-psicose pretreatment, H₂O₂ treatment significantly decreased the MMP in C2C12 cells (Fig. 2J and K).




Figure 4. Co-treatment of D-psicose and H₂O₂ increases ROS generation in C2C12 cells. C2C12 cells were pretreated with 1, 2 or 5 mM D-psicose for 3 h and then 100 μM H₂O₂ for 2 h, after which the medium was changed. Following incubation for 30 min, ROS generation was analyzed using dihydroethidium staining and flow cytometry (A) Representative flow cytometry plots and (B) quantification of ROS generation. (C) C2C12 cells were pretreated with 5 mM D-psicose + 100 μM H₂O₂ or 5 mM D-psicose + 5 mM NAC + 100 μM H₂O₂ for 1 h. The expression levels of p-p38, p-JNK, p-ERK, cleaved caspase-3, Bcl-2 and Bax were analyzed using western blotting. (D) C2C12 cells were pretreated with 1, 2 or 5 mM D-psicose for 3 h and with 100 μM H₂O₂ for 2 h, after which the medium was changed. Following incubation for 3 h, the expression levels of SIRT3 and SOD2 were analyzed using western blotting. Data are presented as the mean ± SEM of independent experiments. [#]P<0.05 and ^{###}P<0.001 vs. NT; ^{*}P<0.05 vs. H₂O₂; [§]P<0.05 vs. D-psicose + H₂O₂ (n=3). H₂O₂, hydrogen peroxide; ROS, reactive oxygen species; p-, phosphorylated; SIRT3, sirtuin 3; SOD2, superoxide dismutase 3; NAC, N-acetylcysteine; NT, No treatment.

D-psicose exacerbates H₂O₂-induced apoptosis in C2C12 cells through the MAPK signaling pathway. Exercising is an intermittent form of cellular stress that was discovered to activate proteins of the MAPK signaling pathways in the skeletal muscle of rats (20). In addition, the H₂O₂-induced

upregulation of ERK, p38 and JNK expression was reported to regulate cell death in C2C12 cells (21). To investigate the role of MAPKs in the damaging effect of D-psicose in exercising skeletal muscle cells, the protein expression levels of ERK, p38 and JNK were analyzed in H₂O₂-stimulated C2C12 cells.




Figure 5. Co-treatment with D-psicose and H_2O_2 induces G₂/M phase cell cycle arrest in C2C12 myoblasts. C2C12 cells were pretreated with 1, 2 or 5 mM D-psicose for 3 h and then with 100 μ M H_2O_2 for 2 h, after which the medium was changed. Following incubation for 24 h, the SubG₁ cell population was analyzed using propidium iodide staining and flow cytometry. (A) Representative flow cytometry plots and (B) quantification of cells in each state of the cell cycle. (C) Colony formation assays were performed using cells seeded at a density of 1×10^3 cells/well into 6-well plates for 24 h, treated as described in part (A) and incubated for 5 days. (D) C2C12 cells were pretreated with 1, 2 or 5 mM D-psicose for 3 h and then with 100 μ M H_2O_2 for 2 h, after which the medium was changed. Following incubation for 3 h, the expression levels of p-Chk1, CDC25C and p-CDC2 were analyzed using western blotting. The relative expression levels of (E) p-Chk1, (F) CDC25C and (G) p-CDC2 were quantified relative to the value of β -actin. Data are presented as the mean \pm SEM of independent experiments. $^{##}P < 0.01$ and $^{###}P < 0.001$ vs. NT; $^*P < 0.05$ and $^{***}P < 0.001$ vs. H_2O_2 (n=3). H_2O_2 , hydrogen peroxide; p-, phosphorylated; Chk1, checkpoint kinase 1; CDC25C, cell division cycle 25 C; CDC2, DNA polymerase δ 1, catalytic subunit; NT, No treatment.

The expression levels of p-JNK and p-p38 were significantly upregulated following the pretreatment with increasing concentrations of D-psicose in H_2O_2 -treated C2C12 cells; however, the expression levels of p-ERK were downregulated (Fig. 3A-D). These data suggested that JNK and p38 may have a role in the regulation of D-psicose-induced apoptosis in muscle cells under oxidative stress conditions.

To validate these findings, muscle cells were exposed to JNK and p-38 inhibitors (JNK inhibitor VIII and SB203580, respectively). Following the pre-incubation of the C2C12 cell culture with JNK inhibitor VIII or SB203580, followed by treatment with D-psicose and H_2O_2 , the expression levels of p-JNK and p38 were downregulated. Concurrently, both of the inhibitors also reversed D-psicose-induced apoptotic effects by upregulating




Figure 6. Proposed molecular mechanism through which D-psicose and H₂O₂ may induce apoptosis and G₂/M cell cycle arrest. H₂O₂, hydrogen peroxide; CDC25C, cell division cycle 25 C; CDC2, DNA polymerase δ 1, catalytic subunit; Chk1, checkpoint kinase 1; H₂O₂, hydrogen peroxide; MMP, membrane mitochondrial potential; ROS, reactive oxygen species; NAC, N-acetylcysteine; p-, phosphorylated.

Bcl-2 expression levels and inhibiting the cleavage of caspase-3 (Fig. 3E and F).

D-psicose pretreatment enhances H₂O₂-induced oxidative stress via ROS generation. The generation of ROS and activation of apoptosis-related proteins serves a key role in triggering apoptosis. Previously, it was reported that H₂O₂ induced high levels of oxidative stress and apoptosis in proliferating skeletal myoblasts (22). Hence, the present study sought to determine whether the same was true for the effect of D-psicose-induced apoptosis in C2C12 cells. The results revealed that pretreatment with increasing concentrations of D-psicose increased the production of ROS in H₂O₂-treated C2C12 cells, while this effect was significantly reversed upon exposure to NAC (Fig. 4A and B). In addition, MAPK signaling molecules were observed to be activated by D-psicose, while the effects of H₂O₂ treatment were markedly blocked by NAC. Furthermore, the phosphorylation levels of JNK and p38 were reduced, while the phosphorylation levels of ERK were increased. Concurrently, the levels of apoptosis were inhibited, which was evident by the upregulated expression levels of Bcl-2, and downregulated expression levels of Bax and cleaved caspase-3 (Fig. 4C). As downregulated expression levels of SIRT3 were previously reported to promote the generation of ROS and inhibit the SIRT3/SOD2-mediated deactivation process of superoxide radicals (23), the present study also investigated the effect of D-psicose and H₂O₂ treatment on the expression levels of SIRT3 and SOD2 in C2C12 cells. The results revealed that the pretreatment with D-psicose significantly downregulated the expression levels of SIRT3 and SOD2 in H₂O₂-treated C2C12 cells in a dose-dependent manner (Fig. 4D).

D-psicose and H₂O₂ induce cell cycle arrest and inhibit the proliferation of C2C12 cells. The effect of D-psicose on

muscle cell proliferation under conditions of oxidative stress were subsequently investigated. The combined treatment with D-psicose and H₂O₂ increased the percentage of cells in the sub-G₁ phase of the cell cycle compared with cells in other phases of the cell cycle (Fig. 5A and B). This finding suggested the potential inhibitory effect of D-psicose on cell viability and proliferation. Furthermore, a colony formation assay was performed for 5 days post-treatment to determine the long-term effects of D-psicose and H₂O₂ treatment on muscle cell proliferation. Both the number and size of colonies were decreased in the treatment groups, while few colonies formed from H₂O₂-stimulated C2C12 cells pretreated with 5 mM D-psicose (Fig. 5C). In addition, the expression levels of several proteins associated with the cell cycle were analyzed using western blotting. The results revealed that, in C2C12 cells, D-psicose treatment in the presence of H₂O₂ downregulated the expression levels of CDC25C and the phosphorylation of CDC2 (Fig. 5D-G). These findings indicated that D-psicose may potentiate cell cycle arrest and decrease the long-term survival of muscle cells under oxidative stress conditions.

Discussion

The present study aimed to determine the effect of D-psicose on ROS generation and H₂O₂-induced apoptosis in C2C12 myogenic cells. H₂O₂ is a widely used method to establish exercise-induced oxidative stress in cells (11,12,22). The present results revealed that the treatment of C2C12 cells with D-psicose under oxidative stress conditions elevated the levels of ROS, decreased the MMP and apoptosis, and induced cell cycle arrest. Moreover, the lethal effects of D-psicose on exercising muscle cells were found to be mediated via the MAPK family molecules, p-JNK and p-p38 (Fig. 6). Notably, the observed effects were reversed following treatment with the ROS inhibitor, NAC, which is consistent with the findings of a previous study (24).

Oxidative stress is the production of excessive reactive oxidants that exceeds the cellular antioxidant capability. In 1992, it was reported that contracting rodent skeletal muscle released superoxide radicals into the interstitial spaces (25). Bailey *et al.* (26) also demonstrated that exercise resulted in a greater production and accumulation of intramuscular free radicals in human skeletal muscle cells. ROS produced in contracting skeletal muscle facilitates the strength of the muscle contraction; however, if ROS concentrations exceed a certain level, it reduces force generation and causes fatigue (27). The findings of the present study revealed that low doses of H₂O₂ induced oxidative stress in muscle cells, which may be due to exercise-induced ROS generation. However, the concomitant exposure to D-psicose further increased the production of ROS, leading to cell death. On the other hand, pretreatment with NAC reversed the increased production of ROS, returning the levels of ROS to almost homeostatic levels, thereby providing a protective effect. These results were consistent with previous studies, which showed that antioxidant infusion in humans during voluntary exercise substantially enhanced performance during prolonged exercise (28,29).

The balance between antioxidants and pro-oxidants is crucial for the survival of aerobic organisms. A previous experiment in rats demonstrated that exercise promoted the expression

and activity of SOD2 in skeletal muscle mitochondria, which subsequently maintained cellular oxidant-antioxidant homeostasis (30). Therefore, as the exercise-induced upregulation of SOD2 in skeletal muscle served a key role in the prevention of superoxide accumulation, it could be hypothesized that the loss of SOD2 may have devastating effects on muscle cell survival. SIRT3 was discovered to regulate SOD2 via deacetylation and activation of FOXO3a, and SOD2 can catalyze the neutralization process involving the conversion of superoxide into water and hydrogen peroxide (23,31). The results of the present study revealed that the treatment with D-psicose alone, in addition to in the presence of H₂O₂, exerted inhibitory effects on the expression levels of SIRT3 and SOD2.

D-psicose was found to suppress post-prandial serum glucose levels and reduce the accumulation of body fat, which indicates its beneficial effects on obesity-related metabolic disturbances (8). In the past, this type of sugar supplement was used in tabletop packets in cafés and restaurants. Currently, these sugars are added to everyday food products to promote the sales of food items as healthy alternatives. In both children and adults, the consumption of low-calorie sweeteners has significantly increased, primarily due to an increased awareness of their non-nutritive value and potential effects in weight loss (32). Hence, future studies should aim to enhance the existing knowledge on the metabolic and health effects of low-calorie sweeteners. A long-term study on D-psicose toxicity in rats revealed that the final body weight and weight gain in D-psicose-fed animals were significantly decreased compared with the control group (33). On the one hand, these findings may be considered as beneficial for weight loss; however, D-psicose may be reducing the body weight by decreasing the number of muscle cells. Yagi and Matsuo (33) investigated the long-term (18 months) toxicity of D-psicose in rats and found a significant increase in organ weight of the liver, kidneys, brain, lungs and pancreas, which suggested that the accumulation of fat, and that muscle proteins may be broken down to promote toxic effects. The present data suggested that the treatment with D-psicose was safe for muscle cells; however, under stressful conditions, D-psicose could exert deleterious effects on the survival of the muscle cells.

Previous studies have shown that D-psicose increased G₂/M phase cell cycle arrest and decreased the population of liver cells in the S phase (5). Cell cycle progression is tightly regulated by a complex network of positive and negative cell cycle regulatory molecules, such as cyclins and CDKs. CDC25C is involved in the DNA damage checkpoints by activating CDK complexes that drive the cell cycle and is a key mediator of cell cycle progression (34). The current findings demonstrated that the treatment with D-psicose in H₂O₂-stimulated muscle cells led to the accumulation of cells in the sub-G₁ phase and the downregulation of the expression levels of the cell cycle-regulating protein, CDC25C.

In conclusion, the findings of the present study suggested that under physiological oxidative stress conditions, D-psicose may exert negative effects on muscle cells by promoting the production of ROS and triggering cell cycle arrest and apoptosis. The results further indicated that the negative effects of D-psicose on exercising muscle cells may be mediated by MAPK and may be partly due to slow cell

cycle progression and the failure to repair DNA damage, which ultimately leads to apoptosis.

Acknowledgements

Not applicable.

Funding

This work was supported by the National Natural Science Foundation of China (grant no. 81330043) and Beijing Municipal Health Commission (grant nos. BMHC-2019-9 and BMHC-2018-4).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

ZW and LS designed the study. ZW and YL performed the experiments and collected the data. ZC and CW confirm the authenticity of all the raw data. ZW interpreted the data with help from LS, YL, JM, YW, QF, YZ, HI, ZC and CW. ZW, LS and YL analyzed the results. ZW and JM wrote the manuscript with help from ZC and CW. ZC and CW were responsible for funding acquisition. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Chu DT, Minh Nguyet NT, Dinh TC, Thai Lien NV, Nguyen KH, Nhu Ngoc VT, Tao Y, Son LH, Le DH, Nga VB, *et al*: An update on physical health and economic consequences of overweight and obesity. *Diabetes Metab Syndr* 12: 1095-1100, 2018.
2. Sylvetsky AC and Rother KI: Trends in the consumption of low-calorie sweeteners. *Physiol Behav* 164: 446-450, 2016.
3. Swithers SE: Artificial sweeteners produce the counterintuitive effect of inducing metabolic derangements. *Trends Endocrinol Metab* 24: 431-441, 2013.
4. He W, Mu W, Jiang B, Yan X and Zhang T: Construction of a food grade recombinant bacillus subtilis based on replicative plasmids with an auxotrophic marker for biotransformation of d-Fructose to d-Allulose. *J Agric Food Chem* 64: 3243-3250, 2016.
5. Sui L, Dong Y, Watanabe Y, Yamaguchi F, Hatano N, Tsukamoto I, Izumori K and Tokuda M: The inhibitory effect and possible mechanisms of D-allose on cancer cell proliferation. *Int J Oncol* 27: 907-912, 2005.
6. Sui L, Dong Y, Watanabe Y, Yamaguchi F, Hatano N, Izumori K and Tokuda M: Growth inhibitory effect of D-allose on human ovarian carcinoma cells in vitro. *Anticancer Res* 25: 2639-2644, 2005.

7. Mitani T, Hoshikawa H, Mori T, Hosokawa T, Tsukamoto I, Yamaguchi F, Kamitori K, Tokuda M and Mori N: Growth inhibition of head and neck carcinomas by D-allose. *Head Neck* 31: 1049-1055, 2009.
8. Tsukamoto I, Hossain A, Yamaguchi F, Hirata Y, Dong Y, Kamitori K, Sui L, Nonaka M, Ueno M, Nishimoto K, *et al.*: Intestinal absorption, organ distribution, and urinary excretion of the rare sugar D-psicose. *Drug Des Devel Ther* 8: 1955-1964, 2014.
9. Qaisar R, Bhaskaran S and Van Remmen H: Muscle fiber type diversification during exercise and regeneration. *Free Radic Biol Med* 98: 56-67, 2016.
10. Derbre F, Ferrando B, Gomez-Cabrera MC, Sanchis-Gomar F, Martinez-Bello VE, Olaso-Gonzalez G, Diaz A, Gratas-Delamarche A, Cerda M and Viña J: Inhibition of xanthine oxidase by allopurinol prevents skeletal muscle atrophy: role of p38 MAPKinase and E3 ubiquitin ligases. *PLoS One* 7:e46668, 2012.
11. Fan X, Hussien R and Brooks GA: H₂O₂-induced mitochondrial fragmentation in C2C12 myocytes. *Free Radic Biol Med* 49: 1646-1654, 2010.
12. Siu PM, Wang Y and Alway SE: Apoptotic signaling induced by H₂O₂-mediated oxidative stress in differentiated C2C12 myotubes. *Life Sci* 84: 468-481, 2009.
13. Sandri M and Carraro U: Apoptosis of skeletal muscles during development and disease. *Int J Biochem Cell Biol* 31: 1373-1390, 1999.
14. Jackson MJ, Edwards RH and Symons MC: Electron spin resonance studies of intact mammalian skeletal muscle. *Biochim Biophys Acta* 847: 185-190, 1985.
15. Goldar S, Khaniani MS, Derakhshan SM and Baradaran B: Molecular mechanisms of apoptosis and roles in cancer development and treatment. *Asian Pac J Cancer Prev* 16: 2129-2144, 2015.
16. Zakki SA, Muhammad JS, Li JL, Sun L, Li ML, Feng QW, Li YL, Cui ZG and Inadera H: Melatonin triggers the anticancer potential of phenylarsine oxide via induction of apoptosis through ROS generation and JNK activation. *Metallomics* 12: 396-407, 2020.
17. Hertveldt K, Philippe J, Thierens H, Cornelissen M, Vral A and De Ridder L: Flow cytometry as a quantitative and sensitive method to evaluate low dose radiation induced apoptosis in vitro in human peripheral blood lymphocytes. *Int J Radiat Biol* 71: 429-433, 1997.
18. Wu J, Liu T, Rios Z, Mei Q, Lin X and Cao S: Heat shock proteins and cancer. *Trends Pharmacol Sci* 38: 226-256, 2017.
19. Li M, Kondo T, Zhao QL, Li FJ, Tanabe K, Arai Y, Zhou ZC and Kasuya M: Apoptosis induced by cadmium in human lymphoma U937 cells through Ca²⁺-calpain and caspase-mitochondria-dependent pathways. *J Biol Chem* 275: 39702-39709, 2000.
20. Goodyear LJ, Chang PY, Sherwood DJ, Dufresne SD and Moller DE: Effects of exercise and insulin on mitogen-activated protein kinase signaling pathways in rat skeletal muscle. *Am J Physiol* 271: E403-E408, 1996.
21. Ronda AC, Vasconsuelo A and Boland R: Extracellular-regulated kinase and p38 mitogen-activated protein kinases are involved in the antiapoptotic action of 17beta-estradiol in skeletal muscle cells. *J Endocrinol* 206: 235-246, 2010.
22. Caporossi D, Ciafre SA, Pittaluga M, Savini I and Farace MG: Cellular responses to H₂O₂ and bleomycin-induced oxidative stress in L6C5 rat myoblasts. *Free Radic Biol Med* 35: 1355-1364, 2003.
23. Lu J, Cheng K, Zhang B, Xu H, Cao Y, Guo F, Feng X and Xia Q: Novel mechanisms for superoxide-scavenging activity of human manganese superoxide dismutase determined by the K68 key acetylation site. *Free Radic Biol Med* 85: 114-126, 2015.
24. Coffey VG, Zhong Z, Shield A, Canny BJ, Chibalin AV, Zierath JR and Hawley JA: Early signaling responses to divergent exercise stimuli in skeletal muscle from well-trained humans. *FASEB J* 20: 190-192, 2006.
25. Reid MB, Shoji T, Moody MR and Entman ML: Reactive oxygen in skeletal muscle. II. Extracellular release of free radicals. *J Appl Physiol* (1985) 73: 1805-1809, 1992.
26. Bailey DM, Lawrenson L, McEneny J, Young IS, James PE, Jackson SK, Henry RR, Mathieu-Costello O, McCord JM and Richardson RS: Electron paramagnetic spectroscopic evidence of exercise-induced free radical accumulation in human skeletal muscle. *Free Radic Res* 41: 182-190, 2007.
27. Nemes R, Koltai E, Taylor AW, Suzuki K, Gyori F and Radak Z: Reactive oxygen and nitrogen species regulate key metabolic, anabolic, and catabolic pathways in skeletal muscle. *Antioxidants (Basel)* 7: 85, 2018.
28. Medved I, Brown MJ, Bjorksten AR and McKenna MJ: Effects of intravenous N-acetylcysteine infusion on time to fatigue and potassium regulation during prolonged cycling exercise. *J Appl Physiol* (1985) 96: 211-217, 2004.
29. McKenna MJ, Medved I, Goodman CA, Brown MJ, Bjorksten AR, Murphy KT, Petersen AC, Sostaric S and Gong X: N-acetylcysteine attenuates the decline in muscle Na⁺,K⁺-pump activity and delays fatigue during prolonged exercise in humans. *J Physiol* 576: 279-288, 2006.
30. Hollander J, Fiebig R, Gore M, Ookawara T, Ohno H and Ji LL: Superoxide dismutase gene expression is activated by a single bout of exercise in rat skeletal muscle. *Pflugers Arch* 442: 426-434, 2001.
31. Kim YS, Gupta Vallur P, Phaeton R, Mythreye K and Hempel N: Insights into the Dichotomous Regulation of SOD2 in Cancer. *Antioxidants (Basel)* 6: 86, 2017.
32. Magnuson BA, Carakostas MC, Moore NH, Poulos SP and Renwick AG: Biological fate of low-calorie sweeteners. *Nutr Rev* 74: 670-689, 2016.
33. Yagi K and Matsuo T: The study on long-term toxicity of d-psicose in rats. *J Clin Biochem Nutr* 45: 271-277, 2009.
34. Cho YC, Park JE, Park BC, Kim JH, Jeong DG, Park SG and Cho S: Cell cycle-dependent Cdc25C phosphatase determines cell survival by regulating apoptosis signal-regulating kinase 1. *Cell Death Differ* 22: 1605-1617, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.