

Isofraxidin enhances hyperthermia-induced apoptosis via redox modification in acute monocytic leukemia U937 cells

PENG LI¹, QING-LI ZHAO¹, MATI UR REHMAN^{1,2}, PARAS JAWAID¹, ZHENG-GUO CUI³,
KANWAL AHMED⁴, TAKASHI KONDO¹, JUN-ICHI SAITOH¹ and KYO NOGUCHI¹

¹Department of Radiology, Graduate School of Medicine and Pharmaceutical Sciences, University of Toyama, Toyama 930-0194, Japan; ²Department of Biological & Biomedical Sciences, Aga Khan University, Karachi 74800, Pakistan; ³Department of Environmental Health, University of Fukui School of Medical Sciences, Fukui 910-8505, Japan; ⁴Department of Basic Medical Sciences, College of Medicine, King Saud Bin Abdulaziz University of Health Sciences, Jeddah 9515, Kingdom of Saudi Arabia

Received January 17, 2022; Accepted September 2, 2022

DOI: 10.3892/mmr.2023.12928

Abstract. The cell-killing potential of most chemotherapeutic agents is enhanced by a temperature elevation. Isofraxidin (IF) is a coumarin compound widely found in plants, such as the Umbelliferae or Chloranthaceae families. IF induces anticancer effects in lung and colorectal cancer. To the best of our knowledge, the combined effects of hyperthermia (HT) and IF on heat-induced apoptosis have not been reported. Acute monocytic leukemia U937 cells were exposed to HT with or without IF pre-treatment. Apoptosis was measured by Annexin V-FITC/PI double staining assay using flow cytometry and cell viability was observed by cell counting kit assay, DNA fragmentation. The mechanism involved in the combination was explored by measuring changes in the mitochondrial membrane potential, (MMP), intracellular ROS generation, expression of apoptosis related protein, and intracellular calcium ion level. It was demonstrated that IF enhanced HT-induced apoptosis in U937 cells. The results demonstrated that combined treatment enhanced mitochondrial membrane potential loss and transient superoxide generation increased protein expression levels of caspase-3, caspase-8 and phosphorylated-JNK and intracellular calcium levels. Moreover, the role of caspases and JNK was confirmed using a pan caspase inhibitor (zVAD-FMK) and JNK inhibitor (SP600125) in U937 cells. Collectively, the data demonstrated that IF enhanced HT-induced apoptosis via a reactive oxygen species mediated mitochondria/caspase-dependent pathway in U937 cells.

Introduction

Isofraxidin (IF) is a coumarin compound widely found in plants, such as those in the Umbelliferae or Chloranthaceae families. It induces numerous biological effects including, anti-bacterial, antioxidant, anti-depressive and anti-inflammatory effects (1-4). However, studies have also reported that IF induces anticancer effects in lung and colorectal cancer cell lines via inhibition of the epidermal growth factor receptor and Akt signaling pathways (5,6). Moreover, IF is reported to exhibit anti-tumor effects on human hepatoma cell lines HuH-7 and HepG2 at a non-toxic concentration by inhibiting matrix metalloproteinase-7 expression and ERK1/2 phosphorylation (7).

Hyperthermia (HT) has been recognized as an effective tool to halt tumor growth, especially in combination with conventional therapies. The advantage of this type of combination therapy is the possibility of using lower doses of chemotherapy or radiation leading to more effective treatment with fewer unwanted side effects (8,9). HT induces apoptosis in cancer cells. Apoptosis is a tightly regulated cell suicide program that serves essential roles in development and maintenance of tissue homeostasis by eliminating unnecessary and harmful cells (10). Reactive oxygen species (ROS) serve an important role as intracellular mediators of HT-induced apoptosis. It has been previously reported that in acute monocytic leukemia U937 cells, HT-induced apoptosis is associated with generation of intracellular superoxide (11). Although cancer cells are sensitive to intracellular ROS, due to biological and technical difficulties, complete eradication of the tumor is usually not achieved with a single modality. Therefore, to overcome these challenges, a multimodal strategy is usually adopted in which a combination of chemotherapeutic drugs or chemotherapeutic drugs combined with a physical modality such as HT is utilized (12). An ideal chemical sensitizer for HT should be non-toxic at normal temperature but cytotoxic at hyperthermic temperatures. To the best of our knowledge, the anticancer effects of the IF + HT combination have not been reported before. Therefore, the present study evaluated the role of IF in U937 cells under HT conditions. In the present study

Correspondence to: Dr Qing-Li Zhao, Department of Radiology, Graduate School of Medicine and Pharmaceutical Sciences, University of Toyama, 2630 Sugitani, Toyama 930-0194, Japan
E-mail: zhao@med.u-toyama.ac.jp

Key words: isofraxidin, hyperthermia, apoptosis, reactive oxygen species, anti-cancer

the ability of IF to enhance HT-induced apoptosis in U937 cells was assessed and its associated mechanism was investigated.

Material and methods

Chemicals. IF (7-hydroxy-6,8-dimethoxy-coumarin) was purchased from Wako Pure Chemical Industries, Ltd. Stock solutions were prepared using dimethyl sulfoxide (DMSO) as a solvent and further dissolved to make the desired concentrations for experimental use. The JNK inhibitor SP600125 was purchased from Calbiochem Biosciences, Inc. (Merck KGaA). The pan caspase-3 inhibitor Z-VAD-FMK was purchased from Promega Corporation.

Cell culture. The human acute monocytic leukemia U937 cell line was purchased from the Human Sciences Research Resource Bank (Japan Human Sciences Foundation) and was maintained in RPMI-1640 medium (Wako Pure Chemical Corporation) supplemented with 10% heat-inactivated fetal bovine serum (SIGMA Life Science) at 37°C in humidified air with 5% CO₂.

Drug treatment and HT. A cell suspension containing 1x10⁶ cells/ml was pre-treated with IF for 1 h at 37°C. For HT treatment, 1x10⁶/ml U937 cells were transferred to plastic tubes and exposed to 37 or 44°C for 20 min by immersing tubes containing 1 ml cell suspension into a water bath. Following treatment, the cells were incubated at 37°C for 6 h.

DNA fragmentation. The quantitative DNA fragmentation assay was performed according to the method reported by Sellins and Cohen (13). Briefly, U937 cells were lysed using a lysis buffer (Tris, 10 mM; EDTA, 1 mM; 0.2% Triton X-100; pH, 7.5) and centrifuged at 13,000 x g for 10 min at 4°C. Subsequently, each DNA sample in the supernatant was precipitated by adding lysis buffer to the pellet and then added 25% TCA to both pellet and supernatant to attain 12.5% trichloroacetic acid (TCA) concentration. Incubate at 4°C overnight in the dark and quantified using diphenylamine reagent following hydrolysis in 5% TCA at 90°C for 20 min. The percentage of DNA fragmentation for each sample was calculated as follows: DNA fragmentation (%) = amount of DNA in the supernatant/total DNA in the sample x100.

Apoptosis assay. Apoptotic assay was determined by apoptotic cells percent in both early and late apoptosis. Apoptotic cells were assessed using Annexin V-FITC/propidium iodide (PI) apoptosis detection kit (Immunotech; Beckman Coulter, Inc.). Briefly, following IF pre-treatment (500 µM) for 1 h at 37°C, the U937 cells were exposed to HT (44°C; 20 min). Following 6 h incubation at 37°C, cells were collected, washed with PBS and centrifuged at 160 x g for 3 min at 4°C. The resulting pellet was mixed with binding buffer of Annexin V-FITC kit. Annexin V-FITC (5 µl) and PI (5 µl) were added to 490 µl each cell suspension, followed by gentle mixing. Following incubation at 4°C for 30 min in the dark, the cells were analyzed using flow cytometry (Epics XL Flow Cytometer; Coulter Epics XL-MCL Flow Cytometer System II Software, version 3.0, Beckman-Coulter, Inc.

Cell counting assay. The cell counting assay was performed using a Cell Counting Kit-8 (CCK-8; Dojindo Laboratories, Inc.) assay according to the manufacturer's protocol. Briefly, following IF pre-treatment (500 µM) for 1 h at 37°C, cells were exposed to HT (44°C; 20 min). Following 6 or 24 h incubation at 37°C, cells were incubated in 100 µl RPMI medium (containing 10 µl CCK-8 reagent) in a 96-well plate and incubated for 2 h at 37°C in 5% CO₂. The absorbance at 450 nm was assessed using an iMark™ Microplate Absorbance Reader (Bio-Rad Laboratories, Inc.).

Cell cycle analysis. Following the drug alone or in combination treatment 1x10⁶ cells/ml were harvested and washed with Dulbecco's phosphate-buffered saline (PBS) and fixed in 10 ml 70% cold ethanol overnight at 4°C. After washing with PBS, cells were incubated with RNase solution (0.25 mg/ml; Wako Pure Chemical Industries, Ltd.) containing PI (50 µg/ml; Wako Pure Chemical Industries, Ltd.) at 37°C for 30 min in the dark. The percentage of subG1 cells was assessed using flow cytometry, as aforementioned. A total of 10,000 events was analyzed for each sample using an Epics XL Flow Cytometer, Coulter Epics XL-MCL Flow Cytometer System II Software, version 3.0, Beckman-Coulter, Inc.

Morphological changes of apoptosis. The morphological changes in the cells both treated and untreated were evaluated using Giemsa staining. The cells were harvested following 6 h incubation at 37°C by centrifugation at 160 x g for 3 min at 4°C and washed with PBS. Then cells were fixed with methanol and acetic acid (3:1) at 4°C, 60 min and spread on glass slides. The slides were incubated for 60 min at room temperature. After drying, staining was performed using 5% Giemsa solution (pH 6.8) for 10 min at room temperature. Cells were observed using an Olympus IX70 light microscope (Olympus Corporation) at a magnification of x400. Cells that demonstrated shrinkage and nuclear changes (condensation, fragmentation and apoptotic bodies) were considered to be apoptotic cells (14).

Assessment of intracellular ROS production. To assess intracellular ROS production, both treated and untreated U937 cells were incubated at 37°C for 15 min with 5 µM hydroxyphenyl fluorescein (HPF; Sekisui Medical Co., Ltd.) to assess intracellular hydroxyl radical and peroxynitrite levels and 5 µM hydroethidine (HE; Molecular Probes; Thermo Fisher Scientific, Inc.) to detect intracellular superoxide levels. The fluorescence emission of all probes was analyzed using flow cytometry as aforementioned. A total of 10,000 events was analyzed for each sample using an Epics XL Flow Cytometer, Coulter Epics XL-MCL Flow Cytometer System II Software, version 3.0, Beckman-Coulter, Inc.

Evaluation of mitochondrial membrane potential (MMP). To assess changes in MMP, acute monocytic leukemia U937 cells were stained with 10 nM tetramethylrhodamine methyl ester (TMRM; Molecular Probes; Thermo Fisher Scientific, Inc.) for 15 min at 37°C in PBS. The fluorescence of red TMRM was analyzed using flow cytometry (excitation, 488 nm; emission, 575 nm), as aforementioned. The percentage of low-MMP cells was determined from cell counts in the 0.1-12.1 low

window of the TMRM log scale. A total of 10,000 events was analyzed for each sample using an Epics XL Flow Cytometer, Coulter Epics XL-MCL Flow Cytometer System II Software, version 3.0, Beckman-Coulter, Inc.

Western blotting. Both treated and untreated U937 cells were collected and lysed in lysis buffer [Tris-HCl, 1 M; NaCl, 5 M; 1% Nonidet P-40 (v/v); 1% sodium deoxycholate; 0.05% SDS; phenylmethylsulfonyl fluoride, 1 mM] for 20 min at 4°C. After sonication at 20 kHz for 5 sec, the lysates were centrifuged at 12,000 x g for 10 min at 4°C and the protein content in the supernatant was assessed using a Bio-Rad Protein Assay kit (Bio-Rad Laboratories, Inc.). The protein lysates were denatured at 96°C for 5 min; after mixing with SDS-loading buffer, 20–30 µg protein/lane was loaded on a 5–20% SDS polyacrylamide gel (SuperStep™ Ace) for electrophoresis and transferred to nitrocellulose membranes. Blots were blocked using 5% skimmed milk for 1 hr at room temperature. Western blotting was performed using primary antibodies as follows: Anti-Bax polyclonal antibody (pAb) (1:1,000; cat. no. 2772; Cell Signaling Technology, Inc.), anti-caspase-3 pAb (1:1,000; cat. no. 9662; Cell Signaling Technology, Inc.), anti-Bcl-xL pAb (1:1,000, cat. no. 2762; Cell Signaling Technology, Inc.), anti-JNK monoclonal antibody (mAb) (1:1,000, cat. no. 9255; Cell Signaling Technology, Inc.), anti-phospho-JNK mAb (1:1,000; cat. no. 9255; Cell Signaling Technology, Inc.), anti-phospho-p38 mAb (1:1,000; cat. no. 9211; Cell Signaling Technology, Inc.), anti-p38 pAb (1:1,000; cat. no. sc-728 Santa Cruz Biotechnology, Inc.). All primary antibodies were incubated overnight at 4°C. Anti-β-actin mAb (1:1,000; cat. no. A5441; MilliporeSigma) was incubated for 1 hr at room temperature. Horseradish peroxidase (HRP) conjugated anti-rabbit IgG (1:1,000, cat. no. 7074; Cell Signaling Technology, Inc.) and HRP conjugated anti-mouse IgG (1:1,000; cat. no. 7076; Cell Signaling Technology, Inc.) were used as secondary antibodies, incubated for 1 hr at room temperature.

Band signals were visualized using a luminescent image analyzer (LAS4000, Fujifilm Co.) using ECL system (Amersham; Cytiva). Band density was semi-quantified using Image J software (ver 1.5.3; National Institutes of Health).

For the preparation of cytosolic and mitochondrial fractions, both untreated and treated U937 cells were collected, washed with PBS and suspended in 300 µl extraction buffer (D-mannitol, 210 mM; sucrose, 70 mM; EDTA, 5 mM; aprotinin, 1 µg/ml; pepstatin, 1 µg/ml; leupeptin, 1 µg/ml; PMSF, 1 mM; HEPES-KOH, 10 mM; pH 7.4). Following 5 min incubation on ice, cells were homogenized and centrifuged at 1,500 x g for 5 min at 4°C to remove nuclei and debris. The supernatant was centrifuged at 105,000 x g for 30 min at 4°C. The resulting supernatant was used as the soluble cytosolic fraction (S100) and the pellet was resolved in mitochondrial buffer as the mitochondrial fraction. Blots were blocked using 5% skimmed milk for 1 hr at room temperature. Western blotting was performed to detect cytochrome c and Bax released to the cytosol and mitochondria using anti-cytochrome c pAb (1:1,000, cat. no. 4272; Cell Signaling Technology, Inc.), anti-Bax pAb (1:1,000, cat. no. 2762; Cell Signaling Technology, Inc.), anti-β-actin mAb (1:1,000; cat. no. A5441; MilliporeSigma) and anti-HSP60 mAb (1:1,000; cat.

no. 611562; BD Biosciences) primary antibodies. All primary antibodies were incubated overnight at 4°C except anti-β-actin mAb, incubated for 1 h at room temperature.

Evaluation of intercellular free calcium ions. To assess the effect of IF treatment on intracellular calcium homeostasis, intracellular free Ca²⁺ levels were assessed using Fluo-3/AM (Dojindo Laboratories Inc.) calcium probe. Cells were treated with IF (500 µM) and incubated for 1 h at 37°C, and then exposed to 44°C for 20 min. Following 6 h incubation at 37°C, the cells were harvested and incubated with 5 µM Fluo-3/AM for 30 min at 37°C. Excess Fluo-3/AM was removed by washing three times with PBS. The fluorescence intensity was assessed using Epics XL Flow Cytometer, Coulter Epics XL-MCL Flow Cytometer System II Software, version 3.0, Beckman-Coulter, Inc.

Statistical analysis. Data are presented as the mean ± standard deviation. Statistical analysis was performed using one-way ANOVA followed by Tukey's multiple comparison test using GraphPad Prism version 5 (GraphPad Software, Inc.). P<0.05 was considered to indicate a statistically significant difference. All experiments were performed in triplicate.

Results

Effect of IF on HT-induced apoptosis. U937 cells were treated with HT exposure with or without IF pre-treatment. DNA fragmentation, cell survival and cell cycle assay were performed to evaluate the effects of IF on HT-induced apoptosis in U937 cells. The DNA fragmentation assay demonstrated that IF markedly enhanced HT-induced apoptosis in a dose-dependent manner (Fig. 1A). IF at a dose of 200 and 500 µM induced significant increase in DNA fragmentation % compared to control. However, 500 µM was used for subsequent experiments because it demonstrated the greatest significant difference in DNA fragmentation following HT exposure compared with 0 µM IF control. The cell survival assay demonstrated that IF in combination with HT significantly decreased cell survival percentage following 6 h incubation compared with the HT group (Fig. 1B). The fraction of cells in subG1 also significantly increased following combined treatment compared with the HT group (Fig. 1C). Similarly, a marked increase in typical apoptotic features, such as cytoplasmic aggregation, nuclear condensation and fragmentation were observed in the combined treatment cells than IF and HT alone treatment, as assessed using Giemsa staining (Fig. 1D). Taken together, the data indicated that IF markedly sensitized U937 cells to HT exposure.

Effect of IF on HT-induced Annexin V-FITC/PI double staining. To assess the effect of combined IF/HT treatment *in vitro*, U937 cells were incubated with or without IF for 1 h and then subjected to HT with subsequent recovery at 37°C for 6 h. Cell death (early apoptosis and late apoptosis) was assessed using flow cytometry via annexin V-FITC/PI double staining. The IF treatment alone did not induce apoptosis compared with control. HT treatment alone markedly increased the proportion of apoptotic cells compared with control and IF treatment alone, however when cells were treated with IF + HT, the




Figure 1. IF increases HT-induced apoptosis in U937 cells. (A) Cells were treated with HT (44°C; 20 min) with or without IF (50, 100, 200 and 500 μM) pre-treatment for 1 h and incubated at 37°C. DNA fragmentation assay was performed after 6 h. * $P < 0.05$. (B) Cells were pre-treated with 500 μM IF for 1 h and then exposed to HT (44°C; 20 min). Cell survival analysis was performed using Cell Counting Kit-8 assay after 6 h. ** $P < 0.01$. (C) Cells were collected after incubation for 6 h and incubated with RNase solution and propidium iodide at 37°C for 30 min. The percentage of sub-G1 cells was determined using flow cytometry. The results are presented as mean \pm SD ($n=3$); significance was assessed using one-way ANOVA with Tukey's multiple comparison test. *** $P < 0.001$. (D) Apoptotic features of U937 cells in response to the combined treatment with HT and IF were assessed using Giemsa staining after 6 h incubation and evaluated under a microscope at x400 magnification. Scale bar 10 μm. IF, isofraxidin; HT, hyperthermia.

proportion of apoptotic U937 cells was significantly increased compared with either treatment alone (Fig. 2A and B).

Effect of IF on HT-induced MMP loss and ROS generation. It has been reported that apoptosis involves a disruption of mitochondrial membrane integrity, which is key for the cell death process (15,16). To determine whether MMP change was involved in IF-enhanced apoptosis, the change in the MMP was assessed 6 h after HT exposure. Loss of MMP was markedly induced by HT and was significantly elevated in the presence of IF at 6 h compared with the HT group (Fig. 3A).

Excess generation of intracellular ROS causes oxidative stress, which serves a key role in the induction of apoptosis (17,18). To assess whether IF increased oxidative stress induced by HT, intracellular ROS generation was evaluated using HE and HPF staining. The results demonstrated that the intensity of HE fluorescence increased markedly in HT-treated cells at 30 min than untreated control and IF treatment alone, which was further significantly enhanced in the combination-treated cells compared with the HT group (Fig. 3B). Furthermore, the intensity of HPF fluorescence was also markedly elevated at 30 min after HT exposure than untreated control and IF treatment, and was significantly enhanced following combination treatment with IF and HT compared with the HT group (Fig. 3C).

Effect of IF on intracellular Ca^{2+} levels. Calcium homeostasis is essential for cellular functions, such as protein folding, processing and transport and signal transduction (19). Increased levels of Ca^{2+} overexcite cells and cause the generation of harmful chemicals such as free radicals (20). Therefore, intracellular Ca^{2+} levels were assessed following treatment. It was demonstrated that the combined treatment significantly enhanced intracellular Ca^{2+} concentration compared with either treatment alone (Fig. 3D).

Effect of IF on HT-induced caspase activation. Caspases, a family of cysteine proteases, are activated in the execution phase of the apoptotic process. Once activated, caspases activate downstream caspases, leading to apoptosis (21). To assess protein expression levels of caspases western blotting was performed. Cells were pre-treated with IF for 1 h, followed by HT exposure and then collected for protein extraction. The results demonstrated that protein expression levels of cleaved caspase-3 and cleaved caspase-8 were markedly increased in the combined treatment cells than HT treatment alone (Fig. 4A). To verify the role of caspases, DNA fragmentation is performed using the pan caspase inhibitor Z-VAD-FMK. The results demonstrated that both HT and combined treatment-mediated apoptosis was significantly suppressed in




Figure 2. IF enhances HT-induced early apoptosis and secondary necrosis. (A) Representative flow cytometry dot plots of annexin V-FITC and PI-stained cells. (B) The cells were treated with HT with or without IF. The percentage of cells in early apoptosis and secondary necrosis were analyzed using flow cytometry 6 h after combined treatment. The results were presented as the mean \pm SD (n=3), significance was assessed using one-way ANOVA with Tukey's multiple comparison test. ***P<0.001 control and IF vs HT. . IF, isofoxadidin; HT, hyperthermia; PI, propidium iodide.




Figure 3. IF enhances HT-induced MMP loss and reactive oxygen species generation in U937 cells. Cells were pre-treated with 500 μ M IF for 1 h and then treated with HT (44°C; 20 min). (A) Following 6 h incubation, cells were collected and analyzed using flow cytometry using tetramethylrhodamine methyl ester staining. (B) Following 30 min incubation, cells were stained using HE for 15 min. Fluorescence intensity was detected using flow cytometry. (C) Cells were stained with HPF for 15 min. Fluorescence intensity was detected using flow cytometry. (D) IF promotes HT-elevated intracellular Ca²⁺ levels in U937 cells. Cells were harvested and stained using 5 μ M Fluo-3/AM for 30 min and intracellular Ca²⁺ levels were assessed using flow cytometry. The results are presented as mean \pm SD (n=3); significance was assessed using one-way ANOVA with Tukey's multiple comparison test. *P<0.05, ***P<0.001 HT vs. HT + IF. **P<0.01 IF vs. HT. HE, hydroethidine; HPF, hydroxyphenyl fluorescein; MMP, mitochondrial membrane potential; IF, isofoxadidin; HT, hyperthermia.

the presence of the inhibitor compared with the uninhibited group, which suggested that caspase cascades were involved

in IF-mediated apoptotic enhancement in response to HT exposure (Fig. 4B).




Figure 4. IF promotes HT-induced apoptosis via a caspase-dependent pathway in U937 cells. (A) Cells were pre-treated with 500 μ M IF for 1 h and then treated with HT (44°C; 20 min). Following 6 h incubation, protein was extracted from cells and western blotting was performed to assess protein expression levels of caspase-3 and caspase-8. (B) Cells were pre-treated with 10 μ M zVAD-FMK, a pan-caspase inhibitor, for 1 h and exposed to HT (44°C; 20 min) in the presence of IF (500 μ M). DNA fragmentation was performed following 6 h incubation. The results are presented as mean \pm SD (n=3); significance was assessed using one-way ANOVA with Tukey's multiple comparison test. * P <0.05 HT vs. HT + zVAD-FMK, *** P <0.001 IF + HT vs. IF + HT + zVAD-FMK. IF, isofraxidin; HT, hyperthermia.

Effect of IF treatment on apoptosis-associated protein. To assess the molecular mechanism of IF in response to HT treatment, western blotting was performed. Bcl-2 family proteins with anti- or pro-apoptotic functions control release of mitochondrial apoptosis factors including cytochrome c and apoptosis-inducing factor (22). Markedly decreased Bcl-xL protein expression levels were demonstrated 6 h following combined treatment compared to either treatment alone (Fig. 5A). No changes in protein expression levels of Bax were demonstrated in whole cell lysate 6 h after HT or combined treatment, however translocation of Bax from cytosol to mitochondria was observed in HT-treated cells and further augmented in combined treatment cells (Fig. 5B). Western blotting also demonstrated that IF markedly enhanced release of cytochrome c from mitochondria to the cytosol compared to control and IF treatment alone, which was induced by HT treatment. Furthermore, induction of heat shock protein is the most well-characterized heat shock response. HSP 70 has been reported to serve a critical role in cell survival and thermotolerance, potentially by preventing recruitment of procaspase-9 to the apoptosome by binding to apoptotic protein activation factor-1 (Apaf-1), thereby inhibiting activation of caspase-3 (23). Therefore, expression of HSP 70 was also assessed using western blotting. HT-induced HSP70 expression was not significantly affected by combined treatment. These results indicated that a mitochondria-dependent pathway was involved in IF-mediated enhancement of apoptosis induced by HT treatment.

Effect of IF on MAPK-associated pathway. Activation of MAPKs serves a key role in apoptosis induced by cellular stresses and oxidative stress is known to activate MAPK family members by protein phosphorylation (24). To assess whether the MAPK signaling pathway was involved in IF-mediated

enhancement of apoptosis, protein expression levels of JNK and p38 were assessed using western blotting at 1 h after HT or combined IF + HT treatment. The results demonstrated that JNK phosphorylation was markedly induced at 1 h following HT treatment compared with the untreated control and IF, and substantially further, markedly elevated by IF + HT compared with the HT group (Fig. 6A). P38 phosphorylation was also markedly induced at 1 h after HT treatment; however, IF did not significantly promote phosphorylation of P38. The protein expression levels of JNK and p38 remained unchanged in cells with HT and/or IF treatment. To evaluate the role of JNK signaling pathway in IF-mediated enhancement of apoptosis induced by HT, a JNK inhibitor, SP10025, was used. The inhibitor did not suppress HT-induced apoptosis; however, it partially suppressed HT + IF-induced apoptosis, which indicated that JNK was, in part, involved in IF-mediated apoptotic enhancement following HT treatment (Fig. 6B).

Discussion

Heat stress induces formation of ROS, such as superoxide anions ($O_2^{\cdot-}$), hydrogen peroxide (H_2O_2), hydroxyl radicals ($\cdot OH$), nitric oxide ($\cdot NO$) and peroxynitrite ($ONOO^{\cdot-}$) by cells. Several studies have reported a link between oxidative stress and heat stress and synergistic augmentation of cell death as increased ROS generation is observed in HT-exposed cells (25-27). $O_2^{\cdot-}$ is the precursor of most ROS, a mediator in oxidative chain reactions and react with other radicals, including $\cdot NO$ (28) to form $ONOO^{\cdot-}$, which is also a potent oxidant (29). HE indicates the presence of superoxide anions and HPF indicates the presence of $\cdot OH$ and $ONOO^{\cdot-}$. Therefore, in the present study, both HE and HPF staining were used to assess the effect of IF on HT-induced ROS formation. The intensity of HE and HPF fluorescence was significantly increased in the presence




Figure 5. IF elevates HT-induced apoptosis in U937 cells via the mitochondrial pathway. Cells were pre-treated with 500 μ M IF for 1 h and exposed to HT (44°C; 20 min). Following 6 h incubation, cells were collected for use in experiments. (A) Western blotting demonstrated changes in expression levels of cell death-associated proteins Bax, Bcl-xL and HSP 70. (B) Changes in protein expression levels of cytochrome c and Bax in the Mito and S100 fractions. Band density was semi-quantified using Image J software (ver 1.5.3) and presented as fold-change vs. control (untreated). Semi-quantified data are presented as mean \pm SD (n=3); significance was assessed using one-way ANOVA with Tukey's multiple comparison test. **P<0.01 HT vs. HT + IF (Bcl-XL, Cyto c) and ***P<0.001 HT vs. HT + IF (HSP70, Cyto c and Bax). HSP70, heat shock protein 70; Mito, mitochondrial; S100, cytosolic; IF, isofradixin; HT, hyperthermia; Cyt c, Cytochrome c.

of IF following HT treatment compared with the HT-alone group. IF was previously reported to protect U937 cells against radiation-induced apoptosis via scavenging of \cdot OH generation (3). The reason for this discrepancy in IF effects may be because in irradiated cells, \cdot OH radicals are generated directly and transiently and IF quenches radiation-induced \cdot OH production, immediately suppressing radiation-induced apoptosis. However, HT generates $O_2^{\cdot-}$ heterogeneously and chronically and does not directly generate \cdot OH production within cells (30). SOD-mimic nitroxides have been reported to cause dismutation of $O_2^{\cdot-}$ in the presence of H^+ and produce H_2O_2 and O_2 by the subsequent chemical reaction *in vitro* (30). Moreover, H_2O_2 participates in the Fenton reaction in the presence of transition metals, particularly Fe^{2+} , to release \cdot OH (31). In the present study, IF transiently increased the intensity of

HE and HPF fluorescence 30 min after HT treatment. This transient elevation of ROS by IF may have resulted in apoptotic enhancement in response to HT. Furthermore, HT has an advantage to sensitize cells because cells in late S phase, which are resistant to X-rays, are most heat sensitive, cells deficient in nutrients are heat sensitive and cells subject to chronic hypoxia demonstrate slightly enhanced heat sensitivity (12). HT also changes cellular status, such as causing increases in membrane fluidity and permeability, alteration of protein structure and function and, increase in cellular metabolism. HT is also selective towards cancer cells; our previous studies demonstrated that HT is well-tolerated by non-tumor cells (12,32,33). Therefore, the cell killing potential of most chemotherapeutic agents may be substantially enhanced by temperature elevation.




Figure 6. Role of IF in the MAPK signaling pathway in HT-treated U937 cells. (A) Cells were pre-treated with 500 μ M IF for 1 h and exposed to HT (44°C; 20 min). Following 1 h incubation, cells were collected for use in experiments. Changes in expression levels of MAPK signaling pathway-associated proteins such as p-JNK, JNK, p-p38 and p38 were assessed using western blotting. Band density was semi-quantified using Image J software and were presented as fold change versus control (untreated). Semi-quantified data was presented as mean \pm SD (n=3). *P<0.05 control vs. HT and IF vs. HT (pJNK/JNK), **P<0.01 control vs. HT + IF and IF vs. HT + IF (pJNK/JNK). *P<0.05 control vs. HT and IF vs. HT (p-p38/p38), **P<0.01 control vs. HT + IF and IF vs. HT + IF (p-p38/p38). (B) Cells were pre-treated with 10 μ M SP600125 for 1 h, then exposed to HT (44°C; 20 min) with IF (500 μ M). DNA fragmentation was performed following 6 h incubation. The results were presented as mean \pm SD (n=3); significance was assessed using one-way ANOVA with Tukey's multiple comparison test. *P<0.05 HT vs. HT + IF, **P<0.01 HT + SP600125 vs. HT + IF, ***P<0.001 HT vs IF and SP600125 (either treatment alone or in combination). IF, isofraxidin; HT, hyperthermia; p, phosphorylated.




Figure 7. Isofraxidin enhances hyperthermia-induced apoptosis via a ROS mediated mitochondria-caspase-dependent signaling pathway in U937 cells. ROS, reactive oxygen species; MMP, mitochondrial membrane potential; Cyt c, cytochrome c; p, phosphorylated.

The apoptotic process occurs via two distinct yet tightly interconnected pathways: Intrinsic pathway (mitochondrial) and extrinsic pathway (death receptors). There are specific proteins in each of these pathways. The key step of the intrinsic apoptosis pathway is permeabilization of the mitochondrial outer membrane (MOMP), which results in MMP loss, then release of certain apoptogenic factors from mitochondria. Cytochrome c, one such protein, binds to the adapter Apaf-1, which recruits cytosolic pro-caspase-9 into a heptameric complex, called the apoptosome. Active caspase-9 cleaves and activates effector caspase-3 and caspase-7 (34,35). In the present study, HT-induced MMP loss was further promoted in IF + HT combination treatment cells. Moreover, the release of cytochrome c from mitochondria to cytosol was clearly observed in HT-treated cells and expression of cytosolic cytochrome c was further enhanced in IF + HT combination treatment cells. In the extrinsic apoptosis pathway, stimulation of death receptors such as CD95 (APO-1/Fas) and TNF-associated apoptosis-inducing ligand receptors results in formation of the death-inducing signaling complex, following activation of the initiator caspase-8. Activated caspase-8 propagates the apoptotic signal either by direct activation of downstream

effector caspases such as caspase-3 or by truncation of BH3 interacting-domain death agonist (Bid) (36). Bid assists in activation and translocation of Bax to the mitochondria, thus triggering the intrinsic pathway (37). In the present study, protein expression levels of cleaved caspase-3 and caspase-8 were induced in HT-treated cells and markedly enhanced by IF pre-treatment following HT exposure. Taken together, these data demonstrated that IF may elevate HT-induced apoptosis in a caspase-dependent mitochondrial signaling pathway.

MOMP leads to both caspase-dependent and -independent cell death signaling and is therefore a key cell life-or-death decision checkpoint. This checkpoint is controlled by the Bcl-2 family of proteins (30). This family is subdivided into three key groups on the basis of their function: Pro-apoptotic BH3-only (Noxa, Bid, Bim, p53 upregulated modulator of apoptosis, Bad, Bcl-2-modifying factor, Hrk and Bcl-2-interacting killer), pro-survival (Mcl-1, Bcl-2, Bcl-xL) and pore-forming proteins (Bax and Bcl-2 homologous antagonist/killer). In the apoptotic process, large amounts of Bax/Bak are bound by Bcl-2 survival factors, BH3-only proteins displace large numbers of Bax/Bak molecules from Bcl-2 and therefore increase the number of pore-forming proteins available for activation of MOMP (38,39). The mitochondrial translocation of Bax is key to the release of cytochrome c during the apoptotic process (40,41). In the present study, the protein expression levels of Bcl-xL were significantly decreased in IF + HT combination treatment cells compared with the HT group. The protein expression levels of Bax in the total cell lysate was not significantly affected by IF in response to HT; however, translocation of Bax from the cytosol to the mitochondria was observed in HT-treated cells and IF + HT combination-treated cells. The data demonstrated that IF pre-treatment potentiated HT-triggered translocation of Bax. Taken together, these data demonstrated that IF enhanced HT-induced apoptosis by decreasing protein expression levels of anti-apoptotic Bcl-xL and increasing translocation of Bax and the release of cytochrome c.

MAPKs are a family of protein kinases that are responsible for transduction of intracellular signaling. They also participate in a range of physiological regulatory activities. Activation of MAPK signaling pathways, including JNK, p38 and ERK, serves a key role in modifying cell death induced by cellular stresses. It has been reported that stress-activated JNK and p38 kinase phosphorylate Bax; phosphorylation of Bax leads to mitochondrial translocation of activated Bax and initiation of mitochondria-dependent apoptosis in cells treated with cell death stimulants (42). To assess the molecular mechanism of IF in response to HT-induced apoptosis, protein expression levels of JNK and p38 were assessed. IF + HT combination treatment markedly increased phosphorylation levels of JNK and p38 compared with the HT group. JNK activation has been reported to serve an essential role in HT-induced apoptosis (43). To evaluate the role of JNK in IF-mediated apoptotic enhancement, a JNK inhibitor (SP600125) was used in U937 cells. The JNK inhibitor markedly decreased IF-mediated apoptotic enhancement following HT treatment, indicating that the JNK pathway may be involved in the enhancement of apoptosis by IF.

Ca²⁺ is a second messenger in cells that translates extracellular stimuli into intracellular activities, which are important for the regulation of physiological processes (44). Excessive elevation or prolonged activation of Ca²⁺ signaling

causes cell death (45,46). It has been reported that oxidative stress and intracellular Ca²⁺ overload are connected in HT-mediated cell death (45,46). It is well established that intracellular Ca²⁺ overload triggers apoptosis via the mitochondria-dependent pathway (47). The mitochondria-dependent apoptosis pathway involves numerous events, such as generation of ROS, the opening of permeability transition pores, MMP loss, release of cytochrome c, expression of Bcl-2 family members and activation of caspase-9 and caspase-3 (48,49). Moreover, calcium increase is secondary to oxidative stress-induced mitochondrial damage. Therefore, the association between increased calcium release and ER stress should be part of future study of the combination of drugs and HT. In the present study, the data supported the classic mitochondria-dependent apoptotic pathway, which was promoted by HT with or without IF pre-treatment in U937 cells. The enhanced intracellular Ca²⁺ overload in HT + IF-treated cells triggered apoptosis via the mitochondria-dependent pathway.

To the best of our knowledge, the present study is the first to demonstrate the effect of IF on HT-induced apoptosis. These findings suggested that IF significantly enhanced HT-induced apoptosis via caspase-mitochondria dependent pathways such as loss of MMP, excessive ROS generation, mitochondrial translocation of Bax, release of cytochrome c and caspase-3 activation. HT was not only cytotoxic to cancer cells but also enhanced fluidity and permeability of the plasma membrane, which promotes drug delivery (50). Therefore, increased IF delivery into cells by HT may be a possible mechanism underlying the enhancement of apoptosis demonstrated by combined treatment (Fig. 7). In the present study, data were collected mainly using U937 cells; lack of evaluation using a normal cell line is as a limitation of the present study. However, it is important to note that a previous study also reported that IF exhibits low toxicity towards BEAS-2B normal lung epithelial cells compared with the lung cancer A549 cell line (6). Furthermore, our previous study also demonstrated that IF serves as an antioxidant and protects cells against radiation-induced cell death (3). These reports suggested that IF does not induce cytotoxic effects in normal cells. These data suggested that IF might be an effective thermo-sensitizer for cancer therapy.

Acknowledgements

Not applicable.

Funding

The present study was supported by JSPS KAKENHI (grant no. JP21K07614).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

Hypotheses and experiments were conceived by QLZ. Experiments were performed by PL, MUR, PJ, KA, ZGC and QLZ. Data analysis and manuscript preparation were

performed by TK, JIS, KN, MUR and QLZ. QLZ and PL confirm the authenticity of all the raw data. All authors have read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Deyama T, Nishibe S and Nakazawa Y: Constituents and pharmacological effects of *Eucommia* and Siberian ginseng. *Acta Pharmacol Sin* 22: 1057-1070, 2001.
- Khan S, Riaz N, Afza N, Malik A, Aziz ur R, Iqbal L and Lateef M: Antioxidant constituents from *Cotoneaster racemiflora*. *J Asian Nat Prod Res* 11: 44-48, 2009.
- Li P, Zhao QL, Wu LH, Jawaid P, Jiao YF, Kadowaki M and Kondo T: Isofraxidin, a potent reactive oxygen species (ROS) scavenger, protects human leukemia cells from radiation-induced apoptosis via ROS/mitochondria pathway in p53-independent manner. *Apoptosis* 19: 1043-1053, 2014.
- Niu X, Xing W, Li W, Fan T, Hu H and Li Y: Isofraxidin exhibited anti-inflammatory effects in vivo and inhibited TNF- α production in LPS-induced mouse peritoneal macrophages in vitro via the MAPK pathway. *Int Immunopharmacol* 14: 164-171, 2012.
- Shen P, Wang HG, Li MM, Ma QY, Zhou CW, Pan F and Xie R: Isofraxidin inhibited proliferation and induced apoptosis via blockage of Akt pathway in human colorectal cancer cells. *Biomed Pharmacother* 92: 78-85, 2017.
- Zhang H, Feng QQ, Gong JH and Ma JP: Anticancer effects of isofraxidin against A549 human lung cancer cells via the EGFR signaling pathway. *Mol Med Rep* 18: 407-414, 2018.
- Yamazaki T and Tokiwa T: Isofraxidin, a coumarin component from *Acanthopanax senticosus*, inhibits matrix metalloproteinase-7 expression and cell invasion of human hepatoma cells. *Biol Pharm Bull* 33: 1716-1722, 2010.
- Datta NR, Ordonez SG, Gaip US, Paulides MM, Crezee H, Gellermann J, Marder D, Puric E and Bodis S: Local hyperthermia combined with radiotherapy and/or chemotherapy: Recent advances and promises for the future. *Cancer Treat Rev* 41: 742-753, 2015.
- Mantso T, Vasileiadis S, Anastopoulos I, Voulgaridou GP, Lampri E, Botaitis S, Kontomanolis EN, Simopoulos C, Goussestis G, Franco R, *et al*: Hyperthermia induces therapeutic effectiveness and potentiates adjuvant therapy with non-targeted and targeted drugs in an in vitro model of human malignant melanoma. *Sci Rep* 8: 10724, 2018.
- Plati J, Bucur O and Khosravi-Far R: Apoptotic cell signaling in cancer progression and therapy. *Integr Biol (Camb)* 3: 279-296, 2011.
- Zhao QL, Fujiwara Y and Kondo T: Mechanism of cell death induction by nitroxide and hyperthermia. *Free Radic Biol Med* 40: 1131-1143, 2006.
- Tabuchi Y, Ahmed K and Kondo T: Induction of oxidative stress by hyperthermia and enhancement of hyperthermia-induced apoptosis by oxidative stress modification. In: *Hyperthermic Oncology from Bench to Bedside*. Springer Singapore, Singapore, pp 7-18, 2016.
- Sellins KS and Cohen JJ: Gene induction by gamma-irradiation leads to DNA fragmentation in lymphocytes. *J Immunol* 139: 3199-3206, 1987.
- Zhao QL, Fujiwara Y and Kondo T: Synergistic induction of apoptosis and caspase-independent autophagic cell death by a combination of nitroxide Tempo and heat shock in human leukemia U937 cells. *Apoptosis* 15: 1270-1283, 2010.
- Antonsson B, Montessuit S, Sanchez B and Martinou JC: Bax is present as a high molecular weight oligomer/complex in the mitochondrial membrane of apoptotic cells. *J Biol Chem* 276: 11615-11623, 2001.
- Narvaez CJ and Welsh J: Role of mitochondria and caspases in vitamin D-mediated apoptosis of MCF-7 breast cancer cells. *J Biol Chem* 276: 9101-9107, 2001.
- Ott M, Gogvadze V, Orrenius S and Zhivotovsky B: Mitochondria, oxidative stress and cell death. *Apoptosis* 12: 913-922, 2007.
- Tripathi P and Hildeman D: Sensitization of T cells to apoptosis-a role for ROS? *Apoptosis* 9: 515-523, 2004.
- Zhao S, Xiong Z, Mao X, Meng D, Lei Q, Li Y, Deng P, Chen M, Tu M, Lu X, *et al*: Atmospheric pressure room temperature plasma jets facilitate oxidative and nitrate stress and lead to endoplasmic reticulum stress dependent apoptosis in HepG2 cells. *PLoS One* 8: e73665, 2013.
- Emam H, Zhao QL, Furusawa Y, Refaat A, Ahmed K, Kadowak M and Kondo T: Apoptotic cell death by the novel natural compound, cinobufotalin. *Chem Biol Interact* 199: 154-160, 2012.
- McArthur K and Kile BT: Apoptotic caspases: Multiple or mistaken identities? *Trends Cell Biol* 28: 475-493, 2018.
- Tsujimoto Y and Shimizu S: Bcl-2 family: Life-or-death switch. *FEBS Lett* 466: 6-10, 2000.
- Beere HM, Wolf BB, Cain K, Mosser DD, Mahboubi A, Kuwana T, Taylor P, Morimoto RI, Cohen GM and Green DR: Heat-shock protein 70 inhibits apoptosis by preventing recruitment of procaspase-9 to the Apaf-1 apoptosome. *Nat Cell Biol* 2: 469-475, 2000.
- Lu TH, Hsieh SY, Yen CC, Wu C, Chen KL, Hung DZ, Chen CH, Wu CC, Su YC, Chen YW, *et al*: Involvement of oxidative stress-mediated ERK1/2 and p38 activation regulated mitochondria-dependent apoptotic signals in methylmercury-induced neuronal cell injury. *Toxicol Lett* 204: 71-80, 2011.
- Gu ZT, Li L, Wu F, Zhao P, Yang H, Liu YS, Geng Y, Zhao M and Su L: Heat stress induced apoptosis is triggered by transcription-independent p53, Ca(2+) dyshomeostasis and the subsequent Bax mitochondrial translocation. *Sci Rep* 5: 11497, 2015.
- McAnulty SR, McAnulty L, Pascoe DD, Gropper SS, Keith RE, Morrow JD and Gladden LB: Hyperthermia increases exercise-induced oxidative stress. *Int J Sport Med* 26: 188-192, 2005.
- Skibba JL, Powers RH, Stadnicka A, Cullinane DW, Almagro UA and Kalbfleisch JH: Oxidative stress as a precursor to the irreversible hepatocellular injury caused by hyperthermia. *Int J Hyperthermia* 7: 749-761, 1991.
- Turrens JF: Mitochondrial formation of reactive oxygen species. *J Physiol* 552: 335-344, 2003.
- Beckman JS and Koppenol WH: Nitric oxide, superoxide, and peroxynitrite: The good, the bad, and the ugly. *Am J Physiol* 271: C1424-C1437, 1996.
- Krishna MC, Russo A, Mitchell JB, Goldstein S, Dafni H and Samuni A: Do nitroxide antioxidants act as scavengers of O₂⁻ or as SOD mimics? *J Biol Chem* 271: 26026-26031, 1996.
- Uchiyama H, Zhao QL, Hassan MA, Andocs G, Nojima N, Takeda K, Ishikawa K, Hori M and Kondo T: EPR-spin trapping and flow cytometric studies of free radicals generated using cold atmospheric argon plasma and X-ray irradiation in aqueous solutions and intracellular milieu. *PLoS One* 10: e0136956, 2015.
- Rohan M, Rehman MU, Zhao QL, Jawaid P, Mitsuhashi Y, Sakurai K, Heshiki W, Ogawa R, Tomihara K, Saitoh JI, *et al*: Combination of 5-aminosalicylic acid and hyperthermia synergistically enhances apoptotic cell death in HSC-3 cells due to intracellular nitric oxide/peroxynitrite generation. *Cancer Lett* 451: 58-67, 2019.
- Rohan M, Rehman MU, Zhao QL, Jawaid P, Takeda K, Ishikawa K, Hori M, Tomihara K, Noguchi K, Kondo T and Noguchi M: Cold atmospheric plasma causes synergistic enhancement in cell death with hyperthermia and an additive enhancement with radiation. *Sci Rep* 7: 11659, 2017.
- Elmore S: Apoptosis: A review of programmed cell death. *Toxicol Pathol* 35: 495-516, 2007.
- Neumann S, El Maadidi S, Faletti L, Haun F, Labib S, Schejtman A, Maurer U and Borner C: How do viruses control mitochondria-mediated apoptosis? *Virus Res* 209: 45-55, 2015.
- Fulda S and Debatin KM: Extrinsic versus intrinsic apoptosis pathways in anticancer chemotherapy. *Oncogene* 25: 4798-4811, 2006.
- Bruin JE, Gerstein HC, Morrison KM and Holloway AC: Increased pancreatic beta-cell apoptosis following fetal and neonatal exposure to nicotine is mediated via the mitochondria. *Toxicol Sci* 103: 362-370, 2008.

38. Borner C and Andrews DW: The apoptotic pore on mitochondria: Are we breaking through or still stuck? *Cell Death Differ* 21: 187-191, 2014.
39. Youle RJ and Strasser A: The BCL-2 protein family: Opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* 9: 47-59, 2008.
40. Haneef J, Parvathy M, Thankayyan RS, Sithul H and Sreeharshan S: Bax translocation mediated mitochondrial apoptosis and caspase dependent photosensitizing effect of *Ficus religiosa* on cancer cells. *PLoS One* 7: e40055, 2012.
41. Wang Z, Lu W, Li Y and Tang B: Alpinetin promotes Bax translocation, induces apoptosis through the mitochondrial pathway and arrests human gastric cancer cells at the G2/M phase. *Mol Med Rep* 7: 915-920, 2013.
42. Kim BJ, Ryu SW and Song BJ: JNK- and p38 kinase-mediated phosphorylation of Bax leads to its activation and mitochondrial translocation and to apoptosis of human hepatoma HepG2 cells. *J Biol Chem* 281: 21256-21265, 2006.
43. Gaitanaki C, Mastri M, Aggeli IK and Beis I: Differential roles of p38-MAPK and JNKs in mediating early protection or apoptosis in the hyperthermic perfused amphibian heart. *J Exp Biol* 211: 2524-2532, 2008.
44. Speidel D: Transcription-independent p53 apoptosis: An alternative route to death. *Trends Cell Biol* 20: 14-24, 2010.
45. Wold LE, Ceylan-Isik AF, Fang CX, Yang X, Li SY, Sreejayan N, Privratsky JR and Ren J: Metallothionein alleviates cardiac dysfunction in streptozotocin-induced diabetes: Role of Ca²⁺ cycling proteins, NADPH oxidase, poly(ADP-Ribose) polymerase and myosin heavy chain isozyme. *Free Radic Biol Med* 40: 1419-1429, 2006.
46. Yu T, Robotham JL and Yoon Y: Increased production of reactive oxygen species in hyperglycemic conditions requires dynamic change of mitochondrial morphology. *Proc Natl Acad Sci USA* 103: 2653-2658, 2006.
47. Orrenius S, Zhivotovsky B and Nicotera P: Regulation of cell death: The calcium-apoptosis link. *Nat Rev Mol Cell Biol* 4: 552-565, 2003.
48. Crompton M, Barksby E, Johnson N and Capano M: Mitochondrial intermembrane junctional complexes and their involvement in cell death. *Biochimie* 84: 143-152, 2002.
49. Toledo FD, Perez LM, Basiglio CL, Ochoa JE, Pozzi EJ and Roma MG: The Ca(2)(+)-calmodulin-Ca(2)(+)/calmodulin-dependent protein kinase II signaling pathway is involved in oxidative stress-induced mitochondrial permeability transition and apoptosis in isolated rat hepatocytes. *Arch Toxicol* 88: 1695-1709, 2014.
50. Seynhaeve ALB, Amin M, Haemmerich D, van Rhoon GC and Hagen TLM: Hyperthermia and smart drug delivery systems for solid tumor therapy. *Adv Drug Deliv Rev* 163: 125-144, 2020.