

# Strong association between lung cancer and the *AXIN2* polymorphism

EMINE GULSEN GUNES<sup>1</sup>, ERGUN PINARBASI<sup>1</sup>, HATICE PINARBASI<sup>2</sup> and YAVUZ SILIG<sup>2</sup>

Departments of <sup>1</sup>Medical Biology and Genetics, and <sup>2</sup>Biochemistry,  
Faculty of Medicine, Cumhuriyet University, 58140 Sivas, Turkey

Received June 12, 2009; Accepted August 10, 2009

DOI: 10.3892/mmr\_00000210

**Abstract.** Accumulated evidence suggests that alterations due to mutations or genetic polymorphisms in the *AXIN2* tumor suppressor gene, a component of the Wnt signaling pathway, contributes to carcinogenesis. The effect of the *AXIN2* exon 1 148 C→T polymorphism was recently investigated in a Japanese population, but has not been investigated in other populations. Additionally, other common polymorphisms of this gene have not been studied. In the present study, 8 polymorphisms of the *AXIN2* gene, including exon 1 148 C→T, were investigated in a Turkish population of 100 lung cancer patients using PCR-RFLP methods. For the exon 1 432 C→T, exon 5 1365 G→A, exon 5 1386 C→T, intron 5 1712+19 G→T, exon 7 2062 C→T and intron 7 2141+73 G→A single nucleotide polymorphisms of *AXIN2*, no significant association was found between the controls and the lung cancer patients. For exon 1 148 C→T, a statistically significant association between the controls and lung cancer patients was found. For this region, lung cancer patients with the TT genotype showed a decreased risk [odds ratio (OR<sub>TT</sub>) 0.33, 95% confidence interval (CI) 0.12-0.89; p=0.032 (adjusted for age, gender and smoking status)] as compared with the controls with the CC genotype. Concerning histological tumor type, it has been found that exon 1 148 C→T SNP is associated with a significant decreased risk in squamous cell carcinoma patients (OR<sub>TT</sub> 0.16; 95% CI 0.03-0.79; p=0.014). Male (OR<sub>TT</sub> 0.19; 95% CI 0.04-0.77; p=0.015) and smoker (OR<sub>TT</sub> 0.11; 95% CI 0.01-0.71; p=0.019) lung cancer patients with the TT genotype showed a decreased risk for the same region. Our results suggest that the risk of lung cancer in a Turkish population is related to polymorphisms of the *AXIN2* gene.

## Introduction

Lung cancer is one of the leading causes of cancer-related death in males and females worldwide (1,2). Epidemiologic

studies show that approximately 80% of all lung cancer cases are attributable to cigarette smoking (3), while the combined effect of environment and genetic factors is responsible for 20% of cases. Among genetic factors, molecular abnormalities such as mutations and deletions have been identified in oncogenes and tumor suppressor genes, and loss of heterozygosity has been detected in a number of chromosomal regions (4-6). In addition, single nucleotide polymorphism (SNP) studies have been carried out in a large number of populations in order to show the susceptibility of an individual to lung cancer. These studies focused mainly on genes encoding xenobiotic metabolizing and DNA repair enzymes, including NAT1 and 2, cytochrome P450 dependent monooxygenases, glutathione S transferases, XRCC1 and XPD (7-9). The association between tumor suppressor gene SNPs and lung cancer risk has also been studied (10-12).

*AXIN2*, a negative regulator of Wnt/β-catenin signaling, acts as a tumor suppressor gene (13,14). The loss of heterozygosity in the genomic locus containing *AXIN2* (15-17) and mutations in the gene itself have been observed in certain types of cancer, including hepatocellular (18), ovarian (19) and colorectal carcinomas (20) and medullablastomas (21), suggesting that this gene plays a role in carcinogenesis. Polymorphisms in the *AXIN2* gene, including exon 1 P50S, have recently been shown to be associated with lung (22) and breast (23) cancers, suggesting they play a role in lung cancer susceptibility.

To our knowledge, there have been no prior studies on the association between *AXIN2* SNPs and lung cancer susceptibility in a Turkish population. We therefore genotyped 200 Turkish individuals (100 controls and 100 lung cancer patients) for 8 SNPs of *AXIN2* in order to investigate their association with susceptibility to lung cancer. The SNPs included the exon 1 148 C→T (Pro50Ser) variant, exon 1 432 T→C, intron 2 956+16 A→G, exon 5 1365 G→A, exon 5 1386 C→T, intron 5 1712+19 G→T, exon 7 2062 C→T and intron 7 2141+73 G→A.

## Materials and methods

**Subjects.** The study population consisted of 100 lung cancer patients admitted to the Respiratory Disease Department at Cumhuriyet University Hospital in Sivas (Central Anatolia) in 2006. Only patients with newly diagnosed lung cancer and no previous diagnosis of cancer or previous radiotherapy or chemotherapy were included in the study. There were no gender, age, histologic type or stage restrictions. The diagnosis of

*Correspondence to:* Dr Ergun Pinarbasi, Department of Medical Biology and Genetics, Faculty of Medicine, Cumhuriyet University, 58140 Sivas, Turkey  
E-mail: epinar@cumhuriyet.edu.tr

**Key words:** *AXIN2*, polymorphism, lung cancer, cancer predisposition, Turkish population

Table I. Conditions for the identification of AXIN2 polymorphisms.

Polymorphism	Marker	Protein effect	Location	Primer sequence (5'-3') (forward and reverse)	Product (bp)	RFLP	Allele (bp)
148 C/T	rs2240308	Pro50Ser	exon 1	F: CCACGCCGATTGCTGAGAGG R: TTCCGCCTGGTGTGGAAGAGACAT	242	<i>Mph1103I</i>	C (218, 24) T (242)
432 T/C	rs2240308	Ile144Ile	exon 1	F: CCTGGAGAGGGAGAAATGC R: CATCACCGACTGGATCTCG	253	<i>Esp3I</i>	T (253) C (145, 108)
956+16 A/G	-	-	intron 2	F: GGGTTTCTGGTGAGGGTCAGT R: CAAGCAAGCCCACGGAAGG	228	<i>MlsI</i>	A (205, 23) G (228)
1365 G/A	rs9915936	Pro455Pro	exon 5	F: ACGTCTTCCCTTTCAGGATG R: GGTA CTGCGAATGGTGGTG	248	<i>MspI</i>	G (100, 80, 36, 32) A (116, 100, 32)
1386 C/T	rs1133683	Pro462Pro	exon 5	F: TGCGTAGGGAGCCGAATGTTG R: GTGGTCCGGGAGCGGGATC	294	<i>TaqI</i>	C (294) T (274, 20)
1712+19 G/T	-	-	intron 5	F: CACCACC ACTACATCCACCAC R: GCTCCACCTCACACCTG	249	<i>CpoI</i>	G (184, 65) T (249)
2062 C/T	-	Leu688Leu	exon 7	F: TCCTTCTGTTTCTCTCTGCTCATTCC R: AGCGTGTGGGTGGGGTCC	209	<i>TaqI</i>	C (209) T (191, 18)
2141+73 G/A	rs4072245	-	intron 7	F: GTGTGCAAGCCCCCAAAG R: CTTGATCCTCCATCTCACAGC	370	<i>BseRI</i>	G (370) A (260, 110)

lung cancer was histologically confirmed. The control group consisted of 100 healthy individuals. All cases and controls were born in Turkey of native Turkish parents, and were interviewed by means of a questionnaire including questions about family history of cancer, smoking and other lifestyle habits, and lifetime occupational history. The study was approved by the local university ethics committee on human research.

**Genotyping.** Prior to the initiation of chemotherapy or radiotherapy, 2 ml of blood was collected per sample. DNA extraction was performed as soon as the blood samples reached the laboratory using a genomic DNA isolation kit (MBI Fermentas, Lithuania) according to manufacturer's instructions. The samples were assayed without knowledge of the case-control status.

A total of 8 regions including 5 exons and 3 introns of the AXIN2 gene were amplified by PCR over 30 cycles (Table I) using previously reported primers (24). PCR was performed in a reaction volume of 25  $\mu$ l containing 100 ng of genomic DNA, 10 pmol of the appropriate amplification primers, 5 nmol each of 4 deoxynucleotide triphosphates (Fermentas), 2.5 Units of Taq DNA polymerase (Fermentas), 10 mmol/l Tris-HCl (pH 8.3 at 25°C), 50 mmol/l KCl and 1.5 mmol/l MgCl<sub>2</sub>. PCR conditions consisted of an initial denaturation at 94°C for 5 min followed by 30 cycles of 94°C for 30 sec, 60°C for 30 sec and 72°C for 60 sec, and lastly 1 cycle at 72°C for 5 min.

In order to detect the 8 AXIN2 gene SNPs, PCR products were digested with 5 Units of appropriate restriction endonuclease enzymes (Fermentas) (Table I) in a total reaction volume of 10  $\mu$ l containing 1X reaction buffer (supplied with the enzyme) for 4 h at optimum temperature according to the manufacturer's instructions. Amplified and digested DNA fragment sizes are shown in Table I. Following digestion, DNA fragments were separated on 2.5% agarose gels.

**Statistical analysis.** Statistical analysis was performed using the Statistical Package for Social Sciences Program (SPSS, version 16). Statistically significant departures from Hardy-Weinberg equilibrium for controls were assessed using the  $\chi^2$  test. For each polymorphism, unconditional logistic regression was used to calculate the odds ratio (OR) and 95% confidence interval (CI) for the lung cancer and tumor types, adjusted for the study matching factors of age, gender and smoking habits. The relationship between genotypes and clinopathological characteristics was examined by the  $\chi^2$  test. A p-value <0.05 was considered to be statistically significant.

## Results

A total of 200 individuals (100 lung cancer patients and 100 healthy controls) were genotyped for 8 different SNPs, including exon 1 148 C→T of the AXIN2 gene. The results of SNP analysis in the control population did not deviate from Hardy-Weinberg equilibrium. The distribution of demographic characteristics in the genotyped cancer cases as similar to the demographic distribution of the controls. The principal characteristics of the study population are presented in Table II. The mean ages of patients and controls were 59.22±9.63 and 57.01±7.89 years, respectively. Relative to the controls, there were proportionally more males among the lung cancer patients (56%). The proportion of smokers was higher among patients (40%) than among controls (30%).

Table III summarizes the main effects of the 8 SNPs of AXIN2 that are involved in lung cancer. Genotype analysis carried out on exons and the exon-intron boundaries of AXIN2 in lung cancer patients indicated that there was no significant association between genotype and risk of lung cancer for the following SNPs: exon 1 432 T→C, intron 2 956+16 A→G, exon 5 1365 G→A, exon 5 1386 C→T, intron 5 1712+19 G→T, exon 7

Table II. Characteristics of lung cancer patients and healthy controls.

	Patients (n=100) n (%)	Controls (n=100) n (%)	p-value <sup>a</sup>
Gender			0.157
Male	56 (56)	46 (46)	
Female	44 (44)	54 (54)	
Age (years $\pm$ SD)	59.22 $\pm$ 9.63	57.01 $\pm$ 7.89	
Smoking habit			0.138
Non-smoker	60 (60)	70 (70)	
Smoker	40 (40)	30 (30)	

<sup>a</sup>P-values were calculated using the  $\chi^2$  test.

2062 C→T and intron 7 2141+73 G→A. However, individuals carrying the 148 TT (Ser/Ser) genotype were found to have a decreased risk of lung cancer ( $p < 0.05$ ). For this region of *AXIN2*, the frequencies of the Pro50Pro (CC), Pro50Ser (C→T) and Ser50Ser (TT) genotypes were 32, 52 and 16% in the controls, respectively, and 45, 47 and 8% in the lung cancer patients, respectively. The calculated values for the 148 C→T polymorphism were  $OR_{TT}$  0.35, 95% CI 0.13-0.93 and  $p = 0.032$ . When age, gender and smoking status were included in the analysis, the estimated OR was found to be 0.33 and the 95% CI was 0.12-0.89.

The  $\chi^2$  test was used to evaluate the association between the *AXIN2* polymorphisms and smoking habit, gender and histopathologic type of lung cancer (Table IV). The most prevalent type of tumors were squamous cell carcinoma (46%) and small cell carcinoma (27%). Others included lung adenocarcinoma, and large cell and adenosquamous carcinoma. With the exception of the exon 1 148 C→T SNP, none of the remaining 7 *AXIN2* SNPs exhibited statistically significant results. For the 148 C→T SNP, squamous cell carcinoma patients ( $OR_{TT}$  0.16, 95% CI 0.03-0.79;  $p = 0.014$ ), males ( $OR_{TT}$  0.19, 95% CI 0.04-0.77;  $p = 0.015$ ) and smokers ( $OR_{TT}$  0.11, 95% CI 0.01-0.71;  $p = 0.019$ ) with the TT genotype were found to exhibit a statistically significant decreased risk of lung cancer.

For the intron 2 956+16 A→G SNP, the p-value was found to be 0.044 for small cell carcinoma ( $OR_{AG}$  0.28, 95% CI 0.80-1.02). However, as it was close to 0.05 it did not reach statistical significance, since the OR value was in the range of 95% CI.

## Discussion

Molecular epidemiology studies indicate that genetic polymorphisms and mutations of xenobiotic metabolizing enzymes and DNA repair genes as well as tumor suppressor genes and oncogenes are among the risk factors for lung cancer (25). Polymorphisms of xenobiotic enzymes have been demonstrated to be a major risk factor for lung cancer by several case-control studies; however, the results of these studies are inconsistent (25). Several lines of evidence indicate that events that alter the tumor suppressor pathways affect lung and other types of cancer (26-31). These events include polymorphisms of *AXIN2*, which

play an important role in Wnt/ $\beta$ -catenin signaling and function as a scaffolding protein for a protein complex including *AXIN2*, adenomatous polyposis coli (APC), glycogen synthase kinase 3 (GSK3) and  $\beta$ -catenin. This complex is required for the phosphorylation of  $\beta$ -catenin (32,33). In this complex is localized human chromosome 17q21-25, a region highly susceptible to allelic losses and chromosomal re-arrangements involved in several types of cancer (24,34,35). A recent study by Kanzaki *et al* in a Japanese population showed that the *AXIN2* SNP at codon 50 is associated with lung cancer (22). In another study, 5 *AXIN* SNPs were investigated in breast cancer patients and found to be associated with the disease (23). In this study, the effect of the *AXIN2* SNP at codon 50 as well as 7 other significant SNPs of *AXIN2* were investigated in a Turkish population in order to further analyse their role in lung cancer susceptibility.

The study comprised 100 lung cancer patients and 100 hospital-based controls. Although the frequency of males and smokers was higher among the lung cancer patients, there was no significant relationship between lung cancer and these factors. No association was observed between lung cancer and the 7 following SNPs of *AXIN2*: exon 1 432 T→C, intron 2 956+16 A→G, exon 5 1365 G→A, exon 5 1386 C→T, intron 5 1712+19 G→T, exon 7 2062 C→T and intron 7 2141+73 G→A.

In the study by Kanzaki *et al*, the SNP at codon 50 of the *AXIN2* gene encoding either proline (CCT) or serine (TCT) was found to be associated with lung cancer, but not with colorectal and head and neck cancers (22). The authors found that patients with the TT genotype showed a reduced risk of cancer ( $OR_{TT}$  0.31, 95% CI 0.12-0.79). The results of the present study also indicate a decreased risk for lung cancer patients with the TT genotype ( $OR_{TT}$  0.35, 95% CI 0.13-0.93). The CC, CT and TT allele frequencies were 45, 47 and 8% in lung cancer patients and 32, 52 and 16% in controls, respectively. These results are similar to those of Kanzaki *et al*, who found a CC, CT and TT genotype distribution of 50, 44 and 8% in lung cancer patients and 42, 52 and 15% in controls, respectively. The 17q21-25 region of *AXIN2* is very close to the functional motif where APC binds. *AXIN2* binding site disruption or mutations in APC result in human cancer (36). It is possible that this SNP of *AXIN2* changes the binding efficiency of APC, though this requires further investigation.

No prior study has examined lung cancer risk associated with 7 of the *AXIN2* polymorphic sites investigated in the present study; however, the intron 2 956+16 A→G and exon 7 2062 C→T polymorphisms were found to be associated with tooth agenesis (24). It has been proposed that this variant of *AXIN2* may exert a weak effect on splicing, since it creates an additional donor splicing site within the sequence of exon 2. This was not observed in the present study. Additionally, the SNP at exon 5 1365 G→A of the *AXIN2* gene has been reported to be A→G (dbSNP ID: rs9915936), but we did not identify any AA genotypes in our control and study groups (24). The GG genotype was therefore selected as the wild-type genotype for statistical testing.

This is the first study carried out in a Turkish population investigating the *AXIN2* polymorphism and its association with lung cancer. The data indicate that there is a significant association between *AXIN2* SNPs and lung cancer in this population. Further investigation in different populations with large case numbers is needed to support these results.

Table III. *AXIN2* genotypes in healthy controls and lung cancer patients.

<i>AXIN2</i> genotype	Patients n (%)	Controls n (%)	p-value <sup>a</sup>	OR (95% CI)	
				Crude	Adjusted <sup>b</sup>
Exon 1 148 C/T					
CC	45 (45.0)	32 (32.0)	1	1 (Reference)	1 (Reference)
CT	47 (47.0)	52 (52.0)	0.148	0.64 (0.35-1.17)	0.62 (0.33-1.16)
TT	8 (8.0)	16 (16.0)	0.032	0.35 (0.13-0.93)	0.33 (0.12-0.89)
Allele frequencies					
C	137 (61.5)	116 (58.0)			
T	63 (38.5)	84 (42.0)			
Exon 1 432 T/C					
TT	96 (96.0)	95 (95.0)	1	1 (Reference)	1 (Reference)
TC	4 (4.0)	5 (5.0)	1.0	1.0 (0.24-4.11)	1.2 (0.28-5.12)
CC	0	0			
Allele frequencies					
T	196 (98.0)	195 (97.5)			
C	4 (2.0)	5 (2.5)			
Intron 2 956+16 A/G					
AA	77 (77.0)	64 (64.0)	1	1 (Reference)	1 (Reference)
AG	20 (20.0)	28 (28.0)	0.121	0.59 (0.30-1.15)	0.57 (0.29-1.13)
GG	3 (3.0)	8 (8.0)	0.08	0.31 (0.07-1.22)	0.30 (0.07-1.22)
Allele frequencies					
A	174 (87.0)	156 (78.0)			
G	26 (13.0)	44 (22.0)			
Exon 5 1365 G/A					
GG	91 (91.0)	88 (88.0)	1	1 (Reference)	1 (Reference)
GA	9 (9.0)	12 (12.0)	0.489	0.72 (0.29-1.80)	0.73 (0.29-1.86)
AA	0	0			
Allele frequencies					
G	191 (95.5)	188 (94.0)			
A	9 (4.5)	12 (6.0)			
Exon 5 1386 C/T					
CC	52 (52.0)	42 (42.0)	1	1 (Reference)	1 (Reference)
CT	38 (38.0)	50 (50.0)	0.102	0.61 (0.34-1.10)	0.57 (0.31-1.05)
TT	10 (10.0)	8 (8.0)	0.985	1.01 (0.36-2.78)	0.80 (0.27-2.30)
Allele frequencies					
C	142 (71.0)	134 (67.0)			
T	58 (29.0)	66 (33.0)			
Intron 5 1712+19 G/T					
TT	97 (97.0)	96 (96.0)	1	1 (Reference)	1 (Reference)
GT	3 (3.0)	4 (4.0)	0.352	1.80 (0.51-6.37)	1.98 (0.55-7.12)
GG	0	0			
Allele frequencies					
T	193 (96.5)	196 (98.0)			
G	7 (3.5)	4 (2.0)			
Exon 7 2062 C/T					
CC	91 (91.0)	86 (86.0)	1	1 (Reference)	1 (Reference)
CT	9 (9.0)	14 (14.0)	0.268	0.60 (0.25-1.47)	0.61 (0.24-1.52)
TT	0	0			
Allele frequencies					
C	191 (95.5)	186 (93.0)			
T	9 (4.5)	14 (7.0)			
Intron 7 2141+73 G/A					
GG	73 (73.0)	80 (80.0)	1	1 (Reference)	1 (Reference)
GA	27 (27.0)	20 (20.0)	0.243	1.47 (0.76-2.86)	1.63 (0.82-3.21)
AA	0	0			
Allele frequencies					
G	173 (86.5)	180 (90.0)			
A	27 (13.5)	20 (10.0)			

<sup>a</sup>p-values were calculated using the  $\chi^2$  test. <sup>b</sup>Adjusted for age, gender and smoking status.

Table IV. Association between the *AXIN2* genotype and the clinicopathological characteristics of the lung cancer patients.

Region	Genotype			p-value	Odds ratio (95% CI)	P-value	Odds ratio (95% CI)
Exon 1 148 C/T	CC	CT	TT		CT		TT
Controls	32	52	16				
Patients	45	47	8	0.148	0.64 (0.35-1.17)	0.032	0.35 (0.13-0.93)
Male	25	27	4	0.082	0.45 (0.18-1.11)	0.015	0.19 (0.04-0.77)
Female	20	20	4	0.620	0.80 (0.34-1.88)	0.463	0.60 (0.15-2.36)
Non-smoker	28	26	6	0.243	0.64 (0.30-1.34)	0.380	0.59 (0.18-1.90)
Smoker	17	21	2	0.268	0.54 (0.18-1.61)	0.019	0.11 (0.01-0.71)
SQC	24	20	2	0.075	0.51 (0.24-1.07)	0.014	0.16 (0.03-0.79)
SCC	10	13	4	0.639	0.80 (0.31-2.03)	1.000	0.80 (0.21-2.95)
Other	11	14	2	0.596	0.78 (0.31-1.93)	0.310	0.36 (0.07-1.84)
Exon 1 432 T/C	TT	TC			TC		
Controls	96	4					
Patients	96	4		0.640	1.00 (0.24-4.11)		
Male	55	1		1.000	0.81 (0.05-13.4)		
Female	41	3		1.000	1.20 (0.23-6.49)		
Non-smoker	56	4		0.703	1.59 (0.34-7.42)		
Smoker	40	0		0.429	0.96 (0.90-1.03)		
SQC	44	2		1.000	1.09 (0.20-6.18)		
SCC	26	1		1.000	0.92 (0.10-8.61)		
Other	26	1		1.000	0.92 (0.09-8.61)		
Intron 2 956+16 A/G	AA	AG	GG		AG		GG
Controls	64	28	8				
Patients	77	20	3	0.121	0.59 (0.30-1.15)	0.080	0.31 (0.79-1.22)
Male	43	11	2	0.196	0.54 (0.21-1.37)	1.000	0.69 (0.10-5.23)
Female	34	9	1	0.367	0.64 (0.24-1.68)	0.113	0.16 (0.02-1.45)
Non-smoker	47	12	1	0.055	0.45 (0.20-1.02)	0.059	0.14 (0.01-1.25)
Smoker	30	8	2	0.747	1.20 (0.35-4.24)	1.000	0.76 (0.10-5.86)
SQC	31	13	2	0.916	0.95 (0.43-2.10)	0.500	0.51 (0.10-2.57)
SCC	24	3	0	0.044	0.28 (0.80-1.02)	0.195	0.88 (0.82-0.96)
Other	22	4	1	0.127	0.41 (0.13-1.31)	0.447	0.36 (0.04-3.07)
Exon 5 1365 G/A	GG	GA			GA		
Controls	88	12					
Patients	91	9		0.489	0.72 (0.29-1.80)		
Male	52	4		0.216	0.42 (0.11-1.56)		
Female	39	5		0.750	1.20 (0.33-4.65)		
Non-smoker	53	7		0.837	0.89 (0.31-2.56)		
Smoker	38	2		0.645	0.47 (0.07-3.03)		
SQC	43	3		0.390	0.51 (0.13-1.90)		
SCC	22	5		0.357	1.66 (0.53-5.22)		
Other	26	1		0.297	0.28 (0.03-2.27)		
Exon 5 1386 C/T	CC	CT	TT		CT		TT
Controls	42	50	8				
Patients	52	38	10	0.102	0.61 (0.34-1.10)	0.985	1.01 (0.36-2.78)
Male	31	22	3	0.201	0.59 (0.26-1.32)	1.000	0.96 (0.14-6.31)
Female	21	16	7	0.318	0.64 (0.27-1.52)	0.752	1.22 (0.35-4.23)
Non-smoker	28	23	9	0.491	0.77 (0.36-1.61)	0.386	1.66 (0.52-5.25)
Smoker	24	15	1	0.072	0.40 (0.14-1.09)	0.265	0.22 (0.02-2.80)
SQC	27	18	1	0.115	0.56 (0.27-1.15)	0.146	0.19 (0.02-1.64)
SCC	12	11	4	0.575	0.77 (0.38-1.92)	0.417	1.75 (0.45-6.82)
Other	13	9	5	0.257	0.58 (0.22-1.49)	0.306	2.01 (0.56-7.24)
Intron 5 1712+19 G/T	GG	GT			GT		
Controls	96	4					
Patients	93	7		0.352	1.80 (0.51-6.37)		
Male	52	4		0.688	1.69 (0.29-9.68)		
Female	41	3		0.654	1.90 (0.30-11.9)		
Non-smoker	53	7		0.080	4.50 (0.90-22.5)		
Smoker	40	0		0.180	0.93 (0.84-1.02)		
SQC	42	4		0.261	2.28 (0.54-9.5)		
SCC	27	0		0.578	0.96 (0.92-1.00)		
Other	24	3		0.165	3.00 (0.62-14.3)		



Table IV. Continued.

Region	Genotype		p-value	Odds ratio (95% CI)	P-value	Odds ratio (95% CI)
Exon 7 2062 C/T	CC	CT		CT		
Controls	86	14				
Patients	91	9	0.268	0.60 (0.25-1.47)		
Male	51	5	1.000	1.02 (0.26-4.07)		
Female	40	4	0.250	0.44 (0.12-1.51)		
Non-smoker	56	4	0.350	0.55 (0.15-1.93)		
Smoker	35	5	0.511	0.57 (0.15-2.08)		
SQC	42	4	0.365	0.58 (0.18-1.88)		
SCC	24	3	1.000	0.76 (0.20-2.89)		
Other	25	2	0.520	0.49 (0.10-2.30)		
Intron 7 2141+73 G/A	GG	GA		GA		
Controls	80	20				
Patients	73	27	0.243	1.47 (0.76-2.86)		
Male	43	13	0.469	1.40 (0.53-3.87)		
Female	30	14	0.285	1.60 (0.66-4.02)		
Non-smoker	40	20	0.183	1.68 (0.77-3.66)		
Smoker	33	7	0.747	1.37 (0.36-5.22)		
SQC	31	15	0.097	1.93 (0.88-4.25)		
SCC	22	5	0.864	0.90 (0.30-2.69)		
Other	20	7	0.504	1.40 (0.52-3.76)		

SQC, squamous cell carcinoma; SCC, small cell carcinoma.

## Acknowledgements

This study was supported by Grant no. T-299 from Cumhuriyet University CUBAP, Sivas, Turkey. The technical assistance of Ms. Meral Yilmaz is gratefully acknowledged.

## References

- Ginsberg RJ, Vokes EE and Raben A: Cancer of the lung. In: *Cancer: Principles and Practice of Oncology*. DeVita VT Jr, Hellman S and Rosenberg SA (eds). 5th edition. Lippincott-Raven, Philadelphia, pp858-865, 1997.
- Marby M, Nelkin BD and Baylin SB: Lung cancer. In: *The Genetic Basis of Human Cancer*. Vogelstein B and Kinzler KW (eds). 2nd edition. McGraw-Hill, New York, pp671-679, 1998.
- Khuder SA: Effect of cigarette smoking on major histological types of lung cancer: a meta-analysis. *Lung Cancer* 31: 139-148, 2001.
- Sekido Y, Fong KM and Minna JD: Progress in understanding the molecular pathogenesis of human lung cancer. *Biochim Biophys Acta* 1378: 21-59, 1998.
- Salgia R and Skarin AT: Molecular abnormalities in lung cancer. *J Clin Oncol* 16: 1207-1217, 1998.
- Reissmann PT, Koga H, Figlin RA, Holmes EC and Slamon DJ: Amplification and overexpression of the cyclin D1 and epidermal growth factor receptor genes in non-small cell lung cancer. *Lung Cancer Study Group. J Cancer Res Clin Oncol* 125: 61-70, 1999.
- Pinarbasi H, Silig Y, Cetinkaya O, Seyfikli Z and Pinarbasi E: Strong association between the GSTM1-null genotype and lung cancer in a Turkish population. *Cancer Genet Cytogenet* 146: 125-129, 2003.
- Bartsch H, Nair U, Risch A, Rojas M, Wikman H and Alexandrov K: Genetic polymorphism of CYP genes, alone or in combination, as a risk modifier of tobacco-related cancers. *Cancer Epidemiol Biomarkers Prev* 9: 3-28, 2000.
- Schwartz AG, Prysak GM, Bock CH and Cote ML: The molecular epidemiology of lung cancer. *Carcinogenesis* 28: 507-518, 2007.
- Kohno T and Yokota J: How many tumor suppressor genes are involved in human lung carcinogenesis? *Carcinogenesis* 20: 1403-1410, 1999.
- Ohta M, Inoue H, Cotticelli MG, *et al*: The FHIT gene, spanning the chromosome 3p14.2 fragile site and renal carcinoma-associated t(3;8) breakpoint is abnormal in digestive tract cancers. *Cell* 84: 587-597, 1996.
- Sard L, Accornero P, Tornielli S, *et al*: The tumor-suppressor gene FHIT is involved in the regulation of apoptosis and in cell cycle control. *Proc Natl Acad Sci USA* 96: 8489-8492, 1999.
- Kikuchi A: Modulation of Wnt signaling by axin and axil. *Cytokine Growth Factor Rev* 10: 255-265, 1999.
- Hughes TA and Brady HJM: Cross-talk between pRb/E2F and Wnt/b-catenin pathways: E2F1 induces AXIN2 leading to repression of Wnt signalling and to increased cell death. *Exp Cell Res* 303: 32-46, 2005.
- Phelan CM, Borg A, Cuny M, *et al*: Consortium study on 1280 breast carcinomas: allelic loss on chromosome 17 targets sub-regions associated with family history and clinical parameters. *Cancer Res* 58: 1004-1012, 1998.
- Plummer SJ, Paris MJ, Myles J, Tubbs R, Crowe J and Casey G: Four regions of allelic imbalance on 17q12-qter associated with high-grade breast tumors. *Genes Chromosomes Cancer* 20: 354-362, 1997.
- Barlund M, Tirkkonen M, Forozan F, Tanner MM, Kallioniemi O and Kallioniemi A: Increased copy number at 17q22-q24 by CGH in breast cancer is due to high-level amplification of two separate regions. *Genes Chromosomes Cancer* 20: 372-376, 1997.
- Taniguchi K, Roberts LR, Aderca IN, *et al*: Mutational spectrum of beta-catenin, AXIN1 and AXIN2 in hepatocellular carcinomas and hepatoblastomas. *Oncogene* 21: 4863-4871, 2002.
- Wu R, Zhai Y, Fearon ER and Cho KR: Diverse mechanisms of beta-catenin deregulation in ovarian endometrioid adenocarcinomas. *Cancer Res* 61: 8247-8255, 2001.
- Liu W, Dong X, Mai M, *et al*: Mutations in AXIN2 cause colorectal cancer with defective mismatch repair by activating beta-catenin/TCF signalling. *Nat Genet* 26: 146-147, 2000.
- Yokota N, Mainprize TG, Taylor MD, *et al*: Identification of differentially expressed and developmentally regulated genes in medulloblastoma using suppression subtraction hybridization. *Oncogene* 23: 3444-3453, 2004.
- Kanzaki H, Ouchida M, Hanafusa H, *et al*: Single nucleotide polymorphism of the AXIN2 gene is preferentially associated with human lung cancer risk in a Japanese population. *Int J Mol Med* 18: 279-284, 2006.

23. Wang X, Goode EL, Fredericksen ZS, *et al*: Association of genetic variation in genes implicated in the beta-catenin destruction complex with risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 17: 2101-2108, 2008.
24. Mostowska A, Biedziak B and Jagodzinski PP: Axis inhibition protein 2 (AXIN2) polymorphisms may be a risk factor for selective tooth agenesis. *J Hum Genet* 51: 262-266, 2006.
25. Risch A and Plass C: Lung cancer epigenetics and genetics. *Int J Cancer* 123: 1-7, 2008.
26. Yokobori T, Mimori K, Iwatsuki M, *et al*: p53-altered FBXW7 expression determines poor prognosis in gastric cancer cases. *Cancer Res* 69: 3788-3794, 2009.
27. Miyaki M, Yamaguchi T, Iijima T, Takahashi K, Matsumoto H and Mori T: Somatic mutations of the CDC4 (FBXW7) gene in hereditary colorectal tumors. *Oncology* 76: 430-434, 2009.
28. Liu JN, Zhang XM, Guo YL, Sun T, Lin DX and Wen T: Genetic polymorphism in MDM2 is associated with susceptibility to colorectal cancer in a Chinese population. *Zhonghua Zhong Liu Za Zhi* 30: 335-338, 2008.
29. Ding L, Getz G, Wheeler DA, *et al*: Somatic mutations affect key pathways in lung adenocarcinoma. *Nature* 455: 1069-1075, 2008.
30. Shinmura K, Suzuki M, Yamada H, *et al*: Characterization of adenocarcinoma of the lung in a familial adenomatous polyposis patient. *Pathol Int* 58: 706-712, 2008.
31. Mechanic LE, Marrogi AJ, Welsh JA, *et al*: Polymorphisms in XPD and TP53 and mutation in human lung cancer. *Carcinogenesis* 26: 597-604, 2005.
32. Akiyama T: Wnt/beta-catenin signaling. *Cytokine Growth Factor Rev* 11: 273-282, 2000.
33. Bienz M and Clevers H: Linking colorectal cancer to Wnt signaling. *Cell* 103: 311-320, 2000.
34. Salahshor S and Woodgett JR: The links between axin and carcinogenesis. *J Clin Pathol* 58: 225-236, 2005.
35. Hart MJ, De los Santos R, Albert IN, Rubinfeld B and Polakis P: Down-regulation of beta-catenin by human Axin and its association with the APC tumor suppressor, beta-catenin and GSK3 beta. *Curr Biol* 8: 573-581, 1998.
36. Rubinfeld B, Robbins P, El-Gamil M, Albert I, Porfiri E and Polakis P: Stabilization of beta-catenin by genetic defects in melanoma cell lines. *Science* 275: 1790-1792, 1997.