

Programmed cell death 4 (Pdcd4) expression in colorectal adenocarcinoma: Association with clinical stage

SUNG-CHUL LIM^{1,2} and RAN HONG^{1,2}

Department of ¹Pathology and ²Research Center for Resistant Cells,
Chosun University School of Medicine, Gwangju, Republic of Korea

Received June 22, 2011; Accepted July 21, 2011

DOI: 10.3892/ol.2011.365

Abstract. The aim of this study was to examine the role of Programmed cell death 4 (Pdcd4) in colorectal adenocarcinoma (CRA). Pdcd4 expression was observed in both the nucleus and cytoplasm in colorectal adenocarcinoma, whereas Pdcd4 was expressed in the nucleus in normal colonic epithelial cells. Loss or weak expression of Pdcd4 was identified in 44 cases (40.7%) of cancer cells. Pdcd4 expression was associated with an increase in the nodal and clinical stage ($p=0.022$ and $p=0.016$, respectively). Nuclear staining was identified in 66 cases (61.15%), with no correlation with clinicopathological factors. Conversely, cytoplasmic staining for Pdcd4 was observed in 45 cases (41.7%), and increased according to nodal and clinical stage ($p=0.011$ and $p=0.009$, respectively), indicating that aberrant Pdcd4 expression leads to tumor progression. However, Pdcd4 expression was not correlated to disease-free survival time. This study demonstrated that during the tumorigenesis of CRA, loss of nuclear Pdcd4 expression occurs, and during tumor progression, aberrant cytoplasmic expression is present, suggesting a higher clinical stage. Although loss of Pdcd4 was not significantly correlated with survival time, as the prognosis of colorectal cancer varies depending on clinical stage including invasion depth, nodal status and metastatic status, cytoplasmic Pdcd4 expression may be a favorable prognostic marker in CRA.

Introduction

Programmed cell death 4 (Pdcd4) is a ubiquitously expressed, novel tumor suppressor gene localized in chromosome 10q24, whose protein product plays a role in the suppression of tumorigenesis and tumor progression and invasion (1-4). Although Pdcd4 was first identified as being differentially upregulated during apoptosis, experimental evidence established it as

a novel tumor suppressor (5-8). Previous studies indicated that the overexpression of Pdcd4 inhibited tumor promoter TPA-induced neoplastic transformation in JB6 and P-positive cells (4,9,10). Pdcd4 expression is often decreased in progressed carcinomas of the lung, ovary, breast, colon and prostate (7,11-13). Pdcd4 protein acts as an inhibitor that suppresses protein translation through its binding to and inhibition of the helicase activity of eukaryotic translation initiation factor 4A (eIF4A), the latter being a component of the protein translation complex (4,9,14). Certain studies investigating cellular functions of Pdcd4 demonstrated that it was capable of regulating molecules such as p21 (7), Cdk3, ornithine decarboxylase (1), carbonic anhydrase II (15) and JNK/c-Jun/AP-1 (6). It has also been shown to suppress the expression of the invasion-related urokinase receptor (u-PAR) gene, and to suppress invasion/intravasation via the Sp1/Sp3 promoter motifs in cancer (8). However, the physiological role of Pdcd4 is not clearly understood. The purpose of the present study was to examine the role of Pdcd4 in colorectal adenocarcinoma (CRA). The prognosis of colorectal cancer patients varies depending on the depth of the invasion of tumor cells and the lymph node status, and these factors are assessed only by a histopathological examination of surgically resected tissues (16). Pdcd4 expression and its subcellular localization were investigated using immunohistochemistry, and Pdcd4 expression was correlated with clinical and pathological parameters including histology, grade, stage and overall survival.

Materials and methods

Patients. Among patients who underwent curative surgery for CRA at Chosun University Hospital between January 1992 and December 2001, the present study was conducted on a non-consecutive series of 108 patients from whom paraffin-embedded tissues were obtained and relatively well preserved, medical records were complete and the patient status had been followed up. Patient survival was confirmed through telephone interviews and by mail. Patients who underwent preoperative chemoradiotherapy and emergency surgery, and patients who had evidence of hereditary non-polyposis colorectal cancer or familial adenomatous polyposis were excluded from the study.

The various clinicopathological parameters of the patients were confirmed by reviewing the patient medical records

Correspondence to: Dr Sung-Chul Lim, Department of Pathology, Chosun University Hospital, 588 Seosuk-dong, Dong-gu, Gwangju 501-140, Republic of Korea
E-mail: nanih@chosun.ac.kr

Key words: programmed cell death 4, colorectal adenocarcinoma

and pathology files. The correlation of clinicopathological parameters and immunohistochemical findings with survival was investigated for all 108 patients. Informed consent was obtained from each patient according to the institutional guidelines, and the research protocols were approved by the Ethics Committee of the University Hospital.

Histopathological analysis

Microscopic examination. Each tumor was re-evaluated by retrospective analysis of the medical records and the tissue slide files of the Department of Pathology. Age, gender, tumor size, histological subtype, degree of differentiation, depth of tumor invasion, status of lymph node metastasis and presence of distant metastasis were assessed. Stage was defined according to the TNM staging system of the American Joint Committee on Cancer (17). The examined tissues were fixed in 10% neutral formalin, and the prepared paraffin-embedded tissues were sectioned at a thickness of 4-5 μ m. Hematoxylin and eosin staining was performed, and the sections were examined under a light microscope. A representative area suitable for the study purpose was selected, and slides were prepared for immunohistochemical analysis.

Immunohistochemical staining. Specimens in this study were tested using a rabbit polyclonal antibody against Pdc44 (Abcam, Cambridge, MA, USA) according to the manufacturer's instructions. Immunolocalization was performed using the mouse ImmunoCruz™ Staining System sc-2050 (Santa Cruz Biotechnology, Santa Cruz, CA, USA), according to the manufacturer's instructions. The staining process was performed according to a standard protocol. Briefly, the 4- μ m sections that were obtained following formalin fixation and paraffin embedding were deparaffinized in xylene and rehydrated with distilled water through a graded series of ethanol solutions. The sections were then placed in a glass jar with 10 mM citrate buffer (pH 6.0) and were irradiated in a microwave oven for 15 min. The sections were allowed to cool in the jar at room temperature for 20 min. The slides were rinsed with Tris-buffered saline (TBS). A blocking reagent was added for 10 min after quenching the endogenous peroxidase activity in 0.3% hydrogen peroxide for 10 min. The slides were then washed as previously described, and the slides were subsequently subjected to the primary antibody reaction. Immunohistochemistry was performed using the Nexes ES (Ventana, Tucson, AZ, USA). Slides were incubated with the antibodies for 32 min. The Ventana basic DAB detection kit (catalog no. 760-001) was the secondary detection method. This kit includes biotinylated immunoglobulin secondary antibody, containing affinity-purified goat anti-mouse IgG and IgM (b200 Ig/ml) and goat anti-rabbit IgG (b200 Ig/ml) in phosphate buffer with preservative. Incubation was performed for 8 min, and was followed by conjugated streptavidin horseradish peroxidase. Slides were counterstained with hematoxylin (Ventana catalog no. 760-2021).

Analysis and interpretation of staining. Two pathologists, who were unaware of the clinical course of the subjects in order to exclude subjectivity, evaluated the results of the staining. Since Pdc44 expression was identified in both the nucleus and the cytoplasm, nuclear and cytoplasmic staining was evaluated separately. Nuclear and cytoplasmic scoring was determined according to the percentage of positive

Table I. Summary of clinicopathological factors.

Characteristics	n (%)
Age	
≤40	14 (13.0)
51-59	28 (25.9)
60-69	38 (35.2)
≥70	28 (25.9)
Gender	
Male	55 (50.9)
Female	53 (49.1)
Pathological tumor classification (pT)	
pT1	2 (1.9)
pT2	21 (19.4)
pT3	80 (74.1)
pT4	5 (4.6)
Pathological lymph node classification	
pN0	73 (67.6)
pN1	26 (24.1)
pN2	9 (8.3)
Metastasis classification (M)	
M0	104 (96.3)
M1	4 (3.7)
AJCC classification	
I	21 (19.4)
IIA	49 (45.3)
IIB	2 (1.9)
IIIA	2 (1.9)
IIIB	22 (20.4)
IIIC	8 (7.4)
IV	4 (3.7)
Differentiation	
Well/Moderate	88 (81.5)
Poor	7 (6.5)
Mucinous	13 (12.0)

AJCC, American Joint Committee on Cancer.

nuclear staining and the intensity of cytoplasmic staining in cells, respectively (18). Nuclear scoring was as follows (18): score 1, negative; score 2, <30%; score 3, 30 to 70%; and score 4, >70%. Cytoplasmic scores were determined according to staining intensity: score 1, negative; score 2, weak staining intensity; score 3, intermediate staining intensity and score 4, strong staining intensity, in reference to a strongly stained tissue as a control. The overall Pdc44 score was calculated by adding the nuclear and cytoplasmic scores, and 4 groups were defined: negative (total score, 1 and 2), weak staining (total score, 3 and 4), intermediate staining (total score, 5 and 6) and strong staining (total score, 7 and 8).

Statistical analysis. For the statistical analysis of Pdc44 in CRA, its association with various clinicopathological factors,

Table II. Summary of Pdc4 expression and correlation with clinicopathologic parameters.

		Pdc4 expression					Nuclear expression					Cytoplasmic expression				
n		0 ^A	1 ^A	2 ^A	3 ^A	P-value	1	2	3	4	P-value	1	2	3	4	P-value
T stage																
1	2	0	1	1	0	0.443	1	0	0	1	0.018 ^d	1	0	1	0	0.903
2	21	3	8	9	1		5	2	10	4		13	4	3	1	
3	80	8	20	46	6		17	14	9	40		45	20	13	2	
4	8	0	4	1	0		1	2	1	1		4	1	0	0	
N stage																
N0	73	10	24	36	3	0.133	17	12	14	30	0.961	49	16	6	2	0.011 ^d
N1	35	1	9	21	4		7	6	16	16		14	9	11	1	
M stage																
M0	104	11	33	53	7	0.294	24	18	20	42	0.133	60	24	17	3	0.808
M1	4	0	0	4	0		0	0	0	4		3	1	0	0	
Clinical stage																
High ^d	72	10	24	35	3	0.114	17	12	14	29	0.906	49	15	6	2	0.009 ^d
Low ^d	36	1	9	22	4		7	6	6	17		14	10	11	1	
Diff. ¹																
W/M ^a	88	11	26	46	5	0.359	22	14	15	37	0.724	54	17	15	2	0.440
P ^b	7	0	4	2	1		1	2	1	3		3	3	1	0	
M ^c	13	0	3	9	1		1	2	4	6		6	5	1	1	
Diff. ²																
Non-M ¹	93	10	29	48	6	0.925	22	16	15	40	0.419	55	20	16	2	0.438
M ²	15	1	4	9	1		2	2	5	6		9	5	1	1	

Pdc4 expression: 0^A, negative staining (score 1-2); 1^A, weak staining (score 3-4); 2^A, intermediate staining (score 5-6); 3^A, strong staining (score 7-8). N stage - N0, N1. Clinical stage: High, clinical stage I and II; Low, clinical stage III and IV. Diff¹: W/M^a, well to moderate differentiation; P^b, poor differentiation; M^c, mucinous carcinoma. Diff²: Non-M¹, non-mucinous carcinoma; M², mucinous carcinoma. ^dStatistically significant, $p < 0.05$.

and survival rate or survival time, we used Pearson's Chi-square test and linear-by-linear association to compare the categorical data. The life table and statistical significance were computed using the Kaplan-Meier method and the log-rank test. $P < 0.05$ was considered to be statistically significant. The Stat View software package was used for all statistical analyses (Abacus Concepts, Berkeley, CA, USA).

Results

The clinical characteristics of the patients are shown in Table I. The average age at the time of surgery was 62.1 years and the ratio of male to female participants was 55:53 (50.9:49.1%). Tumors were mostly well- to moderately differentiated (88 cases, 81.5%), although 7 (6.5%) cases exhibited poor differentiation, and 13 (12.0%) cases were mucinous adenocarcinomas. Pdc4 was usually detected in the nuclei of non-neoplastic epithelial cells (Fig. 1A). However, in the cancer cells, Pdc4 was observed in both the nuclei and the cytoplasm, and nuclear expression was downregulated. The nuclear, cytoplasmic and overall Pdc4 scores were evaluated, respectively. An overall Pdc4 score of 0, 1, 2 and 3 was identified in 11 (10.2%), 33

(30.5%), 57 (52.8%) and 7 (6.5%) cases, respectively. In contrast to normal epithelial cells, the majority of tumor cells had a weak to moderate cytoplasmic staining pattern for Pdc4. A cytoplasmic expression score of 1, 2, 3 and 4 was observed in 63 (58.3%), 25 (23.1%), 17 (15.8%) and 3 (2.8%) cases, respectively (Table II). The cytoplasmic and overall score revealed a tendency to increase gradually according to the nodal status and AJCC clinical stage (linear-by-linear association, $p < 0.05$). The cytoplasmic score also revealed significant difference among nodal status and AJCC clinical stages (Pearson's Chi-square test, $p < 0.05$). Moreover, nuclear expression scores of 1, 2, 3 and 4 were identified in 24 (22.2%), 18 (16.7%), 20 (18.5%) and 46 (42.6%) cases, respectively, and showed no tendency or significant difference with any of the clinicopathological parameters (Table II, Fig. 1B-F).

Discussion

Pdc4 was first identified as being differentially upregulated during apoptosis; however, experimental evidence has established it as a novel tumor suppressor (5-8). Pdc4 has been identified as a suppressor of transformation (5,9,19), tumori-




Figure 1. Various expression patterns of programmed death cell 4 (Pdc44) in colorectal adenocarcinoma and normal cells. (A) Most normal colorectal epithelial cells revealed strong nuclear staining (upper left) compared with the staining in adenocarcinoma. (B-F) In carcinoma, various expression patterns were identified. (B) Moderate nuclear and weak cytoplasmic expression, (C) strong nuclear and weak cytoplasmic expression, (D) weak nuclear with moderate cytoplasmic expression, (E) extremely strong cytoplasmic expression, and (F) no expression in either the nuclei or cytoplasm.

genesis, progression (1), invasion, matrix-metalloproteinase activation (3,8) and tumor growth (20). In the present study, the Pdc44 expression pattern was investigated in CRA (Fig. 1). Nuclear expression exhibited a tendency to be lost or weak in tumor cells, whereas cytoplasmic expression was increased. The transfer of Pdc44 between the nucleus and cytoplasm is understood to have a significant effect on the regulation of its function. Pdc44 is localized predominantly in the nucleus under normal growth conditions but is exported to the cytoplasm upon serum withdrawal (14). A number of authors have studied the localization of Pdc44 in various cell types. Yang *et al* (19) reported that in the mouse JB6 preneoplastic cell line, both nuclear and cytoplasmic localization of Pdc44 were identified. In their study, Yoshinaga *et al* (21) found that Pdc44 accumulated in the nucleus at the G0 phase of asynchronous cultures of human normal fibroblasts but was localized in the cytoplasm during the cell cycle in tumor cell lines. Goke *et al* (7) reported that 6 of 7 colon carcinomas examined revealed a complete absence of nuclear Pdc44 staining and additional cytoplasmic staining in 4 tumors. According to this study, epithelial cells of the prostate, breast and lung exhibited intense nuclear but no cytoplasmic staining; a clear shift from nuclear localization to cytoplasmic staining was observed in all colonic adenomas investigated. In the case of invasive breast cancer tissues, both nuclear and cytoplasmic localization were observed. In an investigation conducted into colorectal cancer, in addition to a decreased expression level of Pdc44, a significant loss of nuclear Pdc44 from normal tissues to colonic adenomas and carcinomas was also observed (18), supporting the hypothesis that the intracellular localization of Pdc44 plays a significant role in the regulation of tumor cell progression. In the present study, similar results to other authors were also demonstrated, in that a differential expression pattern of Pdc44 was found between normal and carcinoma cells by IHC analysis. Additionally, the cytoplasmic

expression level was shown to increase according to the AJCC clinical stage. We suggested that Pdc44 translocates from the nucleus to the cytoplasm during colonic cancer development. As proposed by Zhang (23), the accumulation of Pdc44 in the nuclei is crucial for apoptosis, and the regulatory mechanisms of the localization of Pdc44 protein may play a significant role in the induction of apoptosis in hepatocellular carcinoma cells. Wei *et al* (13) speculated that the accumulation of Pdc44 in the nucleus negatively regulates cell proliferation while the cytoplasmic sequestration of Pdc44 may abolish its involvement in ovarian cancer cells. Palamarchuk *et al* (22) transfected 293 cells that stably expressed wild-type Pdc44 and observed a primarily nuclear Pdc44 staining pattern. However, an S457A mutant of Pdc44 that was unable to be phosphorylated by Akt, in contrast to the wild-type, was not localized in the nucleus but in the cytoplasm, indicating the significance of this amino acid and Akt phosphorylation site in the localization of Pdc44. However, this phenomenon requires further confirmation using a larger patient cohort, and *in vitro* studies are necessary in order to understand its regulatory mechanism. In conclusion, both the localization and expression level of Pdc44 may be potential indicators of colonic cancer progression.

The present study investigated the Pdc44 expression pattern in CRA using IHC, and analyzed the correlation between Pdc44 expression and clinical and pathological parameters, including histology, grade, stage and overall survival. In conclusion, in CRA, nuclear Pdc44 expression was decreased and inversely aberrant cytoplasmic staining was identified, which was significantly increased according to the nodal status and AJCC clinical stage. We suggest that Pdc44 translocates from the nucleus to cytoplasm in CRA progression, and that increased cytoplasmic expression of Pdc44 is a potential prognostic marker in CRA. However, the regulatory mechanism of alteration of subcellular localization remains to be further investigated.

Acknowledgements

This study was supported by the National Research Foundation of Korea (NRF) Grant funded by the Ministry of Education, Science and Technology (MEST) through the Research Center for Resistant Cells (R-13-1003-09).

References

- Jansen AP, Camalier CE and Colburn NH: Epidermal expression of the translation inhibitor programmed cell death 4 suppresses tumorigenesis. *Cancer Res* 65: 6034-6041, 2005.
- Hilliard A, Hilliard B, Zheng SJ, *et al*: Translational regulation of autoimmune inflammation and lymphoma genesis by programmed cell death 4. *J Immunol* 177: 8095-8102, 2006.
- Yang HS, Matthews CP, Clair T, *et al*: Tumorigenesis suppressor Pdc4 down-regulates mitogen-activated protein kinase kinase 1 expression to suppress colon carcinoma cell invasion. *Mol Cell Biol* 26: 1297-1306, 2006.
- Soejima H, Miyoshi O, Yoshinaga H, *et al*: Assignment of the programmed cell death 4 gene (Pdc4) to human chromosome band 10q24 by in situ hybridization. *Cytogenet Cell Genet* 87: 113-114, 1999.
- Cmarik JL, Min H, Hegamyer G, *et al*: Differentially expressed protein Pdc4 inhibits tumor promoter-induced neoplastic transformation. *Proc Natl Acad Sci USA* 96: 14037-14042, 1999.
- Bitomsky N, Bohm M and Klempnauer KH: Transformation suppressor protein Pdc4 interferes with JNK-mediated phosphorylation of c-Jun and recruitment of the coactivator p300 by c-Jun. *Oncogene* 23: 7484-7493, 2004.
- Goke R, Barth P, Schmidt A, Samans B and Lankat-Buttgereit B: Programmed cell death protein 4 suppresses CDK1/cdc2 via induction of p21(Waf1/Cip1). *Am J Physiol Cell Physiol* 287: 1541-1546, 2004.
- Leupold JH, Yang HS, Colburn NH, Asangani I, Post S and Allgayer H: Tumor suppressor Pdc4 inhibits invasion/intravasation and regulates urokinase receptor (u-PAR) gene expression via Sp-transcription factors. *Oncogene* 26: 4550-4562, 2007.
- Yang HS, Jansen AP, Nair R, *et al*: A novel transformation suppressor, Pdc4, inhibits AP-1 transactivation but not NF- κ B or ODC transactivation. *Oncogene* 20: 669-676, 2001.
- Yang HS, Knies JL, Stark C and Colburn NH: Pdc4 suppresses tumor phenotype in JB6 cells by inhibiting AP-1 transactivation. *Oncogene* 22: 3712-3720, 2003.
- Li T, Li D, Sha J, Sun P and Huang Y: MicroRNA-21 directly targets MARCKS and promotes apoptosis resistance and invasion in prostate cancer cells. *Biochem Biophys Res Commun* 383: 280-285, 2001.
- Wen YH, Shi X, Chiriboga L, Matsubashi S, Yee H and Afonja O: Alterations in the expression of Pdc4 in ductal carcinoma of the breast. *Oncol Rep* 18: 1387-1393, 2007.
- Wei NA, Liu SS, Leung TH, *et al*: Loss of Programmed cell death 4 (Pdc4) associates with the progression of ovarian cancer. *Mol Cancer* 8: 70, 2009.
- Bohm M, Sawicka K, Siebrasse JP, Brehmer-Fastnacht A, Peters R and Klempnauer KH: The transformation suppressor protein Pdc4 shuttles between nucleus and cytoplasm and binds RNA. *Oncogene* 22: 1905-1910, 2003.
- Lankat-Buttgereit B, Gregel C, Knolle A, Hasilik A, Arnold R and Goke R: Pdc4 inhibits growth of tumor cells by suppression of carbonic anhydrase type II. *Mol Cell Endocrinol* 214: 149-153, 2004.
- Park JK, Hong R, Kim KJ, Lee TB and Lim SC: Significance of p-STAT3 expression in human colorectal adenocarcinoma. *Oncol Rep* 20: 597-604, 2008.
- Greene FL, Page DL, Fleming ID, *et al* (eds.): *AJCC Cancer Staging Manual*. 6th edition. Springer-Verlag, New York, pp157-164, 2002.
- Mudduluru G, Medved F, Grobholz R, *et al*: Loss of programmed cell death 4 expression marks adenoma-carcinoma transition, correlates inversely with phosphorylated protein kinase B, and is an independent prognostic factor in resected colorectal cancer. *Cancer* 110: 1697-1707, 2007.
- Yang HS, Jansen AP, Komar AA, *et al*: The transformation suppressor Pdc4 is a novel eukaryotic translation initiation factor 4A binding protein that inhibits translation. *Mol Cell Biol* 23: 26-37, 2003.
- Chen Y, Knosel T, Kristiansen G, *et al*: Loss of Pdc4 expression in human lung cancer correlates with tumour progression and prognosis. *J Pathol* 200: 640-646, 2003.
- Yoshinaga H, Matsubashi S, Fujiyama C and Masaki Z: Novel human Pdc4 (H731) gene expressed in proliferative cells is expressed in the small duct epithelial cells of the breast as revealed by an anti-H731 antibody. *Pathol Int* 49: 1067-1077, 1999.
- Palamarchuk A, Efanov A, Maximov V, Aqeilan RI, Croce CM and Pekarsky Y: Akt phosphorylates and regulates Pdc4 tumor suppressor protein. *Cancer Res* 65: 11282-11286, 2005.
- Zhang H, Ozaki I, Mizuta T, *et al*: Involvement of programmed cell death 4 in transforming growth factor-beta1-induced apoptosis in human hepatocellular carcinoma. *Oncogene* 25: 6101-6112, 2005.