

Isolated metastases of hepatocellular carcinoma in the right atrium: Case report and review of the literature

MANRI KAWAKAMI¹, MASAHIKO KODA¹, MARI MANDAI¹, KEIKO HOSHO¹,
YOSHIKAZU MURAWAKI¹, WAKAKO ODA² and KAZUHIKO HAYASHI²

¹Department of Multidisciplinary Internal Medicine, Division of Medicine and Clinical Science, School of Medicine;

²Division of Molecular Pathology, Tottori University, Yonago 683-8504, Japan

Received November 28, 2012; Accepted February 21, 2013

DOI: 10.3892/ol.2013.1240

Abstract. The aim of this study was to clarify the clinical features of patients with isolated HCC metastases to the heart. A 66-year-old female hospitalized with a hepatocellular carcinoma (HCC) ranging from the right to the left lobe and with a tumor thrombus in the main portal vein, was treated with intraarterial cisplatin, 5-fluorouracil, adriamycin and mitomycin. Computed tomography (CT) one month later revealed that the HCC had progressed with multiple lung metastases and moderate ascites. The patient had no symptoms. Magnetic resonance imaging (MRI) and echocardiography revealed a round, movable tumor with a diameter of 2 cm in the right atrium (RA). The patient succumbed to HCC five months later. An autopsy revealed HCC with portal tumor thrombi and metastases to the lungs, inferior vena cava (IVC) and RA. The metastases in the RA and IVC were not continuous with the intrahepatic tumor and were histologically attached to the endocardium and endothelium, respectively. An isolated metastasis of a HCC of the RA and IVC is extremely rare. In conclusion, although the majority of isolated metastases of HCC to the heart were diagnosed by echocardiography and were treated with mainly surgery, they had poor prognosis. The echocardiography should be performed for patients with advanced HCC. A novel treatment including molecular targeted drugs is required.

Introduction

Hepatocellular carcinomas (HCCs) frequently invade the vascular system at points such as the portal and hepatic veins. The results of autopsy studies indicate a 2.7-4.1% incidence of atrial metastases of HCC (1,2). A correct diagnosis is important in the clinical setting since cardiac metastases are able to induce sudden cardiac arrest. The majority of metastases develop as continuous extensions of a tumor thrombus in the hepatic vein. However, isolated cardiac metastases are extremely rare. The present study describes a 66-year-old female with an isolated right atrial metastasis of a HCC and reviews previous published studies, treatments and outcomes in similar patients. Written informed consent was obtained from the patients' family.

Case report

A 66-year-old female was diagnosed with chronic hepatitis type B and HCC in May 2004. The patient had no jaundice, vascular spiders, palmar erythema or cardiac murmurs. A hard mass was palpable from the right hypochondrium to the epigastrium. The laboratory results (normal ranges in parentheses) were as follows: total bilirubin, 0.9 mg/dl (0.2-1.2 mg/dl); aspartate aminotransferase, 115 IU/l (5-40 IU/l); alanine aminotransferase, 105 IU/l (5-47 IU/l); alkaline phosphatase 491 IU/l (111-295 IU/l); and lactate dehydrogenase, 199 IU/l (100-225 IU/l). The patient was positive for the hepatitis B surface antigen and e-antibodies. The serum α -fetoprotein and des- γ -carboxy prothrombin levels were elevated to 687,460 ng/ml (normal range, <13.2 ng/ml) and 1037 mAU/ml (normal range, <40 mAU/ml), respectively.

Abdominal sonography and computed tomography (CT) imaging revealed a large mass reaching from the right to the left lobe and a tumor thrombus in the main portal vein (Fig. 1A and B). Angiography revealed a hypervascular tumor in the right lobe exhibiting the thread and streak sign. No metastases were identified in the right atrium (RA) or inferior vena cava (IVC) prior to starting intraarterial chemotherapy with cisplatin, 5-fluorouracil, adriamycin and mitomycin.

An enhanced CT in July, 2004, showed that the HCC had progressed and that multiple lung metastases had developed with moderate ascites. Magnetic resonance imaging (MRI)

Correspondence to: Dr Masahiko Koda, Department of Multidisciplinary Internal Medicine, Division of Medicine and Clinical Science, Tottori University School of Medicine, 36-1 Nishicho, Yonago 683-8504, Japan
E-mail: masakoda@grape.med.tottori-u.ac.jp

Abbreviations: CT, computed tomography; HCC, hepatocellular carcinoma; MRI, magnetic resonance imaging

Key words: hepatocellular carcinoma, isolated atrial metastasis, prognosis




Figure 1. Imaging findings. (A) Abdominal sonographic image showing ill-defined large isoechoic mass with portal thrombus (arrow). (B) Computed tomography (CT) image showing ill-defined large heterogeneous mass occupying the bilateral liver lobes, and portal thrombus accompanying a cavernous transformation. (C) Transthoracic echocardiography image showing a round, movable tumor of a 2-cm diameter in the right atrium (RA; arrow).

and echocardiography revealed a round, movable tumor with a diameter of 2 cm in the RA (Fig. 1C), but no tumor thrombus in the IVC. The atrial tumor was not continuous with the intrahepatic HCC. Anticoagulation therapy with warfarin was administered, however the patient succumbed to hepatic failure five months later (Fig. 2).

An autopsy revealed diffuse-type HCC in the bilateral lobes of the liver that weighed 1,365 g, with a tumor thrombus in the main trunk of the portal vein, bilateral multiple lung metastases and tumor thrombi in the artery of the right upper lung. A yellowish irregular-surfaced mass with a diameter of 10 mm located in the RA, and a similar small independent mass in the IVC (Fig. 3), were discontinuous with the intrahepatic HCC. Histologically, the intrahepatic and right atrial




Figure 2. Clinical course of the patient. ADR, adriamycin; AFP, α -fetoprotein; AG, angiography; CDDP, cisplatin; DCP, des- γ -carboxy prothrombin; 5-FU, 5-fluorouracil; IFN- β , interferon β ; MMC, mitomycin C.




Figure 3. Appearance and diagnosis of autopsy specimen. (A) Yellowish tumor with irregular surface (arrow) located in the right atrium (RA) of the heart specimen at autopsy. (B) Right atrial tumor attached to the endocardium diagnosed as a moderately differentiated HCC of trabecular type (hematoxylin-eosin stain). (C) Greenish tumor with irregular surface (arrow) located in inferior vena cava (IVC), discontinuous with the intrahepatic HCC. (D) Tumor attached to the endothelium in the IVC, pathologically diagnosed as a moderately differentiated HCC of trabecular type (hematoxylin-eosin stain). HCC, hepatocellular carcinoma.

Table I. Patients with isolated HCC metastases to the heart.

Case	Ref.	Age (years), gender	Diagnosis stage	Symptoms	Pretreatment for cardiac metastases	Pathology	Location of initial HCC	Treatment for cardiac metastase	Prognosis
1	3	52M	Autopsy	Dyspnea, heart murmur	None	NS	RA, endocardium	Supportive care	Succumbed after 1 month
2	4	55M	Autopsy	Dyspnea, cyanosis	None	Ed II-III	RA, LA, endocardium	Supportive care	Succumbed after 1 month
3	5	67M	Autopsy	Fever, cough, chest discomfort	None	Ed III	LV, epicardium	Supportive care	Succumbed after 2 months
4	6	73M	Autopsy	Heart murmur	None	Ed I-II	RA, RV	TAE	Succumbed after 1 month
5	7	51M	CT	Dyspnea, palpitation	Hepatectomy	NS	RV, myocardium	Surgery	Succumbed
6	7	29M	UCG	Dyspnea	Hepatectomy	Ed II	LA	Chemotherapy	Succumbed
7	8	71M	Autopsy	Consciousness disorder	Chemotherapy	Ed II	RV, LV, myocardium	NS	Succumbed after 0.5 months
8	9	76M	UCG	Dyspnea	Hepatectomy, MCT, chemotherapy	Ed I-II	RV, myocardium	Surgery	Succumbed after 6 months
9	10	49F	UCG	Dyspnea, palpitation, heart murmur	Hepatectomy	Ed I-II	RV, myocardium	Surgery	Alive at 21 months
10	11	67F	UCG	Dyspnea, heart murmur	Hepatectomy	Ed II	RV	TAE	Alive at 3 months
11	12	43M	UCG, CT	Dyspnea, cough	Hepatectomy	Ed I-II	RV	Supportive care	Succumbed after 20 days
12	13	45M	UCG, CT	Dyspnea, dizziness	Hepatectomy	NS	RV	Surgery	Alive at 3 months
13	14	65F	CT	Dyspnea	Hepatectomy	NS	RV	Surgery	Succumbed after 3 months
14	15	45F	UCG	Dyspnea, syncope	Hepatectomy, TAE	Ed II	RV	Surgery	Succumbed after 4 months
15	16	63F	UGC	Dyspnea	TAE	Ed IV	RA	Surgery	NS
16	17	74F	UCG	Dyspnea, syncope	TAE	NS	RV	Surgery	Succumbed after 4 months
17	Present study	66F	UCG	None	Chemotherapy	Ed II	RA, endocardium	TAI	Succumbed after 5 months

UCG, ultrasound cardiography; MCT, microwave coagulation therapy; TAE, transcatheter arterial embolization; Ed, Edmondson; NS, not stated; RA, right atrium; RV, right ventricle; LA, left atrium; LV, left ventricle; TAI, transcatheter arterial infusion chemotherapy; HCC, hepatocellular carcinoma.

tumors were moderately differentiated HCCs. The right atrial tumor was fixed to the atrial wall and arose from sites on the endocardium of the RA (Fig. 3A and B). The small tumor in the IVC was similarly fixed to the endothelium (Fig. 3C and D).

Discussion

The mechanism of cardiac metastases is as a contiguous extension from the intrahepatic HCC via a tumor thrombus to the IVC or by lymphatic or hematogenous spread. The majority of cardiac metastases are direct and contiguous extensions of the intrahepatic HCC. Isolated cardiac metastases that are discontinuous with an intrahepatic HCC are extremely rare. The literature was searched for descriptions of isolated cardiac metastases of HCC and 17 patients were identified (Table I) with a mean age of 58 ± 13 years. In total, 15 patients (88.2%) had symptoms, with 13 (76.5%) suffering dyspnea. Intermittent obstruction by tumors in the cardiac cavity led to

the symptoms of ball valve thrombus syndrome, which is able to induce sudden cardiac arrest (18). However, in the present study, the patient had no symptoms since the diameter of the tumor was 2 cm.

Extracorporeal echocardiography was useful for detecting the atrial tumor in the present case. Kanematsu *et al* (19) described a tumor thrombus of HCC in the IVC detected by CT and MRI. Yoshitomi *et al* (20) and Van Camp *et al* (21) found transesophageal echocardiography useful. In theses studies, cardiac metastases were diagnosed by UCG and CT in 10 (58.8%) and 4 (23.5%) patients, respectively. Cardiac metastases were identified at autopsy in the early cases. The HCC was initially treated in 13 (76.5%) of the 17 patients [hepatectomy, n=9; transcatheter arterial embolization (TAE), n=3; chemotherapy, n=3] and then the cardiac metastases were identified.

In the present study, the tumors in the RA and IVC were not continuous growths from the intrahepatic HCC in

the patient. The two tumors were individually isolated and attached to the endocardium or endothelium, respectively. This metastatic pathway was considered to be comprised of two mechanisms. The first was the seeding of a blood flow onto the endocardium or endothelium from the intrahepatic HCC and the second was cancer embolisms of the small vessels under the endocardium and endothelium via arterial systemic spread. However, in the published cases, the mechanism associated with the metastases could not be clarified. Cardiac metastases were notably located in the right ventricle (RV), RA and left ventricle (LV) in 10 (58.8%), 5 (29.4%) and 2 (11.8%) patients, respectively. A greater number of metastases had invaded the RV than the RA. These findings indicate that the main mechanism is embolism via arterial systemic spread as massive myocardial involvement was described in 4 patients.

Surgical (22) or non-surgical approaches, including TAE (23), transcatheter arterial infusion chemotherapy (TAI) (24) and radiotherapy (25), have been undertaken to treat IVC/RA metastases. In the selected published cases, eight (47.1%) out of 17 patients underwent surgery to relieve symptoms. In the present study, the patient underwent TAI only, but succumbed to HCC progression five months later. Overall, all the patients had an extremely poor prognosis regardless of the treatment strategy. Chang *et al* (26) reported that thalidomide is a useful angiogenesis inhibitor for IVC/RA metastases and new molecular target drugs, such as Sorafenib, have since emerged (27). However, further studies involving a larger cohort of patients with IVC/RA tumor thrombi are required.

References

1. Kojiro M, Nakahara H, Sugihara S, Murakami T, Nakashima T and Kawasaki H: Hepatocellular carcinoma with intra-atrial tumor growth. A clinicopathologic study of 18 autopsy cases. *Arch Pathol Lab Med* 108: 989-992, 1984.
2. MacDonald RA: Primary carcinoma of the liver; a clinicopathologic study of one hundred eight cases. *AMA Arch Intern Med* 99: 266-279, 1957.
3. Kato Y, Kurosaki M, Kobayashi K, Sugimoto T and Takada A: Liver cancer associated with a large spheric thrombus caused by tumor in the right atrium of the heart. *Naika* 28: 349-353, 1971 (In Japanese).
4. Takigami S, Komai Y, Lai Y-S, *et al*: An autopsy case of minute hepatoma accompanied by a large intraatrial tumor embolus. *Acta Hepat Jap* 21: 478-486, 1980 (In Japanese).
5. Yoshioka K, Izuchi M, Okayama M, *et al*: An autopsied case of hepatocellular carcinoma with metastases to the heart and spleen. *Acta Hepat Jap* 24: 909-913, 1983 (In Japanese).
6. Miura Y, Kondo K, Watanabe T, *et al*: An autopsy case of hepatocellular carcinoma with tumor thrombus in the right atrium. *Nihon Shokakibyo Gakkai Zasshi* 86: 2460-2463, 1989.
7. Lei MH, Ko YL, Kuan P, Lien WP and Chen DS: Metastases of hepatocellular carcinoma to the heart: unusual patterns in three cases with antemortem diagnosis. *J Formos Med Assoc* 91: 457-461, 1992.
8. Kimitsuki H, Nagasaki Y, Yamashita Y, Kurohiji T, Kakegawa T and Nakashima T: An autopsied case of hepatocellular carcinoma with cardiac metastasis. *Acta Hepat Jap* 35: 629-632, 1994 (In Japanese).
9. Shiota S, Konishi I, Sato N, *et al*: A case of hepatocellular carcinoma with metastases to the right ventricle of the heart. *Jpn J Clin Surg* 61: 1542-1546, 2000 (In Japanese).
10. Murakawa T, Takamoto S, Ezure M, Ono M, Kawauchi M and Tanaka O: Metastatic hepatocellular carcinoma obstructing the right ventricular outflow tract. *Jpn J Thorac Cardiovasc Surg* 48: 516-519, 2000.
11. Kotani E, Kiuchi K, Takayama M, *et al*: Effectiveness of transcatheter chemoembolization for metastatic right ventricular tumor derived from hepatocellular carcinoma. *Chest* 117: 287-289, 2000.
12. Longo R, Mocini D, Santini M, *et al*: Unusual sites of metastatic malignancy: case 1. Cardiac metastasis in hepatocellular carcinoma. *J Clin Oncol* 22: 5012-5014, 2004.
13. Lin TY, Chiu KM, Chien CY, Ming C, Wang MJ and Chu SH: Case 1. Right ventricular outflow obstruction caused by metastatic hepatocellular carcinoma. *J Clin Oncol* 22: 1152-1153, 2004.
14. Chiang SH, Lin CH, Lu MJ and Hung CR: Intracavitary metastatic hepatocellular carcinoma of the right ventricle. *Thorac Cardiovasc Surg* 53: 123-125, 2005.
15. Liu YC, Hung CS, Chan SY, Chen YW, Shun CT and Lai LP: Asymptomatic metastasis of hepatocellular carcinoma into the right ventricular cavity presenting with electrocardiographic changes. *Acta Cardiol Sin* 22: 180-183, 2006.
16. Sekine Y, Kitano M, Akimoto T and Matsuda K: Isolated intracavitary metastatic hepatocellular carcinoma of the right atrium without inferior vena cava involvement: report of a case. *Kyobu Geka* 60: 504-507, 2007 (In Japanese).
17. Kan CB, Chang RY and Chen CK: Isolated right ventricular intracavitary metastasis of hepatocellular carcinoma in a 74-year-old woman. *J Clin Med Assoc* 71: 318-320, 2008.
18. Sung AD, Cheng S, Moslehi J, Scully EP, Prior JM and Loscalzo J: Hepatocellular carcinoma with intracavitary cardiac involvement: a case report and review of the literature. *Am J Cardiol* 102: 643-645, 2008.
19. Kanematsu M, Imaeda T, Minowa H, *et al*: Hepatocellular carcinoma with tumor thrombus in the inferior vena cava and right atrium. *Abdom Imaging* 19: 13-316, 1994.
20. Yoshitomi Y, Kojima S, Sugi T, *et al*: Echocardiography of a right atrial mass in hepatocellular carcinoma. *Heart Vessels* 13: 45-48, 1998.
21. Van Camp G, Abdulsater J, Cosyns B, Liebens I and Vandenbossche JL: Transesophageal echocardiography of right atrial metastasis of a hepatocellular carcinoma. *Chest* 105: 945-947, 1994.
22. Yoshidome H, Takeuchi D, Kimura F, *et al*: Treatment strategy for hepatocellular carcinoma with major portal vein or inferior vena cava invasion: a single institution experience. *J Am Coll Surg* 212: 796-803, 2011.
23. Mizuno S, Kato H, Azumi Y, *et al*: Total vascular hepatic exclusion for tumor resection: a new approach to the intra-thoracic inferior vena cava through the abdominal cavity by cutting the diaphragm vertically without cutting the pericardium. *J Hepatobiliary Pancreat Sci* 17: 197-202, 2010.
24. Murakami E, Aikata H, Miyaki D, *et al*: Hepatic arterial infusion chemotherapy using 5-fluorouracil and systemic interferon- α for advanced hepatocellular carcinoma in combination with or without three-dimensional conformal radiotherapy to venous tumor thrombosis in hepatic vein or inferior vena cava. *Hepatol Res* 42: 442-453, 2012.
25. Komatsu S, Fukumoto T, Demizu Y, *et al*: The effectiveness of particle radiotherapy for hepatocellular carcinoma associated with inferior vena cava tumor thrombus. *J Gastroenterol* 46: 913-920, 2011.
26. Chang JY, Ka WS, Chao TY, Liu TW, Chuang TR and Chen LT: Hepatocellular carcinoma with intra-atrial tumor thrombi. A report of three cases responsive to thalidomide treatment and literature review. *Oncology* 67: 320-326, 2004.
27. Llovet JM, Ricci S, Mazzaferro V, *et al*: SHARP Investigators Study Group: Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 359: 378-390, 2008.