

Cell death pathway induced by resveratrol-bovine serum albumin nanoparticles in a human ovarian cell line

LIYUAN GUO^{1*}, YAN PENG^{2*}, YULIAN LI¹, JINGPING YAO², GUANGMEI ZHANG³,
JIE CHEN¹, JING WANG¹ and LIHUA SUI¹

¹Department of Obstetrics and Gynecology, Third Affiliated Hospital, Harbin Medical University, Harbin 150081;

²Department of Health, First Affiliated Hospital, Heilongjiang University of Chinese Medicine, Harbin 150040;

³Department of Health, First Affiliated Hospital, Harbin Medical University, Harbin 150001, P.R. China

Received November 21, 2013; Accepted August 14, 2014

DOI: 10.3892/ol.2015.2851

Abstract. Resveratrol-bovine serum albumin nanoparticles (RES-BSANP) exhibit chemotherapeutic properties, which trigger apoptosis. The aim of the present study was to investigate the caspase-independent cell death pathway induced by RES-BSANP in human ovarian cancer SKOV3 cells and to analyze its mechanism. Morphological changes were observed by apoptotic body/cell nucleus DNA staining using inverted and fluorescence microscopy. The cell death pathway was determined by phosphatidylserine translocation. Western blot analysis was conducted to detect the activation of apoptosis-inducing factor (AIF), cytochrome *c* (Cyto *c*) and B-cell lymphoma 2-associated X protein (Bax). Apoptotic body and nuclear condensation and fragmentation were observed simultaneously following treatment with RES-BSANP. RES-BSANP induced apoptosis in a dose-dependent manner in the human ovarian cancer SKOV3 cells. The translocation of AIF from the mitochondria to the cytoplasm occurred earlier than that of Cyto *c*. In addition, Bax binding to the mitochondria was required for the release of AIF and Cyto *c* from the mitochondria. The AIF apoptosis pathway may present an alternative caspase-dependent apoptosis pathway in human ovarian cell death induced by RES-BSANP. Elucidation of this pathway may be critical for the treatment of cancer using high doses of RES-BSANP.

Introduction

Resveratrol (RES) exhibits anti-inflammatory and anti-oxidant effects (1,2) and is a potential chemopreventive

agent, which is able to inhibit various stages of carcinogenesis, including the initiation, promotion and progression of tumors (3). RES has been found to induce apoptotic cell death in various cancer cell lines and experimental tumor models (4-7).

Previously, it has been reported that high doses of RES (3 g/kg/day) cause moderate liver toxicity (8). However, the absorption of RES was shown to be poor due to its water insoluble properties (9). Furthermore, the insolubility of RES has been demonstrated to hamper the *in vitro* and *in vivo* biological induction of RES activity (10). Research in nanomedicine has not only become a frontier movement, but is also a revolutionary drug delivery field. Bovine serum albumin (BSA) has been used as a vehicle used for diagnostic and therapeutic agents (11,12), as it is non-toxic, safe, economical and exhibits good biocompatibility and no immunogenicity. Previous studies have explored BSA as a drug carrier (13,14).

RES-BSA nanoparticles (RES-BSANP) are synthesized by protein desolvation chemical crosslinking. Previous studies have demonstrated the antiproliferative effects of RES on the growth of human SKOV3 cell lines in *in vivo* and *in vitro* studies (15-17). The prepared RES-BSANP exhibited an altered distribution. In addition, it was demonstrated that RES-BSANP-treated tumors exhibited a similar apoptotic index to RES control tumors. The cells in the therapeutic groups exhibited apoptotic and necrotic characteristics. The mechanism resulting in these properties may be the release of cytochrome *c* (Cyto *c*) and the upregulation of the dynamic expression of caspase-9 and -3, indicating that the mitochondrial apoptotic pathway was activated (16,17).

Programmed cell death (PCD) is a specific mechanism that initiates cell death. Caspases are involved in PCD. PCD may be divided into two types, caspase-dependent and caspase-independent PCD, according to the involvement of caspase in PCD. Caspase-dependent PCD presents typical apoptosis. Caspase-independent PCD includes autophagy, paraptosis, mitotic catastrophe, apoptosis-like PCD and necrosis-like PCD.

RES-BSANP has been found to inhibit various stages of tumor growth, however, the molecular mechanism of its anti-cancer activity remains unclear, particularly in ovarian cancer. The aim of the present study was to elucidate the molecular

Correspondence to: Dr Jing Wang or Dr Lihua Sui, Department of Obstetrics and Gynecology, Third Affiliated Hospital, Harbin Medical University, 194 Xuefu Road, Harbin 150081, P.R. China
E-mail: wangjing_doctor@163.com
E-mail: ysbj2012@163.com

*Contributed equally

Key words: resveratrol-bovine serum albumin nanoparticles, apoptosis, programmed cell death, caspase-independent

events that occur during RES-BSANP-induced apoptotic cell death in human ovarian SKOV3 cells.

Materials and methods

Reagents. The human ovarian cancer SKOV3 cell line was obtained from the Tumor Research Institute of Harbin Medical University (Harbin Medical University, Harbin, China). RES was purchased from Xian Huacui Biology Co., Ltd. (Xian, China), (purity, $\geq 99.9\%$) and dissolved in dimethyl sulfoxide (DMSO) as a stock solution of 100 mmol/l. RES was further diluted in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum (FBS) to the appropriate final concentrations. RES-BSANP was prepared at the Life Science Laboratory of Northeast Forestry University (Harbin, China).

The general caspase inhibitor, Z-VAD-FMK [Z-Val-Ala-Asp(OMe)-CH₂F], and caspase-9 [Z-LEHD-FMK, Z-Leu-Glu(OMe)-His-Asp(OMe)-CH₂F] were obtained from Calbiochem (La Jolla, CA, USA). Stock solutions of the caspase inhibitors (10 mmol/l each) were prepared in DMSO and diluted in DMEM with 10% FBS to a final concentration of 100 μ mol/l. Polyclonal rabbit anti-rat antibodies against apoptosis-inducing factor (AIF), Cyto *c* and B-cell lymphoma 2 (Bcl-2)-associated X protein (Bax) were obtained from Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd. (Beijing, China). DMEM, penicillin (1 mg/ml) and streptomycin (1 mg/ml) were obtained from Invitrogen Life Technologies (Carlsbad, CA, USA) and FBS was purchased from HyClone Laboratories, Inc. (Logan, UT, USA). The 2.5% trypsin/EDTA solution was purchased from Invitrogen Life Technologies and diluted to 0.5% for trypsinizing attached cells.

Cell culture. The ovarian cancer SKOV3 cell line was cultured in RPMI 1640 (Gibco-BRL, Carlsbad, CA, USA) supplemented with 10% heat-inactivated fetal calf serum (FCS; Gibco-BRL) in a humidified incubator at 37°C, with an atmosphere of 5% CO₂. A total of 2.5 g/l trypsin and 0.2 g/l EDTA were used for subculture.

Morphological study. To evaluate apoptotic cell death, the cells were seeded at a density of 5×10^5 cells/ml in 15-mm diameter wells and incubated for 12 h until the cells had adhered to the bottom of the wells. DNA ladder formation was examined an hour following the start of treatment with the compounds. For morphological examination of the apoptotic changes, the cells were stained with Hoechst 33342 (5 μ g/ml; Invitrogen Life Technologies) at 37°C for 30 min, washed twice with phosphate-buffered saline, pipetted drop-wise onto a glass slide, and examined by fluorescence microscopy using an Olympus microscope (Olympus Corporation, Tokyo, Japan) equipped with an epi-illuminator and appropriate filters.

DNA fragmentation. DNA fragmentation was conducted as previously described (18). Following various treatments, the cells were collected and lysed with lysis buffer [0.5% Triton X-100, 5 mmol/l Tris Buffer (pH 7.4), 20 mmol/l EDTA]. RNA was removed by incubation with RNase A (0.8 mg/ml) at 37°C for 30 min. DNA was extracted by phenol/chloroform and precipitated with 1/10 volume of 3 mol/l sodium acetate

(pH 5.2) along with 20 μ l of 100% ethanol. DNA pellets obtained by centrifugation at 10,000 \times g for 20 min at 4°C, were dried and re-suspended in 25 ml 1xTAE (40 mmol/l Tris-acetate and 1 mmol/l EDTA). Samples were then separated on 2% agarose gels and the DNA ladder was detected by incubation of the gels with ethidium bromide (1 g/ml) for 20 min followed by de-staining with distilled H₂O.

Inhibition of apoptosis by the pan-caspase inhibitor. The tripeptide pan-caspase inhibitor, Z-VAD-FMK was added 12 h prior to treatment with RES-BANP. The optimal concentration of the inhibitor was determined from a dose-response curve using the extent of cell death. The inhibition of apoptosis by Z-VAD-FMK was evaluated by investigating the inhibition of nucleosomal DNA fragmentation, which was observed as DNA ladder formation.

Western blot analysis. Western blot analysis was performed to detect the protein expression of AIF, Cyto *c* and Bax. Cell lysate was prepared by lysis buffer protein extraction [40 mmol/l Tris-Cl (pH 8.0), 120 mmol/l NaCl and 0.1% NP40] supplemented with protease inhibitors. Proteins were separated by SDS-PAGE and transferred to nitrocellulose membranes (Bio-Rad, Hercules, CA, USA). The membranes were blocked with 5% skimmed milk in Tris-buffered saline and incubated with the appropriate primary antibodies for 1 h at room temperature. The blots were developed with peroxidase-conjugated secondary antibody, and the proteins were visualized with an enhanced chemiluminescence kit (GE Healthcare, Piscataway, NJ, USA) according to the manufacturer's instructions.

Statistical analysis. Results are expressed as the mean \pm standard deviation. For multiple comparisons, results were analyzed by one-way analysis of variance. The least significant difference was analyzed by the Bonferroni correction test to identify significant differences between the individual cell groups. $P < 0.05$ was considered to indicate a statistically significant difference. All statistical analyses were performed using SPSS software, version 11.5 (SPSS, Inc., Chicago, IL, USA).

Results

Cell detection via microscopic observations. Following observation under an inverted phase contrast microscope (IX73-F22FL/PH, Olympus Corp., Tokyo, Japan) and a fluorescence microscope (CKX41, Olympus Corp.), numerous cells were found in the culture flask of the control group. The cells had adhered to the flask wall. The level of cell latency and refraction was high and the proliferation rate of the tumor cells was also high. Treatment with 10, 50 and 80 μ mol/l RES-BSANP for 2, 4, 6, 8, 12 and 24 h resulted in morphological cell changes, as observed by apoptotic body/cell nucleus DNA staining. The size of the cell bodies in the experimental groups treated with 50 μ mol/l RES-BSANP for 8 h were reduced significantly and had become round in shape. The latency and refractivity of the cells was weakened, with vacuoles appearing in the cytoplasm. As the concentration of RES-BSANP and the exposure time increased, the cells began to shrink and rupture. The morphology of the

SKOV3 cells became irregular, with surrounding debris, and the culture medium became turbid. A large number of cells collapsed, floated and died. The difference between apoptosis and necrosis induced by 50 $\mu\text{mol/l}$ RES-BSANP is shown in Table I.

DNA fragmentation. To further verify the apoptotic response, DNA fragmentation was examined. DNA ladder formation was observed in SKOV3 cells treated with 50 $\mu\text{mol/l}$ RES and 20 $\mu\text{mol/l}$ RES-BSANP for 4 h. Treatment duration was then extended to 6 h. Large scales of DNA ladder were detected, in contrast to the 50 $\mu\text{mol/l}$ Z-VAD-FMK control group. These results indicated that the marked suppression of cell growth by RES and RES-BSANP was attributable to apoptotic cell death. Gel electrophoresis exhibited DNA ladders and smears in the high-dose RES-BSANP groups (80 and 100 $\mu\text{mol/l}$ RES-BSANP) and exhibited only a smear in the Z-VAD-FMK control group (Fig. 1).

Western blot analysis of AIF, Cyto *c* and Bax expression in SKOV3 cells. Induction of apoptosis by RES-BSANP may involve the translocation of Cyto *c* and AIF. This translocation was investigated by biochemically fractionating different subcellular compartments and quantifying the expression of Cyto *c* and AIF by western blot analysis. Cells treated with 80 $\mu\text{mol/l}$ RES-BSANP for 2 h showed a spatiotemporal release of AIF from the mitochondria into the nucleus (Fig. 2). Similarly, it appeared that Cyto *c* was also released from the mitochondria, however, notably, this was not accompanied by a concomitant cytoplasmic increase at 4 h of treatment. The translocation of AIF from mitochondria to cytoplasm occurred earlier than that of Cyto *c*. These results indicated that RES-BSANP-elicited cell death may not occur via a classical Cyto *c* mitochondria cytosol translocation mechanism but rather, a caspase-independent mechanism of cell death via the nucleus directed shuttling of AIF and Cyto *c*.

Activation of Bax and its mitochondrial translocation are the key regulatory events during induction of mitochondrial membrane depolarization. Treatment of macrophages with macrophage inflammatory protein supernatant resulted in mitochondrial translocation of Bax as early as 2 h following treatment. Reduction of AIF and Cyto *c*, resulted in increased mitochondrial Bax. The binding of Bax to the mitochondria was required for the release of AIF and Cyto *c* from mitochondria.

Discussion

Previous studies have identified two primary forms of cell death, namely apoptosis and necrosis. Apoptosis was initially considered the primary physiological and programmed form of cell death. The apoptotic proteins involved with the caspase-dependent pathway include Cyto *c*, caspase-9 and caspase-3 (16,17). Cell death via this pathway is associated with the activation of caspases. However, it is now widely recognized that PCD may also occur in the absence of caspase activation (18-22). The existence of non-caspase PCD pathways has been found to be associated with caspase-independent elimination, including the use of mitochondrial

protein AIF (19,20,23,24). The current study revealed that the caspase-independent pathway, mediated by AIF, may be involved in the necrotic PCD pathway, particularly following treatment with high doses of RES-BSANP in cell culture.

Caspase activation is regarded as a vital factor in RES-BSANP-induced cell death in *in vitro* and *in vivo* studies. However, previous studies have indicated that RES-induced cell death in human ovarian cancer cells is caspase-independent (17,18); the treatment of these cells with RES-BSANP resulted in the release of Cyto *c* and the activation of caspase-3. Apoptotic cell death is induced by RES-BSANP and appears to be caspase-independent, as caspase inhibitors fail to attenuate induced cell death by RES treatment. It has previously been reported that AIF mediates cell death via a caspase-independent pathway (25). Mitochondrial AIF translocates to the nucleus as a result of cell death stimuli and thus initiates nuclear condensation (26,27). Once the nucleus condenses, this leads to large-scale chromatin fragmentation, followed by cell death. Consistent with these observations, the current study demonstrated translocation of AIF from the mitochondria to the nucleus following RES-BSANP treatment of the SKOV3 cells. AIF translocation and nuclear condensation were detectable within 24 h of RES-BSANP treatment. Whilst AIF protein is considered to be a potent factor in caspase-independent apoptosis, the mechanism by which AIF causes DNA ladder formation remains unclear.

It has been reported that AIF protein release is localized around the cell nuclei and is partly translocated into these nuclei following treatment with the apoptogenic dolichylmonophosphate in SKOV3 cells. Caspase-3 and -8 inhibitors prevent not only DNA fragmentation, but also AIF migration and chromatin condensation (28). In the current study, the pan-caspase inhibitor prevented induced apoptotic cell death following RES-BSANP treatment, indicating that AIF protein release, rather than caspase release, may be pivotal in DNA ladder formation. These results are consistent with previous studies of RES-BSANP-induced apoptosis.

To the best of our knowledge, this is the first study to investigate the caspase-independent signaling mechanism of RES-BSANP-induced apoptosis in SKOV3 cells. Further studies are required to elucidate the precise signaling pathways involved in the RES-BSANP-induced apoptotic death of ovarian cancer SKOV3 cells. Notably, RES-BSANP exhibited a potent effect on the SKOV3 cells, which indicates that polyphenol compounds may also be a candidate for a chemo-preventive and chemotherapeutic agent, as the doses of the compounds used can be considerably lower than the chemotherapy drugs commonly in use currently to exhibit the same activity as RES.

It has been reported that a member of the Bcl-family, Bax, is associated with the release of AIF from the mitochondria (29,30), and the release of Ca^{2+} from the endoplasmic reticulum appears to be important in this process (31). Results of previous studies indicate that the release of AIF, with a decrease in membrane potential and modulation of the expression of Bax may be partly responsible for RES-BSANP-induced apoptosis in SKOV3 cells. The polyphenol compounds have been shown to affect activation of caspase-independent cell death pathways and thus exhibit an anti-cancer activity. Recently, the understanding of how RES induces cell death in

Table I. Effects of resveratrol-bovine serum albumin nanoparticles (80 $\mu\text{mol/l}$) on apoptotic body/cell nucleus DNA staining.

Cell count	Duration of treatment, h					
	2	4	6	8	12	24
Apoptotic bodies	22 \pm 0.21	25 \pm 2.6	47 \pm 3.51	56 \pm 0.8	59 \pm 2.6	63 \pm 6.4
Necrosis	0.44 \pm 0.5	64 \pm 0.6 ^a	87 \pm 5.6 ^a	102 \pm 4.5 ^b	130 \pm 4.7 ^a	178 \pm 1.3 ^b

^aP<0.05, ^bP<0.01, compared with apoptotic bodies. Data are presented as the mean \pm standard deviation.




Figure 1. DNA Fragmentation. A, control; B, 20 $\mu\text{mol/l}$ RES-BSANP; C, 50 $\mu\text{mol/l}$ RES-BSANP; D, 80 $\mu\text{mol/l}$ RES-BSANP; E, 100 $\mu\text{mol/l}$ RES-BSANP; F, 50 $\mu\text{mol/l}$ Z-VAD-FMK and RES-BSANP; G, 50 $\mu\text{mol/l}$ Z-VAD-FMK; H, $\lambda\text{DNA}/\text{HindIII}$; I, 100-base pair DNA ladder. RES-BSANP, resveratrol-bovine serum albumin nanoparticles.




Figure 2. Western blot analysis of AIF, Cyto *c* and Bax protein expression following treatment with 50 $\mu\text{mol/l}$ resveratrol-bovine serum albumin nanoparticles for 0, 2 and 4 h. AIF, apoptosis-inducing factor; Cyto *c*, cytochrome *c*; Bax, B-cell lymphoma 2-associated protein X.

human ovarian cancers has markedly improved. RES usually inhibits signaling via the mitogen-activated protein kinase and phosphatidylinositol 3-kinase/AKT pathways (32-36). Consistently, RES has been found to suppress the activity of the downstream transcription factors AP-1 and nuclear factor- κB (RelA/p65) (35-38). Genes that are considered to be transcriptionally affected by RES with an impact on apoptosis include cyclins, cyclin-dependent kinases, caspases, p53, p21 (Cip1/WAF1), p300, NF- κB , Bcl-2, Bax and inhibitors of apoptosis (39,40). Consequently, further studies investigating RES-BSANP's role in cell death,

excluding the caspase-dependent and caspase-independent pathways, are required. A great deal remains to be determined with regard to the mechanism of cell necrosis or cell death by treatment with RES-BSANP.

Taken together, these results indicate that the main signal transduction pathway of RES-BSANP induced apoptosis in SKOV3 cells is mediated by the activation of caspase. Concurrently, additional intracellular signaling pathways may also be involved. Further research is required to clarify the association between the early response signal and the apoptotic signal. Thus, RES-BSANP, which is a constituent

of an anti-tumor compound, may be a potentially effective candidate for chemoprevention.

Acknowledgements

This study was supported by The Third Hospital of Harbin Medical University (Harbin, China). The authors would like to thank the Tumor Research Institute of Harbin Medical University for providing technical assistance. The study was funded by the State Traditional Chinese Medicine Foundation (grant no. 06-07ZP15), the Harbin Youth Scientific and Technological Innovation Foundation (grant no. 2011RFQYS087) and the China Postdoctoral Science Foundation (grant no. 20110491102).

References

- Koul D, Shen R, Bergh S, Sheng X, Shishodia S, Lafortune TA, Lu Y, de Groot JF, Mills GB and Yung WK: Inhibition of Akt survival pathway by a small-molecule inhibitor in human glioblastoma. *Mol Cancer Ther* 5: 637-644, 2006.
- Harikumar KB and Aggarwal BB: Resveratrol: a multitargeted agent for age-associated chronic diseases. *Cell Cycle* 7: 1020-1035, 2008.
- Delmas D, Lançon A, Colin D, Jannin B and Latruffe N: Resveratrol as a chemopreventive agent: a promising molecule for fighting cancer. *Curr Drug Targets* 7: 423-442, 2006.
- Kundu JK and Surh YJ: Cancer chemopreventive and therapeutic potential of resveratrol: mechanistic perspectives. *Cancer Lett* 269: 243-261, 2008.
- Yang CS, Landau JM, Huang MT and Newmark HL: Inhibition of carcinogenesis by dietary polyphenolic compounds. *Annu Rev Nutr* 21: 381-406, 2001.
- Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, Beecher CW, Fong HH, Farnsworth NR, Kinghorn AD, Mehta RG, *et al*: Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 275: 218-220, 1997.
- Tessitore L, Davit A, Sarotto I and Caderni G: Resveratrol depresses the growth of colorectal aberrant crypt foci by affecting bax and p21(CIP) expression. *Carcinogenesis* 21: 1619-1622, 2000.
- Hebbbar V, Shen G, Hu R, Kim BR, Chen C, Korytko PJ, Crowell JA, Levine BS and Kong AN: Toxicogenomics of resveratrol in rat liver. *Life Sci* 76: 2299-2314, 2005.
- Soleas GJ, Angelini M, Grass L, Diamandis EP and Goldberg DM: Absorption of trans-resveratrol in rats. *Methods Enzymol* 335: 145-154, 2001.
- Singh G and Pai RS: In-vitro/in-vivo characterization of trans-resveratrol-loaded nanoparticulate drug delivery system for oral administration. *J Pharm Pharmacol* 66: 1062-1076, 2014.
- Croy SR and Kwon GS: Polymeric micelles for drug delivery. *Curr Pharm Des* 12: 4669-4684, 2006.
- Escorcia FE, McDevitt MR, Villa CH and Scheinberg DA: Targeted nanomaterials for radiotherapy. *Nanomedicine (Lond)* 2: 805-815, 2007.
- Li F, Zhang X, Li H, Xiang L and Chen Y: Preparation of self-assembled nanoparticles of chitosan oligosaccharide-graft-polycaprolactone as a carrier of bovine serum albumin drug. *Biomed Mater Eng* 24: 2041-2048, 2014.
- Yu Z, Yu M, Zhang Z, Hong G and Xiong Q: Bovine serum albumin nanoparticles as controlled release carrier for local drug delivery to the inner ear. *Nanoscale Res Lett* 9: 343, 2014.
- Guo LY, Yao JP and Sui LH: Preparation and Effects of Resveratrol-Bovine Serum Albumin Nanoparticles on Proliferation of Human Ovarian Carcinoma Cell SKOV3. *Chemical Journal of Chinese Universities* 30: 474-477, 2009.
- Guo LY, Peng Y, Li YL, Yao JP, Wang J, Zhang GM, Chen J and Sui LH: Mechanisms of resveratrol-bovine serum albumin nanoparticle-induced cell death in human ovarian cancer SKOV3 cells. *Nan Fang Yi Ke Da Xue Xue Bao* 30: 2440-2442, 2010 (In Chinese).
- Guo L, Peng Y, Yao J, Sui L, Gu A and Wang J: Anticancer activity and molecular mechanism of resveratrol-bovine serum albumin nanoparticles on subcutaneously implanted human primary ovarian carcinoma cells in nude mice. *Cancer Biother Radiopharm* 25: 471-477, 2010.
- Anastasiadis PZ, Jiang H, Bezin L, Kuhn DM and Levine RA: Tetrahydrobiopterin enhances apoptotic PC12 cell death following withdrawal of trophic support. *J Biol Chem* 276: 9050-9058, 2001.
- Jäättelä M: Programmed cell death: many ways for cells to die decently. *Ann Med* 34: 480-488, 2002.
- Jäättelä M and Tschopp J: Caspase-independent cell death in T lymphocytes. *Nat Immunol* 4: 416-423, 2003.
- Lorenzo HK and Susin SA: Mitochondrial effectors in caspase-independent cell death. *FEBS Lett* 557: 14-20, 2004.
- Blagosklonny MV: Cell death beyond apoptosis. *Leukemia* 14: 1502-1508, 2000.
- Moubarak RS, Yuste VJ, Artus C, Bouharrou A, Greer PA, Menissier-de Murcia J and Susin SA: Sequential activation of poly(ADP-ribose) polymerase 1, calpains, and Bax is essential in apoptosis-inducing factor-mediated programmed necrosis. *Mol Cell Biol* 27: 4844-4862, 2007.
- Zanna C, Ghelli A, Porcelli AM, Martinuzzi A, Carelli V and Rugolo M: Caspase-independent death of Leber's hereditary optic neuropathy cybrids is driven by energetic failure and mediated by AIF and Endonuclease G. *Apoptosis* 10: 997-1007, 2005.
- Miyake K, Bekisz J, Zhao T, Clark CR and Zoon KC: Apoptosis-inducing factor (AIF) is targeted in IFN- α 2a-induced Bid-mediated apoptosis through Bak activation in ovarian cancer cells. *Biochim Biophys Acta* 1823: 1378-1388, 2012.
- Zhan ZL and Chen LY: Changes in the expression of the apoptosis of esophageal cancer EC9706 cells during nuclear matrix proteins induced by curcumin. *Zhong Guo Sheng Wu Hua Xue Yu Fen Zi Sheng Wu Xue Bao* 546-555, 2010.
- Wu B, Gao Q: Microscope laser cytoskeleton changes in the process of cell apoptosis confocal. *Dian Zi Xian Wei Xue Bao* 361-366, 2010.
- Byun HS, Song JK, Kim YR, Piao L, Won M, Park KA, Choi BL, Lee H, Hong JH, Park J, *et al*: Caspase-8 has an essential role in resveratrol-induced apoptosis of rheumatoid fibroblast-like synoviocytes. *Rheumatology (Oxford)* 47: 301-308, 2008.
- Arnoult D, Parone P, Martinou JC, Antonsson B, Estaquier J and Ameisen JC: Mitochondrial release of apoptosis-inducing factor occurs downstream of cytochrome c release in response to several proapoptotic stimuli. *J Cell Biol* 159: 923-929, 2002.
- Cregan SP, Fortin A, MacLaurin JG, Callaghan SM, Cecconi F, Yu SW, Dawson TM, Dawson VL, Park DS, Kroemer G and Slack RS: Apoptosis-inducing factor is involved in the regulation of caspase-independent neuronal cell death. *J Cell Biol* 158: 507-517, 2002.
- Bröker LE, Kruyt FA and Giaccone G: Cell death independent of caspases: a review. *Clin Cancer Res* 11: 3155-3162, 2005.
- Hall JB, Dobrovolskaia MA, Patri AK and McNeil SE: Characterization of nanoparticles for therapeutics. *Nanomedicine (Lond)* 2: 789-803, 2007.
- Dorrie J, Gerauer H, Wachter Y and Zunino SJ: Resveratrol induces extensive apoptosis by depolarizing mitochondrial membranes and activating caspase-9 in acute lymphoblastic leukemia cells. *Cancer Res* 61: 4731-4739, 2001.
- Tinholer I, Bernhard D, Senfter M, Anether G, Loeffler M, Kroemer G, Kofler R, Csordas A and Greil R: Resveratrol, a tumor-suppressive compound from grapes, induces apoptosis via a novel mitochondrial pathway controlled by Bcl-2. *FASEB J* 15: 1613-1615, 2001.
- Howells LM, Moiseeva EP, Neal CP, Foreman BE, Andreadi CK, Sun YY, Hudson EA and Manson MM: Predicting the physiological relevance of in vitro cancer preventive activities of phytochemicals. *Acta Pharmacol Sin* 28: 1274-1304, 2007.
- Athar M, Back JH, Tang X, Kim KH, Kopelovich L, Bickers DR and Kim AL: Resveratrol: a review of preclinical studies for human cancer prevention. *Toxicol Appl Pharmacol* 224: 274-283, 2007.
- Pervaiz S: Chemotherapeutic potential of the chemopreventive phytoalexin resveratrol. *Drug Resist Updat* 7: 333-344, 2004.
- Signorelli P and Ghidoni R: Resveratrol as an anticancer nutrient: molecular basis, open questions and promises. *J Nutr Biochem* 16: 449-466, 2005.
- Le Corre L, Chalabi N, Delort L, Bignon YJ and Bernard-Gallon DJ: Resveratrol and breast cancer chemoprevention: molecular mechanisms. *Mol Nutr Food Res* 49: 462-471, 2005.
- Sheridan C, Delivani P, Cullen SP and Martin SJ: Bax- or Bak-induced mitochondrial fission can be uncoupled from cytochrome C release. *Mol Cell* 31: 570-585, 2008.