

Pretreatment levels of serum osteoprotegerin and p53 protein and urine telomerase as prognostic factors affecting survival in Egyptian bladder cancer patients

RANIA BAKRY¹, MOHAMED I. EL-SAYED², HESHAM M. HAMZA³ and KHALED H. HASSAN⁴

Departments of ¹Clinical Pathology, ²Radiotherapy and ³Surgical Oncology, South Egypt Cancer Institute, Assiut University, Assiut 71111; ⁴Department of Biochemistry, Faculty of Pharmacy (Boys), Al-Azhar University, Cairo 11566, Egypt

Received October 28, 2014; Accepted August 27, 2015

DOI: 10.3892/ol.2015.3925

Abstract. A non-invasive marker is required for the diagnosis and follow-up of patients with bladder cancer. The aim of the current study was to evaluate the potential prognostic significance of serum osteoprotegerin (OPG), p53 protein and urine telomerase in patients with bladder cancer. For all patients, serum levels of OPG and p53 protein were determined using enzyme-linked immunosorbent assay (ELISA), and urine telomerase was assessed using a polymerase chain reaction ELISA technique. Patients were assigned into group 1 (cystectomy and adjuvant radiotherapy) or group 2 (transurethral resection and chemoradiotherapy). The results revealed that serum OPG and p53, and urine telomerase levels were significantly higher in bladder cancer patients compared with in healthy individuals ($P<0.0001$). High serum OPG was associated with significantly lower overall survival and disease-free survival rates (both $P=0.001$), and was correlated with advanced tumor stages ($P<0.0001$), high tumor grades ($P<0.0001$) and the occurrence of disease relapse ($P=0.001$). Serum p53 and urine telomerase did not demonstrate prognostic significance. These findings indicate that serum OPG level may be used as a diagnostic tool and a prognostic variable for patients with muscle invasive bladder cancer. Future trials are required to elucidate its therapeutic role in such patients.

Introduction

Bilharziasis (schistosomiasis) has been endemic in Egypt, with the intensity of infection correlated to the development of bladder cancer (1). Bladder cancer constitutes 30% of all cases of cancer attended at the National Cancer

Institute in Egypt (2), with an overall incidence rate of 13.5/100,000 individuals in Egypt (3). The overall prevalence of *Schistosoma haematobium* (SH) infection in Egypt was 37-48% in the mid-20th century, however, this decreased to 3% by 2002 following the implementation of an antibilharzial campaign (4). In infected patients, the ova deposited in the bladder induces an inflammatory reaction, with the production of oxygen-derived free radicals. Subsequently, this may result in the production of carcinogenic compounds, including N-nitrosamines and polycyclic aromatic hydrocarbons, leading to malignant transformation (5,6). The International Agency for Research on Cancer found that the severity of infection was determined by the urinary egg count and was aggravated by smoking (7). Furthermore, a clear association between bladder cancer and SH infection was identified, with an odds ratio of between 2 and 14 (7). Hematuria, dysuria and necroturia are the main symptoms of infection, which occur as a result of due schistosomal cystitis (8). The median age of patients with schistosoma-associated bladder cancer is lower than patients with schistosoma non-associated bladder cancer. Additionally, schistosomal bladder cancer patients generally present at a later stage of disease, and represent a higher percentage of squamous cell carcinoma (SCC) cases than non-schistosomal bladder cancer patients (9). The majority of tumors associated with bilharziasis are bulky, appear fungating or ulcerative (10), and present at an advanced stage (11), due to the overlap of symptoms of early bilharzial cystitis with those of early malignant cystitis (12). In a report from the Urology and Nephrology center of Mansoura (13), SCC accounted for 59% of 1026 cystectomy specimens in a bilharzial series. This high frequency may be related to squamous metaplasia and dysplasia, which are common in chronic bilharzial cystitis (12).

Bladder cancer possesses diverse biological characteristics resulting from various molecular alterations that lead to uncontrolled cell proliferation and metastasis; such characteristics affect the patient's prognosis (14). Patients with non-invasive bladder cancer must be closely monitored following the termination of treatment. At present, this follow-up is conducted by means of cystoscopy (an invasive procedure) and urine cytology every 3 months during the first two years, every 6 months during third and fourth years, and every 12 months

Correspondence to: Dr Mohamed I. El-Sayed, Department of Radiotherapy, South Egypt Cancer Institute, Assiut University, El-Nasr Towers, Building A2, Flat 907, 7156 Ahmed Hassan Al-Bakory Street, Assiut 71111, Egypt
E-mail: mohebel@yahoo.co.uk

Key words: biomarkers, bladder cancer, overall survival rate, disease-free survival rate

thereafter, unless local recurrence is detected (15). A non-invasive marker is, therefore, required for the follow-up and diagnosis of bladder cancer in order to increase early detection, to decrease costs and to reduce rates of morbidity and mortality (16). Osteoprotegerin (OPG), a cytokine of the tumor necrosis factor (TNF) receptor super-family that is responsible for inhibiting osteoclastogenesis (17,18), may have clinical value as a prognostic marker for patients with bladder carcinoma (19).

Destruction of tumor cells by the immune system is prevented by the development of resistance to TNF-related apoptosis-inducing ligand (TRAIL)-induced toxicity through expression of functional receptors or overexpression of decoy receptors on tumor cell membranes (20). OPG, a soluble decoy receptor for TRAIL, protects tumor cells from TRAIL-mediated apoptosis through binding to TRAIL and inhibiting its activity. As a result, OPG may contribute to tumor progression (19).

The tumor suppressor p53 is crucial in the regulation of cell cycle progression and apoptosis under genotoxic conditions (21). In human cancers, p53 mutations are the most frequent genetic defect (22). Mutations in the p53 protein often result in a prolonged half-life of the protein compared with wild-type p53, leading to its accumulation in cell nuclei, which may then be detected by immunohistochemistry (23).

Telomeres are regions located at the end of human chromosomes that function to stabilize and protect chromosomes (24,25). The ends of the telomeres shorten with successive cell cycles until a critical length is reached, after which cell division results in their breakdown (26). The ribonucleoprotein enzyme telomerase is responsible for maintaining the length of telomeres by adding telomere repeats. This enzyme is inactivated in normal human epithelial tissue; however, it is reactivated in neoplasia (27). The enzymatic activity of telomerase may be measured using a telomeric repeat amplification protocol (TRAP) assay (24).

The aim of the current study was to evaluate the role of pretreatment serum OPG and p53 levels and urine telomerase levels as prognostic factors in patients with muscle-invasive bladder cancer.

Materials and methods

Patients. The current prospective study was conducted in the Departments of Radiotherapy and Clinical Pathology of the South Egypt Cancer Institute, Assiut University (Assiut, Egypt), in the period between June 1st, 2007, and June 1st, 2014. A total of 2,100 patients with non-metastatic muscle-invasive bladder cancer [clinical tumor-node-metastasis (TNM) stage T2-4, N0, M0] were selected. However, due to limited funding, only 65 of these patients and 15 healthy individuals were included. The study protocol was approved by the local institutional review board of South Egypt Cancer Institute (study ID no. IORG0006563; approval no. 190), Assiut University. Written informed consent was obtained from all patients.

For each patient, details of medical history were documented. In addition, all patients were subjected to physical examination, routine laboratory investigations, abdominopelvic computed tomography (CT) scans with contrast, chest X-ray and, if indicated, bone scans. A histopathological diagnosis

Table I. Characteristics of bladder cancer patients (n=65).

Variable	Value
Gender, n (%)	
Female	10 (15.4)
Male	55 (84.6)
Tumor stage, n (%)	
T2	22 (33.9)
T3	35 (53.8)
T4	8 (12.3)
Pathological type, n (%)	
Transitional cell carcinoma	49 (75.4)
Squamous cell carcinoma	16 (24.6)
Bilharzial status and grade, n (%)	
NB-LG	19 (29.2)
B-LG	11 (16.9)
NB-HG	17 (26.2)
B-HG	18 (27.7)
Laboratory tumor markers (mean \pm SD)	
OPG (pg/ml)	211.5 \pm 80.8
p53 (μ g/ml)	27.3 \pm 11.6
Telomerase (μ g protein/assay)	13.9 \pm 6.2
Treatment modality, n (%)	
Total cystectomy and postoperative RT	46 (70.8)
Trimodal therapy (TUR+chemoradiation)	19 (29.2)

NB-LG, non-bilharzial low grade; B-LG, bilharzial low grade; NB-HG, non-bilharzial high grade; B-HG, bilharzial high grade; SD, standard deviation; OPG, osteoprotegerin; RT, radiotherapy; TUR, transurethral resection.

of the bladder mass was determined through cystoscopy and biopsy. Schistosomiasis was diagnosed via cystoscopy based on the presence of schistosoma eggs and peri-oval granulomas in histopathological specimens. According to World Health Organization System for tumor grading (2004) (28) and a previous Malaysian study (29), patients were subdivided into non-bilharzial low-grade (n=19), bilharzial low-grade (n=11), non-bilharzial high-grade (n=17) and bilharzial high-grade (n=18) bladder cancer groups.

Laboratory assays. Samples from all patients and healthy individuals underwent laboratory assays as follows. Serum was isolated from blood samples and subjected to OPG and p53 detection by enzyme-linked immunosorbent assay (ELISA) using a Human Osteoprotegerin (OPG) ELISA kit (TNFRSF 11B; RayBiotech, Inc., Norcross, GA, USA) and p53 ELISA kit (p53 Instant One™ ELISA; eBioscience, Inc., San Diego, CA, USA) respectively. An EVOLIS System (Bio-Rad Laboratories, Inc., Hercules, CA, USA) was used to measure absorbance at 450 nm.

Urine samples obtained from patients and healthy individuals were subjected to telomerase activity measurement using a polymerase chain reaction (PCR)-ELISA technique with a thermal cycler PTC-100 Thermal Cycler

Table II. Laboratory biomarkers in cancer patients and healthy individuals.

Variable	Level, mean \pm standard deviation		P-value
	Healthy individuals (n=15)	Bladder cancer patients (n=65)	
Serum OPG (pg/ml)	62.3 \pm 15.7	214.9 \pm 80.4	<0.0001
Serum p53 (μ g/ml)	11.4 \pm 1.9	27.3 \pm 11.5	<0.0001
Urine telomerase (μ g protein/assay)	3.4 \pm 0.6	13.9 \pm 6.1	<0.0001

OPG, osteoprotegerin.

Table III. Univariate analysis of prognostic factors affecting OAS and DFS rates.

Variable	3-year OAS		3-year DFS	
	Rate, %	P-value	Rate, %	P-value
Age, years		0.280		0.180
<60	66.6		66.6	
\geq 60	52.7		50.9	
Gender		0.240		0.210
Female	78.8		78.8	
Male	56.0		54.4	
Bilharzial status and grade		0.032		0.015
NB-LG	84.2		84.2	
B-LG	68.2		68.2	
NB-HG	52.9		52.9	
B-HG	42.9		28.6	
Tumor stage		<0.001		<0.001
T2	95.5		95.5	
T3	53.8		50.1	
T4	25.0 ^a		25.0 ^a	
Pathological type		0.190		0.260
Transitional cell carcinoma	65.7		61.1	
Squamous cell carcinoma	48.6		48.6	
Treatment modality		0.720		0.890
RC and postoperative RT	60.9		59.3	
TUR and CRT	54.0		54.0	

^a2-year survival rate. OAS, overall survival; DFS, disease-free survival; NB-LG, non-bilharzial low grade; B-LG, bilharzial low grade; NB-HG, non-bilharzial high grade; B-HG, bilharzial high grade; RC, radical cystectomy; RT, radiotherapy; TUR, transurethral resection; CRT, chemoradiotherapy.

(Bio-Rad Laboratories, Inc.) and EVOLIS system (Bio-Rad Laboratories, Inc.). A TRAP kit (TeloTAGGG Telomerase PCR ELISA^{PLUS}; #12013789001) was purchased from Roche Diagnostics GmbH (Mannheim, Germany) and used according to manufacturer's instructions. Telomeric repeats (TTAGGG) were added to the 3' end of a biotin-labelled synthetic P1-TS primer. These elongation products, as well as the internal standard (IS) in the same reaction vessel to avoid false negative results, were amplified by PCR using the primers P1-TS and P2. The Telo TAGGG Telomerase

PCR ELISA^{PLUS} kit contained an IS that produces a 216-bp PCR product, long enough not to interfere with making telomerase ladder visible (30,31). The primer sequences were as follows: P1-TS, anchored complementary telomerase, 5'-GCGCGG[CTAACC]₃-3'; P2, 5'-AATCCGTCGAGCAGAGTTAG[GGTTAG]₄₋₇-3' (where subscript numbers represent maximum telomeric repeats). Sample, negative control (NC) and control template (CT) were added to separate tubes at a volume of 1 μ l. For each, 25 μ l reaction mixture and 5 μ l IS were added. The applied PCR protocol was one cycle for




Figure 1. OAS according to histological grade and bilharzial status. OAS, overall survival; NB-LG, non-bilharzial low grade; B-LG, bilharzial low grade; NB-HG, non-bilharzial high grade; B-HG, bilharzial high grade; FU, follow-up.




Figure 2. DFS according to histologic grade and bilharzial status. DFS, disease-free survival; NB-LG, non-bilharzial low grade; B-LG, bilharzial low grade; NB-HG, non-bilharzial high grade; B-HG, bilharzial high grade; FU, follow-up.




Figure 3. OAS according to disease stage. OAS, overall survival; T, tumor stage; FU, follow-up.




Figure 4. DFS according to disease stage. DFS, disease-free survival; T, tumor stage; FU, follow-up.

Table IV. Cox regression stepwise OAS and DFS analyses according to laboratory markers.

Variable	OAS rate	DFS rate
OPG	P=0.001; HR, 1.01; 95% CI, 1-1.01	P=0.001; HR, 1.01; 95% CI, 1-1.01
p53	P=0.97	P=0.82
Telomerase	P=0.61	P=0.72

OAS, overall survival; DFS, disease free survival; OPG, osteoprotegerin; HR, hazard ratio; CI, confidence interval.

Table V. Disease relapse.

Disease relapse	n	%
Local recurrence	4	6.2
Distant metastases	7	10.8
Mixed relapse	1	1.5
Total	12	18.5

primer elongation (10-30 min at 25°C) followed by one cycle of telomerase inactivation (5 min at 94°C). The subsequent amplification scheme was 30 cycles of denaturation for 30 sec at 94°C, annealing for 30 sec at 50°C, and polymerization for 90 sec at 72°C. Finally, an additional extension was performed for 10 min at 72°C. The denatured and hybridized products were added to streptavidin-coated plates (Roche Diagnostics GmbH). The immobilized amplicons were detected using horseradish peroxidase (HRP)-conjugated anti-digoxigenin (DIG) antibody with DIG-labelled detection probes from the Telo TAGGG Telomerase PCR ELISA^{PLUS} kit (P3 T and P3 IS for the sample and internal standard, respectively; Roche Diagnostics GmbH) for 1 h at 37°C, which were complementary to the telomeric repeat sequence. Sheep polyclonal antibody conjugated to HRP was used for 1 h at 37°C to prepare conjugate (working) solution (dilution, 1:20). Following preparation, anti-DIG-HRP conjugate working solution (10 mU/ml) was added for 30 min at room temperature, followed by washing five times with washing buffer (250 µl). Tetramethylbenzidine substrate was added then 100 µl stopping solution (Roche Diagnostics GmbH) was added. Absorbance was measured at 450 nm and relative telomerase activity (RTA) was determined using the following formula:

$$RTA = \frac{(A_s - A_{s,0}) / A_{s,IS}}{(A_{TS8} - A_{TS8,0}) / A_{TS8,IS}}$$

where A_s was the absorbance of the sample, $A_{s,0}$ was the absorbance of the RNase-treated sample, $A_{s,IS}$ was the absorbance of the IS of the sample, A_{TS8} was the absorbance of the control template (TS8), $A_{TS8,0}$ was the absorbance of the lysis buffer, and $A_{TS8,IS}$ was the absorbance of the IS of the control template.

Table VI. Association between serum OPG level and different prognostic variables.

Variable	OPG, pg/ml; mean \pm SD	P-value
Bilharzial status and grade		<0.0001
NB-LG	128.4 \pm 25.3	
B-LG	163.2 \pm 17.6	
NB-HG	244.1 \pm 20.9	
B-HG	310.3 \pm 49.9	
Tumor stage		<0.0001
T2	157.1 \pm 48.6	
T3	231.4 \pm 77.1	
T4	301.8 \pm 50.9	
Histopathological type		0.37
Transitional cell carcinoma	220.1 \pm 83.1	
Squamous cell carcinoma	199.1 \pm 71.5	
Age, years		0.092
<60	196.2 \pm 84.8	
\geq 60	230.0 \pm 74.4	
Gender		0.72
Female	206.5 \pm 67.7	
Male	216.5 \pm 82.9	
Disease relapse		0.001
Relapse	285.0 \pm 60.8	
No relapse	199.1 \pm 76.1	

OPG, osteoprotegerin; SD, standard deviation; NB-LG, non-bilharzial low-grade; B-LG, bilharzial low-grade; NB-HG, non-bilharzial high-grade; B-HG, bilharzial high-grade.

Radiotherapy. Patients were non-randomly assigned into one of two groups. Group 1 included patients who underwent radical cystectomy (anterior pelvic exenteration in female patients) and postoperative radiotherapy using 2-dimensional planning and a 3-field technique, with a radiation dose of 50 Gy/25 fractions over 5 weeks using 15 MV photons. Group 2 included patients who underwent trimodal therapy [complete transurethral resection (TUR), and concurrent chemo-radiotherapy]. Radiotherapy was initiated 4-6 weeks after TUR using 3-dimensional planning in two phases. In phase I, the clinical target volume (CTV) included the urinary bladder, proximal urethra (and prostate in male patients) and regional lymph nodes (hypogastric, external iliac and obturator nodes). The planning target volume (PTV), including the CTV plus a 1 cm margin, was treated with total dose of 46 Gy in 23 fractions, with 5 fractions administered per week. In phase II, the PTV included the whole bladder plus 2 cm margins, and received an additional total dose of 20 Gy in 10 fractions with 15 MV photons.

Chemotherapy. Gemcitabine was administered at 30 mg/m² by a 30-min intravenous infusion prior to radiation therapy sessions twice weekly. Follow-up examinations were performed routinely every 3 months following treatment. Abdominopelvic CT scans, chest radiography and, for group 2 patients, cystoscopy were performed annually.

Statistical methods. The study cut-off point was June 1st, 2014. Overall survival (OAS) and disease-free survival (DFS) rates were estimated by the Kaplan-Meier method using the GraphPad Prism program (version 5; GraphPad Software, Inc., La Jolla, CA, USA). The log-rank test was used to examine differences in OAS and DFS rates. Comparison of mean values of OPG in different groups was conducted using a one-way analysis of variance. $P < 0.05$ was considered to indicate statistical significance.

Results

Study subjects. A total of 65 patients were recruited into the study. Patient characteristics are listed in Table I. The median age of the patients was 60 years. The majority of patients were male [55 patients; 84.6%; male to female ratio, 5.5:1] and presented with T3 disease stage (35 patients; 54%), transitional cell carcinoma pathological type (49 patients; 75.5%), and high-grade disease (35 patients; 54%). A positive history of bilharziasis was documented in 29 patients (44.5%).

Radical cystectomy followed by adjuvant radiotherapy was conducted in the vast majority of patients (71%), while trimodal therapy was conducted in the remaining 29%. The median follow-up time from the date of enrollment was 30 months, and ranged from 3 to 60 months.

Biomarker levels. Laboratory biomarkers were assessed in cancer patients and healthy individuals (Table II). The mean

values of serum OPG, serum p53 and urine telomerase were significantly higher in bladder cancer patients than in healthy individuals ($P<0.0001$).

Survival rates. After a median follow-up period of 30 months, the 3-year OAS and DFS rates were 59 and 58%, respectively. There were no significant differences between OAS and DFS rates according to age, gender, pathological type, treatment modality, serum p53 level or urine telomerase level (Tables III and IV). However, significant differences in 3-year OAS and DFS rates were associated with T stage (both $P<0.0001$), histological grade and bilharzial status (OAS, $P=0.032$; DFS, 0.015), and serum OPG level (both $P=0.001$).

As indicated in Table V, only 4 patients (6%) developed local recurrence, 7 patients (11%) developed distant metastases (4 in the bones, 2 in the lungs and 1 in the bones and liver) and 1 patient (1.5%) developed mixed local and distant (bones and lungs) metastases.

Association between serum OPG level and prognostic variables. As shown in Table VI, high serum OPG levels were associated with advanced T stage ($P<0.0001$), high histological grade ($P<0.0001$) and occurrence of disease relapse ($P=0.001$). Figs. 1-4 show OAS and DFS rates according to histological grade and disease stage.

Discussion

Identification of biomarkers for bladder cancer may improve the screening and diagnosis for this disease, and aid in determining its prognosis. Ideal biomarkers would be those that may be detected non-invasively and rapidly, that are easy to obtain, use and interpret, are inexpensive and have high accuracy (32,33). A large number of molecular markers that have potential prognostic value have been determined in previous studies of molecular biology and genetics (19,23,34-38), and include serum OPG, serum p53 and urine telomerase.

In the present study, levels of serum OPG and p53 protein, and urine telomerase were identified to be significantly higher ($P<0.0001$) in patients with bladder carcinoma than in healthy individuals. This is in agreement with the findings of a previous study by Mizutani *et al* (19), in which it was reported that the mean serum OPG concentration in patients with bladder carcinoma was ~3 times greater than the mean concentration in healthy volunteers. Shimada *et al* (39) reported that surveillance of serum p53 antibodies may be useful in detecting various types of malignant tumors, including bladder cancer. Telomerase is activated in cancer cells, but not in normal somatic cells; therefore, its detection may be used as a diagnostic marker for cancer (40). Patients with urothelial tumors of all grades exhibit telomerase activity in voided urine (41).

With a median follow-up of 30 months, the 3-year OAS and DFS rates were 59 and 58%, respectively, in the present study. Survival rates in the current study are comparable to those reported in previous studies (42,43). The postoperative radiotherapy (in cystectomy group) and the trimodal therapy (i.e. TUR, and chemoradiation, in bladder preservation group) resulted in a relatively low relapse rate in the present study (18.5%). Zaghloul *et al* (44) demonstrated that adjuvant radiation therapy improve local control rate, and that the therapeutic

benefit of postoperative irradiation was consistent for all tumor types, histological grades and pathological stages in terms of DFS and local control.

Among the investigated variables in the current study, only T stage, bilharzial status and histological grade, and serum OPG biomarker level significantly influenced the rates of OAS ($P<0.0001$, $P=0.032$ and $P=0.001$, respectively) and DFS ($P<0.0001$, $P=0.015$ and $P=0.001$, respectively). Numerous studies have previously analyzed prognostic factors in patients with bladder cancer treated by radical cystectomy, and pathological stage is considered to be the most important prognostic factor in such patients (45-51).

In contrast to OPG in the present study, serum p53 and urine telomerase were not significantly associated with survival rates. Regarding p53 status as a prognostic factor in muscle-invasive bladder cancer, results are contradictory (23). Schmitz-Dräger *et al* (52) conducted a meta-analysis of all published research on the association between p53 positivity and the prognosis of patients with bladder cancer. In the majority of trials (5 of 7) of muscle-invasive bladder cancer, p53 was not regarded as an independent prognostic marker of disease progression. The findings of the present study are in agreement with the results of this meta-analysis. It was also stated that, although urinary telomerase could be a potentially useful urinary tumor marker, its use for diagnosis in symptomatic patients or its impact during surveillance is still unknown (40). Furthermore, normalization and standardization of the assays used to evaluate this is required before they may become valuable in clinical practice (40).

In previous studies investigating the use of serum OPG level as a prognostic marker, elevated levels were reported to be associated with poor prognosis (19,53-55). The 5-year disease-specific survival rate of patients with muscle-invasive bladder carcinoma was determined to be greater for those with low serum OPG levels compared with those exhibiting high serum OPG levels. Such findings indicate that OPG concentration may be a significant prognostic factor in patients with this disease, with low serum OPG levels indicating a favorable prognosis.

In the current study, serum OPG levels in patients with bladder carcinoma were analyzed according to patient's age and gender, disease stage, tumor grade, pathological type and occurrence of disease relapse. High serum OPG was identified to be associated with advanced T stage, high histological grade ($P<0.0001$) and the presence of relapse ($P=0.001$). This is in agreement with studies by Mizutani *et al* (19) and Holen and Shipman (56), who reported high serum OPG was associated with advanced disease stage, high grade and low 5-year disease-specific survival rate. One limitation of the present study is that the analysis was restricted as patient selection for treatment in the two treatment groups was not randomized.

Despite the advent of effective chemotherapy and modern radiation therapy techniques, treatment outcome of bladder cancer is unsatisfactory as local recurrence and distant metastases are major problems. Therefore, treatment protocols inhibiting OPG production may increase TRAIL-mediated toxicity of malignant cells and thus may improve treatment results (19). This necessitates future trials in larger patient populations with longer follow-up, to clarify the prognostic role of OPG in cancers of different stages, pathological types and grades.

References

- Mostafa MH, Sheweita SA and O'Connor PJ: Relationship between schistosomiasis and bladder cancer. *Clin Microbiol Rev* 12: 97-111, 1999.
- Sherif M and Ibrahim AS: The incidence of cancer in Egypt. In: *The Profile of Cancer in Egypt*. Arab World Printing House, Cairo, pp179-190, 1987.
- Ibrahim AS, Khaled HM, Mikhail NH, Baraka H and Kamel H: Cancer incidence in Egypt: Results of the national population-based cancer registry program. *J Cancer Epidemiol* 2014: 1-18, 2014.
- Prevalence of schistosomiasis in Egypt over time. Department of Endemic Diseases, Ministry of Health and Population, Egypt, 2004.
- Rosin MP, Saad el Din Zaki S, Ward AJ and Anwar WA: Involvement of inflammatory reactions and elevated cell proliferation in the development of bladder cancer in schistosomiasis patients. *Mutat Res* 305: 283-292, 1994.
- Marletta MA: Mammalian synthesis of nitrite, nitrate, nitric oxide, and N-nitrosating agents. *Chem Res Toxicol* 1: 249-257, 1988.
- International Agency for Research on Cancer: IARC Monographs on the Evaluation of Carcinogenic Risks to Humans (Vol 61): Schistosomes, Liver flukes and *Helicobacter pylori*. IARC Press, Lyon, 1994.
- Zaghloul MS: Bladder cancer and schistosomiasis. *J Egypt Natl Canc Inst* 24: 151-159, 2012.
- Zaghloul MS: Radiation as adjunctive therapy to cystectomy for bladder cancer: Is there a difference for bilharzial association? *Int J Radiat Oncol Biol Phys* 28: 783, 1994.
- Khalaf I, El-Mallah E, Elstouhi I, Abu-Zeid H and Elmeligy A: Pathologic pattern of invasive bladder carcinoma: Impact of bilharziasis. *Afr J Urol* 14: 90-97, 2010.
- el-Bolkainy MN, Ghoneim MA and Mansour MA: Carcinoma of the bilharzial bladder in Egypt. Clinical and pathological features. *Br J Urol* 44: 561-570, 1972.
- Abol-Enein H, Kava BR and Carmack AJ: Nonurothelial cancer of the bladder. *Urology* 69: 93-104, 2007.
- Ghoneim MA, el-Mekresh MM, el-Baz MA, el-Attar IA and Ashamallah A: Radical cystectomy for carcinoma of the bladder: Critical evaluation of the results in 1,026 cases. *J Urol* 158: 393-399, 1997.
- Ha YS, Kim JS, Yoon HY, Jeong P, Kim TH, Yun SJ, Lee SC, Kim GY, Choi YH, Moon SK, *et al*: Novel combination markers for predicting progression or nonmuscle invasive bladder cancer. *Int J Cancer* 131: E501-E507, 2012.
- Parmar MK, Freedman LS, Hargreave TB and Tolley DA: Prognostic factors for recurrence and followup policies in the treatment of superficial bladder cancer. Report from the british medical research council subgroups on superficial bladder cancer (Urological Cancer Working Party). *J Urol* 142: 284-288, 1989.
- Wadhwa N, Jatava SK and Tiwari A: Non-invasive urine based tests for the detection of bladder cancer. *J Clin Pathol* 65: 970-975, 2012.
- Golledge J, McCann M, Mangan S, Lam A and Karan M: Osteoprotegerin and osteopontin are expressed at high concentrations within symptomatic carotid atherosclerosis. *Stroke* 35: 1636-1641, 2004.
- Hellenthal FA, Buurman WA, Wodzig WK and Schurink GW: Biomarkers of abdominal aortic aneurysm progression. Part 2: Inflammation. *Nat Rev Cardiol* 6: 543-552, 2009.
- Mizutani Y, Mutsuura H, Yamamoto K, Nan Li Y, Mikami K, Okihara K, Kawauchi A, Bonavida B and Miki T: Prognostic significance of serum osteoprotegerin levels in patients with bladder carcinoma. *Cancer* 101: 1794-1802, 2004.
- Makimovic-Ivanic D, Stolic-Grubic S, Nicoletti F and Mijatovic S: Resistance to TRAIL and how to surmount it. *Immunol Res* 52: 157-168, 2012.
- Spike BT and Wahl GM: p53, stem cells, and reprogramming: Tumor suppression beyond guarding the genome. *Genes Cancer* 2: 404-419, 2011.
- Hollstein M, Sidransky D, Vogelstein B and Harris CC: p53 mutations in human cancers. *Science* 253: 49-53, 1991.
- Habuchi T, Marberger M, Droller MJ, Hemstreet GP III, Grossman HB, Schalken JA, Schmitz-Dräger BJ, Murphy WM, Bono AV, Goebell P, *et al*: Prognostic markers for bladder cancer: International consensus panel on bladder tumor markers. *Urology* 66 (6 Suppl 1): 64-74, 2005.
- Kim NW, Piatyszek MA, Prowse KR, *et al*: Specific association of human telomerase activity with immortal cells and cancer. *Science* 266: 2011-2015, 1994.
- Yoshida K, Sugino T, Tahara H, Woodman A, Bolodeoku J, Nargund V, Fellows G, Goodison S and Tahara E: Telomerase activity in bladder carcinoma and its implication for non-invasive diagnosis by detection of exfoliated cancer cells in urine. *Cancer* 79: 362-369, 1997.
- Hodes RJ: Telomere length, aging, and somatic cell turnover. *J Exp Med* 190: 153-156, 1999.
- Mu J and Wei LX: Telomere and telomerase in oncology. *Cell Res* 12:1-7, 2002.
- Eble JN, Sauter G, Epstein JI, and Sesterhenn IA: World Health Organization classification of tumors: Pathology and genetics of tumors of the urinary system and male genital organs. In collaboration with the International Academy of Pathology (IAP): 139-390, 2004.
- Abdulmir AS, Hafidh RR, Kadhim HS, Abubakar F: Tumor markers of bladder cancer: The schistosomal bladder tumors versus non-schistosomal bladder tumors. *J Exp Clin Cancer Res* 28: 27, 2009.
- Wright WE, Shay JW and Piatyszek MA: Modifications of a telomeric repeat amplification protocol (TRAP) result in increased reliability linearity and sensitivity. *Nucl Acids Res* 23: 3794-3795, 1995.
- Kim NW and Wu F: Advances in quantification and characterization of telomerase activity by the telomeric repeat amplification (TRAP). *Nucl Acids Res* 25: 2595-2597, 1997.
- Burchardt M, Burchardt T, Shabsigh A, De La Taille A, Benson MC and Sawczuk I: Current concepts in biomarker technology for bladder cancers. *Clin Chem* 46: 595-605, 2000.
- Selmy MA, Ibrahim GH, El Serafi TI and Ghobeish AA: Evaluation of urinary proepithelin as potential biomarker for bladder cancer detection and prognosis in Egyptian patients. *Cancer biomarker* 7: 163-170, 2010.
- El-Kenawy A, El-Kott AF and Khalil AM: Prognostic value of p53 and MDM2 expression in bilharziasis-associated squamous cell carcinoma of the urinary bladder. *Int J Biol Markers* 18: 284-289, 2003.
- Hemal AK, Khaitan A, Dinda AK, Gupta NP, Seth A, Dogra PN and Nabi G.: Prognostic significance of p53 nuclear overexpression in patients of muscle invasive urinary bladder carcinoma treated with cystectomy. *Urol Int* 70: 42-46, 2003.
- Tsai YS, Tzai TS, Chow NH, Yang WH, Tong YC, Lin JS, Chang CC, Cheng HL and Lin YM: Prognostic values of p53 and HER-2/neu coexpression in invasive bladder cancer in Taiwan. *Urol Int* 71: 262-270, 2003.
- Kelsey KT, Hirao T, Schned A, Hirao S, Devi-Ashok T, Nelson HH, Andrew A and Karagas MR: A population-based study of immunohistochemical detection of p53 alteration in bladder cancer. *Br J Cancer* 90: 1572-1576, 2004.
- Shariat SF, Casella R, Khoddami SM, *et al*: Urine detection of survivin is a sensitive marker for the noninvasive diagnosis of bladder cancer. *J Urol* 171: 626-630, 2004.
- Shimada H, Ochiai T and Nomura F: Japan p53 Antibody Research Group: Titration of serum p53 antibodies in 1,085 patients with various types of malignant tumors. A multiinstitutional analysis by the Japan p53 Antibody Research Group. *Cancer* 97: 682-689, 2003.
- Lamarca A and Barriuso J: Urine telomerase for diagnosis and surveillance of bladder cancer. *Adv Urol* 2012: 693631, 2012.
- Kavaler E, Landman J, Chang Y, Droller MJ and Liu BC: Detecting human bladder carcinoma cells in voided urine samples by assaying for the presence of telomerase activity. *Cancer* 82: 708-714, 1998.
- Lee CT, Madii R, Daignault S, Dunn RL, Zhang Y, Montie JE and Wood DP Jr: Cystectomy delay more than 3 months from initial bladder cancer diagnosis results in decreased disease specific and overall survival. *J Urol* 175: 1262-1267, 2006.
- Sonpavde G, Khan MM, Lerner SP, Svatek RS, Novara G, Karakiewicz PI, Skinner E, Tilki D, Kassouf W, Fradet Y, *et al*: Disease-free survival at 2 or 3 years correlates with 5-year overall survival of patients undergoing radical cystectomy for muscle invasive bladder cancer. *J Urol* 185: 456-461, 2011.
- Zaghloul MS, Awwad HK, Akoush HH, Omar S, Soliman O and el Attar I: Postoperative radiotherapy of carcinoma in bilharzial bladder: Improved disease free survival through improving local control. *Int J Radiat Oncol Biol Phys* 23: 511-517, 1992.
- Bassi P, Ferrante GD, Piazza N, Spinadin R, Carando R, Pappagallo G and Pagano F: Prognostic factors of outcome after radical cystectomy for bladder cancer: A retrospective study of a homogeneous patient cohort. *J Urol* 161: 1494-1497, 1999.

46. Harada K, Sakai I, Hara I, Eto H and Miyake H: Prognostic significance of vascular invasion in patients with bladder cancer who underwent radical cystectomy. *Int J Urol* 12: 250-255, 2005.
47. Antunes AA, Nesrallah LJ, Dall'Oglio MF, Ferreira YA, Passerotti CC, Leite KR, Ortiz V and Srougi M: Analysis of prognostic factors in patients with transitional cell carcinoma of the bladder treated with radical cystectomy. *Int Braz J Urol* 32: 35-42, 2006.
48. Madersbacher S, Hochreiter W, Burkhard F, Thalmann GN, Danuser H, Markwalder R and Studer UE: Radical cystectomy for bladder cancer today-a homogeneous series without neoadjuvant therapy. *J Clin Oncol* 21: 690-696, 2003.
49. Skołyszewski J, Reinfuss M and Weiss M: Radical external beam radiotherapy of urinary bladder carcinoma. An analysis of results in 500 patients. *Acta Oncol* 33: 561-565, 1994.
50. Türkölmez K, Tokgöz H, Reşorlu B, Köse K and Bedük Y: Muscle-invasive bladder cancer: Predictive factors and prognostic difference between primary and progressive tumors. *Urology* 70: 477-481, 2007.
51. Takashi M, Sakata T, Murase T, Hamajima N and Miyake K: Grade 3 bladder cancer with lamina propria invasion (pT1): Characteristics of tumor and clinical course. *Nagoya J Med Sci* 53: 1-8, 1991.
52. Schmitz-Dräger BJ, Goebell PJ, Ebert T and Fradet Y: p53 immunohistochemistry as a prognostic marker in bladder cancer. Playground for urology scientists? *Eur Urol* 38: 691-699, 2000.
53. Ito R, Nakayama H, Yoshida K, Kuraoka K, Motoshita J, Oda N, Oue N and Yasui W: Expression of osteoprotegerin correlates with aggressiveness and poor prognosis of gastric carcinoma. *Virchows Arch* 443: 146-151, 2003.
54. Brown JM, Vessella RL, Kostenuik PJ, Dunstan CR, Lange PH and Corey E: Serum osteoprotegerin levels are increased in patients with advanced prostate cancer. *Clin Cancer Res* 7: 2977-2983, 2001.
55. Eaton CL, Wells JM, Holen I, Croucher PI and Hamdy FC: Serum osteoprotegerin (OPG) levels are associated with disease progression and response to androgen ablation in patients with prostate cancer. *Prostate* 59: 304-310, 2004.
56. Holen I and Shipman CM: Role of osteoprotegerin (OPG) in cancer. *Clin Sci (Lond)* 110: 279-291, 2006.