

Construction of recombinant pEGFP-N1-hPer2 plasmid and its expression in osteosarcoma cells

ANYUAN CHENG¹, YAN ZHANG², HONGJUN MEI³, SHUO FANG¹,
PENG JI¹, JIAN YANG¹, LING YU¹ and WEICHUN GUO¹

¹Orthopedic Department, Renmin Hospital of Wuhan University, Wuhan, Hubei 430060;

²Department of Cardiology, The General Hospital of Fengfeng Group Co., Ltd., Handan, Hebei 056200;

³Orthopedic Department, Wuhan No. 5 Hospital, Wuhan, Hubei 430050, P.R. China

Received November 27, 2014; Accepted November 10, 2015

DOI: 10.3892/ol.2016.4291

Abstract. The aim of this study was to construct the eukaryotic expression vector pEGFP-N1-hPer2 and assess its expression in the human osteosarcoma cell line MG63. Total mRNA was extracted from human osteosarcoma MG63 cells, the human period 2 (hPer2) gene was obtained by reverse transcription-polymerase chain reaction (RT-PCR) and cloned into the pEGFP-N1 vector, then the recombinant pEGFP-N1-hPer2 plasmid was constructed and transfected into MG63 cells using Lipofectamine 2000. The expression of hPer2 in MG63 cells was measured by quantitative RT-PCR and western blot analysis. The accurate construction of pEGFP-N1-hPer2 was verified by double enzyme digestion and DNA sequencing. hPer2 gene expression in the transfected cells was assessed by RT-qPCR and western blot analysis. In conclusion, the recombinant pEGFP-N1-hPer2 plasmid was constructed successfully, and expressed effectively in MG63 cells.

Introduction

The daily light/dark cycles of the Earth are responsible for the physiological and behavioral activity of a number of organisms. This temporal activity is known as circadian rhythm, and has a biological molecular basis, namely the circadian gene (1,2). Previous studies have demonstrated that the circadian genes regulate several molecular and biochemical processes as well as having an established role in the mammalian circadian clock. Research suggests that the role of the circadian clock may be a fundamental regulator for tumor suppression in humans (3,4). Several cancer studies indicate that period 2 (Per2), one of the key circadian genes, plays a significant role in growth control and tumor development (5). To understand

the effects of Per2 on osteosarcoma cell growth *in vitro*, the recombinant pEGFP-N1-hPer2 plasmid was constructed using the vector pEGFP-N1 carrying the fluorescent protein gene and transfected into MG63 cells using Lipofectamine 2000, then the expression of hPer2 in MG63 cells was assessed by reverse transcription-polymerase chain reaction and western blot analysis. This research is expected to lay the foundations for research into the circadian gene in osteosarcoma.

Materials and methods

Materials. The MG63 osteosarcoma cell line was purchased from the Cell Bank of Wuhan University (Wuhan, China); the pMD19-T vector, *Escherichia coli* DH5a and DNA marker were purchased from Transgen Biotechnology (Beijing, China); *Pst*I and *Kpn*I restriction enzymes were purchased from NEB (Ipswich, MA, USA); the plasmid pEGFP-N1 was purchased from Clontech Biotechnology (Mountain View, CA, USA); the DNA ligation kit, RevertAid™ First Strand cDNA synthesis kit, dNTPs, RevertAid reverse transcriptase and HiFi DNA polymerase were purchased from Fermentas (Beijing, China); TRIzol™ reagent, Lipofectamine 2000 and radioimmunoprecipitation assay (RIPA) lysis buffer were purchased from Beyotime Institute of Biotechnology (Shanghai, China); AxyPrep DNA gel extraction kit was purchased from Takara Biotechnology, Inc. (Dalian, China); rabbit anti-hPer2 polyclonal antibody was purchased from ProteinTech Group, Inc. (Chicago, IL, USA; catalog no., 20359-1-AP); and Western Lighting Plus Chemiluminescence was purchased from PerkinElmer, Inc. (Waltham, MA, USA).

Methods

Gene amplification, cloning and sequencing. Total RNA was extracted from the osteosarcoma cell line MG63 using TRIzol reagent according to the procedure supplied by the manufacturer. Extracted RNA (1 µg) was used for cDNA synthesis with the RevertAid First Strand cDNA synthesis kit according to the manufacturer's instructions. The reaction system was prepared in a total volume of 20 µl containing 12.5 µl RNA primer mix, 4 µl 5X RT reaction buffer, 2 µl dNTPs, 1 µl RevertAid reverse transcriptase, 0.5 µl RiboLock RNase inhibitor and dH₂O up to 20 µl. A pair of primers was designed based on the hPer2

Correspondence to: Professor Weichun Guo, Orthopedic Department, Renmin Hospital of Wuhan University, 99 Jiefang Road, Wuchang, Wuhan, Hubei 430060, P.R. China
E-mail: whucay@whu.edu.cn

Key words: biological rhythm, circadian gene period 2, osteosarcoma, MG63, eukaryotic expression vector pEGFP-N1

mRNA sequence (Genbank no. NM_022817.2): *Pst*I tailed forward (5'-AACTGCAGATGAATGGATACGCGGAATTCC-3') and *Kpn*I tailed reverse (5'-CGGCTGCAGCGTCTGCTCTTCGATCCTGT-3') primers (restriction sites are underlined) and named as hPer2-F and hPer2-R, respectively. The length of the amplification segment was 3765 bp. The PCR mixture was blended in a total volume of 20 μ l containing 2 μ l cDNA template, 1 μ l each primer (10 μ mol/l), 10 μ l PCR mix and dH₂O up to 20 μ l. The PCR program was started at 94°C for 5 min, followed by 40 cycles at 94°C for 30 sec, 58°C for 30 sec, 72°C for 5 min and completed with a final extension at 72°C for 5 min. The final PCR products were separated by electrophoresis using 1% polyacrylamide gels, and the target fragment was purified and recovered using an agarose gel extraction kit (Watson Biomedical, Inc., Shanghai, China).

The purified target fragments were ligated into the plasmid pMD19-T and then transformed into competent *E. coli* DH5a cells. Recombinant plasmid was extracted from bacterial colonies and 1.0 μ l plasmid solution was subjected to agarose gel electrophoresis to confirm the presence of the correct sequence of hPer2.

Construction of pEGFP-N1-hPer2 expression plasmid. Double restriction enzyme digestion was applied to the recovered target gene fragment and eukaryotic expression vector pEGFP-N1, respectively. The enzyme reaction contained 2 μ l target gene fragment (or vector pEGFP-N1), 2 μ l 10X Fast Digest buffer, 1 μ l *Pst*I, 1 μ l *Kpn*I and dH₂O up to 20 μ l. Under the guidance of the T4 DNA ligase system instructions, the purified target fragment of the hPer2 gene was directionally ligated into pEGFP-N1 vector in a 10- μ l reaction system containing 3 μ l target fragment, 1 μ l pEGFP-N1, 1 μ l T4 DNA ligase, 1 μ l 2X Quick Ligation buffer and dH₂O up to 10 μ l. The reactants were well mixed at 16°C for 2 h, then the ligation was transformed into competent *E. coli* DH5a cells and inoculated into Luria-Bertani culture media containing 100 μ g/ml ampicillin at a volume ratio of 1:100. After amplification by shaking the culture overnight at 37°C, the target plasmids were extracted from the bacterial liquid according to the instructions for the EndoFree Maxi Plasmid kit (Tiangen Biotech Co., Ltd., Beijing, China). The resulting recombinant eukaryotic expression vector was named pEGFP-N1-hPer2, and the construction procedure is shown in Fig. 1.

The pEGFP-N1-hPer2 was digested using *Pst*I and *Kpn*I, and then evaluated by agarose gel electrophoresis. The recombinant plasmid was further sequenced to confirm its sequence by the Beijing Genomics Institute (BGI; Beijing, China).

Transfection of pEGFP-N1-hPer2 into MG63 cells. MG63 cell lines originated from human osteosarcoma were used in this study. The cells were maintained and cultured in RPMI-1640 media supplemented with 10% fetal bovine serum, 100 U/ml penicillin and 100 μ g/ml streptomycin (Biochrom KG, Berlin, Germany) at 37°C in a humidified 5% CO₂ incubator. The cells were divided into three groups: pEGFP-hPer2, pEGFP-N1 and control. Cells were transiently transfected with the DNA construct using Lipofectamine 2000 reagent. In brief, following the manufacturer's instructions the transfection complex was prepared based on the optimized amounts of plasmid and Lipofectamine reagent and transferred to the




Figure 1. Schematic representation of the construction of pEGFP-N1-hPer2 plasmid.

70-80% confluent MG63 cells, then cells were washed with phosphate-buffered saline (PBS) and collected to conduct subsequent assays.

Analysis of transfection efficiency. The expression of enhanced green fluorescence protein (EGFP) was already visualized by fluorescence microscopy (Olympus IX51; Olympus Corporation, Seoul, South Korea) at 24 h post-transfection, and the percentage of fluorescence-emitting cells was determined by flow cytometry (FACSsort; BD Biosciences, Franklin Lakes, NJ, USA).

Quantitative RT-PCR (RT-qPCR) analysis. Total RNA was isolated from MG63 cells using TRIzol reagent, and the concentration of each RNA sample was determined using a NanoVue Plus spectrophotometer (GE Healthcare Bio-Sciences AB, Uppsala, Sweden). All RNA samples were subsequently adjusted to the same concentration. An SYBR PrimeScript RT-PCR kit (Takara Biotechnology, Inc.) was then used for RT-PCR according to the manufacturer's instructions. The relative mRNA expression of hPer2 was analyzed by qPCR using the IQTM 5 System (Bio-Rad Laboratories, Inc.,




Figure 2. Sequencing results of pEGFP-N1-hPer2 plasmid (partial).




Figure 3. hPer2 cDNA amplified by reverse transcription-polymerase chain reaction. M, DL8000 marker; 1, hPer2 PCR products (3765 bp).




Figure 4. Identification of recombinant pEGFP-N1-hPer2 plasmid double and single enzyme digestion. M, DL8000 marker; 1, pEGFP-N1-hPer2 plasmid double digested with *Pst*I and *Kpn*I enzyme; 2, pEGFP-N1-hPer2 plasmid single digested with *Pst*I enzyme; 3, pEGFP-N1 plasmid.

Hercules, CA, USA) with β -actin (Genbank no. NM_001101) serving as the reference gene. The primer information is as follows: hPER2-F, TACACCGTGGAGGAGATGGAGA; hPER2-R, ATATGGATGCAACCTGGTCAGA; β -actin-F, GTCCACCGCAAATGCTTCTA; β -actin-R, TGCTGT CACCTTCACCGTTC. The PCR reactions were carried out in a 96-well plate in a 25 μ l reaction volume. Each reaction mixture contained 12.5 μ l SYBR-Green I PCR Master mix (Takara Biotechnology, Inc.), 2.5 μ l normalized template DNA, 0.5 μ l each primer and 9.5 μ l sterile ultrapure water. The relative expression of hPer2 was calculated using the 'normalized relative quantification' method followed by the $2^{-\Delta\Delta C_t}$ cycle threshold method (6). PCR reactions were performed in triplicate for each sample.

Western blot analysis. For western blot analysis, cells at 90% confluency were washed in PBS before incubation with RIPA lysis buffer consisting of 50 mM Tris, pH 7.4, 150 mM NaCl, 1% Triton X-100, 1% sodium deoxycholate, 0.1% sodium dodecyl sulphate (SDS) on ice for 10 min. The cell lysates were clarified by centrifugation at 9000 x g for 10 min, and the supernatants were collected. Protein concentration was measured by bicinchoninic assay (Aidlab, Beijing, China). Equal amounts of total protein were separated on 10% SDS-polyacrylamide gel, and then transferred electrophoretically to nitrocellulose membranes blocked with TBST buffer (50 mM Tris-HCl, pH 7.5, 0.15 M NaCl, 0.1% Tween-20) containing 5% fat-free dry milk for 2 h and incubated for 3 h with rabbit polyclonal anti-human hPer2 antibody (dilution,

1:500) in TBST. Following incubation with a horseradish peroxidase-conjugated goat anti-rabbit secondary antibody (dilution, 1:2000; catalog no., SA00001-2; ProteinTech Group, Inc.), immunoreactive proteins were visualized with an enhanced chemiluminescence detection system. The western blot experiments were repeated three times. The relative expression of the target protein was calculated as the gray value ratio of target protein content to β -actin content (target protein/ β -actin) using Quantity One version 4.62 image analysis software (Bio-Rad Laboratories, Inc.).

Statistical analysis. Statistical analyses were carried out with the SPSS version 17.0 (SPSS, Inc., Chicago, IL, USA) statistical software package for Windows. All analyses in this study were performed using analysis of variance. All P-values were based on two-tailed tests and $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Agarose gel electrophoresis of RT-PCR product. The results of RT-PCR demonstrated that there was a visible DNA band just below 4 kb which had the same size as the expectant target gene (the length of hPer2 was 3765 bp; Fig. 2).

Identification of recombinant expression vector. The DNA sequencing results revealed that the sequence of the 3765 bp




Figure 5. Green fluorescence protein expression in MG63 cells observed by fluorescence microscope (magnification, x100). (A) pEGFP-N1 group; (B) pEGFP-N1-hPer2 group.




Figure 6. Expression of hPer2 mRNA and hPer2 protein. hPer2, pEGFP-N1-hPer2 group; Blank, pEGFP-N1 group; NC, control group. * $P < 0.05$.

inserted segment was identical to the cDNA sequence of the hPer2 gene. The results of nucleotide-nucleotide BLAST in NCBI also demonstrated that the sequence alignment was completely correct (Fig. 3). Following double enzyme digestion, a 3765-bp insertion segment and a 4.7-kb vector fragment were observed in the pEGFP-N1-hPer2 group following electrophoresis, while only a 4.7-kb vector fragment was observed in the pEGFP-N1 group. The results confirmed that the construction of the pEGFP-N1-hPer2 eukaryotic expression plasmid was successful (Fig. 4).

Transfection efficiency analysis of hPer2 gene in MG63 cells *in vitro*. The expression of the EGFP reporter gene was clearly observed using fluorescence microscopy (Olympus IX51) 48 h after transfection (Fig. 5) in the pEGFP-N1-hPer2 and the pEGFP-N1 group but not in the control group. The results revealed that in the pEGFP-N1-hPer2 and the pEGFP-N1 group, large numbers of MG63 expressed GFP. EGFP was expressed in 70% of cells in the pEGFP-N1-hPer2 group and 75% of cells in the pEGFP-N1 group, suggesting that pEGFP-N1-hPer2 and pEGFP-N1 may be effectively transfected into MG63 cells, resulting in a high level of EGFP expression. It was expected that the pEGFP-N1 group would express a more intense fluorescence signal than the pEGFP-N1-hPer2 group, since the empty pEGFP-N1 vector was smaller and had a higher transfection efficiency. In addition, as shown in Fig. 6, hPer2 mRNA expression was significantly higher in the pEGFP-N1-hPer2 group compared with the pEGFP-N1 or control groups ($P < 0.05$), and no statistical difference existed between pEGFP-N1 and the control group ($P > 0.05$), suggesting that hPer2 was successfully transfected into MG63 cells and efficiently expressed. Western

blot analysis revealed that hPer2 protein exhibited a significant upregulation in the pEGFP-N1-hPer2 group compared with the pEGFP-N1 and control group ($P < 0.05$), and no statistical difference existed between pEGFP-N1 and the control group ($P > 0.05$). These data also demonstrated that human osteosarcoma MG63 cell lines were successfully generated, in which hPer2 was overexpressed.

Discussion

Osteosarcoma is the most common primary bone tumor and mainly affects children and adolescents (7,8). The etiology of osteosarcoma is largely unknown due to the difficulties in understanding the molecular mechanism of tumor development in the complex structure and numerous genomic rearrangements of bone cancer cells (9). Complete radical surgery remains a preferable choice in osteosarcoma treatment, with adjuvant chemotherapy administered prior to surgery (10). If surgical excision is not possible, the addition of radiation therapy may be beneficial to control the local tumor. Still, a number of patients with osteosarcoma risk having local relapse following chemotherapy (11). For this reason, it is crucial to explore novel and alternative strategies for osteosarcoma treatment. Understanding the fundamental molecular mechanisms in the pathogenesis of osteosarcoma may help to develop novel strategies that have a specific molecular target for the treatment of patients with osteosarcoma (12).

Life on earth has evolved in the presence of a rhythmically changing environment. Most eukaryotes, and certain prokaryotes, have developed a molecular time-keeping mechanism that synchronizes itself with the external environment to

ensure optimal timing of cellular functions, metabolism and physiology (13-15). This mechanism is known as the circadian oscillator or clock. Circadian clocks have been identified in the majority of tissues and cells of mammals (16,17). The molecular basis of the circadian clock is the oscillatory transcription and translation of 'clock genes'. Studies suggest that tumorigenesis is associated with altered circadian function, whether causal or symptomatic, and that a dysfunctional circadian clock promotes carcinogenesis (18). However, the effects of clock genes on the biological behavior of osteosarcoma cells are rarely reported. In the current study, we selected hPer2, one of the key circadian genes, and successfully constructed the recombinant pEGFP-N1-hPer2 plasmid, which was transfected into the MG63 osteosarcoma cell line. This study is expected to lay the foundation for research into the circadian gene in osteosarcoma. A gene transformation technique is applied to modify tumor cells with certain functional genes to increase inhibition effects on the growth of tumors. The development of this technique has been extremely rapid in recent years and has become a new method in the precaution and treatment of tumors (19).

GFP easily forms a fusion protein with other target genes and has no influence on the space conformation and function of the target gene product. It has become the optimal reporter gene for the detection of the target gene transfection efficiency and expression modality (20). EGFP is optimized mutant GFP possessing a much higher sensitivity and 35 times stronger fluorescence than the wild type, having no species specificity and a notable influence on the growth and function of cells (21).

Taken together, in this study, in order to detect the transfection efficiency of the hPer2 gene, we constructed a recombinant plasmid using EGFP which exhibits the changes of its expression position and quantity within MG63 cells. We investigated the function of the target gene in the occurrence and development of tumors, as well as the molecular mechanisms involved. Our preliminary studies provide the ground work for further research on the roles of the circadian gene hPer2 in MG63 osteosarcoma cells, and pave the way for future studies of the intracellular downstream signaling mechanisms responsible for hPer2's ability to affect osteosarcoma cells.

Acknowledgements

The authors thank the Central Laboratory of the First Affiliated Hospital of Wuhan University, as well as Jishuang Zhu for his assistance in primer design and Dr Ling Yu for his careful review of the manuscript.

References

1. Partch CL, Green CB and Takahashi JS: Molecular architecture of the mammalian circadian clock. *Trends Cell Biol* 24: 90-99, 2014.
2. Kelleher FC, Rao A and Maguire A: Circadian molecular clocks and cancer. *Cancer Lett* 342: 9-18, 2014.
3. Greene MW: Circadian rhythms and tumor growth. *Cancer Lett* 318: 115-123, 2012.
4. Beckett M and Roden LC: Mechanisms by which circadian rhythm disruption may lead to cancer. *S Afr J Sci* 105: 415-420, 2009.
5. Ishida N: Circadian clock, cancer and lipid metabolism. *Neurosci Res* 57: 483-490, 2007.
6. Schmittgen TD and Livak KJ: Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3: 1101-1108, 2008.
7. Ottaviani G and Jaffe N: The epidemiology of osteosarcoma. *Cancer Treat Res* 152: 3-13, 2009.
8. Admassi D: Osteosarcoma of medial cuneiform bone. *Ethiop Med J* 47: 305-308, 2009.
9. Martin JW, Squire JA and Zielenska M: The genetics of osteosarcoma. *Sarcoma* 13: 627254, 2012.
10. Yamamoto N and Tsuchiya H: Chemotherapy for osteosarcoma - where does it come from? What is it? Where is it going? *Expert Opin Pharmacother* 14: 2183-2193, 2013.
11. Luetke A, Meyers PA, Lewis I and Juergens H: Osteosarcoma treatment - where do we stand? A state of the art review. *Cancer Treat Rev* 40: 523-532, 2014.
12. Yang J and Zhang W: New molecular insights into osteosarcoma targeted therapy. *Curr Opin Oncol* 25: 398-406, 2013.
13. McWatters HG, Roden LC and Staiger D: Picking out parallels: Plant circadian clocks in context. *Philos Trans R Soc Lond B Biol Sci* 356: 1735-1743, 2001.
14. Levine JD: Sharing time on the fly. *Curr Opin Cell Biol* 16: 210-216, 2006.
15. Dong G and Golden SS: How a cyanobacterium tells time. *Curr Opin Microbiol* 11: 541-546, 2008.
16. Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Slepka SM, Hong HK, Oh WJ, Yoo OJ, *et al*: PERIOD 2: LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci USA* 101: 5339-5346, 2004.
17. Peirson SN, Butler JN, Duffield GE, Takher S, Sharma P and Foster RG: Comparison of clock gene expression in SCN, retina, heart and liver of mice. *Biochem Biophys Res Comm* 351: 800-807, 2006.
18. Hanahan D and Weinberg RA: Hallmarks of cancer: the next generation. *Cell* 144: 646-674, 2011.
19. Noda M, Takahashi C, Matsuzaki T and Kitayama H: What we learn from transformation suppressor genes: lessons from RECK. *Future Oncol* 6: 1105-1116, 2010.
20. Hoffman RM: Imaging metastatic cell trafficking at the cellular level in vivo with fluorescent proteins. *Methods Mol Biol* 1070: 171-179, 2014.
21. Sakharova NY, Mezhevnikina LM, Smirnov AA and Vikhlyantseva EF: Analysis of the effects of blue light on morphofunctional status of in vitro cultured blastocysts from mice carrying gene of enhanced green fluorescent protein (EGFP). *Bull Exp Biol Med* 157: 162-166, 2014.