

Giant elephantiasis neuromatosa in the setting of neurofibromatosis type 1: A case report

GIOVANNI PONTI¹, GIOVANNI PELLACANI¹, DAVIDE MARTORANA²,
VICTOR DESMOND MANDEL¹, PIETRO LOSCHI³, ANNAMARIA POLLIO⁴, ANNARITA PECCHI⁵,
CRISTINA DEALIS⁶, STEFANIA SEIDENARI¹ and ALDO TOMASI⁷

¹Department of Surgical, Medical, Dental and Morphological Sciences with interest in Transplant, Oncological and Regenerative Medicine, University of Modena and Reggio Emilia, I-41124 Modena; ²Department of Genetics, University of Parma, I-43125 Parma; ³Department of Plastic Surgery, University of Modena, I-41124 Modena; ⁴Department of Neurosciences, University of Padua, I-43125 Padua; ⁵Department of Radiology, University of Modena and Reggio Emilia, I-41100 Modena; ⁶Department of Medical Oncology, Bolzano Hospital, I-39100 Bolzano; ⁷Department of Diagnostic and Clinical Medicine and Public Health, University of Modena and Reggio Emilia, I-41100 Modena, Italy

Received October 12, 2014; Accepted June 23, 2015

DOI: 10.3892/ol.2016.4469

Abstract. Elephantiasis neuromatosa (EN) can arise from a plexiform neurofibroma of the superficial and deep nerves developing from a hyperproliferation of the perineural connective tissue infiltrating adjacent fat and muscles. To date, the clinical association between EN and neurofibromatosis type 1 (NF1) has been poorly defined, particularly with regard to the role of lymphatic alterations and the consequent lymphedema. The present study reports the clinical and biomolecular features of EN in a NF1 patient with the clear clinical diagnostic criteria of multiple café-au-lait macules, neurofibromas, EN, a positive family history and a novel NF1 germline c.1541_1542del mutation. Lymphoscintigraphy (LS) highlighted marked dermal backflow in the affected limb, hypertrophy of the ipsilateral inguinal and external iliac lymph nodes, and a bilateral lower limb lymph flow delay. These data support the hypothesis that an extensive hyperproliferative process involving perineural connective, limb soft tissues, bones and the lymphatic system can be

responsible for EN in NF1 patients, on the basis of adipocyte metaplasia triggered by lymphostasis and lymphedema, and bone overgrowth and gigantism caused by chronic hyperemia. LS and magnetic resonance imaging can be efficacious tools in the diagnosis and clinical characterization of the early onset of the disease.

Introduction

Neurofibromatosis type 1 (NF1) (MIM no. 162200), also known as von Recklinghausen disease, is clinically characterized by the presence of simple, diffuse and plexiform neurofibromas. Plexiform neurofibromas are unencapsulated, poorly-circumscribed tumors infiltrating the nerves and adjacent fat and muscles (1). The connective overgrowth can be limited to a single nerve or a plexus; in the latter case, when the plexus spreads to the epidermal and dermal tissues, it is termed molluscum fibrosum. This can occur multiple times, covering all body sites (including the forehead, temple, eyes, nape and upper lip) with the exception of the palms and soles (2). The plexiform neurofibroma variant, mixoglioma gelatiniforme, is usually soft and is located in the lower third of the leg, and when associated with lymphangiomas, it can give rise to elephantiasis neuromatosa (EN). EN is characterized by abnormal soft-tissue hypertrophy and bone dysplasia together with early and excessive bone growth of the affected leg compared with the contralateral leg (3,4). Pachidermocele or dermatholysis may be associated with NF1, showing an overlap of skin layers in the thorax, buttocks and roots of the limbs.

The etiology of EN is not yet fully understood, but the association of primary lymphatic dysplasia with a lymphatic proliferative process has been proposed (5-7).

The current study presents a case of NF1-associated EN with typical clinical manifestations. Written informed consent was obtained from the patient.

Correspondence to: Dr Giovanni Ponti, Department of Surgical, Medical, Dental and Morphological Sciences with interest in Transplant, Oncological and Regenerative Medicine, University of Modena and Reggio Emilia, 71 via del Pozzo, I-41124 Modena, Italy
E-mail: giovanni.ponti@unimore.it

Abbreviations: NF1, neurofibromatosis type 1; EN, elephantiasis neuromatosa; LS, lymphoscintigraphy; MRI, magnetic resonance imaging; 3D-CT, three-dimensional computed tomography

Key words: neurofibromatosis type 1, elephantiasis neuromatosa, plexiform neurofibroma, mixoglioma gelatiniforme, primary lymphatic drainage

Case report

Case history. In January 2014, the case of a female patient was brought to our attention at the Department of Dermatology of University of Modena and Reggio Emilia (Modena, Italy) due to a 29-year history of several neurofibromas and multiple (>6) café-au-lait macules. The patient presented with giant elephantiasis of the right leg, which had started to grow during late childhood, and had since accelerated its expansion in the following years (Fig. 1A). At birth, the patient exhibited a café-au-lait macule on the right thigh. By 1 year old, a semi-liquid mass had developed at the same site. The lesion was noticed by the parents and showed indolent growth with no signs of bleeding or pain. Upon histopathological examination, it presented with the aspects of a lymphangioma. Lymphedema of the ipsilateral foot and discrepant leg lengths were noted successively. In the following years, the uneven leg growth was associated with bone proliferation, which required a number of osteotomies with the aim of stopping the growth. Conventional lymphoscintigraphy (LS) was performed to assess the lower limb lymphatic drainage pathway.

The family history was suggestive of NF1, as the patient's father exhibited macrocephaly, hypertelorism, and multiple café-au-lait macules and neurofibromas.

Genetic testing, magnetic resonance imaging (MRI) and three-dimensional computed tomography (3D-CT) were also performed.

MRI and 3D-CT. The MRI and 3D-CT scan study showed a preponderance of adipose tissue in the elephantiasic limb, corresponding to three-quarters of its whole volume, in addition to a severe dorso-lumbar-sacral scoliosis with convexity on the left. Two central nervous system hamartomas of the pallidus nucleus were also identified by MRI. 3D-CT revealed a mild lumbo-sacral meningocele and giant L4 neurofibroma (Fig. 1B).

Germline mutation analysis of the NF1 gene. Genomic DNA was extracted from the peripheral blood of the patient and the patient's father using the QIAamp DNA Blood Mini kit (Qiagen Inc., Valencia, CA, USA), and stored at -20°C until use. All NF1 exons were amplified by polymerase chain reaction with intron spanning primers, as described previously (8), and then analyzed with denaturing high-performance liquid chromatography, as described previously (9). For each abnormal elution profile, genomic DNA was directly sequenced in each direction using a CEQ Dye-Terminator Cycle Sequencing kit (Beckman Coulter Inc., Miami, FL, USA) according to the manufacturer's instructions.

Mutations were checked using the Mutalyzer program (<http://www.lovd.nl/mutalyzer>). NF1 germline deletion g.129042_129043delAG; c.1541_1542del; p.(Gln514Argfs*43) was found in the proband and the patient's father. To the best of our knowledge, this type of mutation has not previously been described.

Lymphedema and limb lymphatic assessment. A common tape measure was used to assess the limb circumferences of the patient. Reference circumferences were the popliteal crease, point zero (K), +30 cm (A), +20 cm (B) and +10 cm (C) in the thigh, -10 cm (D), -20 cm (E) and -30 cm (F) in the lower leg,



Figure 1. (A) Clinical aspect of elephantiasis neuromatosa. (B) Severe dorso-lumbar-sacral scoliosis with convexity on the left and preponderance of adipose tissue in the elephantiasic limb, corresponding to three-quarters of its whole volume, as detected by three-dimensional computed tomography scan and magnetic resonance imaging. (C) Conventional lymphoscintigraphy performed to assess the lower limb lymphatic drainage pathway: Images obtained at 60 and 120 min post-injection showed bilateral lower limb lymph flow delay.

Table I. EN as a clinical manifestation associated with the NF1 phenotype.

| First author/s, year (ref.) | Gender | Age/age of onset | Proband affected region | Family history | Clinical features |
|----------------------------------------|--------|---------------------|-------------------------|---------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Spittel and Fernando, 1929 (12) | M | 20 years/at birth | Left limb | Grandmother, 2,000 minute cutaneous tumours and scattered subcutaneous swellings; mother, achondroplasia. | Gross EN of the left limb. Subcutaneous tumors scattered all over the body; a pachydermatocele of the occipital scalp; cutaneous freckles. Hyperpigmentation of the mucous membrane of the mouth, and muddy hypopigmentation of the conjunctive. Bone development increasing. |
| Westcott and Ackerman, 1947 (13) | M | 40 years/at birth | Neck | His mother had NF-1. | EN of the neck. Asymmetry of the rib cage. Deformity of the cervical column, the left leg and the left clavicle. |
| Lenson, 1956 (14) | F | 22 years/childhood | Chest wall and axilla | No particular familial history of NF. | EN of extensively involved the chest wall and the axilla. |
| Fethiere <i>et al</i> , 1974 (15) | M | 42 years/- | Right leg | The family history is free from NF or any other inherited disease tendency. Her mother had gastric carcinoma. | Soft irregular bluish end mottled mass involving the distal two-thirds of the right lower extremity. Cavernous haemangioma with the consistency of a lipoma. Numerous femoral cysts. |
| Yaghmai and Tafazoli, 1977 (16) | M | 16 years/ childhood | Penis | Not reported. | EN of the penis. Syringomyelia associated with cervical ependymoma. Facial paralysis on the left side and paralysis of the left eye muscle. Multiple schwannomas involving certain cranial nerves, and numerous dorsal and ventral spinal roots with large tumor formations in cauda equina roots. |
| Harris <i>et al</i> , 1982 (18) | M | 11 years/3 years | Left thigh | Not reported. | Elephantiasis neuromatosa and overgrowth of abnormal bones with subperiosteal haemorrhage. Numerous café-au-lait spots and scoliosis of the dorsolumbar spine. |
| Sty <i>et al</i> , 1981 (17) | F | 9 years/2 years | Right leg | Not reported. | Elephantiasis neuromatosa, multiple café-au-lait spots and overgrowth of abnormal bones with subperiosteal haemorrhage. |
| Holck <i>et al</i> , 1984 (19) | M | 4 years/- | Right lower extremity | NF in 11 of his relatives. | EN involving the right lower extremity. Numerous café-au-lait spots on the lower abdomen and groin. The right lower extremity was longer than the left. Dilation of the medial lymphatic channels in the lower extremity and enlarged right iliac nodes. |
| Birch and Davies, 1988 (20) | F | 18 years/at birth | Right hallux | No familial history of NF. | Only one large café-au-lait spot on the right hallux. Soft-tissue hypertrophy with massive enlargement of the proximal and distal phalanges. |
| Hertzanu <i>et al</i> , 1989 (21) | F | 17 years/childhood | Right gluteal sulcus | There was no predisposition to either familial or hereditary disorders. | EN of the right gluteal sulcus, coexistent or with lipomatosis. No skeletal lesion. Nor were other stigmata of von Recklinghausen's disease apparent. |
| Bardelli and Hadjistilianou, 1989 (22) | F | 30 years/at birth | Left lower limb | Not reported. | EN of the left lower limb maximal in the calf. Dysplastic bones of the left hemipelvis and leg with florid osseous striations. The skin of the thigh was loose and inelastic. |
| | F | 52 years/childhood | Trunk | The patient denied any manifestations of NF in her family. | Symmetric EN of the trunk. At 32 years: Hypovolemic shock due to massive hemorrhage into the mass. |
| | M | Few months/at birth | Facial hemi-hypertrophy | No family history of congenital glaucoma or buphthalmos. | Buphthalmos and regional gigantism with evolution in EN. Corneal diameter 13 mm, rubeosis iridis, ectropion uveae. An increase of optic nerve cupping. A swelling of the right upper lid and preauricular region. Hypertrophy of the soft periorbital tissues. Small nodules on conjunctiva of affected eye. |

Table I. Continued.

| First author/s, year (ref.) | Gender | Age/age of onset | Proband affected regions | Family history | Clinical features |
|-------------------------------------------|--------|--------------------------|-------------------------------|---------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Bardelli and Hadjistilianou, 1989 (22) | M | 13 months/at birth | Right orbital region | Not reported. | Buphthalmos. An enlarged optic canal, a fibrous dysplasia of the greater wing of sphenoid bone. Longer right fibula and a slight deformity of the vertebrae of the lumbar tract. |
| | F | 11 years/at birth | Temporal and zygomatic region | Not reported. | Facial elephantiasis without buphthalmos. Greater wing of sphenoid bone; right optic foramen larger than the left. Iris nodules, ectropion, uveae, increased optic nerve cupping. Small neurofibromas of the zygomatic and temporal region. |
| Kuo and Kuo, 1990 (23) | F | 34 years/8 years | Right limb and pelvis | His mother, grandmother, sister and nephews had NF. | EN involving the right lower limb and pelvis, leading to a right hip disarticulation. Grossly enlarged lower limb complicated by infected decubitus ulcers. A leg length discrepancy. (After surgery a large neurofibroma of the sciatic nerve). |
| Roy and Ghosh, 1992 (24) | F | 22 years/at birth | Right foot | No family history of NF1 was presented. | Enlargement of the whole right foot with conspicuous gigantism of the 2nd to 5th toes. The overlying skin was coarse, dry, thick. Overgrowth and elongation of all the bones of the foot (2nd and 3rd metatarsals, and their phalanges). A huge soft-tissue mass with no bone involvement. |
| | M | 21 years/at birth | Right foot | Not reported. | Gigantism of the right foot and elongation of all the bones of the foot (metatarsals and a huge soft-tissue mass). |
| | M | 18 years/early childhood | Right foot | His father had EN. One of his three brothers has a few cutaneous nodules affecting the right lower limb only. | Gigantism of the right foot with swelling at the heel. Elongation of all the bones (metatarsals and a soft-tissue mass affecting mainly the hind foot). |
| | M | 6 years/at birth | Right foot | No family history of NF. | Enlargement mainly of the forefoot and the swelling was more prominent between the great and 2nd toe. No café-au-lait spots. |
| Kokandkar <i>et al.</i> , 1994 (25) | M | 3 months/at birth | Neck and back | Family history did not reveal occurrence of similar illness in any member. | Congenital plexiform neurofibroma involving the neck with EN with a sarcomatous nodule. The skin covering the tumor was hairy, redundant and dark. The skin covering the nodules was thinned out and ulcerated. |
| Münte <i>et al.</i> , 1996 (26) | F | 33 years/6 years | Right leg | Family history of NF. | EN of the right leg. Severe dysaesthesia in the right lower leg not confined to a single nerve. |
| Stevens <i>et al.</i> , 1998 (27) | M | 43 years/childhood | Left leg | Not reported. | Gross EN of the left leg. Large synovial cyst arising from the synovium of the patello-femoral joint. |
| Akyol <i>et al.</i> , 1999 (28) | M | 20 years/6 years old | Left shoulder and arm | No family history of NF. | EN with Becker's melanosis. Hairy and brown-black hyperpigmented patches on left shoulder, left upper back and left arm. Lisch nodules. |
| Lorberbom <i>et al.</i> , 2000 (29) | M | 35 years/- | Right thigh and sacral region | Not reported. | A soft-tissue mass and enlargement of the right upper leg. |
| Steenbrugge <i>et al.</i> , 2001 (30) | F | 13 years/at birth | Left limb | Her mother had NF-1. | Left leg elephantiasis with recurrent massive subperiosteal hematoma. |
| Hourani <i>et al.</i> , 2006 (31) | F | 41 years/childhood | Right limb | Not reported. | EN of the right leg. Optic chiasm glioma. Right tibia and fibula marrow and cortices hypertrophy. |
| Martínez-García <i>et al.</i> , 2008 (32) | F | 14 years/at birth | Right limb | Not reported. | EN involving the right lower limb. Anaemia and hepatitis B. |

Table I. Continued.

| First author/s, year (ref.) | Gender | Age/age of onset | Proband affected regions | Family history | Clinical features |
|---------------------------------|--------|--------------------|--------------------------|-----------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|
| Hoshi <i>et al.</i> , 2009 (33) | M | 56 years/childhood | Right leg | There was no particular family history of NF. | A huge mass of EN in the right leg. |
| Bano <i>et al.</i> , 2010 (1) | M | 15 years/childhood | Right limb | No family history of NF1 in first-degree relatives. | EN of the right leg. Osseous abnormalities including thinning of bones, erosion of distal articular surfaces and periosteal dysplasia. |

NF1, neurofibromatosis type 1; EN, elephantiasis neuromatosa.

and 10 cm proximal from the tip of the first toe (G). Leg volumes were calculated according to these measurements (10).

Lower limb lymphatic function was assessed by LS, using injections of 37-MBq ^{99m}Tc-labeled human serum albumin. Injection points were in the first, second and fourth interdigital and retromalleolar spaces of the affected and contralateral foot. Image acquisition was obtained after 60 and 240 min using a dual head γ -camera (Philips Healthcare, Andover, MA, USA) equipped with a low-energy and high-resolution collimator, and an energy peak centered on 140 KeV (window of 20%).

The criteria to define lymphatic dysfunction include delay, asymmetric or absent visualization of regional lymph nodes, and the presence of dermal backflow. Other observations include the visualization of asymmetric lymphatic channels, collateral lymphatic channels, interrupted vascular structures, and lymph nodes of the deep lymphatic system (i.e., popliteal lymph nodes following web space injection in the lower extremities) (11).

In LS, the images obtained 60 min after the injections showed bilateral lower limb lymph flow delay. Mild dermal backflow in the absence of tracer migration was observed in the affected lower limb, whilst one inguinal lymph node was observed in the left limb. Posterior images confirmed the findings, also visualizing a popliteal lymph node in the healthy (left) leg. Images acquired at 240 min showed significant dermal backflow in the right limb, and hyperplasia and hypertrophy of the inguinal and external iliac lymph nodes, in comparison to those of the left leg (Fig. 1C).

Treatment and patient outcome. In May 2014, the patient was treated with surgery to reduce the volume of tissue in the leg region; tissue with a mass of 4.3 kg was removed. However, the outcome was not as good as expected due to hemorrhage during surgery, which prevented complete exision. Following surgery, the patient exhibited limited functional improvement and limb lightening. The patient was subsequently administered anticoagulant therapy with enoxaparin (8,000 IU, daily for one year) and underwent regular follow up examinations every three months for the first year and every six months subsequently. At the time of writing, the patient was well with a good prognosis.

Discussion

EN is a rare clinical manifestation associated with the NF1 phenotype. The condition should be defined as early and excessive growth in the width and length of the affected limb due to a neoplastic proliferation of the perineural connective tissue, together with congenital lymphatic insufficiency and chronic hyperemia. While no more than 30 cases are described in the literature (Table I) (12-32), the real incidence is estimated to be higher. The clinical expression is characterized by plexiform neurofibromas located in the superficial or deep nervous system associated with congenital lymphangiomatosis. Signs usually appear during the first years of life, due to lymphostasis and subsequent lymphedema causing adipocyte metaplasia of the adjacent tissues and chronic hyperemia inducing bone overgrowth and focal gigantism (34).

Distinct superficial dysplastic skin alterations known as pachidermocele or dermatholysis, histologically corresponding to mixoglioma gelatiniforme, must be distinguished from EN in patients affected by NF1 (2).

To date, little evidence is available regarding the role of lymphatic alterations in the pathogenesis of EN.

Based on the bilateral lymphatic defect, the presence of a primary lymphatic disease in the current NF1 patient can be hypothesized. The disease is probably supported by a dysplastic-hypertrophic condition as a result of a congenital alteration of the lymphatic network (35,36).

Regarding the lymph to fat transformation, it is known that lymphostasis due to primary and secondary lymphedema determines the transformation of fat cells, resulting in hypertrophied adipose tissue. Several studies (37-39) have suggested that lymphedema leads to adipose tissue accumulation and fibrosis. Moreover, we believe that this process is amplified in NF1 and in EN due to a primary lymphatic disorder, which is at the base of the clinical manifestation induced by the plexiform neurofibroma growth.

Overall, the diagnostic criteria for NF can be improved by the introduction and application of novel criteria based on a wider case series (EN, focal gigantism, mixoglioma gelatiniforme and primary lymphatic disorder), leading to the early diagnosis of NF1, particularly in pediatric patients. LS and MRI can be efficacious tools in the diagnosis and clinical characterization of early onset cutaneous, subcutaneous and skeletal anomalies.

Acknowledgements

The authors would like to thank Dr Federica Arginelli for providing useful comments.

References

- Bano S, Prasad A, Yadav SN, Chaudhary V and Sachdeva N: Elephantiasis neuromatosa of the lower limb in a patient with neurofibromatosis type-1: A case report with imaging findings. *J Pediatr Neurosci* 5: 59-63, 2010.
- Pollock G: Report of a case of mollusum fibrosum or fibroma with observations. *Med Chir Trans* 56: 255-266, 1873.
- Serradell AP: Neurocutaneous syndromes. *Medicine - Programa de formación médica continuada acreditado* 8: 5532-5547, 2003 (In Spanish).
- Ponti G, Martorana D, Pellacani G, Ruini C, Loschi P, Baccarani A, De Santis G, Pollio A, Neri TM, Mandel VD, *et al*: NF1 truncating mutations associated to aggressive clinical phenotype with elephantiasis neuromatosa and solid malignancies. *Anticancer Res* 34: 3021-3030, 2014.
- Flanagan BP and Helwing EB: Cutaneous lymphangioma. *Arch Dermatol* 113: 24-30, 1977.
- Foeldi M: The role of the lymphatic circulation in the fluid circulation of the eye and the central nervous system. *Arch Kreislaufforsch* 41: 186-212, 1963.
- Lohrmann C, Pache G, Felmerer G, Foeldi E, Schaefer O and Langer M: Posttraumatic edema of the lower extremities: Evaluation of the lymphatic vessels with magnetic resonance lymphangiography. *J Vasc Sur* 49: 417-423, 2009.
- De Luca A, Bottillo I, Dasdia MC, Morella A, Lanari V, Bernardini L, Divona L, Giustini S, Sinibaldi L, Novelli A, *et al*: Deletions of NF1 gene and exons detected by multiplex ligation-dependent probe amplification. *J Med Genet* 44: 800-808, 2007.
- De Luca A, Buccino A, Gianni D, Mangino M, Giustini S, Richetta A, Divona L, Calvieri S, Mingarelli R and Dallapiccola B: NF1 gene analysis based on DHPLC. *Hum Mutat* 21: 171-172, 2003.
- International society of lymphology: The diagnosis and treatment of peripheral lymphedema. 2009 consensus document of the international society of lymphology. *Lymphology* 42: 51-60, 2009.
- Pecking AP: Possibilities and restriction of isotopic lymphography for the assessment of therapeutic effects in lymphedema. *Wien Med Wochenschr* 149: 105-106, 1999.
- Spittel RL and Fernando SE: A case of elephantiasis neuromatosa. *Br Med J* 1: 596-597, 1929.
- Westcott RJ and Ackerman LV: Elephantiasis neuromatosa; a manifestation of von Recklinghausen's disease. *Arch Derm Syphilol* 55: 233-241, 1947.
- Lenson N: Neurofibromatosis; a case report of elephantiasis neuromatosa of the right lower extremity, with invasion of the popliteal artery. *AMA Arch Surg* 73: 279-284, 1956.
- Fethiere W, Carter HW and Sturim HS: Elephantiasis neuromatosa of the penis. Light and electron microscopical studies. *Arch Pathol* 97: 326-330, 1974.
- Yaghmai I and Tafazoli M: Massive subperiosteal hemorrhage in neurofibromatosis. *Radiology* 122: 439-441, 1977.
- Sty JR, Starshak RJ and Woods GA: Neurofibromatosis: Lymphoscintigraphic observations. *Clin Nucl Med* 6: 264-265, 1981.
- Harris WC Jr, Alpert WJ and Marcinko DE: Elephantiasis neuromatosa in von Recklinghausen's disease. A review and case report. *J Am Podiatry Assoc* 72: 70-72, 1982.
- Holck S, Medgyesi S, Darre E and Lassen M: Elephantiasis neuromatosa. A light, immunohistochemical and electron microscopic study. *Virchows Arch A Pathol Anat Histopathol* 404: 427-434, 1984.
- Birch PD and Davies AM: The value of computed tomography in elephantiasis neuromatosa. *Br J Radiol* 61: 76-78, 1988.
- Hertzanu Y, Hirsch M, Peiser J and Avinoach I: Computed tomography of elephantiasis neuromatosa. *J Comput Assist Tomogr* 13: 156-158, 1989.
- Bardelli AM and Hadjistilianou T: Buphthalmos and progressive elephantiasis in neurofibromatosis. A report of three cases. *Ophthalmic Paediatr Genet* 10: 279-286, 1989.
- Kuo LA and Kuo RS: Plexiform neurofibromatosis: A difficult surgical problem. *Aust N Z J Surg* 60: 732-735, 1990.
- Roy SM and Ghosh AK: Elephantiasis neuromatosa: A clinicopathologic study of four cases. *J Ind Med Assoc* 90: 185-187, 1992.
- Kokandkar HR, Vyas AS, Kumbhakarna NR and Totala RJ: Congenital plexiform neurofibroma with a sarcomatous nodule in a three month old child. *Indian J Cancer* 31: 130-132, 1994.
- Münste TF, Matzke M, Johannes S, Dietrich B and Dengler R: MRI of elephantiasis neuromatosa. *J Neurol* 243: 619, 1996.
- Stevens KJ, Ludman CN, Sully L and Preston BJ: Magnetic resonance imaging of elephantiasis neuromatosa. *Skeletal Radiol* 27: 696-701, 1998.
- Akyol M, Özçelik S, Marufihah M and Elagöz S: Elephantiasis neuromatosa and Becker's melanosis. *J Dermatol* 26: 396-398, 1999.
- Lorberboym M, Trejo L and Lampl Y: Bone scintigraphy of elephantiasis neuromatosa in Von Recklinghausen's disease. *Clin Nucl Med* 25: 812-813, 2000.
- Steenbrugge F, Poffyn B, Uyttendaele D, Verdonk R and Verstraete K: Neurofibromatosis, gigantism, elephantiasis neuromatosa and recurrent massive subperiosteal hematoma: A new case report and review of 7 case reports from the literature. *Acta Orthop Belg* 67: 168-172, 2001.
- Hourani R, Rizk T, Kung S and Boudghène F: Elephantiasis neuromatosa in neurofibromatosis type I. MRI findings with review of the literature. *J Neuroradiol* 33: 62-66, 2006.
- Martínez-García S, Vera-Casaño A, Eloy-García Carrasco C, del Boz-González J, Martínez-Pilar L and Crespo-Erchiga V: Elephantiasis neuromatosa in a patient with neurofibromatosis type 1. *J Eur Acad Dermatol Venereol* 22: 103-105, 2008.
- Hoshi M, Ieguchi M, Taguchi S and Yamasaki S: A case report of surgical debulking for a huge mass of elephantiasis neuromatosa. *Rare Tumors* 1: e11, 2009.
- Holt JF: 1977 Edward B. D. Neuhauser lecture: Neurofibromatosis in children. *AJR Am J Roentgenol* 130: 615-639, 1978.
- Liu NF, Yan ZX and Wu XF: Classification of lymphatic-system malformations in primary lymphoedema based on MR lymphangiography. *Eur J Vasc Endovasc Surg* 44: 345-349, 2012.
- Preston JM, Starshak RJ and Oechler HW: Neurofibromatosis: Unusual lymphangiographic findings. *AJR Am J Roentgenol* 132: 474-476, 1979.
- Ryan TJ: Lymphatics and adipose tissue. *Clin Dermatol* 13: 493-498, 1995.
- Rosen ED: The molecular control of adipogenesis, with special reference to lymphatic pathology. *Ann N Y Acad Sci* 979: 143-158; discussion 188-196, 2002.
- Brorson H: Liposuction normalizes-in contrast to other therapies-lymphedema-induced adipose tissue hypertrophy. *Handchir Mikrochir Plast Chir* 44: 348-354, 2012.