

Anti-tumoral activity of native compound morelloflavone in glioma

XIANFENG LI¹, HONGYAN AI¹, DEKE SUN¹, TAO WU¹, JIAN HE¹, ZHAI XU², LI DING³ and LING WANG³

¹Neurosurgery Ward 2, Linyi City Yishui Central Hospital, Linyi, Shandong 276400; ²Department of Neurological Surgery Unit 1, The First Affiliated Hospital of Liaoning Medical University, Jinzhou, Liaoning 121001; ³Intensive Care Unit, Linyi Municipal People's Hospital, Linyi, Shandong 276000, P.R. China

Received April 20, 2016; Accepted September 1, 2016

DOI: 10.3892/ol.2016.5094

Abstract. The aim of the study was to investigate the anti-tumoral activity of morelloflavone substances with different structures. We also studied the possible link between morelloflavone structure and its function. Various types of chromatographic techniques were used to isolate and screen morelloflavone substances from the extracts of gambogic tree trunk and the morelloflavone structures were identified by analytical techniques such as high resolution mass spectrometry and nuclear magnetism. Anti-tumoral activities of different compounds were investigated and the link between the antitumor activity and the structure of compound was examined. Our results showed that the isolated morelloflavone substances demonstrated a certain level of antitumor activity. The compound no. 1 had the strongest effect to inhibit glioma U87 and C6 cells followed by compound no. 2 while compound no. 5 was the weakest among them. We conducted a preliminary analysis on the structure-function relationship through the structure comparison and we concluded that the antitumor effects of morelloflavone substances with different structures were significantly different from each other. Thus, the glucose chain in C4 position of biflavone structure can enhance the antitumor activity of the compound in glioma cells. Additionally, the formation of intramolecular hydrogen bonds in biflavone compounds may also play a role in enhancing the antitumor activity and inhibition rate.

Introduction

Glioma originates from nerve epithelium-derived cells (1), accounting for 44.6% of the central nervous system tumors. It is the most common malignant tumor in the central nervous

system with a high incidence and high mortality rate (2). Despite several achievements in cancer therapy in recent years, no effective treatment for glioma has been found. Currently, surgery-based comprehensive treatment is the treatment of choice for glioma, but the prognosis of comprehensive treatment remains poor. Nevertheless, this treatment can improve the recovery rate of glioma to a certain extent with 5-year survival rate of <5% (3,4). Therefore, there is a huge interest in developing a new means of treatment for brain glioma. The application of traditional Chinese medicine for treating the malignant tumors has a long history in China, but traditional Chinese medicine has failed to be recognized in the medical field due to the complexity of Chinese herbal medicinal ingredients and unclear antitumor mechanism. Thus, there are several restrictions in conducting clinical trials for traditional Chinese remedies (5,6). In recent years, several studies have been conducted on the adverse side effects associated with semi-synthetic and synthetic chemotherapy drugs used for treatment of tumors. As a result, there is interest in research on the natural antitumor compounds extracted from Chinese medicines. The *Garcinia* plants are known for their therapeutic effects and there is great interest in the medical effects of *Garcinia*. There are studies on using *Garcinia* for its anti-HIV, antibiosis, antioxidation, anti-inflammation, anti-malarial, insecticide and antitumor activities (7-10). Morelloflavone substances in the *Garcinia* plant have been shown to have broad-spectrum antitumor effects, although their action mechanism remains to be determined (11).

We extracted various plant substances from *Garcinia* plants and then screened the flavonoid substances using LC-MS. The specific structure of these compounds was analyzed using nuclear magnetic resonance and the effects of different compounds on glioma U87 cells and rat glioma C6 cells were investigated. We also analyzed the relationship between the structure of compounds and their anti-glioma effects. This study provides a theoretical reference for further research on anti-glioma agents.

Materials and methods

Experimental materials. The materials used for the present study were: Ethyl alcohol, methyl alcohol (chromatographic grade; Sinopharm Chemical Ltd., Shanghai, China); ultra-pure

Correspondence to: Dr Zhai Xu, Department of Neurological Surgery Unit 1, The First Affiliated Hospital of Liaoning Medical University, 2 Wuduan Renmin Street, Jinzhou, Liaoning 121001, P.R. China
E-mail: zhaixu365@163.com

Key words: biflavone, glioma, structure-function relationship

water (Milli-Q); gambogic tree trunk; deuterated methanol, dimethyl sulfoxide (DMSO); and human glioma U87 and rat glioma C6 cells (ATCC).

Experimental method

Extraction of active substances of gambogic tree trunk. Gambogic tree trunk extract (50.00 g) was broken into pieces and 70% of ethyl alcohol was added. It was transferred to a water bath (80°C) for 8 h, and the leaching liquid was collected and the supernatant after the leaching liquid was centrifuged at 2,000 × g for 15 min and filtrated. A rotary evaporator was used to remove the solvent for freeze-drying, and it was stored at -20°C.

Chromatographic separation of gambogic leaching liquid. Based on Waters preparative high performance liquid chromatography (Pre-HPLC), the XBridge C18 (5 μm, 21.2x250 mm) chromatographic column (Waters, Milford, MA, USA) was used to isolate the gambogic leaching liquid. Conditions for chromatography were as follows: Column temperature was 40°C; flow velocity was 4 ml/min; mobile A phase was methyl alcohol, mobile B phase was 0.1% formic acid aqueous solution, pH 5.0; time 0 min, A (40%), B (60%); 3 min, A (40%), B (60%); 25 min, A (40%), B (60%); 27 min, A (100%), B (0%); 37 min, A (40%), B (60%); and 45 min, A (40%), B (60%). The effluent components were collected according to the chromatographic peak and C18 solid phase extraction column was used to desalt each component. Components were freeze-dried by degreasing solvent and stored at -20°C.

Screening the biflavone substances. High resolution mass spectrometry (HRMS TOF; Waters, Milford, MA, USA) was employed to identify the collected components. Each component was dissolved in the methyl alcohol by connecting the HRMS to the ultra-performance liquid chromatography (UPLC; Waters). Waters BEH C18 (1.7 μm, 2x150 mm) chromatographic column was used for injecting samples directly and each component was identified with HRMS (MassLynx 4.1 analysis method) and by consulting literature and comparing CAS database.

Structure identification. The biflavone substances were prepared again to make the quantity of biflavone substances to reach the concentration necessary for nuclear magnetic resonance. Magnetic resonance spectroscopy (NMR; Bruker ARX 400; Karlsruhe, Germany) was used to confirm and analyze their structure. The sample solvent of nuclear magnetic resonance was CD₃OD and the internal sample standard was TMS. Data acquired were analyzed by MestreNova 4.1 (San Diego, CA, USA).

Inhibition rate of U87 cells of biflavone substances by MTT detection. The biflavone substances with different structures were dissolved in DMSO, and the compound concentration was 0, 10, 20, 40 and 80 μmol/l. The sample of 0 μmol/l was taken as the blank control. The specific experimental operation methods were similar to those reported by Vandamme *et al* (12).

The calculation method for obtaining the inhibition rate was: IR inhibition rate (%) = (1 - OD value of experimental port/OD value of blank control port) × 100%. The mean of




Figure 1. Chromatographic separation of gambogic extraction solution.

the inhibition rate was the end value and the inhibition rate of blank control port was 0 according to the calculation.

Preparation of glioma animal model. To prepare rat glioma cell C6 xenograft tumor model we followed the method reported by Li *et al* (13) with a slight modification. Approval for the animal experiments was received from The First Affiliated Hospital of Liaoning Medical University (Liaoning, China).

Statistical analysis. SPSS 18.0 statistical software (Chicago, IL, USA) was used to conduct statistical analysis. The mean ± standard deviation (SD) was used to record the independent experimental data and the comparison among various groups was analyzed using one-way ANOVA. Inter-group comparisons were tested by t-test. P<0.05 was considered to indicate a statistically significant difference.

Results

Pre-HPLC isolation of gambogic extraction solution. The ethanol water solution was used to extract the active ingredients from the gambogic tree trunk. Due to the fact that flavonoids contain benzene and pyranoid rings in their structure we used 275 nm wavelength to detect the ultraviolet absorption band. To ensure rapid isolation of flavonoids, the gradient elution method was used. The gradient of methanol started from 40% and gradually increased to 100% in order to elute all components. Results for final chromatographic separation are presented in Fig. 1. The chromatogram showed that the separation degree between the chromatographic peaks was improved. Components were eluted from chromatographic column and collected according to the chromatographic peaks (18 components in total). Due to the presence of formic acid in the mobile phase, collected components were desalinated and purified, and C18 solid phase extraction column was used to absorb active components. Ultra-pure water was then used to elute salts and impurities and finally the methanol was used to elute active substances from the solid phase extraction column. Active substances were freeze-dried after the solvent was removed and then stored at -20°C.

Screening of biflavone substances. We used HRMS to analyze the structure of biflavones. Due to the fact that all 18 components mentioned in the present study were isolated and purified by liquid chromatography, the direct injection technique was used for HRMS. Components structures were determined according to the accurate molecular mass and ion fragments information, and finally the biflavone substances

Table I. HRMS information of biflavone substances.

Substances ^a	Retention time	Measured molecular mass	Actual molecular mass	Molecular formula	Error m/z (ppm)	Molecular fragment
1	17.8	751.1505	751.1510	C ₃₆ H ₃₁ O ₁₈	-0.7	151.0031, 285.0392, 435.0708, 437.0866, 445.0556
2	18.6	735.1564	735.1561	C ₃₆ H ₃₁ O ₁₇	+0.4	125.0237, 269.0448, 403.0818, 429.0609, 447.0717
3	24.6	589.0972	589.0982	C ₃₀ H ₂₂ O ₁₃	-1.7	125.0237, 151.0028, 285.0393, 435.0713, 463.0660
4	26.6	573.1043	573.1033	C ₃₀ H ₂₂ O ₁₂	+1.7	125.0237, 296.0313, 419.0761, 447.0711
5	32.7	557.1084	557.1083	C ₃₀ H ₂₂ O ₁₁	+0.1	125.2038, 269.0443, 296.0317, 403.0813, 431.0764

^aCorresponding to the structure of number of substances as shown in Fig. 2. HRMS, high resolution mass spectrometry.

Table II. Proliferation inhibition effect of biflavone substances on glioma U87 cell (% , mean \pm SD).

Concentration (μ mol/l)	Compound 1	Compound 2	Compound 3	Compound 4	Compound 5
0	0.13 \pm 0.11	0.14 \pm 0.09	0.08 \pm 0.12	0.11 \pm 0.10	0.09 \pm 0.15
10	12.24 \pm 0.24	14.10 \pm 0.55	6.75 \pm 0.41	6.12 \pm 0.33	2.23 \pm 0.54
20	19.78 \pm 0.28	17.64 \pm 0.86	8.17 \pm 0.96	8.43 \pm 0.35	1.85 \pm 0.21
40	25.66 \pm 0.68	32.73 \pm 0.94	12.31 \pm 0.68	11.92 \pm 0.68	4.31 \pm 0.35
80	55.64 \pm 0.65 ^a	49.82 \pm 0.91 ^{a,b}	21.17 \pm 0.84 ^{a-c}	22.78 \pm 0.75 ^{a-c}	9.87 \pm 0.66 ^{a-d}

^{a-d}As for the comparison on single factor inter-class variance, the same letters ($P>0.05$), indicate that there is no significant difference; the different letters ($P<0.05$), indicate that there is a significant difference.

were determined by comparing our data with those available in the literature (14,15) and CAS database. Five biflavone substances were finally selected (Table I).

Although HRMS is a fast and effective method to determine the molecular structure of different compounds, structure of biflavones cannot be completely determined just by mass spectrometry. In order to determine the specific structure and position of each functional group they should be analyzed by nuclear magnetic resonance. Therefore, we used a relatively HRMS and prepared large quantities of samples to for nuclear magnetic resonance.

Structure determination of biflavone substances by nuclear magnetic resonance. Five biflavone substances were dissolved in CD₃OD and ¹H and ¹³C spectrum was then analyzed by nuclear magnetic resonance and the specific structures of these compounds were determined by comparing them with literature (16-18) and spectrum database (Fig. 2). After determination of the biflavone molecular structures, we investigated the inhibition effects of these substances on glioma and then analyzed the possible relationship between the structure and the function.

Inhibition effect on U87 cells. Different concentrations of each biflavone substance were added to cells and after 24 h the inhibition rate on glioma U87 cells was determined (Table II). Our results showed that the five biflavone substances demonstrated some inhibitory activities on U87




Figure 2. Structure information of biflavone substances.

glioma cells. The inhibition rate of compound no. 1 was the highest followed by compound no. 2. The inhibition rate of compound no. 3 had no significant difference from that of compound no. 4, while the inhibition rate of compound no. 5 was the lowest.

Inhibition of five biflavone substances on cell C6. To verify the inhibition activity of biflavone substances on xenograft tumor cells, the axillary xenograft tumor model of rat glioma C6 cells was used. Rats were injected with 1 mg/kg of cisplatin and 800 μ g/kg of biflavone on the seventh day after the inoculation of glioma C6 cells. The results are presented in Table III.

Table III. Inhibition effect of biflavone substances on the growth of rat glioma cell C6 xenograft tumors.

Groups	Dosage ($\mu\text{g/kg}$)	Weight (g)	Tumor change (g)	Inhibition rate (%)
Control	0	7.13	2.33 \pm 1.11	0
Compound 1	800	5.62	0.95 \pm 0.55	66.73 \pm 0.12 ^a
Compound 2	800	6.77	1.23 \pm 0.62	65.21 \pm 0.35 ^a
Compound 3	800	4.89	2.28 \pm 0.47	52.12 \pm 0.51 ^{a,b}
Compound 4	800	5.12	2.59 \pm 0.41	53.31 \pm 0.28 ^{a,b}
Compound 5	800	6.08	4.41 \pm 0.53	31.84 \pm 0.33 ^{a-c}

^aCompared with the control, $P < 0.05$. ^bCompared with compound 1, $P < 0.05$. ^cCompared with compound 3, $P < 0.05$.

Our results showed that the inhibition activity of biflavone substances on glioma C6 cells was similar to the inhibition effect in the case of U87 cells. Compound no. 1 was the most potent compound among the five. Compound no. 2 was the second and compound no. 3 had no significant difference compared to compound no. 4. The inhibition rate of compound no. 5 was the lowest.

Discussion

Glioma is one of the most common malignant tumors occurring in the central nervous system, and the incidence of glioma accounts for 50-60% of brain tumors (19,20). Although surgery remains the method of choice for treating malignant glioma, the efficacy of surgery has not greatly improved in recent years. Thus, the trial of drug therapy or assisted surgery therapy has a certain practical significance and broad applications. There are studies on adverse effects associated with synthetic and semi-synthetic drugs used in chemotherapy, therefore people are gradually paying more attention to the research on natural antitumor compounds extracted from traditional Chinese medicine. As type of traditional Chinese medicine, the *Garcinia* has various types of medical effects and the biflavone substances that the carcinia contains have broad-spectrum antitumor effects, but the action mechanism of these biflavone substances are still unclear.

Our results showed that the inhibition effects of biflavone substances on glioma U87 and C6 cells were alike. The inhibition effect of compound no. 1 was the strongest followed by compound no. 2 whose inhibition effect was significantly greater than that of compounds nos. 3-5. From studying the structure of all five compounds we detected the presence of a glucose chain in the structure of compound nos. 1 and 2. The preliminary structure-function relationship showed that the circumscribed glucose chain in the biflavone compounds could enhance the inhibition effect. It is possible that the glucose chain was responsible for these adverse effect on the cell DNA (21).

Compound no. 5, had $R_1=R_2=H$, while compounds nos. 1-4 $R_1=R_2=\text{hydroxyl group}$. It was hypothesized that the reduction of hydroxyl group in biflavone substances may

reduce their inhibition effects. Additionally, the inhibition activity of compound no. 2 was significantly lower than that of compound no. 1. The existence of hydroxyl group can stimulate the formation of external as well as internal hydrogen bonds within and between molecules and indirectly enhance the interaction between molecules and improved the inhibition rate on glioma.

In future, more biflavone substances will be isolated and screened from different parts of *Garcinia* plants and their anti-tumoral activity will be verified. This may lay a more solid foundation for the further verification and exploration of the possible link between biflavone structure and its function.

References

1. Johnson BE, Mazon T, Hong C, Barnes M, Aihara K, McLean CY, Fouse SD, Yamamoto S, Ueda H, Tatsuno K, *et al*: Mutational analysis reveals the origin and therapy-driven evolution of recurrent glioma. *Science* 343: 189-193, 2014.
2. Lee Tittsworth W, Murad GJ, Hoh BL and Rahman M: Fighting fire with fire: the revival of thermotherapy for gliomas. *Anticancer Res* 34: 565-574, 2014.
3. Mirimanoff RO: High-grade gliomas: reality and hopes. *Chin J Cancer* 33: 1-3, 2014.
4. Giovagnoli AR, Meneses RF, Silvani A, Milanesi I, Fariselli L, Salmaggi A and Boiardi A: Quality of life and brain tumors: what beyond the clinical burden? *J Neurol* 261: 894-904, 2014.
5. Ji B, Shi J, Cheng X, Zhou J, Zhou Q, Cao C and Pang J: Association analysis of two candidate polymorphisms in the tumour necrosis factor- α gene with osteoarthritis in a Chinese population. *Int Orthop* 37: 2061-2063, 2013.
6. Lin SZ: Era of stem cell therapy for regenerative medicine and cancers: an introduction for the special issue of Pan Pacific Symposium on Stem Cells and Cancer Research. *Cell Transplant* 24: 311-312, 2015.
7. Boakye PA, Brierley SM, Pasilis SP and Balemba OB: *Garcinia buchananii* bark extract is an effective anti-diarrheal remedy for lactose-induced diarrhea. *J Ethnopharmacol* 142: 539-547, 2012.
8. Castro J, Balemba OB, Blackshaw LA and Brierley SM: *Garcinia buchananii* bark extract inhibits nociceptors, with greater efficacy during inflammation. *Gastroenterology* 140 (Suppl 1): S866, 2011.
9. Kisangau DP, Lyaruu HV, Hosea KM and Joseph CC: Use of traditional medicines in the management of HIV/AIDS opportunistic infections in Tanzania: a case in the Bukoba rural district. *J Ethnobiol Ethnomed* 3: 29-37, 2007.
10. Ye ZN, Yu MY, Kong LM, Wang WH, Yang YF, Liu JQ, Qiu MH and Li Y: Biflavone Ginkgetin, a novel Wnt inhibitor, suppresses the growth of medulloblastoma. *Nat Prod Bioprospect* 5: 91-97, 2015.
11. Pang X, Yi T, Yi Z, Cho SG, Qu W, Pinkaew D, Fujise K and Liu M: Morelloflavone, a biflavonoid, inhibits tumor angiogenesis by targeting rho GTPases and extracellular signal-regulated kinase signaling pathways. *Cancer Res* 69: 518-525, 2009.
12. Vandamme M, Robert E, Pesnel S, Barbosa E, Dozias S, Sobilo J, Lerondel S, Le Pape A and Pouvesle JM: Antitumor effect of plasma treatment on U87 glioma xenografts: preliminary results. *Plasma Process Polym* 7: 264-273, 2010.
13. Li XQ, Ouyang ZG, Zhang SH, Liu H, Shang Y, Li Y and Zhen YS: Synergistic inhibition of angiogenesis and glioma cell-induced angiogenesis by the combination of temozolomide and enediyne antibiotic lidamycin. *Cancer Biol Ther* 15: 398-408, 2014.
14. Stark TD, Losch S, Frank O, Balemba OB and Hofmann T: Purification procedure for (2R,3S,2'R,3'R)-manniflavanone and its minor (2R,3S,2'S,3'S)-isomer from *Garcinia buchananii* stem bark extract. *Eur Food Res Technol* 240: 1075-1080, 2015.
15. Xu G, Kan WL, Zhou Y, Song JZ, Han QB, Qiao CF, Cho CH, Rudd JA, Lin G and Xu HX: Cytotoxic acylphloroglucinol derivatives from the twigs of *Garcinia cowa*. *J Nat Prod* 73: 104-108, 2010.
16. Bing L, Gao-Sheng H, Ling-Ling H, He-Yu L and Jing-Ming J: Extraction and determination of biflavone components in ginkgo leaves. *Chinese Herbal Med* 17: 2014, 2014.

17. Xueping J and Keli C: Research on biflavone compounds in *Selaginella moellendorfii* Hieron. Chin J Pharmaceuticals 1: 2009, 2009.
18. Lianzhu Z: Treatment and possible effect of total flavonoids of *Selaginella tamariscina* and amentoflavone in cognitive disorder model (unpublished PhD dissertation). Jilin University, 2013.
19. Lipeng L: Experimental research on the treatment of C6brain glioma by nimustine combined with cisplatin (unpublished Master dissertation). Tianjin Medical University, 2008.
20. Flavahan WA, Drier Y, Liao BB, Gillespie SM, Venteicher AS, Stemmer-Rachamimov AO, Suvà ML and Bernstein BE: Insulator dysfunction and oncogene activation in IDH mutant gliomas. Nature 529: 110-114, 2016.
21. Bao S, Wu Q, McLendon RE, Hao Y, Shi Q, Hjelmeland AB, Dewhirst MW, Bigner DD and Rich JN: Glioma stem cells promote radioresistance by preferential activation of the DNA damage response. Nature 444: 756-760, 2006.