

The associated pyrazolopyrimidines PP1 and PP2 inhibit protein tyrosine kinase 6 activity and suppress breast cancer cell proliferation

HYUN JAE SHIM¹, HAN IE KIM^{1,2} and SEUNG-TAEK LEE¹

¹Department of Biochemistry, College of Life Science and Biotechnology, Yonsei University, Seoul 03722, Republic of Korea

Received August 12, 2015; Accepted November 17, 2016

DOI: 10.3892/ol.2017.5564

Abstract. Protein tyrosine kinase (PTK)6, also known as breast tumor kinase, is a non-receptor tyrosine kinase. It is closely associated with, but evolutionarily distinct from, the Src family members. PTK6 has a role in proliferation, migration and invasion in various cancers, and therefore has been suggested as a potentially valuable therapeutic target. In an attempt to develop PTK6 inhibitors, chemicals known to inhibit various kinases were screened for their ability to inhibit PTK6. Pyrazolopyrimidine (PP)1, PP2 and a lymphocyte-specific protein tyrosine kinase inhibitor strongly inhibited the catalytic activity of PTK6 *in vitro*. These chemicals suppressed the phosphorylation of PTK6 substrate proteins, including signal transducer and activator of transcription 3, in human embryonic kidney (HEK) 293 cells expressing hyperactive PTK6. They also expressed selectivity towards PTK6 over other PTK members in HEK 293 cells. PP1 and PP2 specifically inhibited the PTK6-dependent proliferation of human breast carcinoma T-47D cells. PP1 and PP2 were more selective for PTK6 than for Src family kinases, and may be useful for the treatment of PTK6-positive malignant diseases such as breast cancer.

Introduction

Protein tyrosine kinase (PTK)6, also known as breast tumor kinase, is a non-receptor type kinase that consists of an Src homology (SH)3 domain, an SH2 domain and a catalytic domain of tyrosine kinase (1,2). PTK6 is overexpressed in >60% of breast carcinoma tissue samples and in the majority of

breast cancer cell lines (3). PTK6 expression is also increased in colon, head and neck, ovary, prostate, lung, bladder, bile duct, pancreas and gastric cancers, and in T- and B-cell lymphomas (4,5).

Expression of PTK6 enhances the proliferation of mammary epithelial and breast cancer cells (6). PTK6 also promotes cell migration and invasion (7). Sublocalization of PTK6 at the plasma membrane is important for its oncogenic potential (8). Activated PTK6 consistently accumulates at the plasma membrane in breast cancer cell lines and tissues (9). Although PTK6 was detected in the nucleus and cytoplasm of normal mammary gland epithelial cells, Tyr342 in the PTK6 activation loop was not phosphorylated, and thus, PTK6 was not active (9).

PTK6 promotes tumorigenicity by enhancing signaling pathways of receptor tyrosine kinases and is particularly well known for sensitizing epidermal growth factor receptor (EGFR) family members (10). Various downstream substrates and interacting proteins, including signal transducing adaptor protein-2, paxillin, Akt, p130 Crk-associated substrate, p190Rho GTPase-activating protein-A and ArfGAP with RhoGAP domain, ankyrin repeat and PH domain 1, contribute to the oncogenic roles of PTK6 (11,12). Similar to other PTKs, mutations of PTK6 identified in different cancer types increase its kinase activity (13).

In view of its oncogenic activity and its presence in various carcinomas such as breast cancer, PTK6 is a potentially valuable therapeutic target for decelerating or arresting tumor growth (14). (E)-5-(Benzylideneamino)-1H-benzo[d]imidazol-2(3H)-one derivatives were previously developed as novel PTK6 inhibitors that exhibited little cytotoxicity, excellent inhibition *in vitro* and at the cellular level, and selectivity for PTK6 (15). Imidazo[1,2-a]pyrazin-8-amines and 4-anilino α -carboline were also identified as PTK6-selective inhibitors that block its catalytic activity (16,17). In the present study, pyrazolopyrimidine (PP)1 [4-amino-5-(4-methylphenyl)-7-(t-butyl) pyrazolo[3,4-d]pyrimidine], PP2 [(4-amino-5-(4-chlorophenyl)-7-(t-butyl) pyrazolo[3,4-d]pyrimidine] and a lymphocyte-specific protein tyrosine kinase (Lck) inhibitor [4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo [3,2-d]pyrimidin-7-yl-cyclopentane] were screened as potent PTK6 inhibitors among the evaluated kinase inhibitors. The selectivity of these compounds for PTK6 and for other PTK family members was analyzed in HEK 293 cells, and

Correspondence to: Professor Seung-Taek Lee, Department of Biochemistry, College of Life Science and Biotechnology, Yonsei University, 50 Yonsei-Ro, Seodaemun-Gu, Seoul 03722, Republic of Korea
E-mail: stlee@yonsei.ac.kr

Present address: ²Department of Life Science and Biotechnology, Shingyeong University, Hwaseong, Gyeonggi-do 18274, Republic of Korea

Key words: PP1, PP2, PTK6 inhibitor, breast cancer cell

it was then examined whether these compounds inhibited PTK6-dependent signaling processes and the proliferation of breast carcinoma T-47D cells.

Materials and methods

Chemicals. PP1, PP2 and the aforementioned Lck inhibitor were purchased from Calbiochem (EMD Millipore, Billerica, MA, USA). Genistein was purchased from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA).

Cell culture. Human embryonic kidney (HEK) 293 cells and human breast cancer T-47D cells (both American Type Culture Collection, Manassas, VA, USA) were maintained in Dulbecco's modified Eagle's medium (DMEM; Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA) containing 10% fetal bovine serum (FBS; HyClone; GE Healthcare Life Sciences, Logan, UT, USA) at 37°C in a CO₂ incubator with a humidified atmosphere of 5% CO₂ and 95% air.

ELISA-based *in vitro* kinase assay for PTK6. ELISA plates (96-well; Greiner Bio-One GmbH, Frickenhausen, Germany) were incubated with 100 μ l of 0.1 mg/ml Poly (Glu, Tyr) (Glu:Tyr, 4:1; Sigma-Aldrich; Merck Millipore, Darmstadt, Germany) in PBS for 16 h at 37°C, and then washed three times with PBS. The Poly (Glu, Tyr)-coated wells were blocked with 1% bovine serum albumin in PBS for 1 h at 37°C, washed three times with PBS, and then incubated for 30 min at room temperature with 10 nM glutathione S-transferase-fused PTK6 catalytic domain (18) in 20 μ l of kinase reaction buffer [20 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, 1 mM MnCl₂ and 50 μ M Na₃VO₄] in the presence of the chemical of interest (Table I) containing a final concentration of 1% dimethyl sulfoxide. The phosphorylation of tyrosine residues was initiated by addition of 300 μ M adenosine triphosphate (ATP) to the reaction mixtures. The wells were washed three times with PBS after incubation for 20 min at room temperature. For the quantification of phosphorylated tyrosines, the wells were incubated with anti-phospho-tyrosine (4G10; 1:1,000; EMD Millipore) and horseradish peroxidase-conjugated anti-mouse immunoglobulin G (K0211589; 1:10,000; Koma Biotech, Seoul, Korea) antibodies for 1 h each at room temperature. The optical density was measured with a 3,3',5,5'-tetramethylbenzidine solution (Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol.

Western blot analysis. HEK 293 cells expressing hyperactive PTK6 (Flag-PTK6-3PA/Y447F) (19), Src, Fyn, Lck, bone marrow tyrosine kinase gene on chromosome X (Bmx) or EGFR were treated with the indicated concentrations of compounds for 2 days. Western blot analysis and immunoprecipitation were performed as previously described (15). Immunoreactive proteins were visualized using anti-phospho-tyrosine (4G10; 1:1,000), anti-PTK6 (sc-1188; 1:2,000; Santa Cruz Biotechnology, Inc.), anti-phospho-signal transducer and activator of transcription (STAT)3 (sc-8059; 1:2,000; Santa Cruz Biotechnology, Inc.), anti-STAT3 (sc-7179; 1:2,000; Santa Cruz Biotechnology, Inc.) and anti-GAPDH (AbC-2003; 1:2,000; AbClon, Inc., Seoul, Korea) primary antibodies overnight at 4°C, followed by incubation with a horseradish

peroxidase-conjugated secondary antibody (K0211589 or K0211708; 1:10,000; Koma Biotech, Seoul, Republic of Korea) for 1 h at room temperature and an enhanced chemiluminescence detection kit (EMD Millipore). For the quantification of the phosphorylation levels in the cell lysates, chemiluminescence was detected using a LAS-3000 imaging system (Fujifilm, Tokyo, Japan) and analyzed using Multi Gauge version 2.2 software (Fujifilm). The half maximal inhibitory concentration (IC₅₀) at the cellular level was determined by quantifying the phosphorylation levels in the HEK 293 cell system. The reference level was the phosphorylation level of the chemical-free control, which was set at 100%.

MTT assay. Subconfluent empty vector-transfected and PTK6-knockdown T-47D cells were incubated for 4 days in DMEM-10% FBS containing various concentrations of the chemicals. Viable cells were measured using MTT assay, as previously described (15). The viability of chemical-free, vector-transfected T-47D cells was set at 100%.

Statistical analysis. All data were expressed as the mean \pm standard deviation of three independent experiments. Statistical analysis was performed using Microsoft Excel (version, 2007; Microsoft Corporation, Redmond, WA, USA). The significant differences between the groups were assessed using a Student's t-test. $P > 0.05$ was considered to indicate a statistically significant difference.

Results

PP1, PP2 and Lck inhibitor inhibit the catalytic activity of PTK6 *in vitro*. Protein kinase inhibitors were analyzed for the inhibition of the PTK6 catalytic activity using an ELISA-based *in vitro* kinase assay system for PTK6. Among the tested kinase inhibitors, PP1, PP2 and the Lck inhibitor exhibited strong inhibition of PTK6 (Table I). The IC₅₀ values for PP1, PP2 and the Lck inhibitor were 230.0, 50.0 and 60.0 nM, respectively (Table II).




PP1, PP2 and Lck inhibitor are highly selective for PTK6 at the cellular level. PP1, PP2 and the aforementioned Lck inhibitor were developed as Src family kinase (SFK) inhibitors (20,21). The present study analyzed the selectivity of each inhibitor for the inhibition of several PTKs, including PTK6, SFK members (Src, Fyn and Lck), a non-receptor type PTK (Bmx) and a receptor type PTK (EGFR). HEK 293 cells expressing hyperactive PTK6, Src, Fyn, Lck, Bmx or EGFR were incubated in the presence of various concentrations of these inhibitors. Inhibition of each PTK activity at the cellular level was assessed for each inhibitor by measuring the decrease in tyrosine phosphorylation levels of cellular proteins via western blot analysis using an anti-phospho-tyrosine antibody. As expected, PP1, PP2 and the Lck inhibitor exhibited strong inhibition of SFK members. The IC₅₀ value of PP1 to Lck was 1.76 μ M; the IC₅₀ value of PP2 to Lck was 4.36 μ M; and the IC₅₀ values of the Lck inhibitor to Lck, Fyn and Src were 0.37, 1.22 and 3.46 μ M, respectively (Table III). Unexpectedly, PP1, PP2 and the Lck inhibitor inhibited PTK6 to a greater degree than SFK members. The IC₅₀ values of PP1, PP2 and the Lck inhibitor to PTK6 were 2.5, 13.0 and 53.0 nM, respectively (Table III and Fig. 1, top panel).

Table I. PTK6 inhibitory activities of protein kinase inhibitors screened from a targeted kinase inhibitor.

Chemical name	PubChem compound identifier	Molecular weight (Da)	Known target	<i>In vitro</i> IC ₅₀ (nM) ^a
Genistein	5280961	270.2	PTKs	>10,000
PP1	1400	281.4	Src, Fyn	<300
PP2	4878	301.8	Lck, Fyn, Hck	<300
Lck inhibitor	6603792	370.5	Lck, Src	<300
LFM-A13	54676905	360.0	Btk	>10,000
JAK inhibitor I	5494425	309.3	JAK1-3	>10,000
Syk inhibitor	6419747	353.4	Syk	>3,000
Emodin	3220	270.2	Lck, ErbB2	>10,000
Erbstatin analog	5353609	194.2	EGFR, Abl	>300
Lavendustin C	3896	275.3	EGFR, Src, CaMKII	>300
Daphnetin	5280569	178.1	EGFR, PKA, PKC	>3,000
PD 157432	10085225	283.3	EGFR, ErbB2	>10,000
Flt-3 inhibitor	1048845	360.4	Flt-3	>10,000
cFMS inhibitor	11617559	366.4	FMS	>10,000
GTP-14564	3385203	234.3	FMS, Kit, Flt-3, PDGFRB	>10,000
PDGFR inhibitor I	5330535	276.3	PDGFRB	>10,000
Oxindole I	5908088	210.2	VEGFR1, PDGFRB	>10,000
VEGFR2 inhibitor II	23301538	343.2	VEGFR2, PDGFRB	>3,000
TrkA inhibitor	5413390	315.4	TrkA	>10,000
Picropodophyllin	72435	414.4	IGF1R	>10,000
JNK inhibitor II	8515	220.2	JNK1-3	>10,000

^aThe IC₅₀ value was the concentration of each inhibitor required for 50% inhibition of the catalytic activity of purified PTK6 in the absence of inhibitors (dimethyl sulfoxide). Data were obtained from three independent experiments, each performed in duplicate. IC₅₀, half maximal inhibitory concentration; PTK, protein tyrosine kinase; PP, pyrazolopyrimidine; Lck, lymphocyte-specific protein tyrosine kinase; LFM-A13, leflunomide metabolite-analog A13; STAT, signal transducer and activator of transcription; JAK, Janus-associated kinase; Syk, spleen tyrosine kinase; PD, Parke-Davis; Flt, Fms-like tyrosine kinase; cFMS, cellular feline McDonough sarcoma; GTP, guanosine triphosphate; PDGFRB, platelet-derived growth factor receptor beta; VEGFR, vascular endothelial growth factor receptor; TrkA, tropomyosin receptor kinase A; JNK, c-Jun N-terminal kinase; Hck, hematopoietic cell kinase; Btk, Bruton's tyrosine kinase; EGFR, epidermal growth factor receptor; Abl, Abelson murine leukemia viral oncogene homolog 1; CaMKII, Ca²⁺/calmodulin-dependent protein kinase II; PK, protein kinase; IGF1R, insulin-like growth factor 1 receptor.

Table II. Inhibition of the catalytic activity of purified PTK6 by selected kinase inhibitors *in vitro*.

Chemical name	Structure	Molecular weight (Da)	<i>In vitro</i> IC ₅₀ (nM) ^a
PP1		301.8	230±20
PP2		317.4	50±10
Lck inhibitor		370.5	60±20

^aThe IC₅₀ value was the concentration of each inhibitor required for 50% inhibition of the catalytic activity of purified PTK6 in the absence of inhibitors (dimethyl sulfoxide). Data are expressed as the mean ± standard deviation values of three independent experiments, each performed in duplicate. IC₅₀, half maximal inhibitory concentration; PTK, protein tyrosine kinase; PP, pyrazolopyrimidine; Lck, lymphocyte-specific protein tyrosine kinase.

Table III. Selectivity of genistein, PP1, PP2 and a Lck inhibitor for various PTKs in the HEK 293 cell system.

PTKs	IC ₅₀ at the cellular level (nM) ^a			
	Genistein	PP1	PP2	Lck inhibitor
PTK6	43,180±1,340	2.5±0.3	13±2	53±7
Src	>100,000	70,300±7,320	53,060±8,230	3,460±850
Fyn	70,640±8,520	13,630±3,160	35,980±1,030	1,220±50
Lck	>100,000	1,760±250	4,360±730	370±40
Bmx	69,550±7,460	2,390±430	16,460±2,780	15,980±3,680
EGFR	24,830±1,590	21,840±1,140	56,800±3,920	>10,000

^aThe IC₅₀ values were determined by quantifying the phosphorylation levels in HEK 293 cells expressing hyperactive PTK6, Src, Fyn, Lck, Bmx or EGFR. Data are expressed as the mean ± standard deviation values of three independent experiments. HEK, human embryonic kidney; IC₅₀, half maximal inhibitory concentration; PTK, protein tyrosine kinase; PP, pyrazolopyrimidine; Lck, lymphocyte-specific protein tyrosine kinase; EGFR, epidermal growth factor receptor; Bmx, bone marrow tyrosine kinase gene on chromosome X.




Figure 1. Effects of selected chemicals on the phosphorylation of cellular proteins and a specific PTK6 substrate, STAT3, in HEK 293 cells. HEK 293 cells stably expressing hyperactive PTK6 were incubated with the indicated concentrations of chemicals for 48 h. Cell lysates were analyzed by western blotting using anti-phospho-tyrosine (4G10), anti-phospho-STAT3, anti-STAT3, anti-PTK6 and anti-GAPDH antibodies. The numbers on the left of the top panel indicate apparent molecular weights in kDa. HEK, human embryonic kidney; PP, pyrazolopyrimidine; Lck, lymphocyte-specific protein tyrosine kinase; PTK, protein tyrosine kinase; STAT, signal transducer and activator of transcription; p-, phosphorylated.

This result suggested that PP1, PP2 and the Lck inhibitor were highly selective for PTK6 compared with other PTKs, including the SFK family members, at the cellular level.

PP1, PP2 and Lck inhibitor inhibit the phosphorylation of PTK6 substrate proteins in HEK 293 cells. To analyze whether

PP1, PP2 and the Lck inhibitor block the PTK6-mediated signaling pathway, HEK 293 cells expressing hyperactive PTK6 were treated with these chemicals at concentrations i) lower than, ii) approximately the same as, and iii) higher than their IC₅₀ values. Phosphorylation of STAT3, which is a known specific substrate of PTK6 (22), was inhibited at




Figure 2. Effects of PTK6 inhibitors on the proliferation of breast cancer T-47D cells. Empty vector-transfected (Vector) and PTK6-knockdown (PTK6-KD) T-47D cells were generated by stable transfection of pLKO.1 and pLKO.1-small hairpin RNA PTK6, respectively (12). (A) Analysis of PTK6 expression levels by western blotting using an anti-PTK6 antibody. (B) Proliferation of Vector and PTK6-KD T-47D cells. (C) Effect of genistein on Vector and PTK6-KD T-47D cells. (D) Effect of PP1 on Vector and PTK6-KD T-47D cells. (E) Effect of PP2 on Vector and PTK6-KD T-47D cells. (F) Effect of a Lck inhibitor on Vector and PTK6-KD T-47D cells. Vector and PTK6-KD cells were incubated in complete medium containing the indicated concentrations of chemicals for 4 days and then subjected to MTT assay. All data are expressed as the mean \pm standard deviation values of three independent experiments, each performed in duplicate. **P<0.01, ***P<0.001 vs. Vector T-47D cells without inhibitors. ††P<0.01, †††P<0.001 vs. PTK6-KD T-47D cells without inhibitors. PTK, protein tyrosine kinase; KD, knockdown; PP, pyrazolopyrimidine; Lck, lymphocyte-specific protein tyrosine kinase.

concentrations equal or greater than the IC_{50} value of each chemical (Fig. 1, mid panel).

PP1 and PP2 inhibit the PTK6-mediated proliferation of T-47D cells. PTK6 is often expressed in breast cancer cell lines (3). Knockdown of PTK6 decreases the proliferation of breast cancer cells (14). Consistent with this observation, the silencing of PTK6 in breast carcinoma T-47D cells using a small hairpin RNA vector suppressed ~24% of cell proliferation, compared with vector transfection (Fig. 2A and B). PP1, PP2, the Lck inhibitor and genistein (used as a control) were applied to T-47D cells at concentrations of 0.33-, 1- and 3-fold their IC_{50} values. PP1 and PP2 inhibited the proliferation of vector-transfected T-47D cells in a dose-dependent manner, but did not affect the proliferation of PTK6-knockdown T-47D cells at values of ≤ 3 -fold their IC_{50} values (Fig. 2D and E). However, the Lck inhibitor and genistein suppressed the

proliferation of both vector-transfected and PTK6-knockdown T-47D cells (Fig. 2C and F). These results suggest that the Lck inhibitor is not specific for PTK6, which is similar to the general PTK inhibitor genistein (23).

Discussion

In our study, PP1, PP2 and a Lck inhibitor were screened as potential inhibitors for PTK6 using an *in vitro* kinase assay. These chemicals were initially developed as ATP-competitive inhibitors of SFKs (21,22). PP1 inhibits Lck, Fyn and Src with IC_{50} values of 5, 6 and 170 nM, respectively, as assessed with an *in vitro* kinase assay (20). PP2 inhibits Lck and Fyn with IC_{50} values of 5 and 6 nM, respectively (20). The Lck inhibitor was developed as a derivative of PP1 and PP2, and inhibits Lck and Src with IC_{50} values of <1 and 70 nM, respectively (21). These chemicals were widely used to investigate

the physiological roles for SFKs, but were not used in human clinical trials. PTK6 is evolutionarily distinct from, but still closely associated with, SFKs (2). Thus, it is expected that PP1, PP2 and the Lck inhibitor inhibit PTK6.

Although PP1 and PP2 are more selective for SFKs than the previous generation of PTK inhibitors (including herbimycin A and genistein), they can inhibit off-target kinases (including C-terminal Src kinase, ephrin type-A receptor 2, platelet-derived growth factor receptor, fibroblast growth factor receptor 1, p21 activated kinase, receptor-interacting protein 2, p38 and casein kinase 1 δ) with sufficient potency (24-27). When the selectivity of these chemicals for various PTKs was analyzed in HEK 293 cells expressing one of the PTKs, they displayed high selectivity for PTK6 over various SFK members, including Src, Fyn, Lck, and other PTK family members such as Bmx and EGFR. In particular, PP1 and PP2 inhibited PTK6 activity at IC₅₀ values of 2.5 and 13.0 nM, respectively. Although they also inhibited the catalytic activities of other PTKs, including SFKs, their IC₅₀ values were mostly at micro-molar concentrations. Incubation of T-47D cells with 0.8-7.5 nM PP1 or 4.0-39.0 nM PP2 reduced the PTK6-dependent proliferation of T-47D cells without a decrease in the proliferation of PTK6-knockdown cells. The Lck inhibitor also exhibited inhibitory selectivity for PTK6, but reduced the proliferation of T-47D cells in a PTK6-independent manner. These results demonstrate that PP1 and PP2 reduce the PTK6-mediated signaling pathways and cell proliferation in PTK6-positive cells. Thus, it can be suggested that PP1 and PP2 could be applied as therapeutic agents in PTK6-positive malignant diseases.

Resistance to chemotherapy and molecularly targeted therapies is a major problem confronting current cancer research. The results of a recent study indicated that PTK6 confers resistance of breast cancer SUM102 cells to cetuximab, an EGFR-blocking antibody that is approved for the treatment of several types of human solid tumors (28). Knockdown of PTK6 sensitized the cells to cetuximab by inducing apoptosis. Assuming that PTK6 catalytic activity is essential for drug resistance (28), PTK6 inhibitors such as PP1 and PP2 may be useful for the treatment of chemotherapy-resistant cancer cells.

Acknowledgements

The present study was supported by a grant from the National Research Foundation of Korea funded by the Ministry of Science, ICT & Future Planning (grant no. 2014M3C9A2064536).

References

- Mitchell PJ, Barker KT, Martindale JE, Kamalati T, Lowe PN, Page MJ, Gusterson BA and Crompton MR: Cloning and characterisation of cDNAs encoding a novel non-receptor tyrosine kinase, brk, expressed in human breast tumours. *Oncogene* 9: 2383-2390, 1994.
- Lee H, Kim M, Lee KH, Kang KN and Lee ST: Exon-intron structure of the human PTK6 gene demonstrates that PTK6 constitutes a distinct family of non-receptor tyrosine kinase. *Mol Cells* 8: 401-407, 1998.
- Barker KT, Jackson LE and Crompton MR: BRK tyrosine kinase expression in a high proportion of human breast carcinomas. *Oncogene* 15: 799-805, 1997.
- Brauer PM and Tyner AL: Building a better understanding of the intracellular tyrosine kinase PTK6-BRK by BRK. *Biochim Biophys Acta* 1806: 66-73, 2010.
- Mizuguchi Y, Specht S, Isse K, Sasatomi E, Lunz JG III, Takizawa T and Demetris AJ: Breast tumor kinase/protein tyrosine kinase 6 (Brk/PTK6) activity in normal and neoplastic biliary epithelia. *J Hepatol* 63: 399-407, 2015.
- Kamalati T, Jolin HE, Mitchell PJ, Barker KT, Jackson LE, Dean CJ, Page MJ, Gusterson BA and Crompton MR: Brk, a breast tumor-derived non-receptor protein-tyrosine kinase, sensitizes mammary epithelial cells to epidermal growth factor. *J Biol Chem* 271: 30956-30963, 1996.
- Chen HY, Shen CH, Tsai YT, Lin FC, Huang YP and Chen RH: Brk activates rac1 and promotes cell migration and invasion by phosphorylating paxillin. *Mol Cell Biol* 24: 10558-10572, 2004.
- Je Kim H and Lee ST: Oncogenic functions of PTK6 are enhanced by its targeting to plasma membrane but abolished by its targeting to nucleus. *J Biochem* 146: 133-139, 2009.
- Peng M, Emmadi R, Wang Z, Wiley EL, Gann PH, Khan SA, Banerji N, McDonald W, Asztalos S, Pham TN, *et al*: PTK6/BRK is expressed in the normal mammary gland and activated at the plasma membrane in breast tumors. *Oncotarget* 5: 6038-6048, 2014.
- Ostrander JH, Daniel AR and Lange CA: Brk/PTK6 signaling in normal and cancer cell models. *Curr Opin Pharmacol* 10: 662-669, 2010.
- Goel RK and Lukong KE: Tracing the footprints of the breast cancer oncogene BRK-Past till present. *Biochim Biophys Acta* 1856: 39-54, 2015.
- Kang SA, Lee ES, Yoon HY, Randazzo PA and Lee ST: PTK6 inhibits down-regulation of EGF receptor through phosphorylation of ARAP1. *J Biol Chem* 285: 26013-26021, 2010.
- Tsui T and Miller WT: Cancer-associated mutations in breast tumor kinase/PTK6 differentially affect enzyme activity and substrate recognition. *Biochemistry* 54: 3173-3182, 2015.
- Harvey AJ and Crompton MR: The Brk protein tyrosine kinase as a therapeutic target in cancer: Opportunities and challenges. *Anticancer Drugs* 15: 107-111, 2004.
- Shim HJ, Yang HR, Kim Hie, Kang SA, No KT, Jung YH and Lee ST: Discovery of (E)-5-(benzylideneamino)-1H-benzo[d]imidazol-2(3H)-one derivatives as inhibitors for PTK6. *Bioorg Med Chem Lett* 24: 4659-4663, 2014.
- Zeng H, Belanger DB, Curran PJ, Shipps GW Jr, Miao H, Bracken JB, Arshad Siddiqui M, Malkowski M and Wang Y: Discovery of novel imidazo[1,2-a]pyrazin-8-amines as Brk/PTK6 inhibitors. *Bioorg Med Chem Lett* 21: 5870-5875, 2011.
- Mahmoud KA, Krug M, Wersig T, Slynko I, Schächtele C, Totzke F, Sippl W and Hilgeroth A: Discovery of 4-anilino α -carboline as novel Brk inhibitors. *Bioorg Med Chem Lett* 24: 1948-1951, 2014.
- Kim Hie and Lee ST: An intramolecular interaction between SH2-kinase linker and kinase domain is essential for the catalytic activity of protein-tyrosine kinase-6. *J Biol Chem* 280: 28973-28980, 2005.
- Kang SA, Cho HS, Yoon JB, Chung IK and Lee ST: Hsp90 rescues PTK6 from proteasomal degradation in breast cancer cells. *Biochem J* 447: 313-320, 2012.
- Hanke JH, Gardner JP, Dow RL, Changelian PS, Brissette WH, Weringer EJ, Pollok BA and Connelly PA: Discovery of a novel, potent and Src family-selective tyrosine kinase inhibitor. Study of Lck- and FynT-dependent T cell activation. *J Biol Chem* 271: 695-701, 1996.
- Arnold LD, Calderwood DJ, Dixon RW, Johnston ND, Kamens JS, Munschauer R, Rafferty P and Ratnofsky SE: Pyrrolo[2,3-d]pyrimidines containing an extended 5-substituent as potent and selective inhibitors of lck I. *Bioorg Med Chem Lett* 10: 2167-2170, 2000.
- Liu L, Gao Y, Qiu H, Miller WT, Poli V and Reich NC: Identification of STAT3 as a specific substrate of breast tumor kinase. *Oncogene* 25: 4904-4912, 2006.
- Smal C, Cardoen S, Bertrand L, Delacauw A, Ferrant A, Van den Berghe G, Van Den Neste E and Bontemps F: Activation of deoxycytidine kinase by protein kinase inhibitors and okadaic acid in leukemic cells. *Biochem Pharmacol* 68: 95-103, 2004.
- Summy JM, Trevino JG, Lesslie DP, Baker CH, Shakespeare WC, Wang Y, Sundaramoorthi R, Metcalf CA III, Keats JA, Sawyer TK and Gallick GE: AP23846, a novel and highly potent Src family kinase inhibitor, reduces vascular endothelial growth factor and interleukin-8 expression in human solid tumor cell lines and abrogates downstream angiogenic processes. *Mol Cancer Ther* 4: 1900-1911, 2005.

25. Ma QL, Yang F, Frautschy SA and Cole GM: PAK in Alzheimer disease, Huntington disease and X-linked mental retardation. *Cell Logist* 2: 117-125, 2012.
26. Bain J, McLauchlan H, Elliott M and Cohen P: The specificities of protein kinase inhibitors: An update. *Biochem J* 371: 199-204, 2003.
27. Bain J, Plater L, Elliott M, Shpiro N, Hastie CJ, McLauchlan H, Klevernic I, Arthur JS, Alessi DR and Cohen P: The selectivity of protein kinase inhibitors: A further update. *Biochem J* 408: 297-315, 2007.
28. Li X, Lu Y, Liang K, Hsu JM, Albarracin C, Mills GB, Hung MC and Fan Z: Brk/PTK6 sustains activated EGFR signaling through inhibiting EGFR degradation and transactivating EGFR. *Oncogene* 31: 4372-4383, 2012.