

Expression of miR-146a in patients with ovarian cancer and its clinical significance

MIŁOSZ WILCZYŃSKI¹, EWELINA ŻYTKO², BOŻENA SZYMAŃSKA³, MONIKA DZIENIECKA⁴,
MAREK NOWAK², JUSTYNA DANIELSKA⁵, GRZEGORZ STACHOWIAK² and JACEK R. WILCZYŃSKI²

Departments of ¹Operative Gynecology, Endoscopy and Gynecologic Oncology, and ²Gynecology and Oncological Gynecology, Polish Mother's Memorial Hospital Research Institute, 93-338 Lodz;

³Central Scientific Laboratory CoreLab, Medical University of Lodz, 92-215 Lodz;

⁴Department of Pathology, Polish Mother's Memorial Hospital Research Institute, 93-388 Lodz;

⁵Radiotherapy Department, Medical University of Lodz, 93-509 Lodz, Poland

Received October 14, 2016; Accepted April 7, 2017

DOI: 10.3892/ol.2017.6477

Abstract. The aim of the present retrospective study was to compare microRNA (miR)-146a expression levels in primary tumors and omental metastases of 48 patients, who had undergone surgery for advanced ovarian serous cancer. Possible correlations between miR-146a expression level and clinicopathological features were investigated, including chemosensitivity and survival. miR-146a was evaluated in formalin-fixed, paraffin-embedded samples. miR-146a expression level in primary tumors was demonstrated to be increased in comparison with normal ovary tissues ($P=0.02$) and metastases ($P=0.01$). A negative correlation was demonstrated between miR-146a expression in primary tumors and serum levels of cancer antigen 125 ($R=-0.37$; $P=0.03$) and Risk of Malignancy Algorithm index ($R=-0.79$; $P=0.0007$). Overall survival positively correlated with miR-146a expression in primary tumor tissue samples ($R=0.38$; $P=0.01$). Probability of survival was decreased in patients with low miR-146a expression levels in primary tumor tissues (hazard ratio=0.21; $P=0.003$). Lower levels of miR-146a in primary tumor tissue samples were correlated with a shorter progression-free

survival ($P=0.04$) and platinum-resistance of metastases ($P=0.006$). In conclusion, miR-146a may be a prognostic marker for serous ovarian cancer.

Introduction

Ovarian cancer is one of the most fatal types of female neoplasms. Despite the availability of extensive management of the disease, the number of patients who survive ≥ 5 years following diagnosis remains low (1). Ultra-radical surgery, platinum- and taxane-based chemotherapy and immunotherapy are all improvements. However, the impact of these treatments on overall survival (OS) remains unsatisfactory (2). One of the main concerns is the low efficacy of chemotherapy due to the primary tumor platinum-refractoriness or acquired chemoresistance during the course of adjuvant treatment (3). It is not possible to predict which patients, subject to standard chemotherapeutic regimen, would respond to the therapy. Inefficient response to the treatment leads to tumor progression or recurrence. Therefore, identification of markers for potentially chemoinensitive tumors may aid in modifying and individualizing treatments prior to the recognition of chemorefractoriness or chemoresistance. MicroRNAs (miRNA/miR), small non-coding RNAs involved in post-transcriptional gene regulation, may be candidates for potential markers (4).

miR-146a has been described as a modulator of differentiation and function of innate and adaptive immunity. In human T cells, miR-146a is expressed abundantly in memory T cells, and its expression is critical for function of T regulatory cells (Tregs). miR-146a was also demonstrated to upregulate the macrophage inflammatory response (5,6). The molecular function of miR-146a in the immune response involves negative regulation of the signal transduction pathway, which leads to activation of nuclear factor- κ B (NF κ B), disruption of downstream T lymphocyte receptor-4 signaling pathway, and modulation of chemokine interleukin-8, RANTES (regulated on activation, normal T-cell expressed and secreted) and CXC chemokine receptor type 4 expression (6). The association between miR-146a and the immune response may have a potential impact in solid tumors.

Correspondence to: Dr Miłosz Wilczyński, Department of Operative Gynecology, Endoscopy and Gynecologic Oncology, Polish Mother's Memorial Hospital Research Institute, 281/289 Rzgowska Street, 93-338 Lodz, Poland
E-mail: jrwil@wp.pl

Abbreviations: Tregs, T regulatory cells; NF κ B, nuclear factor- κ B; RANTES, regulated on activation, normal T-cell expressed and secreted; CXCR4-C-X-C; chemokine receptor type 4; FIGO, International Federation of Gynecology and Obstetrics; CA125, cancer antigen 125; HE4, serum human epididymis antigen-4; FFPE, formalin-fixed, paraffin-embedded; PFS, progression-free survival; OS, overall survival; AUC, area under the curve

Key words: microRNA-146a, ovarian cancer, cancer antigen 125, risk of malignancy algorithm, survival, chemoresistance

Furthermore, it was demonstrated that miR-146a is involved in the regulation of various RNAs encoding a number of proteins involved in cell differentiation, proliferation and migration (7). Consequently, aberrant disturbed miR-146a expression level was observed in numerous types of malignancies, including thyroid, breast, gastric, prostate, pancreatic and ovarian cancer (8-10). miR-146a polymorphism may increase the risk of developing various types of cancer. For example, the G/C polymorphism in the pre-miR-146a sequence was associated with a decrease or an increase of miRNA-146a expression levels, depending on the cancer type. The change in miRNA-146a expression modified the risk of papillary thyroid, hepatocellular, gastric cancer and glioma (6,8,9). The G/C polymorphism (SNP no. rs2910164) may induce the onset of breast and ovarian cancers in breast cancer 1/2 (BRCA1/2) positive cases (11). The level of miR-146a expression may depend on the type of tumor and the aggressiveness of the tumor. Increased expression levels of miR-146a were reported in papillary thyroid and cervical cancer compared with normal tissues (12,13). However, in highly metastatic breast cancer cell lines, the expression of miR-146a was downregulated and exogenous miR-146a expression impaired the invasion and migratory capacity of cancer cells. Breast cancer metastasis suppressor-1, which affects multiple steps of the metastasizing process, may partially function by upregulating miR-146a expression in breast cancer cells (14). A study by Boldin *et al* investigating miR-146a-knock-out mice confirmed human studies and demonstrated that the lack of miR-146a expression favors development of hematologic neoplasms (15). Chang *et al* (16) suggested that downregulation of miR-146a may contribute to the Myc-mediated tumorigenesis. These observations strongly suggested that miR-146a may serve a role as a tumor suppressor.

Various miRs were revealed to be either upregulated or downregulated in patients with ovarian cancer (17,18). The expression of miR-30a-3p was increased in well-differentiated tumors compared with poorly differentiated tumors (19). Detection of high levels of plasma miR-205 and low Let-7f expression levels combined with high serum cancer antigen 125 (CA125) levels improved the accuracy of ovarian cancer detection (20). Let-7f was identified as a predictive factor for ovarian cancer prognosis (20). A predictive model based on the serum expression levels of miR-200b/miR-200c was able to discriminate between normal controls and age-matched patients with high-grade serous ovarian cancer (21). A number of other miRs were studied, and the levels of these miRs were correlated with the hazard ratio for patient survival or tumor recurrence (22). *In vitro* and *in vivo* studies have suggested that the pattern of miR expression may have an impact on the chemosensitivity of ovarian tumors (23-26). Vang *et al* (27) performed a study on a small group of patients with advanced ovarian serous cancer: The study revealed dysregulation of miR-146a and miR-150 in omental metastases and suggested their possible role in increased platinum tolerance (27).

The aim of this retrospective study was to compare expression levels of miR-146a in primary tumor tissues and omental metastases from patients who underwent surgery for advanced ovarian serous cancer. The second aim of the present study

was to investigate an association between miR-146a expression levels and clinicopathological features, including chemosensitivity and survival.

Materials and methods

Patient collection. The present study was approved by the Ethics Committee of the Polish Mother's Memorial Hospital Research Institute (Lodz, Poland; grant no. 37/2014). Written informed consent was obtained from all patients prior to enrolment in the present study. A total of 48 patients with advanced ovarian cancer, who underwent cytoreductive abdominal surgery between March 2006 and December 2010, were included in the present study. The inclusion criteria were serous tumor histology and stage III/IV according to the International Federation of Gynecology and Obstetrics (FIGO) clinical staging system (28). Total hysterectomy with bilateral salpingo-oophorectomy, omentectomy and appendectomy was performed in all cases, supplemented with partial resection of infiltrated intestine or bowel, peritonectomy or splenectomy for the purposes of optimal cytoreduction. Systemic or sampling lymphadenectomy was performed only in cases when optimal cytoreduction was achieved or in the presence of bulky nodes. Adjuvant treatment with platinum-taxane regimen, six standard courses of carboplatin 5-7.5 area under the curve (AUC) and paclitaxel 175 mg/m² and modified according to the patient's general status, was introduced in all cases. Clinical information was acquired from medical records. Serum CA125 levels and Risk of Malignancy Algorithm (ROMA) index calculated based on the levels of serum CA125, serum human epididymis antigen-4 (HE4) and pre-menopausal or menopausal status were acquired prior to cytoreductive surgery. Platinum-sensitive tumors were identified when there was no relapse ≥ 6 months following completion of the chemotherapy. Resistant patients were defined as patients with primary chemo-refractory tumors (progression despite treatment with a first-line chemotherapy). Platinum-resistance was also diagnosed when relapse occurred ≤ 6 months following completion of chemotherapy. A total of 27 patients were identified to be chemosensitive and 21 patients were recognized as chemoresistant. Clinical characterizations of chemosensitive and chemoresistant patients are presented in Table I.

Sample collection. miR-146a expression was evaluated in tissues obtained from archival formalin-fixed (tissues were fixed with 10% formalin for 24-48 h at room temperature) paraffin-embedded (FFPE) serous ovarian cancer samples. Other histological types of ovarian cancer were excluded from the present study. All archival FFPE samples were re-evaluated by an experienced pathologist. Following confirmation of the cancer type, the areas of cancerous tissues were carefully selected and micro-dissected from the samples in order to avoid areas of extensive necrosis and to minimize the risk of contamination with noncancerous tissues. From each patient, two samples were obtained, with one sample from primary ovarian tumor and another sample from omental metastasis. The reference group consisted of 48 normal ovarian tissue samples that were retrieved from peri-menopausal women during hysterectomy with bilateral salpingo-oophorectomy

Table I. Clinicopathological characteristics and outcomes of patients in the platinum-sensitive and platinum-resistant groups.

Parameter	Platinum-sensitive	Platinum-resistant
Case number (n)	27	21
Mean age, years (range)	48 (24-81)	54 (48-75)
FIGO stage		
III (n)	23	20
IV (n)	4	1
Tumor grade ^a		
1 (n)	2	2
2 (n)	8	8
3 (n)	17	11
Recurrence		
No (n)	5	6
Yes (n)	22	15
Median PFS, months (range)	25 (12-67)	4 (0-11)
Median OS, months (range)	33 (13-70)	16 (2-113)
Survival		
No (n)	3	9
Yes (n)	24	12

PFS, progression-free survival; OS, overall survival; FIGO, International Federation of Gynecology and Obstetrics.^a(55).

due to benign uterine disease (uterine leiomyoma). All hysterectomy procedures were performed between January 2014 and December 2014. The mean age of patients was 47 years (range, 39-56 years). Surgery was performed in the Polish Mother's Memorial Hospital Research Institute and informed consent was obtained from all of them.

Total RNA isolation and miRNA expression analysis. Total RNA was extracted from FFPE tissues using the Roche High Pure miR Isolation kit (Roche Diagnostics GmbH, Mannheim, Germany), according to the manufacturer's instructions. In brief, the FFPE microsamples were processed in 2 ml Eppendorf tubes, deparaffinized with 100% xylene, washed in 100% ethanol and dried at 55°C for -10 min. The dried tissue was resuspended in 100 μ l Paraffin Tissue Lysis Buffer (included in the kit) and digested with proteinase K at 55°C overnight. Subsequent steps of RNA purification on columns were performed according to the manufacturer's protocol (Roche Diagnostics GmbH). Briefly, 325 μ l of binding buffer and 325 μ l of binding enhancer was added and the mixture applied on the columns, centrifuged for 30 s at 13,000 x g and washed twice with 500 μ l and 300 μ l of wash buffer. An additional step of centrifugation for 1 min at 13,000 x g was performed to dry the filter fleece completely and RNA was eluted with 50 μ l of Elution Buffer. The yield and quality (260/280 optical density ratios) of the RNA products were determined using a PicoDrop spectrophotometer (PicoDrop Ltd., Hinxton, UK). The purified total RNA was immediately used for cDNA synthesis or stored at -80°C until use.

Quantification of differentially expressed miRNAs. Reverse transcription was carried out using the Universal cDNA Synthesis kit (Exiqon A/S, Vedbaek, Denmark), according to the manufacturer's protocol A for individual assays. A total of 10 ng total RNA were used. The RT reaction was diluted 80 times in nuclease-free water, and 4 μ l aliquots were subsequently used for PCR amplification with 5 μ l ExiLent SYBR® Green Master mix (Exiqon A/S, Vedbaek, Denmark) and 1 μ l commercially available primers (Exiqon A/S, Vedbaek, Denmark): hsa-miR-146a-5p LNA™ PCR primer set (cat. no. 204688; target sequence, UAGCAGCACAAUAUGGUUUGUG); SNORD48 (hsa) PCR primer set (cat. no. 203903; target sequence, AGUGAUGAUGACCCAGGUAACUCUGAGUGUGUCGUGAUGCCAUCACCGCAGCGCUCUGACC); and U6 snRNA (hsa, mmu, rno) PCR primer set (cat. no 203907; target sequence, GUGCUCGCUUCGGCAGCACAUUAACUAAAAUUGGAACGAUACAGAGAAGAUUAGCAUGGCCCCUGCGCAAGGAUGACACGCAAAUUCGUGAAGCGUCCAUUUUUU). U6 small nuclear RNA and small nuclear RNA, C/D box 48 were used as the internal controls. The reactions were incubated in a 96-well plate at 95°C for 10 min, followed by 40 cycles of 95°C for 15s and 60°C for 1 min. All reactions were performed in duplicate using a 7900HT Fast Real-Time PCR System (Applied Biosystems; Thermo Fisher Scientific, Inc., Waltham, MA, USA). Relative expression level was determined using to the $2^{-\Delta\Delta C_q}$ method (29).

Statistical analysis. Kaplan-Meier survival curves were used to evaluate the association between the expression levels of miR-146a and patient survival rate. The differences between the studied groups were determined by using the Mann-Whitney U test or Kruskal-Wallis. Spearman's rank correlation coefficient was used in order to determine the statistical dependence between two variables. Multivariate analysis was used to estimate correlations between ≥ 3 variables. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Patients. Age, FIGO stage and histological grading distribution did not differ significantly between the groups of chemosensitive and chemoresistant patients. The number of patients who relapsed was similar in both groups. Progression-free survival (PFS) was longer for platinum-sensitive patients compared with platinum-resistant patients ($P < 0.05$). OS was not significantly longer for platinum-sensitive patients compared with platinum-resistant patients ($P = 0.07$; Table I).

miR-146a expression in primary tumor tissues, metastases and normal ovarian tissues. miR-146a expression in primary tumor samples was significantly increased in comparison with normal ovarian tissues ($P = 0.02$) and metastases ($P = 0.01$; Figs. 1 and 2). The range of relative quantification (RQ) values for primary tumors was heterogeneous, whereas the range of RQ values for metastases was homogeneous, suggesting stable and low level of miR-146a expression.

miR-146a expression and clinical parameters in ovarian cancer patients. For primary tumor tissues, a negative




Figure 1. miR-146a expression level in primary tumor tissues and in normal ovarian tissues. $P=0.02$. miR, microRNA; RQ, relative quantification; SD, standard deviation.




Figure 2. miR-146a expression level in primary tumor tissues and in metastases ($P=0.01$). miR, microRNA; RQ, relative quantification; SD, standard deviation.

Spearman's correlation was identified between miR-146a expression and serum levels of CA125 ($R=-0.37$; $P=0.03$) and ROMA index ($R=-0.79$; $P=0.0007$; Figs. 3 and 4). Serum CA125 levels and ROMA index were determined prior to cytoreductive surgery. The expression level of miR-146a was not correlated with FIGO stage or histological grading. The OS of the group of patients with ovarian cancer was




Figure 3. Correlation between miR-146a expression in primary tumor tissues and preoperative CA125 serum levels. $P=0.03$. miR, microRNA; RQ, relative quantification; CA125, cancer antigen 125.




Figure 4. Correlation between miR-146a expression in primary tumor tissues and preoperative ROMA index values. $P=0.0007$. miR, microRNA; RQ, relative quantification; ROMA, Risk of Malignancy Algorithm.

positively correlated with the level of miR-146a expression in the primary tumor tissues ($R=0.38$; $P=0.01$; Fig. 5). The Kaplan-Meier analysis revealed that the probability of survival was significantly decreased for patients with lower levels of miR-146a expression ($HR=0.21$; $P=0.003$) in the primary tumor tissues (Fig. 6). In multivariate analysis, lower levels of miR-146a expression in primary tumor tissues were associated with shorter PFS ($P=0.04$). Multivariate analysis indicated that lower expression levels of miR-146a in metastases correlated with platinum-resistance ($P=0.006$). This finding was not demonstrated for primary tumor tissues.

Discussion

The present retrospective study demonstrated that there is a difference in miR-146a expression in primary tumor tissues and omental metastases from patients with advanced serous ovarian cancer, who were subject to routine cytoreductive surgery and standard chemotherapy. Furthermore, an




Figure 5. Correlation between miR-146a expression level in primary tumor tissues and overall survival ($P=0.01$). miR, microRNA.




Figure 6. Kaplan-Meier analysis of survival patients with ovarian cancer and miR-146a expression levels in primary tumor tissues. ($P=0.003$). OS, overall survival; miR, microRNA; RQ, relative quantification.

association was identified between level of miR-146a expression and clinicopathological factors, including platinum resistance, survival rate, CA125 serum levels and ROMA index values.

miR-146a expression was observed to be increased and downregulated in various human cancer types compared with expression in normal tissues (6). Aberrant participation of miR-146a in the regulation of the immune response may result in a local inflammatory environment typical for many solid tumors, including ovarian cancer (18). A number of data studies have demonstrated expression of various miRs in serum or peripheral blood exosomes (17). However, the data concerning the levels of miR-146a expression in ovarian tumors are limited. Wyman *et al* (30) observed decreased expression of miR-146a in ovarian tumor tissue samples compared with normal ovarian cells. However, the tissue samples used in the study contained various histological types of cancer (including clear cell and endometrial cancer) and the reference samples used were cultured human ovarian surface epithelial (HOSE) cells. Similarly, Cui *et al* (31) detected downregulation of miR-146a in OVCAR3, CAOV3 and HEY ovarian cancer epithelial cells compared with HOSE cells. To

the best of our knowledge, the only previous published study investigating the differences between ovarian primary tumor and metastases in the context of miR expression levels, was performed by Vang *et al* (27) on 9 paired tissue specimens of primary lesions and omental metastases in serous cancer: Vang *et al* (27) demonstrated that miR-146a expression level was increased in metastases compared with in primary lesion tissues. It was demonstrated that high miR-146a expression was able to induce the formation of spheroids from cancer cells *in vitro*. This process mimics the *in vivo* intraperitoneal dissemination of cancer cell conglomerates over peritoneal cavities to form metastases (27).

By contrast to the previously mentioned results, the present study revealed that miR-146a expression was increased in primary tumor tissues compared with normal ovarian tissues, and that miR-146a expression level in omental metastases was reduced compared with primary ovarian tumor tissues. Similar inconsistencies in miRNA expression have been also observed by another study (32).

There are several factors that may account for the differences in miR-146a expression identified in the present study when compared with previous studies. First, only tumors of serous histology were included in the present study and tumors of different origin may exhibit different patterns of miRNA expression. Previous profiling studies that included different histological types of ovarian cancer support this observation and reported different miRNA expression in serous, mucinous, endometrial and clear-cell types of cancer (33), as well as in borderline and invasive tumors (19). It has also been demonstrated that miR-146a expression was downregulated in highly metastatic breast cancer. However, in breast tumors originating from BRCA1/2 mutation, miR-146a was overexpressed (34). The cancer tissues samples used in the present study were micro-dissected from the paraffin-embedded specimens. The specimens not only contained cancerous epithelium but also infiltrated stroma. Similarly, the reference samples also comprised normal surface epithelium and ovarian stroma. Heterogeneity of tissue samples may have had an impact on the expression level of miR-146a observed in the present study.

Furthermore, tissue samples extracted *in vivo* from the primary tumor were a heterogeneous population of cells compared with cultured cancer cell lines. An extensive range of miR-146a expression levels observed in primary tumor tissues in the present study may support this hypothesis. Primary tumors are composed of cell clones with different invasive potentials (35). It was demonstrated that the pattern of miRNA expression varied between cultured SKOV-3 and OVCAR-3 cell lines and was associated with invasiveness (36). The present study, which compared primary lesions and metastases, indicated that miR-146a expression levels in metastases were reduced compared with primary lesions. RQ values for metastases were more reproducible and demonstrated a more homogeneous range compared with the primary tumor tissues. Peritoneal metastases in ovarian cancer grow from small cell conglomerates, which originate from primary lesions that spread via lymphatic vessels or ascitic fluid (37). Metastatic cells create a less heterogeneous population and usually represent a more aggressive phenotype (38). Differences in the expression of miRNAs and other

small RNAs may regulate the type of tumor spread (milliary vs. bulky) in serous ovarian cancer (39). In the majority of cancer cases, miR-146a acts as a tumor suppressor, and its expression level was demonstrated to be downregulated in highly metastatic types of cancer (40). This finding suggested that more aggressive metastases may exhibit lower expressions of miR-146a, and the results of the present study supported this finding.

miRNA expression may be associated with clinical outcomes of ovarian cancer, and this has been reported by a number of studies. The level of miR-370 expression was increased in FIGO stage I/II samples compared with FIGO stage III/IV tissue samples. Additionally, upregulated expression of miR-181d, miR-30c, miR-30d and miR-30e-3p significantly improved disease-free survival or OS (19). To the best of our knowledge, miR-146a expression has not been associated with clinical outcomes of ovarian cancer. The results of the present study demonstrated that the levels of miR-146a expression in primary serous ovarian tumor tissues may be negatively correlated with serum levels of CA125 and ROMA index values. Assessment of the serum level of CA125 has been widely used in the diagnosis of ovarian cancer (41). High serum CA125 levels have been associated with FIGO stage, but predominantly with serous histology, low-grade cancer cases and the presence of ascites (42-45). Patients with CA125 levels >100 U/l were demonstrated to have shorter OS compared with patients with CA125 levels ≤100 U/l (46). The CA125 AUC value was also associated with FIGO stage, residual disease, response to chemotherapy and final outcome (47). ROMA index, originally described by Moore *et al* (48), is an algorithm, which takes into consideration the level of HE4 and CA125 together with menopausal status in order to classify patients with adnexal mass into groups of high or low risk of ovarian cancer. In patients with ovarian cancer, increased ROMA values were associated with advanced FIGO stage, undifferentiated tumors, ascites and lymph node involvement, as well as shorter overall, disease-free and progression-free survival rates (49,50). In the present study, high CA125 serum levels and increased ROMA values (reflecting the advancement of ovarian cancer) correlated with low expression of miR-146a in primary tumor tissues. OS was longer for patients with increased miR-146a expression in the primary tumor, and the probability of survival was significantly decreased for patients with low levels of miR-146a expression. In multivariate analysis, lower miR-146a expression levels in primary tumor tissues were associated with shorter PFS. These observations are consistent with previous studies performed in gastric cancer, which demonstrated that patients with high-expression profiles of miR-146a exhibited reduced lymph node metastases and exhibited longer OS compared with patients with low miR-146a expression profiles (9). These observations further supports the hypothesis that miR-146a may act as a tumor suppressor.

In the context of ovarian cancer chemosensitivity, various miRNAs have been studied. For example, the upregulation of miR-125b and miR-106a was previously reported *in vitro* in platinum-resistant ovarian cancer cells (23,51,52). miR-31 was also demonstrated to be downregulated in

taxane-resistant ovarian cancer cell lines (24). Similarly, downregulated expression of miR-9, -22, -129-5p, -155, -320a and -640 was observed in paclitaxel-resistant compared with paclitaxel-sensitive serous ovarian adenocarcinoma tissue samples (26). Validation of miRNA profiles in tumor tissue samples revealed that it was possible to predict chemo-sensitivity of ovarian tumors by detecting expression of miR-484, miR-642 and miR-217 (25). In the present study, a correlation was identified between lower miR-146a expression in metastatic tissue samples and platinum-resistance, but this correlation was not observed in primary tumor tissues. There are a number of potential miRNA-dependent mechanisms underlying chemoresistance in ovarian cancer. Analysis of miR-484 revealed that the chemosensitive tumor phenotype was induced by modulation of vascular endothelial growth factor (VEGF) B and VEGFR2 signaling pathways (25). Another possible mechanism underlying chemoresistance may be dependent on miR-146a and miR-150 (27). The expression levels of miR-146a and miR-150 were studied in SKOV-3, OVCAR-8 and IGROV-1 cell lines, and it was demonstrated that increased platinum tolerance was associated with spheroid formation, a model of peritoneal spread of cancer cells (27). Additionally, it was revealed that spheroid formation correlated with elevated miR-146a expression in metastases (27), contrary to the results of the present study. By contrast, another study reported that miR-146a is able to downregulate the expression of superoxide dismutase-2 and increase reactive oxygen species generation, which leads to increased apoptosis, inhibition of proliferation and increased sensitivity to chemotherapy (20). miR-146a may also mediate its effect via upregulating epidermal growth factor receptor, NFκB, interleukin-1 receptor-associated kinase 1 and metastasis-associated protein-2 (MTA2) (40). MTA2 is a transcription factor that regulates metastasis and may be triggered by a low expression of miR-146a, as it was demonstrated in pancreatic cancer cultured cells (40). Similarly in breast cancer cell lines, downregulation of miR-146a induced NFκB activation and augmented metastatic potential (53). It was revealed that the upregulation of transcription factor NFκB in numerous solid tumors prevents apoptosis induced by stress signals, including chemotherapeutic agents (54).

In conclusion, the results of the present study indicated differences in miR-146a expression levels between primary tumor tissues and in metastases of ovarian cancer. A correlation between the expression of miR-146a and clinicopathological features characterizing cancer advancement and chemoresistance was demonstrated. Low miR-146a expression level was also revealed to be a prognostic factor for an unfavorable outcome in patients with cancer. Further studies are required in order to investigate the precise pattern of miR-146a expression in various types of ovarian cancer, its role as a disease marker and as a potential target for novel therapeutic regimens.

Acknowledgements

The present study was supported by the Polish Mother's Memorial Hospital Research Institute (grant no. 2014/VII/25-GW).

References

- Trétarre B, Molinié F, Woronoff AS, Bossard N, Bessaoud F, Marrer E, Grosclaude P, Guizard AV, Delafosse P, Bara S, *et al*: Ovarian cancer in France: Trends in incidence, mortality and survival, 1980-2012. *Gynecol Oncol* 139: 324-329, 2015.
- Edwards HM, Noer MC, Sperling CD, Nguyen-Nielsen M, Lundvall L, Christensen IJ and Høgdall C: Survival of ovarian cancer patients in Denmark: Results from the Danish gynaecological cancer group (DGCG) database, 1995-2012. *Acta Oncol* 55 (Suppl 2): S36-S43, 2016.
- Cho KR and Shih IeM: Ovarian cancer. *Annu Rev Pathol* 4: 287-313, 2009.
- Huang J, Hu W and Sood AK: Prognostic biomarkers in ovarian cancer. *Cancer Biomark* 8: 231-251, 2010-2011.
- Lu LF, Boldin MP, Chaudhry A, Lin LL, Taganov KD, Hanada T, Yoshimura A, Baltimore D and Rudensky AY: Function of miR-146a in controlling Treg cell-mediated regulation of Th1 responses. *Cell* 142: 914-929, 2010.
- Labbaye C and Testa U: The emerging role of miR-146A in the control of hematopoiesis, immune function and cancer. *J Hematol Oncol* 5: 13, 2012.
- Chen G, Umelo IA, Lv S, Teugels E, Fostier K, Kronenberger P, Dewaele A, Sadones J, Geers C and De Grève J: miR-146a inhibits cell growth, cell migration and induces apoptosis in non-small cell lung cancer cells. *PLoS One* 8: e60317, 2013.
- Xu T, Zhu Y, Wei QK, Yuan Y, Zhou F, Ge YY, Yang JR, Su H and Zhuang SM: A functional polymorphism in the miR-146a gene is associated with the risk for hepatocellular carcinoma. *Carcinogenesis* 29: 2126-2131, 2008.
- Kogo R, Mimori K, Tanaka F, Komune S and Mori M: Clinical significance of miR-146a in gastric cancer cases. *Clin Cancer Res* 17: 4277-4284, 2011.
- Williams AE, Perry MM, Moschos SA, Larner-Svensson HM and Lindsay MA: Role of miRNA-146a in the regulation of the innate immune response and cancer. *Biochem Soc Trans* 36: 1211-1215, 2008.
- Shen J, Ambrosone CB, DiCioccio RA, Odunsi K, Lele SB and Zhao H: A functional polymorphism in the miR-146a gene and age of familial breast/ovarian cancer diagnosis. *Carcinogenesis* 29: 1963-1966, 2008.
- Wang X, Tang S, Le SY, Lu R, Rader JS, Meyers C and Zheng ZM: Aberrant expression of oncogenic and tumor-suppressive microRNAs in cervical cancer is required for cancer cell growth. *PLoS One* 3: e2557, 2008.
- He H, Jazdzewski K, Li W, Liyanarachchi S, Nagy R, Volinia S, Calin GA, Liu CG, Franssila K, Suster S, *et al*: The role of microRNA genes in papillary thyroid carcinoma. *Proc Natl Acad Sci USA* 102: 19075-19080, 2005.
- Hurst DR, Edmonds MD, Scott GK, Benz CC, Vaidya KS and Welch DR: Breast cancer metastasis suppressor 1 up-regulates miR-146, which suppresses breast cancer metastasis. *Cancer Res* 69: 1279-1283, 2009.
- Boldin MP, Taganov KD, Rao DS, Yang L, Zhao JL, Kalwani M, Garcia-Flores Y, Luong M, Devrekanli A, Xu J, *et al*: miR-146a is a significant brake on autoimmunity, myeloproliferation, and cancer in mice. *J Exp Med* 208: 1189-1201, 2011.
- Chang TC, Yu D, Lee YS, Wentzel EA, Arking DE, West KM, Dang CV, Thomas-Tikhonenko A and Mendell JT: Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat Genet* 40: 43-50, 2008.
- Zheng H, Liu JY, Song FJ and Chen KX: Advances in circulating microRNAs as diagnostic and prognostic markers for ovarian cancer. *Cancer Biol Med* 10: 123-130, 2013.
- Di Leva G and Croce CM: The role of microRNAs in the tumorigenesis of ovarian cancer. *Front Oncol* 3: 153, 2013.
- Lee H, Park CS, Deftereos G, Morihara J, Stern JE, Hawes SE, Swisher E, Kiviat NB and Feng Q: MicroRNA expression in ovarian carcinoma and its correlation with clinicopathological features. *World J Surg Oncol* 10: 174, 2012.
- Zheng H, Zhang L, Zhao Y, Yang D, Song F, Wen Y, Hao Q, Hu Z, Zhang W and Chen K: Plasma miRNAs as diagnostic and prognostic biomarkers for ovarian cancer. *PLoS One* 8: e77853, 2013.
- Kan CWS, Hahn MA, Gard GB, Maidens J, Huh JY, Marsh DJ and Howell VM: Elevated levels of circulating microRNA-200 family members correlate with serous epithelial ovarian cancer. *BMC Cancer* 12: 627, 2012.
- Delfino KR and Rodriguez-Zas SL: Transcription factor-microRNA-target gene networks associated with ovarian cancer survival and recurrence. *PLoS One* 8: e58608, 2013.
- Li H, Xu H, Shen H and Li H: microRNA 106a modulates cisplatin sensitivity by targeting PDCD4 in human ovarian cancer cells. *Oncol Lett* 7: 183-188, 2014.
- Mitamura T, Watari H, Wang L, Kanno H, Hassan MK, Miyazaki M, Katoh Y, Kimura T, Tanino M, Nishihara H, *et al*: Downregulation of miRNA-31 induces taxane resistance in ovarian cancer cells through increase of receptor tyrosine kinase MET. *Oncogenesis* 2: e40, 2013.
- Vecchione A, Belletti B, Lovat F, Volini S, Chiappetta G, Giglio S, Sonogo M, Cirombella R, Onesti EC, Pellegrini P, *et al*: A microRNA signature defines chemoresistance in ovarian cancer through modulation of angiogenesis. *Proc Natl Acad Sci USA* 110: 9845-9850, 2013.
- Li X, Lu Y, Chen Y, Lu W and Xie X: MicroRNA profile of paclitaxel-resistant serous ovarian carcinoma based on formalin-fixed paraffin-embedded samples. *BMC Cancer* 13: 216, 2013.
- Vang S, Wu HT, Fischer A, Miller DH, MacLaughlan S, Douglass E, Comisar L, Steinhoff M, Collins C, Smith PJ, *et al*: Identification of ovarian cancer metastatic miRNAs. *PLoS One* 8: e58226, 2013.
- Pecorelli S: Revised FIGO staging for carcinoma of the vulva, cervix, and endometrium. *Int J Gynaecol Obstet* 105: 103-104, 2009.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408, 2001.
- Wyman SK, Parkin RK, Mitchell PS, Fritz BR, O'Brian K, Godwin AK, Urban N, Drescher CW, Knudsen BS and Tewari M: Repertoire of microRNAs in epithelial ovarian cancer as determined by next generation sequencing of small RNA cDNA libraries. *PLoS One* 4: e5311, 2009.
- Cui Y, She K, Tian D, Zhang P and Xin X: miR-146a inhibits proliferation and enhances chemosensitivity in epithelial ovarian cancer via reduction of SOD2. *Oncol Res* 23: 275-282, 2016.
- Chen Y, Zhang L and Hao Q: Candidate microRNA biomarkers in human epithelial ovarian cancer: Systematic review profiling studies and experimental validation. *Cancer Cell Int* 13: 86, 2013.
- Zaman MS, Maher DM, Khan S, Jaggi M and Chauhan SC: Current status and implications of microRNAs in ovarian cancer diagnosis and therapy. *J Ovarian Res* 5: 44, 2012.
- Garcia AI, Buisson M, Bertrand P, Rimokh R, Rouleau E, Lopez BS, Lidereau R, Mikaelian I and Mazoyer S.: Down-regulation of BRCA1 expression by miR-146a and miR-146b-5p in triple negative sporadic breast cancers. *EMBO Mol Med* 3: 279-290, 2011.
- Diaz-Cano SJ: Tumor heterogeneity: Mechanisms and bases for a reliable application of molecular marker design. *Int J Mol Sci* 13: 1951-2011, 2012.
- Kobayashi M, Salomon C, Tapia J, Illanes SE, Mitchell MD and Rice GE: Ovarian cancer cell invasiveness is associated with discordant exosomal sequestration of Let-7 miRNA and miR-200. *J Transl Med* 12: 4, 2014.
- Lengyel E: Ovarian cancer development and metastasis. *Am J Pathol* 177: 1053-1064, 2010.
- Périgny M, Bairati I, Harvey I, Beauchemin M, Harel F, Plante M and Têtu B: Role of immunohistochemical overexpression of matrix metalloproteinases MMP-2 and MMP-11 in the prognosis of death by ovarian cancer. *Am J Clin Pathol* 129: 226-231, 2008.
- Bachmayr-Heyda A, Auer K, Sukhbaatar N, Aust S, Deycmar S, Reiner AT, Polterauer S, Dekan S and Pils D: Small RNAs and the competing endogenous RNA network in high grade serous ovarian cancer tumor spread. *Oncotarget* 7: 39640-39653, 2016.
- Li Y, VandenBoom TG II, Wang Z, Kong D, Ali S, Philip PA and Sarkar FH: MiR-146a suppresses invasion of pancreatic cancer cells. *Cancer Res* 70: 1486-1495, 2010.
- Duffy MJ, Bonfrer JM, Kulpa J, Rustin GJ, Soletormos G, Torre GC, Tuxen MK and Zwierner M: CA125 in ovarian cancer: European group on tumor markers guidelines for clinical use. *Int J Gynecol Cancer* 15: 679-691, 2005.
- But I and Gorisek B: Preoperative value of CA 125 as a reflection of tumor grade in epithelial ovarian cancer. *Gynecol Oncol* 63: 166-172, 1996.
- Topalak O, Saygili O, Soyuturk M, Karaca N, Batur Y, Uslu T and Erten O: Serum, pleural effusion, and ascites CA-125 levels in ovarian cancer and nonovarian benign and malignant diseases: A comparative study. *Gynecol Oncol* 85: 108-113, 2002.
- Rossi AC, Di Vagno G, Cormio G, Cazzolla A, Stefanelli S, D'Elia E and Selvaggi L: A retrospective study of preoperative CA 125 levels in 82 patients with ovarian cancer. *Arch Gynecol Obstet* 269: 263-265, 2004.

45. Ayhan A, Guven S, Guven ES and Kucukali T: Is there a correlation between tumor marker panel and tumor size and histopathology in well-staged patients with borderline ovarian tumors? *Acta Obstet Gynecol Scand* 86: 484-490, 2007.
46. Markmann S, Gerber B and Briese V: Prognostic value of Ca 125 levels during primary therapy. *Anticancer Res* 27: 1837-1839, 2007.
47. Mano A, Falcão A, Godinho I, Santos J, Leitão F, Oliveira C and Caramona M: CA-125 AUC as a new prognostic factor for patients with ovarian cancer. *Gynecol Oncol* 97: 529-534, 2005.
48. Moore RG, McMeekin DS, Brown AK, DiSilvestro P, Miller MC, Allard WJ, Gajewski W, Kurman R, Bast RC Jr and Skates SJ: A novel multiple marker bioassay utilizing HE4 and CA125 for the prediction of ovarian cancer in patients with a pelvic mass. *Gynecol Oncol* 112: 40-46, 2009.
49. Bandiera E, Romani C, Specchia C, Zanotti L, Galli C, Ruggeri G, Tognon G, Bignotti E, Tassi RA, Odicino F, *et al*: Serum human epididymis protein 4 and risk for ovarian malignancy algorithm as new diagnostic and prognostic tools for epithelial ovarian cancer management. *Cancer Epidemiol Biomarkers Prev* 20: 2496-2506, 2011.
50. Steffensen KD, Waldstrøm M, Brandslund I, Petzold M and Jakobsen A: The prognostic and predictive value of combined HE4 and CA-125 in ovarian cancer patients. *Int J Gynecol Cancer* 22: 1474-1482, 2012.
51. Kong F, Sun C, Wang Z, Han L, Weng D, Lu Y and Chen G: miR-125b confers resistance of ovarian cancer cells to cisplatin by targeting pro-apoptotic Bcl-2 antagonist killer 1. *J Huazhong Univ Sci Technolog Med Sci* 31: 543-549, 2011 .
52. Li H, Xu H, Shen H and Li H: microRNA-106a modulates cisplatin sensitivity by targeting PDCD4 in human ovarian cancer cells. *Oncol Lett* 7: 183-188, 2014.
53. Bhaumik D, Scott GK, Schokrpur S, Patil CK, Campisi J and Benz CC: Expression of microRNA-146 suppresses NF-kappaB activity with reduction of metastatic potential in breast cancer cells. *Oncogene* 27: 5643-5647, 2008.
54. Rayet B and Gélinas C: Aberrant rel/nfkb genes and activity in human cancer. *Oncogene* 18: 6938-6947, 1999.
55. Pecorelli S, Ngan HYS and Hacker NF (eds): Staging classifications and clinical practice guidelines for gynaecological cancers. A collaboration between FIGO and IGCS. 3rd edition. London: FIGO, 2006.