

# Expression of circadian clock genes in human colorectal adenoma and carcinoma

TOMOYUKI MOMMA, HIROKAZU OKAYAMA, MASARU SAITOU, HIDEKAZU SUGENO, NOBUHIRO YOSHIMOTO, YUJI TAKEBAYASHI, SHINJI OHKI and SEIICHI TAKENOSHITA

Department of Organ Regulatory Surgery, Fukushima Medical University School of Medicine, Fukushima 960-1295, Japan

Received July 29, 2015; Accepted October 5, 2016

DOI: 10.3892/ol.2017.6876

**Abstract.** Circadian rhythms are fundamental biological systems in most organisms. Epidemiological and animal studies have demonstrated that disruption of circadian rhythms is linked to tumor progression and mammalian tumorigenesis. However, the clinical significance of *in situ* clock gene expression in precancerous and cancerous colorectal lesions remains unknown. The present study aimed to investigate mRNA transcript levels of circadian clock genes within human colorectal cancer and adenoma tissue sections. Using *in situ* hybridization, the expression of key clock genes, including period circadian protein homolog (*Per*) 1 and 2, cryptochrome 1 (*Cry1*), circadian locomotor output cycles protein kaput (*Clock*), brain and muscle ARNT-like protein 1 (*Bmall*) and casein kinase 1 $\epsilon$  (*CK1 $\epsilon$* ) were retrospectively examined in 51 cases of colorectal carcinoma and 10 cases of adenoma. The expression of clock genes was almost undetectable in the majority of adenomas, whereas positive expression of clock genes was observed in 27-47% of carcinomas. Notably, positive *Per1*, *Per2* and *Clock* staining in colorectal carcinomas were each significantly associated with a larger tumor size (P=0.012, P=0.011 and P=0.009, respectively). Tumors with positive *Per2* and *Clock* expression tended to exhibit deeper depth of invasion and were generally more advanced than tumors that did not express these genes (P=0.052 and P=0.064, respectively). However, no statistically significant association was observed between clock gene expression and clinicopathological variables, including histopathological differentiation,

lymph node metastasis, depth of invasion or disease stage, although *Per2*-positive tumors tended to be associated with poorer overall survival (P=0.060). The results of the current study suggest that dysregulated expression of clock genes may be important in human colorectal tumorigenesis.

## Introduction

A number of biochemical, physiological and behavioral processes have demonstrated that an internal time-keeping mechanism, referred to as the biological clock, regulates circadian rhythms. The master circadian clock coordinates peripheral clocks elsewhere in the body and is located in the suprachiasmatic nuclei (SCN) within the anterior hypothalamus (1). The core oscillator driving this clock is intergrated by an auto-regulatory transcription-(post) translation-based feedback loop, which is comprised of genes related to the circadian rhythm (1,2).

Epidemiological studies have suggested that disruption of the circadian clock may increase cancer risk in humans (3-5). In particular, it has been observed that shift workers have an increased risk of developing malignancies, including breast, endometrial, prostate and colorectal cancer, due to their disrupted circadian cycles (5-9). Fu *et al* (10) previously demonstrated that a period circadian protein homolog 2 (*Per2*) mutation induced upregulation of c-Myc and downregulation of p53 transcription in mice; furthermore, the incidence of spontaneous and radiation-induced lymphoma increased, as did lymphoma-associated mortality. Other *in vivo* studies have identified an association between alterations of the circadian rhythm and tumorigenesis (2,9,11). In a number of types of human solid cancer, including breast, endometrial and colorectal cancer, the dysregulated expression of circadian genes has been investigated by immunohistochemistry and/or reverse transcription quantitative polymerase chain reaction (RT-qPCR) (9,12-14).

The aim of the present study was to investigate the clinical significance of the mRNA expression of clock genes in human colorectal carcinoma and adenoma tissues, using *in situ* hybridization.

## Patients and methods

**Patients and tumor samples.** A total of 51 patients (32 males and 19 females) with colorectal carcinoma, and 10 patients with

---

**Correspondence to:** Dr Seiichi Takenoshita, Department of Organ Regulatory Surgery, Fukushima Medical University School of Medicine, 1 Hikarigaoka, Fukushima, Fukushima 960-1295, Japan  
E-mail: takenoss@fmu.ac.jp

**Abbreviations:** SCN, suprachiasmatic nuclei; Per, period circadian protein homolog; Bmall, brain and Muscle ARNT-Like Protein 1; Clock, circadian locomotor output cycles protein kaput; Cry, cryptochrome; CK1 $\epsilon$ , casein Kinase 1 $\epsilon$

**Key words:** circadian rhythm, colorectal cancer, clock gene, *in situ* hybridization

Table I. Associations between clock gene expression and clinicopathological variables.

Variables	Per1			Per2			Cry1			Clock			Bmal1			CKIε			
	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	
	51	27	24	24	25	26	27	24	30	21	31	20	37	14	14	37	14	P-value	
Age																			
Mean	65.1	63.7	67.6	0.242	68.9	62.2	65.1	65.9	65.6	65.4	65.3	65.9	65.8	64.6	0.423	65.8	64.6	0.727	
Range	33-84	33-81	43-84		49-84	33-81	33-84	41-84	41-81	33-84	41-84	33-84	33-84	46-80		33-84	46-80		
Gender																			
Male	32	17	15	1	16	16	16	16	20	12	20	12	25	7	0.563	25	7	0.333	
Female	19	10	9		9	10	11	8	10	9	11	8	12	7		12	7		
Tumor size (mm)																			
<50	29	20	9	0.012 <sup>a</sup>	19	10	18	11	22	7	21	8	23	6	0.009 <sup>a</sup>	23	6	0.342	
≥50	22	7	15		6	16	9	13	8	14	10	12	14	8		14	8		
Tumor location																			
Colon	24	12	12	0.782	12	12	14	10	13	11	16	8	19	5	0.578	19	5	0.363	
Rectum	27	15	12		13	14	13	14	17	10	15	12	18	9		18	9		
Histological differentiation																			
Well	34	17	17	0.636	17	17	18	16	20	14	22	12	27	7	0.991	27	7	0.212	
Moderately	13	7	6		6	7	7	6	7	6	6	7	7	6		7	6		
Poorly	0	0	0		0	0	0	0	0	0	0	0	0	0		0	0		
Mucinous	4	3	1		2	2	2	2	3	1	3	1	3	1		3	1		
Depth of invasion																			
pT1	5	3	2	0.482	4	1	3	2	4	1	2	3	2	3	0.303	2	3	0.240	
pT2	7	4	3		5	2	4	3	5	2	4	3	6	1		6	1		
pT3	31	14	17		11	20	16	15	15	16	21	10	22	9		22	9		
pT4	8	6	2		5	3	4	4	6	2	4	4	7	1		7	1		
Depth of invasion																			
pT1-2	12	7	5	0.749	9	3	7	5	9	3	6	6	8	4	0.315	8	4	0.715	
pT3-4	39	20	19		16	23	20	19	21	18	25	14	29	10		29	10		
Lymph node metastasis																			
Absent	27	14	13	1	13	14	12	15	16	11	16	11	18	9	1	18	9	0.363	
Present	24	13	11		12	12	15	9	14	10	15	9	19	5		19	5		

Table I. Continued.

Variables	Per1			Per2			Cry1			Clock			Bmal1			CKIε		
	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive	Total	Negative	Positive
	51	27	24	25	26	24	27	24	30	21	31	20	37	14	51	27	24	24
			P-value			P-value			P-value				P-value					P-value
Stage			0.723			0.240			0.506						0.180			0.987
I	9	6	3	6	3	4	5	4	8	1	5	4	6	3				
II	16	7	9	6	10	10	6	10	7	9	10	6	10	6				
III	21	11	10	12	9	8	13	8	12	9	13	8	18	3				
IV	5	3	2	1	4	2	3	2	3	2	3	2	3	2				
Stage			0.473			0.291			1						0.064			0.724
I	9	6	3	6	3	4	5	4	8	1	5	4	6	3				
II-IV	42	21	21	19	23	20	22	20	22	20	26	16	31	11				

\*P<0.05, indicates a significant difference. P, P-value; Per, period circadian protein homolog; Cry1, cryptochrome 1; Clock, circadian locomotor output cycles protein kaput; Bmal1, brain and muscle ARNT-like protein 1; CKIε, casein kinase 1ε.

colorectal adenoma were examined. All patients underwent endoscopic or surgical resection to completely remove tumors in the Department of Organ Regulatory Surgery, Fukushima Medical University Hospital (Fukushima, Japan) between April 1999 and July 2005. In several tissue specimens, the surrounding normal mucosa was also examined. None of the patients had received prior chemotherapy or irradiation or had experienced any other form of cancer. The clinicopathological characteristics of the 51 patients with colorectal cancer investigated in this study are summarized in Table I.

All tissue samples were embedded in optimal cutting temperature (OCT) compound (Sakura Finetek USA, Inc., Torrance, CA, USA) and immediately stored at -8°C. Tumors were histopathologically classified as well-differentiated, moderately differentiated, poorly differentiated or mucinous adenocarcinomas (15), and tumor size was defined as the largest diameter of the tumor. Histopathological diagnoses were performed at the Department of Pathology, Fukushima Medical University Hospital following standard procedures. Informed consent was obtained from each patient and the Fukushima Medical University Committee approved the protocol of the present study.

*In situ hybridization.* Digoxigenin (DIG)-UTP labeled cRNA probes were used to evaluate the mRNA expression of clock genes. The DIG-labeled cRNA probes were synthesized using a DIG RNA Labeling kit (Roche Diagnostics, Basel, Switzerland), and were labeled with SP6 or T7 RNA polymerase in the presence of DIG-UTP. Sections 5-μm thick were formed from the embedded tissue specimens, sufficiently dried with cold air and fixed in 4% paraformaldehyde diluted with phosphate-buffered saline for 30 min. Sections were hybridized overnight at 42°C in hybridization buffer containing 1 μg/ml of DIG-labeled probe. The DIG-labeled probes were diluted to 1 μg/ml with hybridization buffer (Nippon Gene Co., Ltd., Tokyo, Japan) and dropped to the sections, which were incubated at 42°C for 16-20 h to hybridize with each probe. Following hybridization, the sections were washed in 2 x standard citrate buffered saline and 0.2 x saline sodium citrate buffer at 5°C for 20 min and treated with 1% blocking solution at room temperature for 30 min, using the DIG Nucleic Acid Detection Kit (Roche Diagnostics). The sections were subsequently incubated at room temperature for 30 min with alkaline phosphatase labeled anti-digoxigenin antibody (Roche) diluted with a blocking solution (1:5,000). Color reaction was conducted using NTB/BCIP at 4°C for 12 h. As a negative control, the serial section was hybridized with a sense probe. Sections were simultaneously evaluated by two investigators. The tumor cells were classified into 4 groups based on intensity of staining (none, weak, moderate or strong) indicating levels of gene expression within the cells.

*Statistical analysis.* Differences between groups were evaluated by the χ<sup>2</sup> test, Fisher's exact test, Student's *t* test or the Mann-Whitney U test. Cumulative survival was estimated by the Kaplan-Meier method and differences were analyzed by the log-rank test. All statistical analyses were two-sided and P<0.05 was considered to indicate a statistically significant difference.




Figure 1. Expression of clock genes detected by *in situ* hybridization in colorectal carcinoma and surrounding normal mucosa. (A) *Per1*, (B) *Per2*, (C) *Cry1*, (D) *Clock*, (E) *Bmal1* and (F) *CK1ε*. Panels 1 and 2 represent carcinoma tissue with anti-sense probe (magnification, x100 and x400, respectively). Panels 3 and 4 represent carcinoma tissues with sense probe (magnification, x100 and x400, respectively). Panels 5 and 6 represent normal mucosa tissues with anti-sense and sense probes, respectively (magnification, x100 for both). *Per*, Period circadian protein homolog; *Cry1*, Cryptochrome 1; *Clock*, Circadian locomotor output cycles protein kaput; *Bmal1*, Brain and Muscle ARNT-like Protein 1; *CK1ε*, Casein Kinase 1ε.

## Results

**Potential involvement of clock genes in colorectal tumor progression.** mRNA expression of the clock genes, including *Per1* and 2, cryptochrome 1 (*Cry1*), circadian locomoter output cycles protein kaput (*Clock*), brain and muscle ARNT-like protein 1 (*Bmal1*) and casein kinase 1ε (*CK1ε*), was examined in colorectal cancer tissues by *in situ* hybridization. As presented in Fig. 1, sense probes as negative controls exhibited no staining (Fig. 1 panels 3-4), whereas various levels of staining were observed in tumor cells detected by anti-sense probes (Fig. 1, panels 1-2). Normal epithelial areas were evaluated in a number of specimens where the surrounding normal mucosa was available, however, no clear staining was detected by any of the probes (Fig. 1, panels 5-6). The levels of expression of clock genes in tumor cells were classified into four groups (none, weak, moderate or strong), based on the intensity of staining (Fig. 2). Tumors with no or weak staining were further defined as a negative group, while tumors with moderate and strong staining were a positive group. Of the 51 colorectal carcinomas evaluated, positive staining for *Per1*, *Per2*, *Cry1*, *Clock*, *Bmal1* and *CK1ε* was observed in 24 (47%), 26 (51%), 24 (47%), 21 (41%), 20 (39%) and 14 (27%) tumors, respectively (Table I and Fig. 3). However, no significant associations

were observed between levels of clock gene expression and histopathological type, depth of invasion, lymph node metastasis or disease stage (Table I). Although positive *Per2* and positive *Clock* groups tended to be associated with a deeper depth of invasion and advanced stage, respectively, these associations were not significant ( $P=0.052$  and  $P=0.064$ , respectively). By contrast, positive-*Per1*, *Per2* and *Clock* groups were each associated with larger tumor size ( $>50$  mm;  $P=0.012$ ,  $P=0.011$  and  $P=0.009$ , respectively; Table I). Similar results were obtained when tumor size was treated as a continuous variable (Fig. 4), therefore, the potential prognostic significance of *Per1*, *Per2* and *Clock* was investigated. No association was observed between *Per1* or *Clock* positive expression and overall survival rates ( $P=0.0599$  and  $P=0.994$ , respectively; Fig. 5A and B). On the other hand, patients with carcinomas exhibiting positive-*Per2* expression tended to have lower rates of survival than patients with negative-*Per2* carcinomas, although this association was not significant ( $P=0.060$ ; Fig. 5C).

**The expression of clock genes in colorectal adenoma.** To investigate the expression of clock genes in precancerous lesions compared with cancer tissues, 10 colorectal adenomas were examined by *in situ* hybridization. In contrast to cancer tissues, the expression of each clock




Figure 2. Classification of the expression of clock genes by *in situ* hybridization. The intensity of staining in tumor cells was classified into four groups as: (A) None, (B) weak, (C) moderate and (D) strong. Magnification, x100.




Figure 3. Proportion of expression levels of clock genes in colorectal carcinoma (left panel) and adenoma (right panel). Differences between positive and negative were analyzed by Fisher's exact test. Per, period circadian protein homolog; Cry1, cryptochrome 1; Clock, circadian locomotor output cycles protein kaput; Bmal1, brain and muscle ARNT-like protein 1; CK1ε, casein kinase 1ε.

gene was undetectable in the majority of adenoma tissue (Fig. 3). No adenomas (0%) exhibited positive staining for *Per1*, *Per2*, *Cry1*, *Bmal1* or *CK1ε* and only 10% of adenomas

exhibited positive expression of *Clock*. Hence, the proportion of tissues indicating positive *Per1*, *Per2*, *Cry1* and *Cry2* expression in colorectal carcinoma was significantly higher




Figure 4. Comparison of tumor size between positive and negative staining of each gene. (A) *Per1*, (B) *Per2*, (C) *Cry1*, (D) *Clock*, (E) *Bmal1* and (F) *CK1ε*. Boxes correspond to the inter-quartile ranges, with the lower boundary of the box representing the 25th percentile and the upper boundary representing the 75th percentile. Differences between groups were analyzed by Student's *t*-test. Per, period circadian protein homolog; *Cry1*, cryptochrome 1; *Clock*, circadian locomotor output cycles protein kaput; *Bmal1*, brain and muscle ARNT-like protein 1; *CK1ε*, casein kinase 1ε.




Figure 5. Kaplan-Meier survival curves for *Per1* (A), *Per2* (B) and *Clock* (C). Survival differences were assessed using the log-rank test. Per, period circadian protein homolog; *Clock*, circadian locomotor output cycles protein kaput.

than in colorectal adenoma ( $P=0.046$ ,  $P=0.004$ ,  $P=0.004$  and  $P=0.015$ , respectively; Fig. 3).

## Discussion

Epidemiological studies have suggested that disruption of the circadian rhythm is associated with increased cancer incidence and poorer disease outcome (3-5,8). Previous studies have indicated that in *Per2* mutant mice, *Bmal1* expression decreased causing an increase in c-Myc transcription, thus disrupting the circadian rhythm and increasing cancer risk (10). In colorectal cancer, animal studies using chemically induced models as well as *APC<sup>Min/+</sup>* mice, have suggested a link between alterations of circadian genes and colorectal tumor development and progression (16-18). Other studies have used RT-qPCR to demonstrate that clock gene expression is dysregulated in human colorectal cancer (9,19,20).

In the present study, unlike previous studies, the *in situ* hybridization technique was utilized to detect clock gene mRNA expression in colorectal tumor tissues, including precancerous and cancerous lesions. The proportion of colorectal carcinomas with positive *Per1*, *Per2*, *Cry1* and *Cry2* expression was observed to be significantly higher than the proportion of adenomas, suggesting that dysregulated clock gene expression may be involved in colorectal tumorigenesis. Colorectal carcinoma tumors exhibiting positive staining of *Per1*, *Per2* and *Clock* were significantly larger than those exhibiting negative staining. Correspondingly, tumors with positive staining of *Per2* and *Clock* tended to be associated with deeper depth of invasion and a more advanced stage of cancer. Furthermore, an association was observed between positive-*Per2* expression and poorer overall survival outcome, though this was not technically significant. However, due to the relatively small sample size and short follow-up time of the

present study, the prognostic impact of positive-*Per2* expression remains to be fully determined.

Clock genes are involved in cell cycle regulation (21). A positive factor, the Clock-Bmal1 dimer, is required for transcription initiation of the *Per* and *Cry* genes oscillating mechanism in the feedback mechanisms of the clock genes (21). By contrast, *Per* and *Cry* proteins are supposed to act as negative factors and promote oscillation (22). The Clock-Bmal1 dimer promotes transcription through the E-box of *Wee1*, suppressing the cell cycle at M-phase (22). The results of previous studies have demonstrated a connection between the alterations of clock genes, and cell cycle progression and proliferation through c-Myc/p21 signaling and the Wnt/ $\beta$ -catenin pathway, which are implicated in the molecular pathogenesis of colorectal cancer (11,17,18). Taken together with the results of the present study, this indicates that the imbalance of clock gene expression levels, which results in the dysregulation of cell cycle, may stimulate the adenoma-carcinoma transition and tumor progression during colorectal carcinogenesis. However, the current study did not address whether clock gene expression directly contributes to cell cycle dysregulation and tumorigenesis. The biological significance of *in situ* clock gene expression remains to be elucidated. Clock gene expression *in situ* may at least in part represent the dysregulated rhythms in carcinomas, therefore, future studies are required to address the molecular mechanisms by which the imbalance of clock genes expression contributes to dysregulated circadian rhythms and consequently, to tumorigenesis.

In conclusion, mRNA expression of key clock genes, including *Per1*, *Per2*, *Cry1* and *Cry2* was frequently found in carcinomas, but not in adenomas, using *in situ* hybridization. Also, the expression of some clock genes were associated with tumor size, and tended to be associated with depth of invasion and survival outcome. Therefore, the present study suggests that dysregulated clock gene expression may serve an important role in human colorectal tumorigenesis.

## References

1. Fu L and Lee CC: The circadian clock: Pacemaker and tumour suppressor. *Nat Rev Cancer* 3: 350-361, 2003.
2. Hunt T and Sassone-Corsi P: Riding tandem: Circadian clocks and the cell cycle. *Cell* 129: 461-464, 2007.
3. Devilee P, Schuurin E, van de Vijver MJ and Cornelisse CJ: Recent developments in the molecular genetic understanding of breast cancer. *Crit Rev Oncog* 5: 247-270, 1994.
4. Ronco A, De Stefani E, Mendilaharsu M and Deneo-Pellegrini H: Meat, fat and risk of breast cancer: A case-control study from Uruguay. *Int J Cancer* 65: 328-331, 1996.
5. Stevens RG, Brainard GC, Blask DE, Lockley SW and Motta ME: Breast cancer and circadian disruption from electric lighting in the modern world. *CA Cancer J Clin* 64: 207-218, 2014.
6. Ambrosone CB, Freudenheim JL, Graham S, Marshall JR, Vena JE, Brasure JR, Michalek AM, Laughlin R, Nemoto T, Gillenwater KA and Shields PG: Cigarette smoking, N-acetyltransferase 2 genetic polymorphisms, and breast cancer risk. *JAMA* 276: 1494-1501, 1996.
7. Magnusson C, Baron J, Persson I, Wolk A, Bergström R, Trichopoulos D and Adami HO: Body size in different periods of life and breast cancer risk in post-menopausal women. *Int J Cancer* 76: 29-34, 1998.
8. Schernhammer ES, Laden F, Speizer FE, Willett WC, Hunter DJ, Kawachi I, Fuchs CS and Colditz GA: Night-shift work and risk of colorectal cancer in the nurses' health study. *J Natl Cancer Inst* 95: 825-828, 2003.
9. Karantanos T, Theodoropoulos G, Pektasides D and Gazouli M: Clock genes: Their role in colorectal cancer. *World J Gastroenterol* 20: 1986-1992, 2014.
10. Fu L, Pelicano H, Liu J, Huang P and Lee C: The circadian gene *Period2* plays an important role in tumor suppression and DNA damage response in vivo. *Cell* 111: 41-50, 2002.
11. Huisman SA, Oklejewicz M, Ahmadi AR, Tamanini F, Ijzermans JN, van der Horst GT and de Bruin RW: Colorectal liver metastases with a disrupted circadian rhythm phase shift the peripheral clock in liver and kidney. *Int J Cancer* 136: 1024-1032, 2014.
12. Yeh KT, Yang MY, Liu TC, Chen JC, Chan WL, Lin SF and Chang JG: Abnormal expression of period 1 (PER1) in endometrial carcinoma. *J Patol* 206: 111-120, 2005.
13. Shih HC, Choo KB, Chang TJ, Yang MY, Shih MC, Yeh KT, Liu TC, Lin SF and Chang JG: Disturbance of circadian gene expression in endometrial cancer: Detection by real-time quantitative RT-PCR. *Oncol Rep* 14: 1533-1538, 2005.
14. Chen ST, Choo KB, Hou MF, Yeh KT, Kuo SJ and Chang JG: Deregulated expression of the PER1, PER2 and PER3 genes in breast cancers. *Carcinogenesis* 26: 1241-1246, 2005.
15. Sobin LH: World Health Organization international histological classification of tumors: Histological typing of intestinal tumors. 2nd edition. New York: Springer-Verlag, 1989.
16. Wood PA, Yang X, Taber A, Oh EY, Ansell C, Ayers SE, Al-Assaad Z, Carnevale K, Berger FG, Peña MM and Hrushesky WJ: Period 2 mutation accelerates *ApcMin/+* tumorigenesis. *Mol Cancer Res* 6: 1786-1793, 2008.
17. Yang X, Wood PA, Ansell CM, Ohmori M, Oh EY, Xiong Y, Berger FG, Peña MM and Hrushesky WJ: Beta-catenin induces beta-TrCP-mediated PER2 degradation altering circadian clock gene expression in intestinal mucosa of *ApcMin/+* mice. *J Biochem* 145: 289-297, 2009.
18. Soťák M, Polidarová L, Ergang P, Sumová A and Pácha J: An association between clock genes and clock-controlled cell cycle genes in murine colorectal tumors. *Int J Cancer* 132: 1032-1041, 2013.
19. Oshima T, Takenoshita S, Akaike M, Kunisaki C, Fujii S, Nozaki A, Numata K, Shiozawa M, Rino Y, Tanaka K, *et al*: Expression of circadian genes correlates with liver metastasis and outcomes in colorectal cancer. *Oncol Rep* 25: 1439-1446, 2011.
20. Karantanos T, Theodoropoulos G, Gazouli M, Vaiopoulou A, Karantanou C, Lymberi M and Pektasides D: Expression of clock genes in patients with colorectal cancer. *Int J Biol Markers* 28: 280-285, 2013.
21. Matsuo T, Yamaguchi S, Mitsui S, Emi A, Shimoda F and Okamura H: Control mechanism of the circadian clock for timing of cell division in vivo. *Science* 302: 255-259, 2003.
22. Yagita K, Tamanini F, Yasuda M, Hoeijmakers JH, van der Horst GT and Okamura H: Nucleocytoplasmic shuttling and mCRY-dependent inhibition of ubiquitylation of the mPER2 clock protein. *EMBO J* 21: 1301-1314, 2002.