

Aberrant expression of miR-153 is associated with the poor prognosis of cervical cancer

LI LIU¹, XIAOJING LAI², CHANGJIN YUAN¹, XIUWEI LV¹, TAO YU¹,
WENYU HE¹, JIAOPING LIU¹ and HAIMING ZHANG¹

¹Department of Oncology, Integrated Traditional Chinese and Western Medicine, The Central Hospital of Wuhan, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430012;

²Applied Biotechnology Research Center, Hubei Engineering Research Center of Viral Vector, Wuhan Institute of Bioengineering, Economic Development Zone of Yangluo, Wuhan, Hubei 430415, P.R. China

Received September 16, 2017; Accepted January 24, 2018

DOI: 10.3892/ol.2018.8475

Abstract. Previous studies have demonstrated that microRNAs (miRNAs) are frequently dysregulated in tumors and are associated with the initiation and progression of various types of cancer. miR-153 has been previously shown to have an anti-tumor effect in the majority of cancer types. However, to date, the expression status and function of miR-153 in cervical cancer (CC) remains unclear. In the present study, the expression of miR-153 in CC tissues and cell lines was examined, revealing that the expression of miR-153 was markedly downregulated in the CC tissues and cell lines investigated, when compared with matched noncancerous tissues and normal cervical epithelial cell line. Furthermore, ectopic expression of miR-153 by miR-153 mimic inhibited cell proliferation; however, transfection with the miR-153 inhibitor promoted the cell proliferation in CC cell lines. Finally, the results showed that the downregulation of miR-153 was associated with poor 5-year over survival in CC patients and it could be regarded as an independent biomarker to predict the prognosis of CC patients. Collectively, these results indicated that miR-153 may function as a tumor suppressor in CC, and it may be a potential novel therapeutic target for CC.

Introduction

Cervical cancer (CC) is one of the most malignant female reproduction cancers in human, as well as the fourth most common cause of cancer-related mortality in the world (1,2). Globally, it was estimated that there were about 530,000 new

diagnosed cases each year (1). In the meantime, approximately 135,000 causes were found in China, accounting for about 25% of all the new CC causes each year (3). Despite advances in the diagnosis and treatment methods for CC in recent years, unfortunately, the prognosis of CC patients remains poor (4-6). Therefore, exploring the molecular mechanisms related to the tumor initiation and progression of CC is of vital importance and might provide new biomarker to predict and improve the prognosis of CC patients.

Increasing researches have demonstrated that microRNAs (miRs) are a class of small, non-coding RNAs that play critical roles in various biological cellular processes, including cell proliferation, migration, differentiation, and apoptosis (7-9). In recent decades, dysregulation of miRs have been implicated in the development and malignant progression of human cancers and might be used as potential diagnostic or therapeutic targets for cancers (10-12). To date, over 2,500 miRs have been identified in human genome (13). Among numerous cancer-related miRs, miR-153 was recently found to be dysregulated in human cancers including breast cancer (14), ovarian cancer (15), osteosarcoma (16), esophageal squamous cell carcinoma (17), gastric cancer (18), pancreatic cancer (19), non-small-cell lung cancer (20), and colorectal cancer (21). Significantly, miR-153 expression was found upregulated in colorectal cancer but downregulated in the rest cancer types mentioned above (14-21), which implies that miR-153 plays a dual role in the progression of human cancers. However, up to now, the role of miR-153 in regulating human CC cells remains unexplored. Importantly, miR-153 was found dysregulated in gynecological tumors including breast cancer and ovarian cancer (14,15). Therefore, it is worthwhile to investigate the clinical significance of miR-153 in CC.

In the present study, we found miR-153 was significantly downregulated in CC tissues and cell lines. Ectopic expression of miR-153 inhibited cell proliferation in CC cell lines. Moreover, the downregulation of miR-153 in CC patients was associated with tumor size and lymph node metastasis. Furthermore, we revealed that the downregulation of miR-153 predicts the poor prognosis of CC patients. Taken together, our data revealed that miR-153 plays a tumor suppressor role

Correspondence to: Dr Li Liu, Department of Oncology, Integrated Traditional Chinese and Western Medicine, The Central Hospital of Wuhan, Tongji Medical College, Huazhong University of Science and Technology, 26 Shengli Street, Jiang'an, Wuhan, Hubei 430012, P.R. China
E-mail: liliu8310@126.com

Key words: microR-153, cervical cancer, proliferation, prognosis

in the progression of CC and may provide a novel potential therapeutic target for CC.

Materials and methods

Clinical tissue samples. This study was approved by the Ethics Committee of The Central Hospital of Wuhan. A total of 93 pairs of CC tissues and matched noncancerous tissues were collected at the Central Hospital of Wuhan between March 2008 and September 2011. The written informed consent has been obtained from all the enrolled patients. These enrolled patients did not receive any anti-tumor treatments prior to surgical resection. Tissues were immediately snap-frozen in liquid nitrogen after surgical resection, and stored in liquid nitrogen before usage. Clinicopathological parameters of the enrolled patients were collected at the beginning of the follow-up period and summarized in Table I. The overall survival time was calculated as the time between the date of surgery and the date of mortality or last follow-up for 60 months.

Cell lines and cell culture. The human CC cell lines including C-33A, HeLa, and SiHa were purchased from Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). These cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS; HyClone; GE Healthcare Life Sciences, Logan, UT, USA), and 100 U/ml penicillin and 0.1 mg/ml streptomycin. The human cervical epithelial cell line End1 was purchased from American Type Culture Collection (ATCC, Manassas, VA, USA) and cultured in Keratinocyte-Serum Free medium (K-SFM; Invitrogen; Thermo Fisher Scientific, Inc.) with 0.1 ng/ml human recombinant EGF, 0.05 mg/ml bovine pituitary extract, and additional calcium chloride 44.1 mg/l (final concentration 0.4 mM). Cells were maintained in humidified incubator at 37°C containing 5% of CO₂.

Cell transfection. The miR-153 mimic (5'-UUGCAUAGU CACAAAAGUGAUC-3'), miR-153 inhibitor (5'-AUCACU UUUGUGACUAUGCA-3') and negative control (miR-con) (5'-UAGCUUAUCAGACUGAUGUUGA-3') were purchased from Guangzhou RiboBio Co., Ltd., (Guangzhou, China). The cells were seeded into 6-well plates at a density of 3x10⁵ cells/well. The synthetic miRNAs were transfected to the cultured cells using Lipofectamine 2000 (Invitrogen; Thermo Fisher Scientific, Inc.) according to the instructions provided by the manufacturer. Further analyses were performed 24 h post transfection.

Reverse transcriptase-quantitative polymerase chain reaction (RT-qPCR). Total RNA from the cultured cell lines was isolated using TRIzol reagent (Beyotime Institute of Biotechnology, Haimen, China). RNA was reverse transcribed to cDNA using TaqMan miRNA reverse transcription kit (Applied Biosystems; Thermo Fisher Scientific, Inc.). RT-PCR was performed using PrimeScript miRNA RT-PCR kit (Takara Biotechnology Co., Ltd., Dalian, China) according to the manufacturer's instructions. The following procedures were used: 1 cycle at 95°C for 2 min, 40 cycles at

Table I. Associations between microRNA-153 expression and clinicopathological features.

Variable	Cases (n)	miR-153 expression level		P-value
		High (n=25) %	Low (n=68) %	
Age (years)				
>50	45	12 (26.7)	33 (72.3)	0.662
<50	48	13 (27.1)	35 (71.9)	
Lymph node metastasis				
Negative	39	11 (28.2)	28 (71.8)	0.039
Positive	54	14 (25.9)	40 (74.1)	
Tumor size				
≥4 cm	55	16 (29.1)	39 (70.9)	0.018
<4 cm	38	9 (23.7)	29 (76.3)	
Clinical stage				
I-II	47	10 (21.3)	37 (78.7)	0.093
III	46	15 (32.6)	31 (67.4)	
miR-153, microRNA-153.				

95°C for 30 sec and 58°C for 40 sec. U6 snRNA was used as an internal control to normalize the expression of miR-153. The primers for miR-153 and U6 snRNA used in this study were purchased from Guangzhou RiboBio Co., Ltd. and the detailed sequences were as follows: miR-153: Forward, 5'-TTGCATAGTCACAAAAGTGAT-3', Reverse, 5'-CAG TCGTGTCTGGAGT-3'; U6 snRNA: Forward, 5'-CTC GCTTCGGCAGCACATATACT-3', Reverse, 5'-ACGCTT CACGAATTTGCGTGTC-3'. The relative expression levels were calculated using the 2^{-ΔΔC_q} method (22).

Cell proliferation assay. To assess the cell proliferation rate, MTT method was employed. Briefly, the cells were cultured in 96-well plate at the density of 3x10³ cells/well. At indicated time points (0, 24, 48, 72 h), 10 μl MTT solution (5 mg/ml; Beyotime Institute of Biotechnology) was added to each well and incubated for additional 3 h. Then, the supernatant was discarded followed by adding 200 μl DMSO to dissolve the violet formazan crystals. After incubation for 2 h, the optical density was measured at 570 nm using the Multiskan Spectrum equipment (Thermo Fisher Scientific, Inc.).

Statistical analysis. SPSS v16.0 software (SPSS, Inc., Chicago, IL, USA) was used for statistical analysis. To determine the significance of two groups and multiple groups, Student's t test and one-way ANOVA were conducted respectively. Chi-square test was used to analyze the correlation between the expression of miR-153 and clinicopathological features. The Kaplan-Meier curve and log-rank test was used to analyze the overall survival of CC patients. Univariate and multivariate analyses with Cox proportional hazards model were used to identify the independent predictors for the prognosis of CC




Figure 1. Downregulation of miR-153 in CC. (A) RT-qPCR was performed to analyze the expression of miR-153 in the CC cell lines C-33A, SiHa and HeLa, and the normal cervical epithelial cell line End1. (B) RT-qPCR was performed to analyze the expression of miR-153 in CC tissues and adjacent noncancerous tissues. **P<0.01 and ***P<0.001 vs. control (End1/Adjacent tissues). miR-153, microRNA-153; RT-qPCR, reverse transcription-quantitative polymerase chain reaction. CC, cervical cancer.




Figure 2. miR-153 overexpression inhibits the proliferation of CC cell lines. (A) Reverse transcription-quantitative polymerase chain reaction was performed to analyze the expression of miR-153 in the CC cell lines C-33A, SiHa and HeLa, following transfection with miR-153 mimic, miR-153 inhibitor and negative control. (B) An MTT assay was performed to analyze the cell proliferation of the CC cell lines (C-33A, SiHa and HeLa) and the normal cervical epithelial cell line End1. (C) An MTT assay was also performed to analyze the cell proliferation of the CC cell lines (C-33A, SiHa and HeLa) following transfection with miR-153 mimic, miR-153 inhibitor and negative control. *P<0.05, **P<0.01 and ***P<0.001 vs. control (End1/miR-con). miR-153, microRNA-153; CC, cervical cancer; miR-con, negative control miRNA; OD 570 nm, optical density at 570 nm.

patients. Data were presented as the mean \pm SD. P<0.05 was considered to indicate a statistically significant difference.

Results

MiR-153 is downregulated in CC tissues and cell lines. We measured the expression of miR-153 in CC cell lines (C-33A, HeLa, and SiHa) and normal cervical epithelial cell line End1 by RT-qPCR. As presented in Fig. 1A, the expression of miR-153 was significantly reduced in CC cell lines investigated compared with that in normal cervical epithelial cell line (all P<0.01). To further confirm the significance of miR-153 expression in CC, the expression level of miR-153 in 93 paired CC tissues and adjacent normal tissues was measured by the same method. We found the miR-153 expression was significantly lower in CC tissues compared with in adjacent normal tissues (P<0.01; Fig. 1B), which was consistent with the observations on CC cell lines and normal cervical epithelial cell line. These 93 CC patients were then classified into two groups based on the expression level of miR-153: Namely high miR-153 expression group and low miR-153

expression group. The 75th percentile of $2^{-\Delta\Delta C_q}$ was used as the cut-off point (0.72) for patients with high or low miR-153 expression (23).

MiR-153 inhibits the proliferation of CC cell lines in vitro. To explore the role of miR-153 in CC cells, miR-153 expression level was altered with miR-153 mimic and inhibitor. RT-qPCR showed that miR-153 expression was significantly enhanced in all the investigated CC cell lines transfected with miR-153 mimic compared with those transfected with miR-con (all P<0.01; Fig. 2A). Conversely, the miR-153 inhibitor transfection could reduce the expression of miR-153 in the CC cell lines investigated (all P<0.01; Fig. 2A). Cell proliferation rate was measured using MTT assay, the cell proliferation of CC cell lines was significantly higher than that of normal cervical epithelial cell (all P<0.01; Fig. 2B). Following, the cell proliferation of CC cell lines transfected with miRNAs was also measured by the same method. As shown in Fig. 2C, we found the CC cells transfected with miR-153 mimic showed obvious growth inhibition, while those transfected with miR-153 inhibitor showed obvious growth stimulation (all P<0.05).

Table II. Univariate and multivariate analyses of overall survival.

Variable	Univariate analysis			Multivariate analysis		
	HR	95% CI	P-value	HR	95% CI	P-value
miR-153	2.062	1.080-3.936	0.028	2.080	1.096-3.949	0.025
Age	1.798	0.870-3.716	0.113	-	-	-
Lymph node metastasis	1.981	1.103-3.873	0.046	2.001	1.030-3.889	0.041
Tumor size	2.021	1.047-3.904	0.036	2.043	1.065-3.922	0.032
Clinical stage	1.896	0.945-3.804	0.072	-	-	-

HR, hazard ratio; CI, confidence interval; miR-153, microRNA-153.

Clinical significance of miR-153 expression in CC. To investigate the clinical significance of miR-153, we assessed the association between miR-153 expression and clinical-pathological features. The results revealed that low miR-153 expression was associated with tumor size ($P=0.018$), lymph node metastasis ($P=0.039$) but not related to age ($P=0.662$), clinical stage ($P=0.093$) in CC patients (Table I). The Kaplan-Meier curve and log-rank test was employed to analyze the effect of miR-153 expression on 5-year overall survival of the 93 enrolled CC patients. We found the patients with high miR-153 expression ($n=25$) live longer than those with low miR-153 expression ($n=68$) ($P=0.037$; Fig. 3). Additionally, multivariate analysis indicated that miR-153 expression was an independent predictor for the overall survival in CC patients ($P=0.025$; Table II). Taken together, there was a strong correlation between low miR-153 expression and poor outcomes in CC patients.

Discussion

miRNAs function mainly through complementary binding to the 3'-UTR of target gene which can induce the degradation of its mRNA, or suppress gene transcription to reduce the expression levels of its target genes (24-26). Many studies have confirmed that miRNAs were critical players in the initiation and progression of human cancers including CC (27,28). For CC patients, the overall survival has significantly improved in the past decade due to the usage of HPV vaccination and improvement of therapeutics methods but it is still undesirable (4,5). Therefore, it is of great importance to understand the molecular mechanisms underlying the progression of CC, which are important for developing novel therapeutic strategies for patients with CC.

Although documented evidence indicates that miR-153 can function as either oncogene or tumor-suppressor in cancers, the role of miR-153 in regulating CC cells remains unexplored (14-21). In the present study, the miR-153 expression was found to be downregulated in CC tissues and cell lines. The cell proliferation rate analysis results demonstrated that the cell proliferation rate in CC cells was higher than in normal cervical epithelial cell line. To further elucidate the effect of miR-153 on the progression of CC, the synthetic miRNAs were introduced into the CC cell lines. We discovered that the overexpression of miR-153 diminished but the




Figure 3. Low expression levels of miR-153 are associated with the poor prognosis of patients with CC. Patients were divided into the miR-153 low group ($n=68$) and high group ($n=25$) to analyze the cumulative survival rate. miR-153, microRNA-153; CC, cervical cancer.

downregulate miR-153 expression increased the proliferation rate in the CC cells investigated. The above findings indicated that miR-153 functions as tumor-suppressor in CC and inhibits CC progression partly through cell proliferation inhibition. The progression of CC is a complex pathological process in which proliferation and apoptosis of CC cells serves an important role (29). The proliferation and apoptosis of CC cells are regulated by multiple molecules, including phosphatase and tension homolog (PTEN) (30), miR-21 (31), miR-940 (32), and DJ-1 (33). Therefore, the aberrant expressed status of miR-153 and the involvement of miR-153 in CC proliferation process highlighted the importance of miR-153 in CC.

Then, we analyzed the 5-year overall survival rate of the recruited CC patients. We found the low expression of miR-153 indicates a poor 5-year survival. Following, we found the low miR-153 expression was related to tumor size and lymph node metastasis. The univariate and multivariate analyses demonstrated that low miR-153 expression was an independent predictor for poor overall survival of patients with CC.

In conclusion, we provided evidence that miR-153 exert its tumor suppressor potential by inhibiting cell proliferation. Meanwhile, there was a strong correlation between low miR-153 expression and poor outcomes in CC patients, indicating miR-153 could potentially serve as a therapeutic target for CC.

Acknowledgements

The present study was supported by National Natural Science Foundation of China (grant no. 31500840); Health and Family Planning Commission of Wuhan Province (grant no. WZ17Z03).

References

- Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J and Jemal A: Global cancer statistics, 2012. *CA Cancer J Clin* 65: 87-108, 2015.
- Siegel RL, Miller KD and Jemal A: Cancer Statistics, 2017. *CA Cancer J Clin* 67: 7-30, 2017.
- Wang JM, Ju BH, Pan CJ, Gu Y, Li MQ, Sun L, Xu YY and Yin LR: MiR-214 inhibits cell migration, invasion and promotes the drug sensitivity in human cervical cancer by targeting FOXM1. *Am J Transl Res* 9: 3541-3557, 2017.
- Ellenson LH and Wu TC: Focus on endometrial and cervical cancer. *Cancer Cell* 5: 533-538, 2014.
- Yoshikawa H: Progress in the World and challenges in Japan on HPV vaccination for cervical cancer prevention. *Gan To Kagaku Ryoho* 37: 971-975, 2010 (In Japanese).
- Ghebrey RG, Grover S, Xu MJ, Chuang LT and Simonds H: Cervical cancer control in HIV-infected women: Past, present and future. *Gynecol Oncol Rep* 21: 101-108, 2017.
- Calin GA and Croce CM: MicroRNA-cancer connection: The beginning of a new tale. *Cancer Res* 66: 7390-7394, 2006.
- Bartel DP: MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell* 116: 281-297, 2004.
- Garzon R, Calin GA and Croce CM: MicroRNAs in cancer. *Annu Rev Med* 60: 167-179, 2009.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA, *et al*: MicroRNA expression profiles classify human cancers. *Nature* 435: 834-838, 2005.
- Iorio MV and Croce CM: MicroRNA dysregulation in cancer: Diagnostics, monitoring and therapeutics. A comprehensive review. *EMBO Mol Med* 9: 852, 2017.
- Esquela-Kerscher A and Slack FJ: Oncomirs-microRNAs with a role in cancer. *Nat Rev Cancer* 6: 259-269, 2006.
- Kozomara A and Griffiths-Jones S: miRBase: Annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res* 42: D68-D73, 2014.
- Wu XW, Li L, Li Y and Liu ZH: MiR-153 promotes breast cancer cell apoptosis by targeting HECTD3. *Am J Cancer Res* 6: 1563-1571, 2016.
- Zhou J, Xie M, Shi Y, Luo B, Gong G, Li J, Wang J, Zhao W, Zi Y, Wu X and Wen J: MicroRNA-153 functions as a tumor suppressor by targeting SET6 and ZEB2 in ovarian cancer cells. *Oncol Rep* 34: 111-120, 2015.
- Niu GF, Li B, Sun L and An CG: MicroRNA-153 inhibits osteosarcoma cells proliferation and invasion by targeting TGF- β 2. *PLoS One* 10: e0119225, 2015.
- Zuo J, Wang D, Shen H, Liu F, Han J and Zhang X: MicroRNA-153 inhibits tumor progression in esophageal squamous cell carcinoma by targeting SNAI1. *Tumor Biol*, 2016 (Epub ahead of print).
- Wang Z and Liu C: MiR-153 regulates metastasis of gastric cancer through SNAI1. *Tumor Biol*, 2015 (Epub ahead of print).
- Liu F, Liu B, Qian J, Wu G, Li J and Ma Z: miR-153 enhances the therapeutic effect of gemcitabine by targeting SNAI1 in pancreatic cancer. *Acta Biochim Biophys Sin (Shanghai)* 49: 520-529, 2017.
- Shan N, Shen L, Wang J, He D and Duan C: MiR-153 inhibits migration and invasion of human non-small-cell lung cancer by targeting ADAM19. *Biochem Biophys Res Commun* 456: 385-391, 2015.
- Zhang L, Pickard K, Jenei V, Bullock MD, Bruce A, Mitter R, Kelly G, Paraskeva C, Strefford J, Primrose J, *et al*: miR-153 supports colorectal cancer progression via pleiotropic effects that enhance invasion and chemotherapeutic resistance. *Cancer Res* 73: 6345-6347, 2013.
- Schmittgen TD and Livak KJ: Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc* 3: 1101-1108, 2008.
- Chen ZL, Zhao XH, Wang JW, Li BZ, Wang Z, Sun J, Tan FW, Ding DP, Xu XH, Zhou F, *et al*: microRNA-92a promotes lymph node metastasis of human esophageal squamous cell carcinoma via E-cadherin. *J Biol Chem* 286: 10725-10734, 2011.
- Wongfieng W, Jumnainsong A, Chamgramol Y, Sripan B and Leelayuwat C: 5'-UTR and 3'-UTR regulation of MICB expression in human cancer cells by novel microRNAs. *Gene (Basel)* 8: E213, 2017.
- Biegel JM, Henderson E, Cox EM, Bonenfant G, Netzband R, Kahn S, Eager R and Pager CT: Cellular DEAD-box RNA helicase DDX6 modulates interaction of miR-122 with the 5' untranslated region of hepatitis C virus RNA. *Virology* 507: 231-241, 2017.
- Xiao Y, Li X, Wang H, Wen R, He J and Tang J: Epigenetic regulation of miR-129-2 and its effects on the proliferation and invasion in lung cancer cells. *J Cell Mol Med* 19: 2172-2180, 2015.
- Gregory RI and Shiekhattar R: MicroRNA biogenesis and cancer. *Cancer Res* 65: 3509-3512, 2005.
- Calin GA and Croce CM: MicroRNA signatures in human cancers. *Nat Rev Cancer* 6: 857-866, 2006.
- Xu L, Xu Q, Li X and Zhang X: MicroRNA-21 regulates the proliferation and apoptosis of cervical cancer cells via tumor necrosis factor- α . *Mol Med Rep* 16: 4659-4663, 2017.
- Feng Y, Zou W, Hu C, Li G, Zhou S, He Y, Ma F, Deng C and Sun L: Modulation of CASC2/miR-21/PTEN pathway sensitizes cervical cancer to cisplatin. *Arch Biochem Biophys* 623-624: 20-30, 2017.
- Du G, Gao D and Meng L: miR-21 inhibitor suppresses cell proliferation and colony formation regulating the PTEN/AKT pathway and improves paclitaxel sensitivity in cervical cancer cells. *Mol Med Rep* 15: 2713-2719, 2017.
- Su K, Wang CF, Zhang Y, Cai YJ, Zhang YY and Zhao Q: miR-940 upregulation contributes to human cervical cancer progression through p27 and PTEN inhibition. *Int J Oncol*, 2017 (Epub ahead of print).
- Wang H and Gao W: DJ-1 expression in cervical carcinoma and its effects on cell viability and apoptosis. *Med Sci Monit* 22: 2943-2949, 2016.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.