MicroRNA-599 suppresses glioma progression by targeting RAB27B

YU JIANG^{1*}, XIAOHUI WANG^{2*}, JI ZHANG³ and RENCHUN LAI¹

¹State Key Laboratory of Oncology in South China, Department of Anesthesiology;

²State Key Laboratory of Oncology in South China, Department of Radiation Oncology;

³State Key Laboratory of Oncology in South China, Department of Neurosurgery, Collaborative Innovation Center for Cancer Medicine, Sun Yat-sen University Cancer Center, Guangzhou, Guangdong 510060, P.R. China

Received July 9, 2017; Accepted January 17, 2018

DOI: 10.3892/ol.2018.8727

Abstract. MicroRNAs (miRNAs/miRs) serve tumor promoting or suppressive roles in different human cancer types, including glioma; however, the regulatory underlying mechanism by which miR-599 affects glioma progression remains largely unknown. The aim of the present study was to investigate the expression of miR-599 in glioma, as well as the underlying regulatory mechanism. Reverse transcription-quantitative polymerase chain reaction and western blot analysis were used to examine mRNA and protein expression, respectively. MTT, wound healing and transwell assays were conducted to study cell proliferation, migration and invasion, respectively. A dual-luciferase reporter gene assay was used to confirm the targeting association between miR-599 and Ras-related protein Rab-27B (hereafter RAB27B). In the present study, miR-599 expression was observed to be significantly downregulated in human glioma tissues and cell lines, when compared with normal brain tissues and normal human astrocyte cells, respectively. Low miR-599 expression was significantly associated with glioma progression. Ectopic expression of miR-599 caused a significant reduction in the proliferation, migration and invasion of U-87MG Uppsala and U251 cells. Bioinformatics analysis and dual-luciferase reporter gene assay data identified that RAB27B was a direct target gene of miR-599. The mRNA and protein expression of RAB27B was significantly downregulated following miR-599 overexpression in U-87MG Uppsala and U251 cells. Rescue experiments demonstrated that RAB27B rescued the miR-599-induced

inhibition of glioma cell growth, migration and invasion. In addition, RAB27B expression was significantly upregulated in glioma tissues and cell lines, and the expression levels of RAB27B were inversely correlated with miR-599 levels in glioma tissues; therefore, the present study demonstrated that miR-599 exerts a tumor-suppressive role in glioma progression by targeting RAB27B, indicating that miR-599 may be used as a potential candidate for glioma treatment.

Introduction

Glioma is the most common type of primary brain tumor, and one of the most aggressive and lethal human malignancies (1). The majority of patients are diagnosed at an advanced stage of the disease, despite great efforts towards the early detection of glioma (1,2). The overall prognosis of patients with advanced glioma is poor (1,2). In recent years, a number of studies have focused on investigating the pathogenesis of glioma (3-5). However, the molecular mechanisms underlying glioma progression remain largely unknown; therefore, there is an urgent requirement to identify novel and effective diagnostic and therapeutic targets for this disease.

MicroRNAs (miRNAs/miRs) are a class of short, endogenous, single-stranded RNAs that have been demonstrated to function as key regulators for gene expression by directly binding to the 3'-untranslational region (3'-UTR) of target mRNAs, causing mRNA degradation or translation repression (6,7). Previous studies have demonstrated that miRNAs are involved in a variety of cellular biological processes, including development and differentiation, cell proliferation, apoptosis and migration, in addition to tumorigenesis (8,9). Complete deregulation of miRNAs has been observed in various human cancer types, indicating that an altered miRNA expression profile may contribute to the development and malignant progression of human cancer (10,11). Recently, Zhang *et al* (12) reported that miR-599 inhibited the proliferation and invasion of glioma by targeting periostin. As one miRNA can have numerous target genes, other target genes of miR-599 may also serve important roles in glioma.

Ras-related protein Rab-27B (hereafter RAB27B) is a member of the Rab protein family, which are prenylated, membrane-bound proteins involved in vesicular fusion and

Correspondence to: Professor Renchun Lai, State Key Laboratory of Oncology in South China, Department of Anesthesiology, Collaborative Innovation Center for Cancer Medicine, Sun Yat-sen University Cancer Center, 651 Dongfeng East Road, Guangzhou, Guangdong 510060, P.R. China
E-mail: lairechunductor@163.com

*Contributed equally

Key words: glioma, microRNA, Ras-related protein Rab-27B, tumor suppressor

trafficking (13). Deregulation of RAB27B has been observed in a number of common cancer types (14,15). For instance, RAB27B was determined to be significantly upregulated in ovarian cancer tissues, where it was associated with distant metastasis and poor prognosis (15). Furthermore, patients with high-grade glioma harboring RAB27B hypomethylation or overexpression exhibited a reduced survival time, indicating that the upregulation of RAB27B contributes to glioma progression and poor patient prognosis (16); however, whether miR-599 regulates RAB27B expression in glioma remains unclear.

Therefore, the present study aimed to investigate the clinical significance of miR-599 and RAB27B expression in glioma. The regulatory mechanism of miR-599 and RAB27B underlying glioma progression was studied.

Materials and methods

Clinical tissue samples. The present study was approved by the Ethics Committee of Sun Yat-sen University Cancer Center (Guangzhou, China). A total of 50 glioma tissues were collected from patients with primary glioma at Sun Yat-sen University Cancer Center between April 2014 and March 2016, as were 13 matched normal brain tissues. The age of patients with normal brain tissue ranged between 34–58 years, with 8 male and 5 female. The 50 patients with glioma included 29 men and 21 women, aged 14–69 years old, with a mean age of 43.3 years old. Written informed consent was obtained from all patients. All tissues were pathologically confirmed at Sun Yat-sen University Cancer Center, and necrosis in glioma tissues was also evaluated pathologically. The patients with glioma were classified according to the World Health Organization criteria and staged according to the Tumor-Node-Metastasis classification (17). Following surgical resection, tissues were immediately snap-frozen in liquid nitrogen. The clinicopathological characteristics of these patients with glioma are summarized in Table I.

Cell culture. Normal human astrocyte (NHA) cells were purchased from the American Type Cell Culture Collection (Manassas, VA, USA). Human glioma cell lines (U-373MG Uppsala, U-87MG Uppsala, U251 and T98G) were obtained from the Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). Cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS) (Thermo Fisher Scientific, Inc.) at 37°C in a humidified incubator containing 5% CO₂.

Cell transfection. For miR-599 and RAB27B functional analysis, U-87MG Uppsala and U251 cells were transfected with 100 nM negative control miRNA (miR-NC), miR-599 mimic, NC inhibitor, or miR-599 inhibitor (all purchased from Guangzhou Fugen Co., Ltd., Guangzhou, China), or co-transfected with miR-599 mimic and RAB27B plasmid (generated by Yearthbio, Changsha, China), or with miR-599 mimic and blank pcDNA3.1 vector, using Lipofectamine® 2000 (Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol. The groups in the present study were as follows: miR-NC, miR-599 mimic, NC inhibitor, miR-599 inhibitor, miR-599 mimic+RAB27B plasmid and miR-599

mimic+blank pcDNA3.1 vector. Subsequent experimentation were conducted 48 h following transfection.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Total RNA was extracted from tissues and cells (NHA, U-373MG Uppsala, U-87MG Uppsala, U251 and T98G) using TRIzol® reagent (Thermo Fisher Scientific, Inc.). RevertAid RT Reverse Transcription kit (Thermo Fisher Scientific, Inc.) was used to convert 1 µg RNA into cDNA. To detect the expression of miR-599, qPCR was conducted using the All-in-One™ miRNA qRT-PCR detection kit (GeneCopoeia, Inc., Rockville, MD, USA), according to the manufacturer's instruction. U6 was used as internal reference. The primers for miR-599 and U6 were directly purchased from Fulengen, Guangzhou, China. For detecting mRNA expression, qPCR was conducted using the SYBR Green qPCR Master mix (Bio-Rad Laboratories, Inc., Hercules, CA, USA), according to the manufacturer's instruction. GAPDH was used as internal reference. The primer sequences were as follows: RAB27B, forward, TAGACTTTCGGGAAAAACGTGTG, and reverse, AGAAGCTCTGTTGACTGGTGA; GAPDH, forward, GGAGCGAGATCCCTCCAAAAT, and reverse, GGCTGTTGTCATACTTCTCATGG. The thermocycling conditions were as follows: Denaturation at 95°C for 5 min followed by 35 cycles of denaturation at 95°C for 15 sec and annealing/elongation at 60°C for 30 sec. The relative expression was analyzed using the 2^{-ΔΔC_q} method (18).

Western blot analysis. Cells (NHA, U-373MG Uppsala, U-87MG Uppsala, U251 and T98G) were solubilized in cold radioimmunoprecipitation lysis buffer (Beyotime Institute of Biotechnology, Shanghai, China). The protein concentration was detected using a BCA Protein Assay kit (Pierce; Thermo Fisher Scientific, Inc.), according to the manufacturer's instruction. Proteins (60 µg/lane) were separated by 12% SDS-PAGE and were then transferred onto a polyvinylidene difluoride (PVDF) membrane (Thermo Fisher Scientific, Inc.). The PVDF membrane was then incubated with PBS (Thermo Fisher Scientific, Inc.) containing 5% non-fat milk at room temperature for 3 h. Following three washes with PBS, the PVDF membrane was incubated with rabbit polyclonal anti-RAB27B antibody (dilution, 1:50; cat. no. ab103418; Abcam, Cambridge, MA, USA) or rabbit polyclonal anti-GAPDH antibody (dilution, 1:50; cat. no. ab9485; Abcam) at room temperature for 3 h. After being washed with PBS three times, the PVDF membrane was then incubated with the horseradish peroxidase-conjugated goat anti-rabbit monoclonal secondary antibody (dilution, 1:5,000; cat. no. ab6721; Abcam) at room temperature for 40 min. An Enhanced Chemiluminescence Western Blotting kit (Pierce; Thermo Fisher Scientific, Inc.) was used to detect immune complex on the PVDF membrane. Protein expression was analyzed using Image-Pro Plus software 6.0 (Media Cybernetics, Inc., Rockville, MD, USA). GAPDH was used as an internal control.

MTT assay. To analyze cellular proliferation, 5×10⁴ U-87MG Uppsala and U251 cells were cultured in a 96-well plate with 100 µl of DMEM containing 0.5 g/l MTT (Thermo Fisher Scientific, Inc.). Following this, U251 cells were cultured at 37°C for 12, 24, 48 or 72 h, after which the medium was

Table I. Association between miR-599 expression and clinicopathological characteristics of patients with glioma.

Variables	Patients, n	High miR-599, n	Low miR-599, n	P-value
Total	50	18	32	
Age, years				0.774
<55	26	10	16	
≥55	24	8	16	
Sex				0.551
Male	29	9	20	
Female	21	9	12	
Tumor size, cm				0.073
<5	31	8	23	
≥5	19	10	9	
Necrosis				0.033 ^a
Yes	19	3	16	
No	31	15	16	
Clinical stage				0.018 ^a
I-II	27	14	13	
III-IV	23	4	19	

^aP<0.05. miR-599, microRNA-599.

removed. Next, 50 μ l dimethyl sulfoxide (Thermo Fisher Scientific, Inc.) was added. Following incubation at 37°C for 10 min, the A570 of each sample was measured at a wavelength of 570 nm using an enzyme immunoassay analyzer (Tecan Infinite® M200; Tecan Group Ltd., Männedorf, Switzerland).

Wound-healing assay. U-87MG Uppsala and U251 cells were cultured to full confluence and a wound of ~1 mm width was created using a plastic scribe. Next, cells were washed in DMEM and incubated in serum-free DMEM at 37°C for 24 h. For the negative control, cells were fixed using 90% ethanol at room temperature for 10 min and observed under an inverted microscope (magnification, x40; Olympus Corporation, Tokyo, Japan). Following this, cells were incubated at 37°C in DMEM supplemented with 10% FBS for 48 h. Next, cells were fixed using 90% ethanol at room temperature for 10 min and observed under an inverted microscope (magnification, x40; Olympus Corporation). This experiment was repeated 3 times.

Invasion assay. Transwell chambers pre-coated with Matrigel (BD Biosciences, Franklin Lakes, NJ, USA) were used to conduct cell invasion analysis. The U-87MG Uppsala and U251 cell suspension (3×10^5 cells/ml) was prepared in DMEM. Next, 300 μ l cell suspension was added into the upper chamber, and 300 μ l DMEM supplemented with 10% FBS was added into the lower chamber. Following incubation at 37°C for 24 h, the cells that did not invade through the membrane were removed using a cotton-tipped swab. Cells that invaded the membrane were fixed in 90% ethyl alcohol at room temperature for 10 min, and then stained with 0.1% crystal violet (Beyotime Institute of Biotechnology) at room temperature for 10 min. The invaded cells were counted under an inverted microscope (magnification, x400).

Bioinformatics prediction. Targetscan 7.1 software (http://www.targetscan.org/vert_71/) was used to predict the potential target genes of miR-599. The search terms 'Human' and 'miR-599' were used.

Dual-luciferase reporter assay. The wild type (WT) RAB27B 3'-UTR containing the predicted binding sequences of miR-599 was constructed using PCR, which was then subcloned downstream of the *Renilla* luciferase gene in the psiCHECK-2 vector (Promega Corporation, Madison, WI, USA). The mutant type (MT) RAB27B 3'-UTR lacking the binding sequences of miR-599 was constructed using the Directed Mutagenesis kit (Agilent Technologies, Inc., Santa Clara, CA, USA), which was also subcloned downstream of the *Renilla* luciferase gene in the psiCHECK-2 vector. U-87MG Uppsala and U251 cells were then co-transfected with WT or MT RAB27B 3'-UTR plasmid, and miR-599 mimic or miR-negative control (NC), using Lipofectamine 2000 according to the manufacturer's instruction. Following transfection for 48 h, the luciferase activity was determined using the Dual-Luciferase Reporter Assay system (Promega Corporation) on an Lmax multiwell luminometer (Molecular Devices, LLC, Sunnyvale, CA, USA), according to the manufacturer's instruction. The *Renilla* luciferase activity was normalized to firefly luciferase activity.

Statistical analysis. The data are presented as the mean $\pm$ standard deviation of three independent experiments. Statistical analysis was conducted using the SPSS Graduate Pack, version 19.0 (IBM Corp., Armonk, NY, USA). Differences were analyzed using a Student's t-test for two-group comparison or a one-way analysis of variance for multiple-group comparison followed by Tukey's post hoc test. The correlation between the expression of miR-599 and

Figure 1. Downregulation of miR-599 expression in glioma. (A) RT-qPCR data indicated that glioma tissues exhibited significantly lower expression of miR-599 than normal brain tissues. ** $P < 0.01$ vs. normal. (B) RT-qPCR data indicated that miR-599 expression was significantly downregulated in glioma cell lines, compared with NHA cells. ** $P < 0.01$ vs. NHA. NHA, normal human astrocyte; miR-599, microRNA-599; RT-qPCR, reverse transcription-quantitative polymerase chain reaction.

RAB27B was analyzed using Pearson's correlation analysis. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Downregulation of miR-599 expression is associated with glioma progression. RT-qPCR analysis was conducted to determine the expression of miR-599 in glioma tissues, which was then compared to that in normal brain tissues. These data indicated that glioma tissues exhibited significantly lower expression levels of miR-599, than normal brain tissues (Fig. 1A). Furthermore, miR-599 expression was also significantly downregulated in glioma cell lines, compared with NHA cells (Fig. 1B).

Next, the clinical significance of miR-599 expression in glioma was studied. Patients with glioma were divided into high-miR-599-expression and low-miR-599-expression groups according to the mean value of miR-599 expression. Low expression of miR-599 was significantly associated with necrosis and advanced clinical stage in glioma (Table I).

miR-599 could inhibit glioma cell proliferation, migration and invasion. As miR-599 expression was significantly downregulated in glioma, U-87MG Uppsala and U251 cells were transfected with miR-599 mimic to upregulate its expression. Following transfection, the miR-599 levels were significantly increased compared with the control group; however, transfection with scramble miRNA mimic did not affect the miR-599 expression in U-87MG Uppsala and U251 cells (Fig. 2A). The proliferative, migratory and invasive capacities of the transfected cells were examined using MTT, wound healing and transwell assays, respectively. As depicted in Fig. 2B-D, the overexpression of miR-599 caused a significant reduction in U-87MG Uppsala and U251 cell proliferation, migration and invasion. These data demonstrated that miR-599 could inhibit glioma cell proliferation, migration and invasion.

RAB27B is a target gene of miR-599 in glioma cells. As miRNAs function by mediating the expression of their target

genes, Targetscan software was used to predict the potential target genes of miR-599. The 3'-UTR of RAB27B mRNA contained the binding sequences of miR-599 (Fig. 3A). To confirm this prediction, the WT RAB27B-3'-UTR and MT RAB27B 3'-UTR luciferase reporter plasmids were generated (Fig. 3A). A dual-luciferase reporter gene assay was then conducted in U-87MG Uppsala and U251 cells. As depicted in Fig. 3B, the luciferase activity was significantly decreased in cells co-transfected with miR-599 mimics and WT RAB27B 3'-UTR luciferase reporter plasmid, which was eliminated by transfection with the WT RAB27B 3'-UTR luciferase reporter plasmid. These results indicated that miR-599 could directly bind to the 3'-UTR of RAB27B mRNA in U-87MG Uppsala and U251 cells. Following this, the regulatory effect of miR-599 on the expression of RAB27B in U-87MG Uppsala and U251 cells was assessed. As depicted in Fig. 3C and D, the mRNA and protein levels of RAB27B were significantly downregulated following overexpression of miR-599. To confirm these findings, U-87MG Uppsala and U251 cells were transfected with miR-599 inhibitor or NC inhibitor. Following transfection, the miR-599 levels were significantly downregulated in the miR-599 inhibitor group, compared with the control group; however, transfection with the NC inhibitor did not affect its expression (Fig. 3E). It was determined that transfection with miR-599 inhibitor caused an upregulation of RAB27B expression, when compared with the NC inhibitor group (Fig. 3F and G). Therefore, the expression of RAB27B was negatively mediated by miR-599 in U-87MG Uppsala and U251 cells.

RAB27B rescues the inhibitory effects of miR-599 on the malignant phenotypes of glioma cells. As it was determined that overexpression of miR-599 resulted in inhibitory effects on the proliferation, migration and invasion of U-87MG Uppsala and U251 cells, accompanied by the significant downregulation of RAB27B, a rescue experiment was conducted to assess whether RAB27B was involved in the miR-599-mediated malignant phenotypes of U-87MG Uppsala and U251 cells. miR-599-overexpressing U-87MG Uppsala and U251 cells were transfected with a RAB27B expression plasmid to upregulate its expression levels. Following transfection, the mRNA and

Figure 2. miR-599 may inhibit glioma cell proliferation, migration and invasion. U-87MG Uppsala and U251 cells were transfected with miR-599 mimic or miR-NC. Non-transfected U-87MG Uppsala and U251 cells were used as the control group. (A) Reverse transcription-quantitative polymerase chain reaction analysis was conducted to examine the expression of miR-599. (B) MTT, (C) wound healing and (D) transwell invasion assays were conducted to examine cellular proliferation, migration and invasion, respectively. Magnification for wound healing assay, x40; magnification for transwell assay: x400. **P<0.01 vs. miR-NC. miR-NC, microRNA-negative control; OD, optical density.

protein levels of RAB27B were significantly increased in the miR-599+RAB27B group, compared with the miR-599+blank

group (Fig. 4A and B). Next, the cell proliferation, migration and invasion capacities was examined using MTT, wound-healing

Figure 3. RAB27B is a target gene of miR-599 in glioma cells. (A) WT RAB27B 3'-UTR and MT RAB27B 3'-UTR luciferase reporter plasmids were generated from the present study. (B) Dual luciferase reporter gene assay was then conducted in U-87MG Uppsala and U251 cells. ** $P < 0.01$ vs. Control. (C) RT-qPCR and (D) western blot analysis were used to determine the mRNA and protein levels of RAB27B in U-87MG Uppsala and U251 cells transfected with miR-599 mimic or scramble miRNA mimic (miR-NC). ** $P < 0.01$ vs. miR-NC. Following this, U-87MG Uppsala and U251 cells were transfected with miR-599 inhibitor or negative control (NC) inhibitor, and non-transfected U-87MG Uppsala and U251 cells were used as the control group. (E) RT-qPCR was conducted to examine the miR-599 expression levels. (F) RT-qPCR and (G) western blot analysis were used to determine the mRNA and protein levels of RAB27B. ** $P < 0.01$ vs. NC inhibitor. 3'-UTR, 3'-untranslated region; WT, wild type; MT, mutant type; RT-qPCR, reverse transcription-polymerase chain reaction; miR-NC, microRNA-negative control.

and transwell assays, respectively. As depicted in Fig. 4C-E, the proliferation, migration and invasion of U-87MG Uppsala and U251 cells were significantly upregulated in the miR-599+RAB27B group, compared with the miR-599+blank group, indicating that RAB27B rescued the inhibition effects of miR-599 on the malignant phenotypes of glioma cells. These data indicated that RAB27B acted as a downstream effector of miR-599 in U-87MG Uppsala and U251 cells.

RAB27B is upregulated in glioma. The expression of RAB27B in glioma tissues and cell lines was studied. As depicted in Fig. 5A, RT-qPCR data demonstrated that RAB27B was significantly upregulated in glioma tissues, compared with normal brain tissues. Similarly, the mRNA and protein expression of RAB27B was also increased in glioma cell lines, compared with NHA cells (Fig. 5B and C). Notably, an inverse correlation was observed between the expression of RAB27B and

Figure 4. RAB27B rescues the inhibitory effects of miR-599 on the malignant phenotypes of glioma cells. miR-599-overexpressing U-87MG Uppsala and U251 cells were transfected with RAB27B expression plasmid or a blank vector. (A) Reverse transcription-quantitative polymerase chain reaction and (B) western blot analysis were used to determine the mRNA and protein levels of RAB27B. (C) MTT, (D) wound healing and (E) transwell assays were conducted to examine cellular proliferation, migration and invasion, respectively. Magnification for wound healing assay, $\times 40$; magnification for transwell assay, $\times 400$. ** $P < 0.01$ vs. miR-599+blank. miR-599, microRNA-599; OD, optical density.

Figure 5. RAB27B is upregulated in glioma. (A) RT-qPCR data demonstrated that RAB27B was significantly upregulated in glioma tissues, compared with normal brain tissues. ** $P < 0.01$ vs. normal. (B) RT-qPCR and (C) western blot analysis were used to determine the mRNA and protein expression levels of RAB27B in the indicated glioma cell lines and NHA cells. ** $P < 0.01$ vs. NHA. (D) An inverse correlation was observed between the expression of RAB27B and miR-599 in glioma tissues. RT-qPCR, reverse transcription-quantitative polymerase chain reaction; NHA, normal human astrocyte.

miR-599 in glioma tissues (Fig. 5D), which indicated that the reduced miR-599 expression may contribute to the increased expression of RAB27B in glioma tissues.

Discussion

The underlying mechanism by which miR-599 regulates glioma progression remains largely unclear. The present study determined that low miR-599 expression was significantly associated with glioma progression. Ectopic expression of miR-599 reduced glioma cell proliferation, migration and invasion. Bioinformatic analysis identified that RAB27B was a direct target gene of miR-599, and its expression was suppressed by miR-599 in glioma cells. Overexpression of RAB27B rescued the miR-599-induced inhibition of glioma cell proliferation, migration and invasion. Additionally, RAB27B expression was significantly upregulated in glioma, and inversely correlated to the miR-599 expression in glioma tissues.

A previous study indicated that miR-599 inhibits the proliferation and migration of vascular smooth muscle

cells by targeting transforming growth factor $\beta 2$ (19); however, there are a limited number of studies regarding the function of miR-599 in human cancer. Tian *et al* (20) reported that miR-599 was upregulated in patients with non-small cell lung cancer, and promoted cancer cell proliferation and invasion by directly targeting special AT-rich sequence-binding protein 2. Chi *et al* (21) determined that miR-599 could directly inhibit the expression of inositol polyphosphate-4-phosphatase type II B, and thus served a tumor-suppressive role in melanoma. Recently, Zhang *et al* (12) reported that the expression of miR-599 was reduced in 33 glioma tissues, compared with adjacent normal brain tissues. The present study also demonstrated that miR-599 was significantly downregulated in 67 glioma tissues, compared with normal brain tissues. Furthermore, the present study and that by Zhang *et al* (12) indicated that the increased expression of miR-599 was significantly associated with the malignant progression of glioma. In addition, Zhang *et al* (12) demonstrated that miR-599 could inhibit glioma cell proliferation and invasion by targeting

periostin expression. The present study also indicated that overexpression of miR-599 could inhibit glioma U251 cellular proliferation, migration and invasion.

Further investigation identified that RAB27B was a novel target gene of miR-599, and the expression levels of RAB27B was negatively affected by miR-599 in U-87MG Uppsala and U251 cells. RAB27B, a member of the Rab protein family, was previously reported to act as an oncogene in several common types of cancer (22,23). High expression of RAB27B is associated with poor prognosis of patients with breast cancer, and it promotes the invasive growth and metastasis in estrogen receptor-positive breast cancer cells (22). Pu *et al* (23) demonstrated that miR-193a-3p and miR-193a-5p suppressed the metastasis of osteosarcoma cells via inhibition of RAB27B expression. Wang *et al* (16) reported that high expression of RAB27B was associated with the poor prognosis of patients with glioma, and knockdown of RAB27B significantly inhibited glioma U-87MG Uppsala and LN229 cell invasion, possibly owing to a reduction in matrix metalloproteinase 9 expression and activation. Additionally, Wang *et al* (16) demonstrated that inhibition of RAB27B expression also reduced the tumor growth *in vivo*; their data also indicated that RAB27B acts as an oncogene in glioma. The present study determined that overexpression of miR-599 caused a reduction in the expression of RAB27B in U251 cells, which further confirmed that miR-599 could directly bind to RAB27B mRNA and repress its translation. RAB27B overexpression rescued the inhibitory effects of miR-599 on the proliferation, migration and invasion of U251 cells, indicating that RAB27B may be involved in the miR-599-mediated malignant phenotypes of glioma cells. Additionally, it was also determined that RAB27B was significantly upregulated in glioma, consistent with the previous report (16). The upregulation of RAB27B may be partly due to the reduced expression of miR-599; an inverse correlation was observed between miR-599 and RAB27B expression in glioma tissues.

In summary, to the best of our knowledge the present study is the first to have demonstrated that miR-599 served a suppressive role in regulating the proliferation, migration and invasion of glioma cells, at least in part, by directly targeting RAB27B. The present study indicated that miR-599 may represent a potential therapeutic candidate for glioma treatment.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

RL designed this study and revised this manuscript. YJ collected clinical tissues, performed statistical analysis and wrote the manuscript. XW and JZ performed the experiments.

Ethics approval and consent to participate

The present study was approved by the Ethics Committee of Sun Yat-sen University Cancer Center (Guangzhou, China).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J and Jemal A: Global cancer statistics, 2012. *CA Cancer J Clin* 65: 87-108, 2015.
2. Anjum K, Shagufta BI, Abbas SQ, Patel S, Khan I, Shah SAA, Akhter N and Hassan SSU: Current status and future therapeutic perspectives of glioblastoma multiforme (GBM) therapy: A review. *Biomed Pharmacother* 92: 681-689, 2017.
3. Liu ZJ, Liu HL, Zhou HC and Wang GC: TIPE2 inhibits hypoxia-induced wnt/beta-catenin pathway activation and EMT in Glioma cells. *Oncol Res* 24: 255-261, 2016.
4. Li Z, Xu C, Gao M, Ding B, Wei X and Ji N: Reduced expression of Jumonji, AT-rich interactive domain 2 (JARID2) in glioma inhibits tumor growth in vitro and in vivo. *Oncol Res* 25: 365-372, 2017.
5. Wang S, Hui Y, Li X and Jia Q: Silencing of lncRNA-CCDC26 restrains the growth and migration of glioma cells in vitro and in vivo via targeting miR-203. *Oncol Res*: Jun 9, 2017 (Epub ahead of print).
6. Ambros V: The functions of animal microRNAs. *Nature* 431: 350-355, 2004.
7. Ji S, Zhang B, Kong Y, Ma F and Hua Y: MiR-326 inhibits gastric cancer cell growth through down regulating NOB1. *Oncol Res* 25: 853-861, 2017.
8. Li H, Xiang Z, Liu Y, Xu B and Tang J: MicroRNA-133b inhibits proliferation, cellular migration, and invasion via targeting LASP1 in hepatocarcinoma cells. *Oncol Res* 25: 1269-1282, 2017.
9. Liu X, Li J, Yu Z, Sun R and Kan Q: MiR-935 promotes liver cancer cell proliferation and migration by targeting SOX7. *Oncol Res* 25: 427-435, 2017.
10. Wang G, Fu Y, Liu G, Ye Y and Zhang X: miR-218 inhibits proliferation, migration, and EMT of gastric cancer cells by targeting WASF3. *Oncol Res* 25: 355-364, 2017.
11. Chen X and Chen J: MiR-3188 regulates cell proliferation, apoptosis, and migration in breast cancer by targeting TUSC5 and regulating the p38 MAPK signaling pathway. *Oncol Res*: May 26, 2017 (Epub ahead of print).
12. Zhang T, Ma G, Zhang Y, Huo H and Zhao Y: miR-599 inhibits proliferation and invasion of glioma by targeting periostin. *Biotechnol Lett* 39: 1325-1333, 2017.
13. Chen D, Guo J, Miki T, Tachibana M and Gahl WA: Molecular cloning and characterization of rab27a and rab27b, novel human rab proteins shared by melanocytes and platelets. *Biochem Mol Med* 60: 27-37, 1997.
14. Yang X, Ye X, Sun L, Gao F, Li Y, Ji X, Wang X, Feng Y and Wang X: Downregulation of serum RAB27B confers improved prognosis and is associated with hepatocellular carcinoma progression through PI3K-AKT-P21 signaling. *Oncotarget* 8: 61118-61132, 2017.
15. Ren P, Yang XQ, Zhai XL, Zhang YQ and Huang JF: Overexpression of Rab27B is correlated with distant metastasis and poor prognosis in ovarian cancer. *Oncol Lett* 12: 1539-1545, 2016.
16. Wang H, Wang Y, Bao Z, Zhang C, Liu Y, Cai J and Jiang C: Hypomethylated Rab27b is a progression-associated prognostic biomarker of glioma regulating MMP-9 to promote invasion. *Oncol Rep* 34: 1503-1509, 2015.
17. Rogers TW, Toor G, Drummond K, Love C, Field K, Asher R, Tsui A, Buckland M and Gonzales M: The 2016 revision of the WHO classification of central nervous system tumours: Retrospective application to a cohort of diffuse gliomas. *J Neurooncol* 137: 181-189, 2018.

18. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
19. Xie B, Zhang C, Kang K and Jiang S: miR-599 inhibits vascular smooth muscle cells proliferation and migration by targeting TGFB2. *PLoS One* 10: e0141512, 2015.
20. Tian W, Wang G, Liu Y, Huang Z, Zhang C, Ning K, Yu C, Shen Y, Wang M, Li Y, *et al*: The miR-599 promotes non-small cell lung cancer cell invasion via SATB2. *Biochem Biophys Res Commun* 485: 35-40, 2017.
21. Chi MN, Guo ST, Wilmott JS, Guo XY, Yan XG, Wang CY, Liu XY, Jin L, Tseng HY, Liu T, *et al*: INPP4B is upregulated and functions as an oncogenic driver through SGK3 in a subset of melanomas. *Oncotarget* 6: 39891-39907, 2015.
22. Hendrix A, Maynard D, Pauwels P, Braems G, Denys H, Van den Broecke R, Lambert J, Van Belle S, Cocquyt V, Gespach C, *et al*: Effect of the secretory small GTPase Rab27B on breast cancer growth, invasion, and metastasis. *J Natl Cancer Inst* 102: 866-880, 2010.
23. Pu Y, Zhao F, Cai W, Meng X, Li Y and Cai S: MiR-193a-3p and miR-193a-5p suppress the metastasis of human osteosarcoma cells by down-regulating Rab27B and SRR, respectively. *Clin Exp Metastasis* 33: 359-372, 2016.

RETRACTED