# The mitochondrial one-carbon metabolic pathway is associated with patient survival in pancreatic cancer

KOZO NOGUCHI<sup>1,2\*</sup>, MASAMITSU KONNO<sup>2\*</sup>, JUN KOSEKI<sup>3</sup>, NAOHIRO NISHIDA<sup>2</sup>, KOICHI KAWAMOTO<sup>1</sup>, DAISAKU YAMADA<sup>1</sup>, TADAFUMI ASAOKA<sup>1</sup>, TAKEHIRO NODA<sup>1</sup>, HIROSHI WADA<sup>1</sup>, KUNIHITO GOTOH<sup>1</sup>, DAISUKE SAKAI<sup>2</sup>, TOSHIHIRO KUDO<sup>2</sup>, TAROH SATOH<sup>2</sup>, HIDETOSHI EGUCHI<sup>1</sup>, YUICHIRO DOKI<sup>1</sup>, MASAKI MORI<sup>1</sup> and HIDESHI ISHII<sup>2,3</sup>

Departments of <sup>1</sup>Gastroenterological Surgery, <sup>2</sup>Frontier Science for Cancer and Chemotherapy, and <sup>3</sup>Cancer Profiling Discovery, Graduate School of Medicine, Osaka University, Osaka 565-0871, Japan

Received February 24, 2016; Accepted March 17, 2017

DOI: 10.3892/ol.2018.8795

Abstract. The expression levels of one-carbon metabolic enzymes were investigated and observed to be correlated with clinicopathological parameters in patients with pancreatic cancer. Mitochondrial one-carbon metabolism comprises a network of biological reactions that integrate nutrient status with nucleotide synthesis, amino acid metabolism, antioxidant reduced nicotinamide adenine dinucleotide phosphate production and epigenetic methylation processes. Previous studies have reported that the hyper-activation of mitochondrial one-carbon metabolism serves a significant role in malignant cancer phenotypes. A total of 103 patients underwent surgical resection of pancreatic ductal adenocarcinomas (PDAC) at Osaka University Hospital between April 2007 and December 2013 and were enrolled in this study. Subsequently, the expression of the one-carbon metabolic enzymes methylenetetrahydrofolate dehydrogenase 2 (MTHFD2), aldehyde dehydrogenase 1 family member L2 (ALDH1L2), and serine hydroxymethyltransferase (SHMT2) was examined using immunohistochemical analysis. The immunohistochemical analyses demonstrated that patients with high expression levels of MTHFD2, ALDH1L2 or SHMT2 had significantly poor overall survival (OS) and disease-free survival (DFS)

*Correspondence to:* Professor Masaki Mori, Department of Gastroenterological Surgery, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan E-mail: mmori@gesurg.med.osaka-u.ac.jp

Professor Hideshi Ishii, Department of Frontier Science for Cancer and Chemotherapy, Graduate School of Medicine, Osaka University, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan E-mail: hishii@cfs.med.osaka-u.ac.jp

\*Contributed equally

*Key words:* pancreatic cancer, one-carbon metabolism, MTHFD2, ALDH1L2, SHMT2, folate pathway

rates, as compared with patients with low expression levels. Furthermore, multivariate Cox proportional hazards analysis indicated that MTHFD2 and ALDH1L2 were independent prognostic factors for OS and DFS, whereas SHMT2 was not predictive of DFS. However, high and low expression levels of all three folate metabolic enzymes were significantly associated with improved OS and DFS, compared with the high expression of one or two folate metabolic enzymes. The expression levels of mitochondrial one-carbon metabolic enzymes are independent prognostic factors and potential therapeutic targets for future pancreatic cancer treatments.

### Introduction

Pancreatic ductal carcinoma is the fourth most common cause of cancer-associated mortality in the United States of America (1). Despite recent advances in treatment modalities, the 5-year overall survival (OS) rate of pancreatic cancer patients is <5% (2), reflecting the aggressive invasion and early metastasis of the disease; the presence of extra-pancreatic dissemination at diagnosis is typical for patients with pancreatic cancer (3-5). Although a number of molecules have been identified to serve roles in the progression and metastasis of pancreatic cancer, the underlying molecular mechanisms remain unclear.

One-carbon metabolism is a network of biological reactions that serve critical roles in DNA methylation and DNA synthesis, and facilitate cross talk between genetic and epigenetic processes (6) (Fig. 1). One-carbon metabolism is also referred to as folate-mediated one-carbon metabolism and can impact genetic and epigenetic pro-carcinogenic processes, reflecting critical roles in DNA methylation and DNA synthesis. Methylenetetrahydrofolate dehydrogenase 2 (MTHFD2) is an enzyme of the mitochondrial folate metabolic pathway that functions as a methylenetetrahydrofolate dehydrogenase and a cyclohydrolase (6). Previous studies demonstrated that MTHFD2 was specifically upregulated in various types of cancer, compared with in normal tissues, and MTHFD2 has been identified as a novel drug target with the potential to block cancer cell migration and invasion in breast cancer and melanoma (7,8). In the present study, the expression of MTHFD2 and the coupling enzymes aldehyde dehydrogenase 1 family member L2 (ALDH1L2) and serine hydroxymethyltransferase (SHMT2), and the clinicopathological significance of these mitochondrial folate metabolic pathway enzymes in pancreatic cancer were investigated.

#### Materials and methods

Patients and specimens. Between April 2007 and December 2013, a total of 103 patients underwent surgical resection for pancreatic ductal adenocarcinoma (PDAC) at the Department of Surgery at Osaka University Hospital (Osaka, Japan) and all patients were enrolled in the present study (mean age,  $67.2\pm9.6$ , range 38 to 84 years; 64 males and 39 females; Table I). The clinicopathological characteristics of the enrolled patients are presented in Table I. Patients who underwent preoperative chemotherapy were included in this study. The tumor stages were determined according to the 7th edition of the American Joint Committee on Cancer/Unio Internationalis Contra Cancrum (UICC) tumor-node-metastasis classification (9). Two pathologists examined all histological slides, and the immunohistochemical (IHC) diagnoses of tumors and degrees of differentiation were determined. The median follow-up time was 60.9 months (range, 2 to 72 months), and the regular follow-up included measurements of carcinoembryonic antigen (CEA) and carbohydrate 19-9 (CA19-9). The present study was approved by an institutional review board of Osaka University School of Medicine (approved by Professor Y. Kaneda) and written informed consent was provided by all patients.

Immunohistochemistry. IHC analyses were performed as described previously (10). Briefly, surgical tissue specimens were fixed in 10% formaldehyde, embedded in paraffin and cut into  $3.5-\mu m$  sections. The sections were then deparaffinized in xylene, incubated at 95°C for 10 min with antigen retrieval buffer (cat no. CTS014; Funakoshi Co., Ltd., Tokyo, Japan) and then incubated with the following specific antibodies overnight at 4°C: Anti-MTHFD2 (cat no. ab176016; rabbit polyclonal; dilution, 1:100), anti-ALDH1L2 (cat no. ab170176; rabbit polyclonal; dilution, 1:200) or anti-SHMT2 (cat no. ab64417; rabbit polyclonal; dilution, 1:300; all Abcam, Cambridge, MA, USA). The sections were subsequently visualized using avidin-biotin complex reagents by direct addition (no dilution; ABC-HRP kit; Vector Laboratories, Inc., Burlingame, CA, USA) and diaminobenzidine. The reaction was performed for 30 min at 23°C. Two reviewers performed IHC analyses independently in a blinded manner. All IHC analyses were performed using a series of tissue sections for each antibody.

*Quantification of immunostaining parameters*. IHC analyses of MTHFD2, ALDH1L2 and SHMT2 were performed at x200 magnification using a light microscope. The samples were scored according to the intensity of cytoplasmic staining for MTHFD2 and were categorized as follows: 0, no staining; 1, weak (weaker staining than the positive control); 2, strong (equal to or stronger than the positive control, which included advanced colorectal cancer tissues or normal pancreatic acini tissues). IHC analyses of ALDH1L2 and MTHFD2 were scored in the same manner. The positively stained cells were counted in four representative fields of tumor regions at x100

magnification. Patients with scores of 1 or 0 were included in the low expression group and those with scores of 2 were included in the high expression group.

Statistical analysis. The associations between MTHFD2, ALDH1L2 and SHMT2 expression levels and other parameters were identified using chi-squared tests, Fisher's exact tests or independent t-tests as appropriate. The OS and disease-free survival (DFS) rates were estimated using the Kaplan-Meier estimator method and were compared using the log-rank test. Variables that were identified as significant in univariate analyses were included in subsequent Cox proportional hazards regression models. All statistical analyses were performed using JMP Statistical Software (version 11; SAS Institute Inc., Cary, NC, USA). P<0.05 was considered to indicate a statistically significant difference.

## Results

IHC analyses of one-carbon metabolism enzymes in PDAC. The clinicopathological characteristics of the patients included in the current study are presented in Table I. IHC analysis demonstrated that MTHFD2, ALDH1L2 and SHMT2 were localized in the cytoplasm of pancreatic cancer cells (Fig. 2). In addition, MTHFD2 expression was greater in pancreatic epithelial cancer tissue specimens compared with in surrounding non-epithelial tissues (Fig. 2). Representative images of ALDH1L2 scoring are demonstrated as: Score 0 in Fig. 2A, score 0 in Fig. 2B, score 2 in Fig. 2C, and score 2 in Fig. 2D. Examples of MTHFD2 scoring are presented as: Score 0 in Fig. 2E, score 1 in Fig. 2F, score 2 in Fig. 2G, and score 2 in Fig. 2H. Representative SHMT2 scoring is demonstrated as: Score 0 in Fig. 2I, score 1 in Fig. 2J, score 2 in Fig. 2K, and score 2 in Fig. 2L. With the exception of pathological N stage, no significant differences were observed in the clinicopathological parameters between the high and low expression groups in all three enzymes (data not presented). High MTHFD2 expression was positively correlated with the incidence of lymph node metastases (P=0.033; data not presented). IHC analyses using advanced colorectal cancer tissues or normal pancreatic acini tissues were used as positive controls as observed by two pathologists. Following IHC analyses of MTHFD2, ALDH1L2 and SHMT2, patients with scores of 1 or 0 were included in the low expression group and those with scores of 2 were included in the high expression group (Fig. 2).

*Expression of MTHFD2, SHMT2 and ALDH1L2 in patients with PDAC correlates with poor outcomes.* A previous report correlated the expression of the mitochondrial folate metabolic pathway enzyme MTHFD2 with poor clinical survival in breast cancer (7). Thus, we determined the association of mitochondrial folate metabolic pathway enzymes, including MTHFD2, ALDH1L2 and SHMT2, with clinical survival in patients with pancreatic cancer using Kaplan-Meier estimator analyses. Kaplan-Meier estimator curves demonstrated significantly lower DFS among patients with high tumor MTHFD2 expression (P<0.001; Fig. 3A). In addition, high ALDH1L2 expression was significantly associated with poor OS and DFS (P<0.001;



Figure 1. Schematic of one-carbon metabolism. Enzymes with polymorphisms investigated in the present study are illustrated in the boxes. ALDH1L1, aldehyde dehydrogenase 1 family member L1; ALDH1L2, aldehyde dehydrogenase 1 family member L2; AMT, aminomethyltransferase; CH2-THF, Methylenetetrahydrofolate; CH0-THF, formyl-tetrahydrofolate; DHFR, dehydrofolate reductase; MTHFD1, methylenetetrahydrofolate dehydrogenase D1; MTHFD2, methylenetetrahydrofolate dehydrogenase D2; HCHO, formaldehyde; NADPH, nicotinamide adenine dinucleotide phosphate; SHMT1, serine hydroxymethyltransferase 1; SHMT2, serine hydroxymethyltransferase 2; THF, tetrahydrofolate.

Fig. 3B). SHMT2 expression was significantly associated with poor OS (P=0.001), but was not associated with DFS (P=0.068; Fig. 3C). These data suggest that one-carbon metabolism serves an important role in the biologically malignant features such as invasion and metastasis of pancreatic cancer. In addition, based on the univariate analysis of OS (Table II), clinicopathological features that correlated with poor patient survival included pathological N stage (P<0.001), pathological UICC stage (P=0.007) and IHC scores for MTHFD2 (P<0.001), ALDH1L2 (P<0.001) and SHMT2 (P=0.002). Subsequent multivariate Cox proportional hazards analyses revealed that pathological N stage and IHC scores for MTHFD2, ALDH1L2 and SHMT2 were independent prognostic factors for OS (P=0.001, P=0.015 and P=0.017, respectively). Similarly, based on the univariate analyses of DFS (Table III), clinicopathological features that correlated with poor patient survival included pathological N stage (P<0.001), pathological UICC stage (P=0.034) and IHC scores for MTHFD2 (P<0.001) and ALDH1L2 (P<0.001); however, SHMT2 expression was not associated with patient survival (P=0.070). Finally, multivariate Cox proportional hazards analyses demonstrated that pathological N stage and IHC scores for MTHFD2 are independent prognostic factors for OS (P=0.025 and P<0.001, respectively).

Low expression of MTHFD2, SHMT2 and ALDH1L2 correlates with improved survival among patients with PDAC. Overexpression of MTHFD2, SHMT2 or ALDH1L2 was associated with poor survival (Fig. 3); however, it remains unclear whether the upregulation of one folate metabolic pathway enzyme directly enhances another folate pathway. As tumors that depend on the folate metabolic pathway may upregulate multiple folate metabolic pathway enzymes for progression and survival, the clinical outcomes of patients with PDAC with high expression of all three enzymes were compared with those of patients with low expression of all three enzymes, or with high expression of one or two of the enzymes. Kaplan-Meier estimator analyses revealed that low expression levels of MTHFD2, ALDH1L2 and SHMT2 correlated with improved OS and DFS in patients with PDAC, as compared with in patients with high expression of one or two of the enzymes (P<0.001; Fig. 4).

#### Discussion

In the present study, overexpression of each of the one-carbon metabolic enzymes, MTHFD2, ALDH1L2 and SHMT2, which are specifically located in the mitochondria, was associated with poor clinical outcomes among patients with PDAC. However, high and low expression levels of all three folate pathway enzymes were associated with improved survival, compared with the high expression of one or two of the three enzymes studied. Previously, a high intake of folate and the upregulation of the folate pathway were considered to reduce the risks of cancer due to the associated antioxidant activities of the folate metabolic pathway (6). However, recent epidemiologic data regarding the association between folate intake and pancreatic cancer are inconsistent (11). In addition, the expression of folate metabolic pathway enzymes, including MTHFD2, was identified to be markedly elevated in numerous types of cancer (12) and MTHFD2 expression has previously been associated with poor clinical outcomes (7). Therefore, one-carbon metabolism is currently considered one of the most important metabolic pathways for cancer progression (6).

	Table I. Clinicopathological	characteristics of 103	patients with	pancreatic ductal	adenocarcinoma.
--	------------------------------	------------------------	---------------	-------------------	-----------------

Variables	n (%)		
Age (years)	67.2±9.6		
Gender (male:female)	64 (62.1):39 (37.9)		
Carcinoembryonic antigen (ng/ml)	6.68±32.9		
Cancer antigen 19-9 (U/ml)	267.7±952.8		
Duke pancreatic monoclonal antigen type 2 (U/ml)	643.6±2014.5		
Operative procedure (PD:DP:others)	64 (62.1):37 (35.9):2 (2.0)		
Preoperative chemoradiotherapy (yes vs. no)	63 (61.2):40 (38.8)		
Adjuvant therapy (yes vs. no)	84 (81.6):19 (18.4)		
T stage (1:2:3:4)	21 (20.2):9 (8.7):73 (71.2):0 (0)		
N stage (0 vs. 1)	68 (66.0): 35 (34.0)		
UICC stage (IA:IB:IIA:IIB:III:IV)	17 (16.5):7 (6.8):44 (42.7):35 (34.0):0 (0):0 (0)		
Histological type (well:moderately:poor:other)	7 (6.8):90 (87.4):5 (4.9):1 (1.0)		
Median follow-up time (months)	60.9		

UICC, American Joint Committee on Cancer/Unio Internationalis Contra Cancrum; N, node; T, tumor; PD, pancreaticoduodenectomy; DP, distal pancreatectomy; others, 2 cases included total pancreatectomy and partial resection; UICC, American Joint Committee on Cancer/Unio Internationalis Contra Cancrum; N, node; T, tumor.



Figure 2. Immunohistochemical analyses of MTHFD2, ALDH1L2 and SHMT2. Representative images of ALDH1L2, (A) score 0; (B) score 0; (C) score 2; (D) score 2. Representative images of MTHFD2, (E) score 0; (F) score 1; (G) score 2; (H) score 2. Representative images of SHMT2, (I), score 0; (J) score 1; (K), score 2; (L) score 2. Acini were used as positive controls. Scale bar, 100  $\mu$ m. A, Acini; T, Tumors; ALDH1L2, aldehyde dehydrogenase 1 family member L2; MTHFD2, methylenetetrahydrofolate dehydrogenase D2; SHMT2, serine hydroxymethyltransferase 2.

Previous studies have revealed a number of molecular mechanisms underlying the effects of the folate metabolic pathway on tumorigenesis and metastasis (6). As a source of one-carbon units that are necessary for DNA replication and repair, folate protects normal tissues from mutations and chromosomal damage. However, folate may enhance the growth of pre-existing neoplastic lesions, as rapidly proliferating tissues such as tumors have increased nucleotide demands. Additionally, antioxidants have been demonstrated to promote distant metastasis in immunodeficient mice, and the inhibition of the folate pathway using low-dose methotrexate, ALDH1L2 knockdown or MTHFD1 knockdown limited distant metastasis without significantly affecting the growth of subcutaneous tumors in mice. Therefore, oxidative stress inhibits the distant metastasis of melanoma cells *in vivo* (8).

In the present study, the systematic downregulation of individual folate metabolic pathway enzymes was demonstrated to significantly correlate with improved survival. Recently, innovative drugs have been developed that target one-carbon metabolism. However, the majority of



Figure 3. Kaplan-Meier estimator analysis. Associations between patient survival and (A) MTHFD2, (B) ALDH1L2 or (C) SHMT2 expression levels. MTHFD2, methylenetetrahydrofolate dehydrogenase D2; SHMT2, serine hydroxymethyltransferase 2; ALDH1L2, aldehyde dehydrogenase 1 family member L2.

	Univariate			Multivariate		
Clinicopathological characteristic	HR	95% CI	P-value <sup>a</sup>	HR	95% CI	P-value
Age (years)	1.05	0.98-1.05	0.262			
Gender (female vs. male)	0.84	0.50-1.42	0.523			
Carcinoembryonic antigen (ng/ml)	0.99	0.98-1.01	0.412			
Cancer antigen 19-9 (U/ml)	0.99	0.99-1.00	0.443			
Duke pancreatic monoclonal antigen type 2 (U/ml)	1.00	0.99-1.00	0.241			
Preoperative chemoradiotherapy (yes vs. no)	0.97	0.58-1.66	0.939			
Adjuvant therapy (yes vs. no)	0.56	0.29-1.05	0.072			
T stage (3+4 vs. 1+2)	1.76	0.94-3.28	0.077			
N stage (1 vs. 0)	2.43	1.42-4.14	0.001	2.49	1.43-4.32	0.001
UICC Stage (IA+IIA vs. IB+IIB)	2.07	1.22-3.51	0.007			
Histological type (moderate vs. well+poor+muc)	0.77	0.24-2.48	0.661			
MTHFD2 (high vs. low)	3.48	2.04-5.91	<0.001 <sup>b</sup>	2.22	1.17-4.20	0.015 <sup>b</sup>
ALDH1L2 (high vs. low)	3.64	1.98-6.70	<0.001 <sup>b</sup>	2.35	1.16-4.74	$0.017^{b}$
SHMTL2 (high vs. low)	2.35	1.38-4.02	0.002 <sup>b</sup>	1.39	0.75-2.57	0.299 <sup>b</sup>

Table II. Multivariate analysis of overall survival.

MTHFD2, methylenetetrahydrofolate dehydrogenase D2; SHMT2, serine hydroxymethyltransferase 2; ALDH1L2, aldehyde dehydrogenase 1 family member L2; HR, hazard ratio; CI, confidence interval; UICC, American Joint Committee on Cancer/Unio Internationalis Contra Cancrum; N, node; T, tumor; muc, mucinous. <sup>a</sup>P<0.05 was considered to indicate a statistically significant difference. <sup>b</sup>The P-values are for the association between high MTHFD2/ALDH1L2/SHMTL2 expression and OS.

Table III. Multivariate analysis of disease-free survival.

	Univariate			Multivariate		
Categories	HR	95% CI	P-value	HR	95% CI	P-value
Age (years)	1.02	0.99-1.05	0.230			
Gender (female vs. male)	0.78	0.48-1.29	0.336			
Carcinoembryonic antigen (ng/ml)	1.00	0.99-1.00	0.370			
Cancer antigen 19-9 (U/ml)	1.00	0.99-1.00	0.242			
Duke pancreatic monoclonal antigen type 2 (U/ml)	1.00	0.99-1.00	0.297			
Preoperative chemoradiotherapy (yes vs. no)	1.28	0.76-2.13	0.353			
Adjuvant therapy (yes vs. no)	1.02	0.52-2.02	0.935			
T stage (3+4 vs. 1+2)	1.48	0.84-2.62	0.175			
N stage (1 vs. 0)	1.95	1.17-3.24	0.010	1.81	1.08-3.04	0.025
UICC Stage (IA+IIA vs. IB+IIB)	1.71	1.04-2.80	0.034			
Histological type (mod vs. well+poor+muc)	1.13	0.41-3.11	0.820			
MTHFD2 (high vs. low)	3.28	1.99-5.41	< 0.001	2.59	1.47-4.55	< 0.001
ALDH1L2 (high vs. low)	2.68	1.50-4.80	< 0.001	1.70	0.88-3.24	0.111
SHMT2 (high vs. low)	1.57	0.96-2.61	0.070			

MTHFD2, methylenetetrahydrofolate dehydrogenase D2; SHMT2, serine hydroxymethyltransferase 2; ALDH1L2, aldehyde dehydrogenase 1 family member L2; HR, hazard ratio; CI, confidence interval; UICC, American Joint Committee on Cancer/Unio Internationalis Contra Cancrum; N, node; T, tumor.

drugs targeted dihydrofolate reductase (DHFR). When the DHFR enzyme was inhibited, there was a metabolic flow of one-carbon metabolism. Therefore, the effect of DHFR targeting drugs may be insufficient. Cytoplasmic one-carbon metabolic enzymes, including MTHFD1, ALDH1L1 and SHMT1, and mitochondrial enzymes, including MTHFD2, ALDH1L2 and SHMT2, constitute the circinate metabolic flow, and this metabolic flow does not have an alternative loophole pathway. Therefore, the effect of targeting these enzymes may be sufficient. However, normal pancreatic



Figure 4. Associations between patient survival and the combined expression patterns of MTHFD2, ALDH1L2 and SHMT2. Associations between (A) OS or (B) DFS and triple low expression of MTHFD2, ALDH1L2 and SHMT2 (solid line; line 1); triple high expression of MTHFD2, ALDH1L2 and SHMT2 (dotted line, line 3); and other groups (broken line, line 2). Survival was estimated using Kaplan-Meier estimator analyses. P-values were calculated using the log-rank test. \*P<0.001. MTHFD2, methylenetetrahydrofolate dehydrogenase D2; SHMT2, serine hydroxymethyltransferase 2; ALDH1L2, aldehyde dehydrogenase 1 family member L2; OS, overall survival; DFS, disease-free survival.

tissues have high expression levels of these one-carbon metabolic enzymes. Thus, the drug targeting these enzymes must be delivered using cancer-specific drug delivery systems. Specifically, data from the present study demonstrated that high expression levels of one or two of the enzymes MTHFD2, ALDH1L2 and SHMT2 are associated with poor prognosis (Figs. 3 and 4), suggesting that the activity of one carbon metabolism plays a role in invasion and metastasis of pancreatic cancer cells and associates with clinical survival of patients with pancreatic cancer. These proteins may be independent prognostic factors and potential therapeutic targets for pancreatic cancer.

## Acknowledgements

Not applicable.

## Funding

The present study was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (Tokyo, Japan; grant nos. 23390199, 25112708, 25134711, 30253420 and 26670604), a Grant-in-Aid from the Ministry of Health, Labour and Welfare (Tokyo, Japan; grant no. H23-003), a grant from the National Institute of Biomedical Innovation (Osaka, Japan; grant no. 12-4) and a grant from the Osaka University Drug Discovery Fund (Osaka, Japan). Partial support was also received from the Takeda Science and Medical Research Foundation (Osaka, Japan; to Dr H Ishii), the Princess Takamatsu Cancer Research Fund (Tokyo, Japan; to Dr H Ishii), the Suzuken Memorial Foundation (to Dr M Konno), the Yasuda Medical Foundation (to Dr M Konno), the Pancreas Research Foundation (to Dr K Kawamoto), the Nakatani Foundation (to Dr H Ishii) and the Nakatomi Foundation, Japan (to Dr M Konno).

### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

#### Authors' contributions

KN, MK, HE, YD, MM, and HI conceived and designed the study. KN, MK and HI developed the methodology. KN, MK, JK, NN, KK, DY, TA, TN, HW, KG, DS, TK, TS, HE and HI acquired the data. KN, MK, JK, NN and KK analyzed and interpreted the data. KN, MK, HE, YD, MM and HI wrote, reviewed and revised the manuscript. MM and HI supervised the study.

### Ethics approval and consent to participate

This study has been approved by the research ethics committee of Osaka University (ID of the approval: 15149-2, by Dr. Y. Kaneda).

#### **Consent for publication**

All patients provided written consent for the publication of results.

#### **Competing interests**

Institutional endowments were received partially from Taiho Pharmaceutical Co., Ltd. (Tokyo, Japan), Evidence-Based Medical Research Center (Osaka, Japan), Chugai Co., Ltd. (London, UK), Yakult Honsha Co., Ltd. (Tokyo, Japan) and Merck & Co., Ltd., Whitehouse Station, NJ, USA). These funding bodies had no role in the main experimental equipment, the study design, data collection and analysis, the decision to publish or the preparation of this manuscript. 1834

#### References

- 1. Howe HL, Wu X, Ries LA, Cokkinides V, Ahmed F, Jemal A, Miller B, Williams M, Ward E, Wingo PA, et al: Annual report to the nation on the status of cancer, 1975-2003, featuring cancer among U.S. Hispanic/Latino populations. Cancer 107: 1711-1742, 2006. 2. Majumder K, Gupta A, Arora N, Singh PP and Singh S:
- Premorbid obesity and mortality in patients with pancreatic cancer: A systematic review and meta-analysis. Clin Gastroenterol Hepatol 14: 355-368.e, 2016.
- 3. Hashimoto D, Chikamoto A, Ohmuraya M, Sakata K, Miyake K, Kuroki H, Watanabe M, Beppu T, Hirota M and Baba H: Pancreatic cancer in the remnant pancreas following primary pancreatic resection. Surg Today 44:
- 1313-1320, 2014. 4. Kanda M, Fujii T, Takami H, Suenaga M, Inokawa Y, Yamada S, Nakayama G, Sugimoto H, Koike M, Nomoto S and Kodera Y: Combination of the serum carbohydrate antigen 19-9 and carcinoembryonic antigen is a simple and accurate predictor of mortality in pancreatic cancer patients. Surg Today 44: 1692-1701, 2014.
- 5. Pandol S, Gukovskaya A, Edderkaoui M, Dawson D, Eibl G and Lugea A: Epidemiology, risk factors, and the promotion of pancreatic cancer: Role of the stellate cell. J Gastroenterol Hepatol 27 (Suppl 2): S127-S134, 2012.

- 6. Konno M, Asai A, Kawamoto K, Nishida N, Satoh T, Doki Y, Mori M and Ishii H: The one-carbon metabolism pathway highlights therapeutic targets for gastrointestinal cancer (Review). Int J Oncol 50: 1057-1063, 2017
- 7. Liu F, Liu Y, He C, Tao L, He X, Song H and Zhang G: Increased MTHFD2 expression is associated with poor prognosis in breast cancer. Tumour Biol 35: 8685-8690, 2014
- 8. Piskounova E, Agathocleous M, Murphy MM, Hu Z, Huddlestun SE, Zhao Z, Leitch AM, Johnson TM, DeBerardinis RJ
- Hasegawa S, Eguchi H, Nagano H, Konno M, Tomimaru Y, Wada H, Hama N, Kawamoto K, Kobayashi S, Nishida N, et al: MicroRNA-1246 expression associated with CCNG2-mediated chemoresistance and stemness in pancreatic cancer. Br J Cancer 111: 1572-1580, 2014.
- 11. Lin HL, An QZ, Wang QZ and Liu CX: Folate intake and pancreatic cancer risk: An overall and dose-response meta-analysis. Public Health 127: 607-613, 2013.
- Nilsson R, Jain M, Madhusudhan N, Sheppard NG, Strittmatter L, Kampf C, Huang J, Asplund A and Mootha VK: Metabolic enzyme expression highlights a key role for MTHFD2 and the mitochondrial folate pathway in cancer. Nat Commun 5: 3128, 2014.