

Influence of *SLCO1B1* in gastric cancer patients treated with EOF chemotherapy

WANJING FENG^{1,2*}, XIN LIU^{1,2*}, XIAOYING ZHAO^{1,2}, MINGZHU HUANG^{1,2},
WEIJIAN GUO^{1,2}, JILIANG YIN^{1,2}, ZHIYU CHEN^{1,2} and XIAODONG ZHU^{1,2}

¹Department of Medical Oncology, Fudan University Shanghai Cancer Center;

²Department of Oncology, Shanghai Medical College, Fudan University, Shanghai 200032, P.R. China

Received November 19, 2017; Accepted June 28, 2018

DOI: 10.3892/ol.2018.9147

Abstract. Cytochrome-P450 enzymes, ATP-binding cassette transporters, and solute carriers mediate drug metabolism as metabolic enzymes and membrane transporters, respectively. The present study investigated whether single nucleotide polymorphisms (SNPs) in genes encoding these proteins were predictive or prognostic factors in patients with metastatic gastric cancer (MGC) undergoing chemotherapy. A retrospective study of 108 MGC patients who received epirubicin, oxaliplatin, and 5-fluorouracil (EOF) as first-line treatment was performed. A total of 13 SNPs were genotyped, including *SLCO1B1* (rs4149056), *SLC2A9* (rs16890979, rs6449213, rs734553), *ABCG2* (rs2231142), *CYP2C9* (rs1057910, rs1799853), *CYP2C19* (rs72552267, rs28399504, rs56337013, rs41291556) and *CYP1A2* (rs12720461, rs56107638). The associations between these genotypes and disease-control rate (DCR), progression-free survival (PFS) and overall survival (OS) were analyzed. Patients with *SLCO1B1* rs4149056 TT genotype had a significantly shorter OS compared with those with a C allele (CC + CT; 312 vs. 565 days, $P=0.039$). Multivariate analysis revealed that the rs4149056 TT homozygous genotype was an independent prognostic factor for shorter OS (hazard ratio: 2.565, 95% confidence interval: 1.215-5.415, $P=0.014$). However, no significant associations between *SLCO1B1* rs4149056 and PFS were observed, between the other 12 SNPs and PFS or OS, or between any of the 13 SNPs and DCR. In conclusion, *SLCO1B1*

rs4149056 TT may be an independent predictor of survival in patients with MGC treated with EOF chemotherapy.

Introduction

Gastric cancer (GC) represents the fourth most common malignant neoplasm and the second leading cause of cancer-related death worldwide. More than 40% of all GC cases diagnosed annually occur in China, of whom almost 50% are diagnosed at an advanced stage and cannot be cured. Treatment of advanced GC is challenging. Epirubicin, cisplatin, and fluorouracil (ECF) and its modified regimens, such as EOF (epirubicin, oxaliplatin, and fluorouracil) and EOX (epirubicin, oxaliplatin, and capecitabine), are widely used to treat GC patients based on the results of phase III clinical trials. However, although these regimens are common first-line treatments, their response rates remain <50%. Biomarkers able to predict chemotherapeutic efficacy are therefore urgently needed.

Both membrane transporters and metabolic enzymes affect cytotoxic-drug metabolism. Solute carrier (SLC) superfamily proteins and ATP-binding cassette (ABC)-transporters are vital membrane transporters. *SLCO1B1* encodes the transporter protein, organic anion-transporting polypeptide-1 (OATP1B1), which mediates liver uptake of a wide variety of drugs, and its role in the efficacy of cytotoxic drugs, including 5-fluorouracil (5FU) (1,2), methotrexate (MTX) (3-5), irinotecan (6,7), and paclitaxel (8) has been widely reported. MTX was the first cytotoxic drug reported to be associated with *SLCO1B1*, and *SLCO1B1* SNPs were shown to affect MTX pharmacokinetics in children with acute lymphoblastic leukemia, particularly in terms of deposition and toxicity (5). These findings have been validated in several later studies (9-11). The breast cancer resistance protein (BCRP/ABCG2) has been reported to affect drug resistance in many cancer types, including colorectal cancer, lymphoblastic leukemia, and breast cancer (12-15), and ABCG2 polymorphisms are prognostic factors in breast cancer patients treated with anthracycline-based neoadjuvant chemotherapy (16). Expression levels of glucose transporters (GLUT/SLC1A) were also shown to be related to response to 5FU chemotherapy in GC cells (17), and glucose transporters were reported to be independent prognostic factors in patients with GC (18,19).

Correspondence to: Dr Xiaodong Zhu or Dr Zhiyu Chen, Department of Oncology, Shanghai Medical College, Fudan University, 270 Dong' An Road, Shanghai 200032, P.R. China
E-mail: xddr001@163.com
E-mail: chanhj75@aliyun.com

*Contributed equally

Key words: gastric cancer, single nucleotide polymorphism, received epirubicin, oxaliplatin, and 5-fluorouracil regimen, *SLCO1B1*

Among metabolic enzymes, the cytochrome-P450 (CYP) enzyme family plays an important role in the metabolism of various anticancer drugs (20). Several studies have shown that SNPs in *CYP2C9* influence disease-free survival in breast cancer patients treated with tamoxifen (21,22). Moreover, *CYP2C9* polymorphism was related to response to fluorouracil-based neoadjuvant chemotherapy in breast cancer (23). *CYP2C19* is involved in the metabolism of cyclophosphamide (24,25) and tamoxifen (26,27), and P450 enzymes in human liver microsomes, including *CYP1A2* have also been reported to catalyze tegafur into 5FU (28).

Metabolism-related genes, including *SLCO1B1*, *ABCG2*, *SLC2A9*, *CYP2C9*, *CYP2C19*, and *CYP1A2*, might thus influence the efficacy of EOF regimens. In the present study, we investigated the associations between metabolism-related genes and the clinical outcomes of GC patients treated with first-line EOF regimens, in terms of disease-control rate (DCR), progression-free survival (PFS), and overall survival (OS).

Materials and methods

Study population. This retrospective study enrolled 108 consecutive Chinese Han patients with metastatic GC (MGC) treated with EOF regimens as first-line chemotherapy at Fudan University Shanghai Cancer Center (Shanghai, China) between May 2009 and June 2012. Their diagnoses were pathologically confirmed as gastric adenocarcinoma. The study was approved by the Ethics Committee of Fudan University Shanghai Cancer and complied with the principles of the Helsinki Accord. This was a retrospective study and patient consent was therefore deemed unnecessary. However, blood samples were collected from all subjects before treatment, with patient consent, and stored in the tissue bank at Fudan University Shanghai Cancer Center.

Treatment. All patients in this study were treated with first-line chemotherapy using an EOF regimen, consisting of epirubicin infusion (50 mg/m²) combined with an intravenous infusion of oxaliplatin (130 mg/m²) for 2 h on day 1, following a 24-h continuous infusion of 5FU (375-425 mg/m²/day) for 5 days, over a 21-day treatment cycle. Tumor responses were evaluated every 6 weeks in accordance to the Response Evaluation Criteria in Solid Tumors 1.0 (RECIST 1.0). Treatment was terminated in the event of disease progression or unacceptable toxicity. If the lesions continued to shrink after six cycles, with no unacceptable toxicity, a further one or two EOF cycles were recommended; otherwise, oral FU was recommended as follow-up treatment.

SNP selection and genotyping. We selected and genotyped 13 drug-metabolism-related genetic polymorphisms located at *SLCO1B1* (rs4149056), *SLC2A9* (rs16890979, rs6449213, rs734553), *ABCG2* (rs2231142), *CYP2C9* (rs1057910, rs1799853), *CYP2C19* (rs72552267, rs28399504, rs56337013, rs41291556), and *CYP1A2* (rs12720461, rs56107638), respectively, from the Hapmap project (www.hapmap.org) and dbSNP databases (www.ncbi.nlm.nih.gov/SLP) (Table I). Genomic DNA was extracted from venous blood leukocytes using a standard phenol-chloroform method. The selected SNPs were genotyped using the TaqMan assay method and an ABI 7900

DNA detection system (Applied Biosystems, Foster City, CA, USA). All the probes and primers were designed using the Assay-on-Design service from Applied Biosystems. The experiments were repeated for 15% of the samples. The genotype error rate was <0.03%.

Statistical analysis. The allelic and genotypic distributions and Hardy-Weinberg equilibrium were analyzed using the online analysis tool, SHEsis. Differences in clinical characteristics among the 108 GC patients were analyzed by χ^2 tests, and differences in allelic and genotypic frequencies between the controlled-disease and progressive-disease groups were compared with χ^2 or Fisher's exact probability tests. Genetic power was calculated using the G*Power program (29). Survival curves were analyzed using the Kaplan-Meier method. Differences in PFS and OS between genotype groups were estimated by log-rank tests. Multivariate analysis of prognostic predictors was carried out using a Cox proportional hazards model. A two-sided P-value <0.05 was considered significant. P-values for association analysis were corrected by a false-discovery rate (FDR) procedure (30). To be detailed, $\text{FDR } p = p^* n/a$. In this formula, n is the number of SNPs in the same gene, and a is the rank of the P-value among the SNPs in the same gene.

Results

Patient characteristics. Most of the 108 GC patients enrolled in the study had ≥ 3 tumor sites, with the most common metastatic organs being the liver and retroperitoneal lymph nodes. Among the clinicopathological features, liver metastasis and pleural effusion were significantly associated with OS, whereas histological grade, number of tumor sites, retroperitoneal lymph node involvement, and ascites were significantly associated with PFS (Table II).

The responses to EOF chemotherapy were CR, $n=1$; PR, $n=41$; SD, $n=47$; and PD, $n=19$. The 89 patients with CR, PR, or SD were classified as the disease-control group and the 19 patients with PD were classified as the disease-progression group.

Genotype frequency and disease control. There was no significant relationship between genotype frequency and disease-control rate for any of the 13 SNPs analyzed in the present study (Table III). There was only one genotype distribution each for *CYP2C19* (rs72552267, rs28399504, rs56337013, rs41291556) and *CYP1A2* (rs12720461, rs56107638) in the 108 patients (data not shown).

Genotype frequency and survival analysis. Univariate analysis of the 13 SNPs in relation to survival (Table IV) showed that patients with *SLCO1B1* rs4149056 CC and CT genotypes had significantly longer median OS than patients with the TT genotype (565 vs. 312 days, log-rank $P=0.039$; Fig. 1). However, there was no significant association between PFS and *SLCO1B1* rs4149056 (log-rank $P=0.956$; Fig. 2), or between any of the other 12 SNPs and OS or PFS (Table IV).

Multivariate PFS and OS analysis with cox regression. Multivariate analysis showed that *SLCO1B1* rs4149056

Table I. SNPs in the *SLCO1B1*, *SLC2A9*, *SLC17A1*, *ABCG2*, *CYP2C9*, *CYP2C19* and *CYP1A2* genes analyzed in the article.

Gene	SNP ID	Chromosome	Function	Allele	HWE test P-value
<i>SLCO1B1</i>	rs4149056	12:21178615	Missense	T/C	0.608
<i>SLC2A9</i>	rs16890979	4:9920543	Missense	C/T	0.903
<i>SLC2A9</i>	rs6449213	4:9992591	Intron	C/T	0.903
<i>SLC2A9</i>	rs734553	4:9921380	Intron	A/C	0.903
<i>ABGC2</i>	rs2231142	4:88131171	Missense	A/C	0.047
<i>CYP2C9</i>	rs1057910	10:94981296	Missense	A/C	0.808
<i>CYP2C9</i>	rs1799853	10:94942290	Missense	C/T	1.000
<i>CYP2C19</i>	rs72552267	10:94775453	Missense	A/G	1.000
<i>CYP2C19</i>	rs28399504	10:94762706	Missense	A/G/T	1.000
<i>CTP2C19</i>	rs56337013	10:94852738	Missense	C/T	1.000
<i>CYP2C19</i>	rs41291556	10:94775416	Missense	C/T	1.000
<i>CYP1A2</i>	rs12720461	15:74749010	Intron	C/T	1.000
<i>CYP1A2</i>	rs56107638	15:74753271	Splice donor	A/C/G	1.000

HWE, hardy-weinberg equilibrium; SNP, single nucleotide polymorphism.

genotype, liver metastasis, ascites, pleural effusion, and number of tumor sites were significantly or borderline-significantly associated with OS (Table V). In addition, histological grade, retroperitoneal lymph node involvement, ascites, pleural effusion, and number of tumor sites were significantly or borderline-significantly associated with PFS. All the above factors were included in a stepwise multivariate Cox regression model, which confirmed that *SLCO1B1* rs4149056 was an independent prognostic factor for shorter OS ($P=0.014$), but not PFS ($P=0.533$).

Linkage disequilibrium analysis. We analyzed linkage disequilibrium (LD) for the *SLC2A9* and *CYP2C9* SNPs in AGC patients HaploView software. The *SLC2A9* SNPs rs16890979 and rs6449213 ($D'=1.000$, $r^2=0.796$), *SLC2A9* SNPs rs16890979 and rs734553 ($D'=1.000$, $r^2=1.000$) and *SLC2A9* SNPs rs6449213 and rs734553 ($D'=1.000$, $r^2=0.796$) all showed strong LD (Fig. 3A). *CYP2C9* SNPs rs1057910 and rs1799853 showed no LD ($D'=1.000$, $r^2=0.000$) (Fig. 3B). Since the most samples showed the same allele and only less than 5 samples showed different allele in the *SLC2A9* SNPs, linkage disequilibrium and haplotype analysis cannot find clinically meaningful result.

Discussion

We investigated the association between SNPs of metabolism-related genes and chemotherapy response in patients with MGC. Patients with *SLCO1B1* rs4149056 CC and CT genotypes had longer OS than patients with TT genotype (312 vs. 565 days, $P=0.039$), and *SLCO1B1* rs4149056 TT was confirmed as an independent prognostic factor for shorter OS in GC patients treated with EOF by Cox regression analysis (hazard ratio: 2.565, 95% confidence interval: 1.215-5.415, $P=0.014$).

SLCO1B1 encodes the SLC family member OATP1B1, which is highly expressed in hepatocytes and associated with

hepatic drug uptake and elimination (31). Among single-nucleotide variants of *SLCO1B1*, c.521T>C rs4149056 had the greatest effect on OATP1B1 activity in this study. *SLCO1B1* c.521T>C decreases OATP1B1 transporting activity, thus increasing plasma concentrations of drugs, including cytotoxic drugs. *SLCO1B1* rs4149056 has been significantly associated with exposure to SN-38, the active metabolite of irinotecan (7). In another study, OATP1B1 was responsible for SN-38 uptake from plasma into hepatocytes (6). Innocenti *et al* also revealed that *SLCO1B1* rs4149056 increased patient exposure to CPT-11 and was associated with an increased risk of severe neutropenia (6). OATP1B1 was shown to transport paclitaxel in an *in vitro* ovarian cancer study, implying that it contributed to the disposition of paclitaxel (8). Huang *et al* reported that another *SLCO1B1* variant, rs2306283, was an independent prognostic factor for longer PFS in patients with metastatic colorectal cancer treated with a fluoropyrimidine plus irinotecan (1). Overall, these studies suggest that *SLCO1B1* affects the metabolism of fluoropyrimidines, CPT-11, and docetaxel, thus influencing their chemotherapeutic efficacy, consistent with the results of the current study.

To the best of our knowledge, the present study provides the first evidence for an association between *SLCO1B1* and GC. A literature search of MEDLINE via PubMed found no similar reports with respect to the efficacy of first-line chemotherapy, GC risk, or prognosis.

Notably, although we identified *SLCO1B1* rs4149056 as a prognostic factor for OS in MGC patients treated with first-line EOF, we failed to detect any correlation between *SLCO1B1* rs4149056 and PFS. There are two possible reasons for this apparent discrepancy. First, *SLCO1B1* rs4149056 may only be a prognostic factor in GC patients, and may not affect the short-term efficacy of first-line chemotherapy. Second, CPT-11 or paclitaxel were administered as the main second- or third-line chemotherapy regimens in the present study; as noted above, *SLCO1B1* rs4149056 has been associated with the metabolism

Table II. Associations between patient characteristics and survival.

Characteristics	Number of patients (%)	Median OS	P-value	Median PFS	P-value
Age (years)					
≤60	80 (74.1)	465	0.916	159	0.176
>60	28 (25.9)	403		187	
Sex					
Male	64 (59.3)	534	0.359	166	0.163
Female	44 (40.7)	372		182	
ECOG performance status					
0	14 (13.0)	704	0.153	240	0.768
1	89 (82.4)	367		167	
2	5 (4.6)	299		237	
Histological grade					
Moderate and high	13 (12.0)	403	0.226	156	0.004
Low/undifferentiated	66 (61.2)	875		380	
Unclassified	29 (26.8)	367		180	
Number of tumor sites					
1	5 (4.6)	984	0.076	545	0.038
2	8 (7.4)	570		411	
≥3	95 (88.0)	372		167	
Metastasis sites					
Liver					
Yes	36 (33.3)	281	0.004	169	0.232
No	72 (66.7)	570		178	
Lung					
Yes	7 (6.5)	252	0.997	169	0.885
No	101 (93.5)	444		187	
Retroperitoneal lymph node					
Yes	45	372	0.710	156	0.033
No	63	534		192	
Ascites					
Yes	32	312	0.101	126	0.015
No	76	534		192	
Pleural effusion					
Yes	9	211	0.001	144	0.054
No	99	475		178	

Log-rank test. PFS, progression free survival; OS, overall survival.

of CPT-11 and paclitaxel, and rs4149056 may thus affect OS by influencing the efficacy of the second- or third-line regimens.

In our study, *ABCG2* (BCRP2) C421A rs2231142 was borderline-significantly associated with PFS ($P=0.083$), though multivariate analysis did not identify it as an independent predictor in GC patients treated with EOF regimen. However, several previous studies reported significant relationships between *ABCG2* (BCRP2) C421A rs2231142 and the toxicity or efficacy of oxaliplatin or anthracyclines. Custodio *et al* revealed that *ABCG2* rs3114018 was associated with oxaliplatin-induced peripheral neuropathy in patients with stage II-III colon cancer (32), while breast cancer patients with an *ABCG2* rs2231142 A allele reportedly showed better responses to anthracycline-based neoadjuvant chemotherapy (16). Ghafouri *et al* also showed that the *ABCG2* rs22311442 A allele

was associated with stronger responses to anthracyclines and paclitaxel (33). Overall, these three studies indicated that patients with an *ABCG2* rs22311442 A allele showed better responses to anthracyclines and paclitaxel. Although the current univariate analysis failed to find an association between *ABCG2* rs22311442 and PFS, there was a nonsignificant tendency towards longer PFS among rs22311442 AA/AC compared with rs22311442 CC patients. There are three possible reasons for this apparent discrepancy. First, it is possible that the sample-size was too small to demonstrate any significant difference between the survival trends of patients with rs22311442 A and C alleles, respectively. Second, the previous studies involved patients with colon cancer or breast cancer, rather than GC, and the results may thus have been affected by this tumor heterogeneity. Third, the EOF regimen contains epirubicin, oxaliplatin, and 5FU.

Table III. Allelic and genotypic distribution of the 15 SNPs in the disease control (CR, PR and SD) and the disease progressive (PD) to chemotherapy.

Gene	SNP	Allele frequency		Chi ² value	P-value	Odds ratio 95% CI		Genotype frequency			P-value
SLCO1B1	Rs4149056 CR+PR+SD PD	C	T	0.372	0.541	0.705 (0.229-2.168)	0 (0.000)	CC	CT	TT	0.700
		4 (0.105) 24 (0.143)	34 (0.895) 144 (0.857)								
SLC2A9	rs16890979 CR+PR+SD PD	C	T	0.185	0.666	0.606 (0.061-6.006)	17 (0.944) 85 (0.966)	CC	CT		0.663
		35 (0.972) 173 (0.983)	1 (0.028) 3 (0.017)								
SLC2A9	rs6449213 CR+PR+SD PD	C	T	0.036	0.847	1.242 (0.134-11.457)	1 (0.056) 4 (0.045)	CT	TT		0.845
		1 (0.028) 4 (0.022)	35 (0.972) 174 (0.978)								
SLC2A9	rs734553 CR+PR+SD PD	A	C	0.168	0.681	0.621 (0.062-6.149)	17 (0.944) 83 (0.965)	AA	AC		0.678
		35 (0.972) 169 (0.983)	1 (0.028) 3 (0.017)								
ABGC2	Rs2231142 CR+PR+SD PD	A	C	2.081	0.149	0.525 (0.217-1.272)	2 (0.111) 11 (0.124)	AA	AC	CC	0.175
		7 (0.194) 56 (0.315)	29 (0.806) 122 (0.685)								
CYP2C9	Rs1057910 CR+PR+SD PD	A	C	2.952	0.085	0.204 (0.027-1.500)	17 (0.895) 87 (0.978)	AA	AC		0.082
		36 (0.947) 176 (0.989)	2 (0.053) 2 (0.011)								
CYP2C9	rs1799853 CR+PR+SD PD	C	T	0.203	0.652		18 (1.000) 88 (0.989)	CC	CT		0.651
		36 (1.000) 177 (0.994)	0 (0.000) 1 (0.006)								

χ² test. SNP, single nucleotide polymorphism.

Table IV. Univariate survival analysis of SNPs and overall survival/progression free survival time.

SNPs	Median OS	95% CI	P-value	Median PFS	95% CI	P-value
SLCO1B1 rs4149056						
TT	312.0	178.0-445.9	0.039	173.0	142.7-203.2	0.760
CC+CT	565.0	108.9-1021.0		179.0	157.6-200.3	
SLC2A9 rs16890979						
CC	465.0	303.1-626.8	0.301	176.0	156.7-195.2	0.427
TC	189.0	0.0-383.6		162.0	0.0-353.6	
SLC2A9 rs6449213						
CT	265.0	94.0-435.9	0.161	162.0	55.8-268.1	0.257
TT	465.0	304.2-625.7		176.0	161.4-190.5	
SLC2A9 rs734553						
AA	444.0	307.7-580.2	0.282	173.0	154.2-191.7	0.489
CA	189.0	0.0-383.6		162.0	0-353.6	
ABGC2 rs2231142						
CC	444.0	293.2-594.7	0.750	167.0	144.1-189.8	0.083
AA+CA	475.0	128.5-821.4		182.0	138.1-225.8	
CYP2C9 rs1057910						
AA	444.0	289.8-598.1	0.382	176	156.8-195.1	0.311
CA	295.0	0.0-727.6		41	0.0-202.7	
CYP2C9 rs1799853						
CC	465.0	272.9-657.0	0.711	173.0	153.6-192.3	0.750
CT	444.0	-		178.0	-	

Log-rank test. PFS, progression-free survival; OS, overall survival; HR, hazards ratio; CI, confidence interval; SNP, single nucleotide polymorphism.




Figure 1. Kaplan-Meier OS curves in patients with different SLCO1B1 rs4149056 genotypes. OS, overall survival.




Figure 2. Kaplan-Meier PFS curves in patients with different SLCO1B1 rs4149056 genotypes. PFS, progression-free survival.

Custodio showed an association between *ABCG2* rs22311442 and oxaliplatin toxicity rather than oxaliplatin response, which differed from our current study.

Evidence suggests that some CYP450 enzyme family members affect 5FU metabolism (34-37). 5FU was reported to reduce the ability to metabolize a CYP2C9 probe drug (38).

Moreover, several studies have indicated that 5FU can down-regulate the expression of CYP enzymes, including CYP2C9 and CYP2C19 (39,40). However, although CYP2C9 and CYP2C19 have been shown to influence 5FU metabolism *in vitro*, these results have not yet been verified in clinical studies. Furthermore, no studies have reported on the relationship

Table V. Multi-factorial analysis of prognostic factors for PFS and OS.

Clinical factor	OS		PFS	
	P-value	HR 95% CI	P-value	HR 95% CI
rs4149056 TT	0.014	2.565 (1.215-5.415)	-	-
Grade	-	-	0.221	-
PLN	-	-	0.003	2.041 (1.271-3.278)
Ascites	0.050	0.508 (0.258-0.999)	0.436	0.818 (0.494-1.355)
Pleural effusion	0.016	0.363 (0.159-0.827)	0.071	0.497 (0.232-1.061)
Liver	0.002	0.372 (0.179-0.703)	-	-
Number of sites	-	-	0.200	-
rs2231142 CC	-	-	0.342	1.239 (0.796-1.929)

Cox proportional hazards model. PFS, progression-free survival; OS, overall survival; HR, hazards ratio; CI, confidence interval; PLN, retroperitoneal lymph node.




Figure 3. Haploview linkage disequilibrium. Linkage disequilibrium for haplotype blocks within (A) the *SLC2A9* SNPs rs16890979, rs734553 and rs6449213, (B) the *CYP2C9* SNPs rs1057910 and rs1799853.

between 5FU efficacy and SNPs in *CYP2C9* or *CYP2C19*. Our results showed that all 108 blood samples had the same genotypic distributions of *CYP2C9* and *CYP2C19* SNPs, except for *CYP2C9* rs1057910, which was not significantly associated with PFS or OS. The roles of *CYP2C9* and *CYP2C19* in 5FU metabolism thus currently remain unclear.

There were several limitations to the present study. First, we only analyzed some of the genes related to the CYP, SLC, and ABC families, and any associations between the remaining untested genes and GC remain unknown. Second, we only detected selected polymorphisms for each target agent, and other potentially related polymorphisms may have been missed. Third, the sample size was relatively small. The conclusions of this study therefore require verification in further studies.

In conclusion, the present study identified an association between the *SLCO1B1* rs4149056 SNP and clinical

outcomes of MGC patients treated with EOF chemotherapy. The resulting risk model successfully divided patients into low-risk and high-risk groups, with a significant difference in OS, but not PFS. *SLCO1B1* rs4149056 is therefore a prognostic, but not a predictive factor in MGC patients treated with EOF. Further studies are required to validate our results and to detect other potential prognostic sites in *SLCO1B1*.

Acknowledgements

Not applicable.

Funding

This study was funded by The National Key Research and Development Program of China (grant no. 2017YFC1308900).

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Authors' contributions

WF and XL performed the statistical analysis and wrote the article. XZhu and ZC designed and managed the research. XZha and MH performed the experiments. WG and JY assisted with the research design.

Ethics approval and consent to participant

Informed consent was been obtained from all participants. The study was approved by the Ethics Committee of Fudan University Shanghai Cancer.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Huang L, Zhang T, Xie C, Liao X, Yu Q, Feng J, Ma H, Dai J, Li M, Chen J, *et al*: SLC01B1 and SLC19A1 gene variants and irinotecan-induced rapid response and survival: A prospective multicenter pharmacogenetics study of metastatic colorectal cancer. *PLoS One* 8: e77223, 2013.
- De Mattia E, Toffoli G, Polesel J, D'Andrea M, Corona G, Zagonel V, Buonadonna A, Dreussi E and Cecchin E: Pharmacogenetics of ABC and SLC transporters in metastatic colorectal cancer patients receiving first-line FOLFIRI treatment. *Pharmacogenet Genomics* 23: 549-557, 2013.
- Ramsey LB, Panetta JC, Smith C, Yang W, Fan Y, Winick NJ, Martin PL, Cheng C, Devidas M, Pui CH, *et al*: Genome-wide study of methotrexate clearance replicates SLC01B1. *Blood* 121: 898-904, 2013.
- Radtke S, Zolk O, Renner B, Paulides M, Zimmermann M, Möricke A, Stanulla M, Schrappe M and Langer T: Germline genetic variations in methotrexate candidate genes are associated with pharmacokinetics, toxicity, and outcome in childhood acute lymphoblastic leukemia. *Blood* 121: 5145-5153, 2013.
- Treviño LR, Shimasaki N, Yang W, Panetta JC, Cheng C, Pei D, Chan D, Sparreboom A, Giacomini KM, Pui CH, *et al*: Germline genetic variation in an organic anion transporter polypeptide associated with methotrexate pharmacokinetics and clinical effects. *J Clin Oncol* 27: 5972-5978, 2009.
- Innocenti F, Kroetz DL, Schuetz E, Dolan ME, Ramírez J, Relling M, Chen P, Das S, Rosner GL and Ratain MJ: Comprehensive pharmacogenetic analysis of irinotecan neutropenia and pharmacokinetics. *J Clin Oncol* 27: 2604-2614, 2009.
- Teft WA, Welch S, Lenehan J, Parfitt J, Choi YH, Winquist E and Kim RB: OATP1B1 and tumour OATP1B3 modulate exposure, toxicity, and survival after irinotecan-based chemotherapy. *Br J Cancer* 112: 857-865, 2015.
- Svoboda M, Wlcek K, Taferner B, Hering S, Stieger B, Tong D, Zeillinger R, Thalhammer T and Jäger W: Expression of organic anion-transporting polypeptides 1B1 and 1B3 in ovarian cancer cells: Relevance for paclitaxel transport. *Biomed Pharmacother* 65: 417-426, 2011.
- Zhang HN, He XL, Wang C, Wang Y, Chen YJ, Li JX, Niu CH and Gao P: Impact of SLC01B1 521T>C variant on leucovorin rescue and risk of relapse in childhood acute lymphoblastic leukemia treated with high-dose methotrexate. *Pediatr Blood Cancer* 61: 2203-2207, 2014.
- Ramsey LB, Bruun GH, Yang W, Treviño LR, Vattathil S, Scheet P, Cheng C, Rosner GL, Giacomini KM, Fan Y, *et al*: Rare versus common variants in pharmacogenetics: SLC01B1 variation and methotrexate disposition. *Genome Res* 22: 1-8, 2012.
- Lima A, Bernardes M, Azevedo R, Monteiro J, Sousa H, Medeiros R and Seabra V: SLC19A1, SLC46A1 and SLC01B1 polymorphisms as predictors of methotrexate-related toxicity in Portuguese rheumatoid arthritis patients. *Toxicol Sci* 142: 196-209, 2014.
- Yuan SQ, Zhou ZW, Liang YJ, Fu LW, Chen G, Qiu HB and Zhang LY: Correlation of chemosensitivity tested using histoculture drug response assay to expression of multidrug resistance genes and proteins in colorectal cancer tissues. *Ai Zheng* 28: 932-938, 2009.
- Cortez MA, Scrideli CA, Yunes JA, Valera ET, Toledo SR, Pavoni-Ferreira PC, Lee ML, Petrilli AS, Brandalise SR and Tone LG: mRNA expression profile of multidrug resistance genes in childhood acute lymphoblastic leukemia. Low expression levels associated with a higher risk of toxic death. *Pediatr Blood Cancer* 53: 996-1004, 2009.
- Zhou S, Liao L, Chen C, Zeng W, Liu S, Su J, Zhao S, Chen M, Kuang Y, Chen X and Li J: CD147 mediates chemoresistance in breast cancer via ABCG2 by affecting its cellular localization and dimerization. *Cancer Lett* 337: 285-292, 2013.
- Wang Z, Wang N, Li W, Liu P, Chen Q, Situ H, Zhong S, Guo L, Lin Y, Shen J and Chen J: Caveolin-1 mediates chemoresistance in breast cancer stem cells via β -catenin/ABCG2 signaling pathway. *Carcinogenesis* 35: 2346-2356, 2014.
- Wu H, Liu Y, Kang H, Xiao Q, Yao W, Zhao H, Wang E and Wei M: Genetic variations in ABCG2 gene predict breast carcinoma susceptibility and clinical outcomes after treatment with anthracycline-based chemotherapy. *Biomed Res Int* 2015: 279109, 2015.
- Won HJ, Ha TK, Kwon SJ, Cho HY, Hur SJ, Baik HH, Suh SI, Ha E and Kim YH: Differential effects of 5-fluorouracil on glucose transport and expressions of glucose transporter proteins in gastric cancer cells. *Anticancer Drugs* 21: 270-276, 2010.
- Berlth F, Mönig S, Pinther B, Grimminger P, Maus M, Schlösser H, Plum P, Warnecke-Eberz U, Harismendy O, Drebbler U, *et al*: Both GLUT-1 and GLUT-14 are independent prognostic factors in gastric adenocarcinoma. *Ann Surg Oncol* 22 (Suppl 3): S822-S831, 2015.
- Schlößer HA, Drebbler U, Urbanski A, Haase S, Baltin C, Berlth F, Neiß S, von Bergwelt-Baildon M, Fetzner UK, Warnecke-Eberz U, *et al*: Glucose transporters 1, 3, 6, and 10 are expressed in gastric cancer and glucose transporter 3 is associated with UICC stage and survival. *Gastric Cancer* 20: 83-91, 2017.
- Dees EC and Watkins PB: Role of cytochrome P450 phenotyping in cancer treatment. *J Clin Oncol* 23: 1053-1055, 2005.
- Jernström H, Bågeman E, Rose C, Jönsson PE and Ingvar C: CYP2C8 and CYP2C9 polymorphisms in relation to tumour characteristics and early breast cancer related events among 652 breast cancer patients. *Br J Cancer* 101: 1817-1823, 2009.
- Boruban MC, Yasar U, Babaoglu MO, Sencan O and Bozkurt A: Tamoxifen inhibits cytochrome P450 2C9 activity in breast cancer patients. *J Chemother* 18: 421-424, 2006.
- Seredina TA, Goreva OB, Talaban VO, Grishanova AY and Lyakhovich VV: Association of cytochrome P450 genetic polymorphisms with neoadjuvant chemotherapy efficacy in breast cancer patients. *BMC Med Genet* 13: 45, 2012.
- Jamieson D, Lee J, Cresti N, Jackson R, Griffin M, Sludden J, Verrill M and Boddy AV: Pharmacogenetics of adjuvant breast cancer treatment with cyclophosphamide, epirubicin and 5-fluorouracil. *Cancer Chemother Pharmacol* 74: 667-674, 2014.
- Melanson SE, Stevenson K, Kim H, Antin JH, Court MH, Ho VT, Ritz J, Soiffer RJ, Kuo FC, Longtine JA and Jarolim P: Allelic variations in CYP2B6 and CYP2C19 and survival of patients receiving cyclophosphamide prior to myeloablative hematopoietic stem cell transplantation. *Am J Hematol* 85: 967-971, 2010.
- Beelen K, Opdam M, Severson TM, Koornstra RH, Vincent AD, Hauptmann M, van Schaik RH, Berns EM, Vermorken JB, van Diest PJ and Linn SC: CYP2C19 2 predicts substantial tamoxifen benefit in postmenopausal breast cancer patients randomized between adjuvant tamoxifen and no systemic treatment. *Breast Cancer Res Treat* 139: 649-655, 2013.
- van Schaik RH, Kok M, Sweep FC, van Vliet M, van Fessem M, Meijer-van Gelder ME, Seynaeve C, Lindemans J, Wesseling J, Van't Veer LJ, *et al*: The CYP2C19*2 genotype predicts tamoxifen treatment outcome in advanced breast cancer patients. *Pharmacogenomics* 12: 1137-1146, 2011.
- Komatsu T, Yamazaki H, Shimada N, Nakajima M and Yokoi T: Roles of cytochromes P450 1A2, 2A6, and 2C8 in 5-fluorouracil formation from tegafur, an anticancer prodrug, in human liver microsomes. *Drug Metab Dispos* 28: 1457-1463, 2000.
- Faul F, Erdfelder E, Lang AG and Buchner A: G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 39: 175-191, 2007.
- Benjamini Y, Drai D, Elmer G, Kafkafi N and Golani I: Controlling the false discovery rate in behavior genetics research. *Behav Brain Res* 125: 279-284, 2001.
- Niemi M, Pasanen MK and Neuvonen PJ: Organic anion transporting polypeptide 1B1: A genetically polymorphic transporter of major importance for hepatic drug uptake. *Pharmacol Rev* 63: 157-181, 2011.
- Custodio A, Moreno-Rubio J, Aparicio J, Gallego-Plazas J, Yaya R, Maurel J, Higuera O, Burgos E, Ramos D, Calatrava A, *et al*: Pharmacogenetic predictors of severe peripheral neuropathy in colon cancer patients treated with oxaliplatin-based adjuvant chemotherapy: A GEMCAD group study. *Ann Oncol* 25: 398-403, 2014.
- Ghafouri H, Ghaderi B, Amini S, Nikkhoo B, Abdi M and Hoseini A: Association of ABCB1 and ABCG2 single nucleotide polymorphisms with clinical findings and response to chemotherapy treatments in Kurdish patients with breast cancer. *Tumour Biol* 37: 7901-7906, 2016.
- Park SR, Kong SY, Nam BH, Choi JJ, Kim CG, Lee JY, Cho SJ, Kim YW, Ryu KW, Lee JH, *et al*: CYP2A6 and ERCC1 polymorphisms correlate with efficacy of S-1 plus cisplatin in metastatic gastric cancer patients. *Br J Cancer* 104: 1126-1134, 2011.

35. Park SR, Hong YS, Lim HS, Seong MW, Kong SY, Kim SY, Park YI and Jung KH: Phase I clinical and pharmacokinetic/pharmacogenetic study of a triplet regimen of S-1/irinotecan/oxaliplatin in patients with metastatic colorectal or gastric cancer. *Cancer Chemother Pharmacol* 72: 953-964, 2013.
36. Kim YW, Kim MJ, Ryu KW, Lim HS, Lee JH, Kong SY, Lee JS, Choi IJ, Kim CG, Lee JY, *et al*: A phase II study of perioperative S-1 combined with weekly docetaxel in patients with locally advanced gastric carcinoma: Clinical outcomes and clinicopathological and pharmacogenetic predictors for survival. *Gastric Cancer* 19: 586-596, 2016.
37. Kong SY, Lim HS, Nam BH, Kook MC, Kim YW, Ryu KW, Lee JH, Choi IJ, Lee JS, Park YI, *et al*: Association of CYP2A6 polymorphisms with S-1 plus docetaxel therapy outcomes in metastatic gastric cancer. *Pharmacogenomics* 10: 1147-1155, 2009.
38. Karadag O, Babaoglu MO, Altundag K, Elkiran T, Yasar U and Bozkurt A: 5-Fluorouracil-induced coronary spasm: May inhibition of hyperpolarization factors produced by CYP2C enzymes be the cause? *Oncology* 66: 510-511, 2004.
39. Helsby NA, Lo WY, Thompson P and Laking GR: Do 5-fluorouracil therapies alter CYP2C19 metaboliser status? *Cancer Chemother Pharmacol* 66: 405-407, 2010.
40. Afsar A, Lee C and Riddick DS: Modulation of the expression on constitutive rat hepatic cytochrome P450 isozymes by 5-fluorouracil. *Can J Physiol Pharmacol* 74: 150-156, 1996.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.