

Stimulated upregulation of *HO-1* is associated with inadequate response of gastric and ovarian cancer cell lines to hyperthermia and cisplatin treatment

VAIDOTAS CESNA¹, ARTURAS SUKOVAS², ALDONA JASUKAITIENE³, GIEDRE SILKUNIENE³, SAULIUS PASKAUSKAS², ZILVINAS DAMBRAUSKAS³ and ANTANAS GULBINAS³

Departments of ¹Surgery and ²Obstetrics and Gynaecology; ³Institute for Digestive Research, Lithuanian University of Health Sciences, LT-50161 Kaunas, Lithuania

Received September 9, 2018; Accepted February 21, 2019

DOI: 10.3892/ol.2019.10489

Abstract. Heme oxygenase (HO)-1 is a heat shock protein induced by hyperthermia, responsible for cellular resistance to temperature. The aim of this *in vitro* study was to clarify the response of gastric and ovarian cancer cells to hyperthermic intraperitoneal chemotherapy, following the modulation of *HO-1* expression. AGS and OVCAR-3 cells were treated with different temperature regimens, either alone or in combination with an IC₅₀ dose of cisplatin for 1 h. Prior to treatment, *HO-1* expression was silenced by short interfering RNA transfection. In OVCAR-3 cells, cisplatin increased *HO-1* mRNA expression by 3.73-fold under normothermia and 2.4-fold under hyperthermia; furthermore, these factors similarly increased *HO-1* protein expression levels. Exposure to cisplatin under hyperthermia reduced the viability of OVCAR-3 cells by 36% and *HO-1*-silencing enhanced this effect by 20%. *HO-1*-silencing under normothermia increased apoptotic rates in cisplatin-treated OVCAR-3 cells by 2.07-fold, and hyperthermia enhanced the effect by 3.09-fold. Semi-quantitative polymerase chain reaction (PCR) cell analysis indicated that exposure to cisplatin decreased the cell index under normothermia, and that hyperthermia boosted this effect in OVCAR-3. In AGS cells, only temperature increased cellular *HO-1* levels. Silencing *HO-1* in AGS cells at 37°C reduced viability by 16% and increased apoptotic rates 2.63-fold. Hyperthermia did not affect AGS viability; however, apoptosis was increased 6.84-fold. PCR analysis indicated no additional effects of hyperthermia on the AGS cell index. *HO-1* is induced in cancer cells by different stressors in a variable manner. In tumors with highly inducible *HO-1*, prior

silencing of this gene could improve the cellular response to hyperthermia and cisplatin.

Introduction

The treatment of cancer in the peritoneal cavity is a major problem in gastrointestinal and gynecological oncology worldwide (1,2). Globally, patients with gastric cancer with peritoneal carcinomatosis have a median survival time of 3-6 months, when the disease is left untreated (3). In the majority of newly diagnosed ovarian cancer cases, peritoneal metastases are already present (4).

Hyperthermic intraperitoneal chemotherapy (HIPEC) is a promising treatment option for intraperitoneally disseminated cancer (5-7). It has been reported to be more efficient compared with intravenous chemotherapy, however, as a treatment of peritoneal metastases it is limited by the plasma-peritoneal barrier (8,9). It has been previously suggested that hyperthermia can enhance the intraperitoneal application of cytotoxic agents (10,11). Residual microtumors, following cytoreductive surgery, are treated by intraperitoneal chemotherapy in combination with hyperthermia up to 42-43°C (12). Overall, hyperthermia as part of HIPEC is thought to boost pharmacokinetics and increase DNA damage induced by the cytostatic agent (13). Cisplatin is an alkylating drug used to treat residual gastric and ovarian cancer in the peritoneal cavity (14). Previous *in vitro* studies produced controversial data about the additivity of hyperthermia to cisplatin (15,16).

The response and resistance of cancer cells to chemotherapy and hyperthermia depend on the induction and expression of a number of cytoprotective proteins, including Hsp70 and Hsp27 (17-19). Therefore, the modulation of cytoprotective proteins may serve a crucial role in cancer treatment. One potential target is heme oxygenase (HO)-1, particularly in HIPEC, since high temperatures are a component of HIPEC, and it has been reported that, under hyperthermia, cells enhance *HO-1* expression for self-protective purposes (19). *HO-1* is normally expressed at low levels in the majority of tissues, including the gastrointestinal tract, female reproductive organs, brain and bone marrow (20); however, it is highly inducible by a variety of stimuli, including cytokines, lipopolysaccharides (21) and

Correspondence to: Mr. Vaidotas Cesna, Department of Surgery, Lithuanian University of Health Sciences, 2 Eiveniu Street, LT-50161 Kaunas, Lithuania
E-mail: vaidotas.cesna@lsmuni.lt

Key words: cisplatin, hyperthermic intraperitoneal chemotherapy, heme oxygenase-1

serine/threonine kinases (22). Cellular levels of *HO-1* are known to be temperature-dependent (23,24). *HO-1* is overexpressed under hyperthermic conditions, exerting a protective function (25,26).

An *in vitro* study was conducted to clarify the underlying mechanism of how cisplatin and hyperthermia induce *HO-1* expression in ovarian and gastric cancer cells. In addition, the present study investigated the response of these cancer cells to cisplatin and hyperthermia following the modulation of *HO-1* protein expression.

Materials and methods

Cell lines and conditions. Human gastric adenocarcinoma, AGS, and ovarian adenocarcinoma, OVCAR-3, cell lines were purchased from the American Type Tissue Culture Collection (Manassas, VA, USA). OVCAR-3 cells were cultivated in RPMI-1640 medium (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA) with 20% fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.), 1% penicillin/streptomycin and 0.01 mg/ml bovine insulin (Gibco; Thermo Fisher Scientific, Inc.). AGS cells were harvested in Ham's F-12K medium with 10% FBS and 1% penicillin/streptomycin at 37°C and 5% CO₂.

Experimental design. Cells were harvested for 24 h in the conditions described previously. The cells were subjected to conditions of normothermia (37°C) or 43°C and/or an IC₅₀ dose of cisplatin. The IC₅₀ dose was determined for each cell line individually in an experimental manner. The IC₅₀ doses of cisplatin for AGS and OVCAR-3 cells were determined (at 37°C) to be 111 and 152 µM, respectively. Hyperthermia and/or cisplatin exposure lasted for 1 h; this step began once the media reached the desired temperature (37 or 43°C), as measured by a digital thermometer in a humid incubator with a set temperature of 43°C. Following treatment, the medium was changed and cells were harvested after 48 h of incubation in a humidified atmosphere at 37°C with 5% CO₂. AGS and OVCAR-3 cell viability, apoptosis, and cell index were all subsequently measured. Additionally, these cell lines were used in real time cell analysis, western blotting and semi-quantitative polymerase chain reaction (qPCR) assays.

Silencing of *HO-1*. *HO-1* small interfering RNA (siRNA; 30 nM HMOX1; sense 5'-UGAACACUCUGGAGAUGAC-3', and antisense 5'-GUCAUCUCCAGAGUGUCCA-3') was obtained from Ambion; Thermo Fisher Scientific, Inc., and negative control (30 nM AllStars Negative Control siRNA; sense 5'-UUCUCCGAACGUGUCACGU-3' and antisense 5'-ACGUGACACGUUCGGAGAA-3') was obtained from Qiagen GmbH (Hilden, Germany). Lipofectamine® RNAiMAX (Invitrogen; Thermo Fisher Scientific, Inc.) and Opti-MEM™ media (Gibco; Thermo Fisher Scientific, Inc.) were used according to the manufacturer's protocols. The efficiency of transfection was verified using BLOCK-iT Alexa Fluor (Invitrogen; Thermo Fisher Scientific, Inc.). The efficiency of knockdown was verified by western blot analysis. *HO-1*-silencing was performed 72 h prior to implementation of the experimental temperature and treatment with cisplatin.

MTT assay. Cell viability was determined using an MTT assay (Invitrogen; Thermo Fisher Scientific, Inc.). Cells were incubated for 4 h at 37°C following the addition of 5 mg/ml MTT reagent. The supernatant was subsequently removed and dimethyl sulfide was added (Carl Roth GmbH Co KG, Karlsruhe, Germany). Absorbance was measured at a wavelength of 570 nm and the reference was measured at 650 nm using a Sunrise spectrophotometer (Tecan Austria GmbH, Grödig, Austria).

qPCR. Cellular RNA was extracted using a PureLink® RNA Mini kit (Ambion; Thermo Fisher Scientific, Inc.), according to the manufacturer's protocols. Purified RNA was measured and verified for purity using ultraviolet (UV) spectrophotometry (NanoDrop; Thermo Fisher Scientific, Inc.). Using the Super Script Vilo Master Mix (Invitrogen; Thermo Fisher Scientific, Inc.) with 2 µg RNA, cDNA was generated, according to the manufacturer's protocols. RNA amplification was performed in a 20 µl reaction volume, which contained 1X PCR Master Mix, primers, and 2 µl cDNA template. Thermocycling conditions were as follows: initial step at 95°C for 10 min (1 cycle), denaturation at 95°C for 15 sec and annealing/extending at 60°C for 1 min (40 cycles), followed by a final extension step at 72°C for 2 min. *HO-1* primers were obtained from Invitrogen (Thermo Fisher Scientific, Inc.): forward, 5'-TGCTCAACATCCAGCTCTTTGAGGA-3'; and reverse, 5'-CAGGCAGAGAATGCTGAGTTC-3'. The products were loaded on 1.5% agarose gels. Ethidium bromide staining and UV light (Gel Doc™ XR+ Gel Documentation System; Bio-Rad Laboratories, Inc., Hercules, CA, USA) were used for visualization. Analysis was performed using ImageLab software (version 6.0.0; Bio-Rad Laboratories, Inc.).

Flow cytometry. Apoptosis was evaluated by flow cytometry using Annexin V-PE and 7-aminoactinomycin D. A Guava Nexin Annexin V Assay kit (Merck KGaA, Darmstadt, Germany) was used according to the manufacturer's protocols. Analysis was performed with the Guava Personal Cell Analysis Flow Cytometer (Guava; EMD Millipore, Billerica, MA, USA) and CytoSoft software (version 2.1.4; Guava; EMD Millipore).

Western blot analysis. Lysates were prepared using radioimmunoprecipitation lysis buffer (Abcam, Cambridge, UK) containing protease inhibitors (Roche Diagnostics, Basel, Switzerland). A bicinchoninic acid protein assay kit (Thermo Fisher Scientific, Inc.) was used to determine the protein concentration, according to the manufacturer's protocols. Following heating at 97°C for 5 min, protein samples (50 µg) were subjected to 4-12% SDS-PAGE and transferred to polyvinylidene fluoride membranes at 30 V for 50 min. Membranes were blocked with a blocking buffer (20% diluent A, 30% diluent B; WesternBreeze Blocker/Diluent; Invitrogen; Thermo Fisher Scientific, Inc.) at room temperature for 1 h and incubated with the primary antibodies rabbit anti-*HO-1* (dilution, 1:2,000; cat. no., EP1391Y; Abcam) and mouse anti-GAPDH (dilution, 1:1,000; cat. no., AM4300; Ambion; Thermo Fisher Scientific, Inc.) at 4°C overnight. The following day, the blots were incubated with ready to use secondary antibodies against rabbit (cat. no. WP20007; Invitrogen, Thermo Fisher Scientific, Inc.)




Figure 1. *HO-1* expression is differentially induced by hyperthermia and cisplatin treatment in ovarian adenocarcinoma, OVCAR-3, and gastric adenocarcinoma, AGS, cell lines. (A) *HO-1* expression was activated by cisplatin in OVCAR-3 cells, whereas hyperthermia was not indicated to have an additional effect. (B) In AGS cells, only hyperthermia indicated to induce *HO-1* expression. * $P < 0.05$, compared with the control group (not treated with cisplatin, normothermia). *HO-1*, heme oxygenase-1.

or mouse immunoglobulin G (cat. no. WP20006; Invitrogen; Thermo Fisher Scientific, Inc.) for 1 h at room temperature. Chemiluminescence substrate (CDP-Star; Invitrogen; Thermo Fisher Scientific, Inc.) was added and the ChemiDoc imaging system (Bio-Rad Laboratories, Inc.) was used for visualization. ImageJ software (version 1.48; National Institutes of Health, Bethesda, MD, USA) was used for quantification of western blots (27).

Real time cell analysis. The xCELLigence® RTCA DP Real-Time Analyzer (ACEA Biosciences Inc., San Diego, CA, USA) was used to investigate the PCR cellular response to treatment. To present the real time cell analysis data obtained by xCELLigence system, the cell index was used. The determination of the cell index parameter is an automatic system feature and is based on the rapid measurements of impedance between gold electrodes in the analyzer wells (28). Cell-free medium (RPMI-1640; Gibco; Thermo Fisher Scientific, Inc.) and medium with cells have different cell impedance and proliferation indices, which is reflected in changes in impedance value and CI; this is calculated as follows: (impedancetime point n -impedancebackground)/nominal impedance value. Delta cell index was measured immediately when cells reached the electronic microplate. Cells were evaluated for 48 h following the experiment and/or 72 h following *HO-1*-silencing.

Statistical analysis. SPSS v.21.0 software (IBM Corp., Armonk, NY, USA) and GraphPad (version 6.01; GraphPad Software Inc., La Jolla, CA, USA) were used for statistical evaluation. The Mann-Whitney test and one-way analysis of variance with a Bonferroni post hoc test were performed to assess clinical significance. Data are presented as the mean \pm standard deviation of three independent experiments, performed in triplicate. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Hyperthermia and cisplatin differentially induce HO-1 mRNA and protein expression in ovarian and gastric cancer cells.

Enhanced expression levels of *HO-1* mRNA were only observed in OVCAR-3 cells. The exposure of OVCAR-3 cells to cisplatin resulted in a significant increase of *HO-1* mRNA expression. Cisplatin induced a 3.75- and 2.4-fold increase of *HO-1* expression in conditions of normothermia (37°C) and hyperthermia (43°C), respectively ($P < 0.05$). While hyperthermia at 43°C boosted *HO-1* expression by 1.34-fold, the addition of cisplatin increased the effect on *HO-1* expression by 3.22-fold ($P < 0.05$; Fig. 1A). In AGS cells, *HO-1* expression was not significantly affected by temperature or cisplatin (Fig. 1B).

Furthermore, cisplatin significantly increased *HO-1* protein expression in OVCAR-3 cells ($P < 0.05$). *HO-1* expression was increased 9.5-fold, following cisplatin treatment under normothermia. At 43°C, this effect was slightly higher, with a 9.77-fold increase ($P < 0.05$). The exposure of OVCAR-3 cells to hyperthermia alone had no effect on *HO-1* protein expression levels. However, the combination of cisplatin and hyperthermia increased *HO-1* protein expression by 11-fold compared with the control (untreated cells in normothermia; $P < 0.01$; Fig. 2A).

In AGS cells, cisplatin had no notable effect on *HO-1* expression. At 37°C, cisplatin increased *HO-1* protein expression in AGS cells by 1.2-fold; however, this increase was not statistically significant, whereas the exposure of cells to 43°C in the absence of cisplatin increased *HO-1* protein expression levels by 2.75-fold ($P < 0.05$). *HO-1* expression dropped slightly when cisplatin was added at 43°C. Therefore, concomitant treatment of AGS cells with cisplatin and hyperthermia at 43°C resulted in a 2.14-fold increase in *HO-1* protein compared with the control (Fig. 2B). Furthermore, *HO-1* knockdown was assessed by western blotting (Fig. 2C and D).

HO-1-silencing does not influence AGS cell viability. The results of the MTT assay revealed that *HO-1*-silencing in OVCAR-3 cells does not affect viability in response to cisplatin at 37°C. The exposure of OVCAR-3 cells to cisplatin and hyperthermia (43°C) resulted in a 36% drop in cell viability ($P < 0.05$). *HO-1*-silencing enhanced this effect by an additional 20% ($P < 0.05$; Fig. 3A).

HO-1-silencing in AGS cells enhanced the cisplatin effect and reduced cell viability by 16% at 37°C ($P < 0.05$).




Figure 2. *HO-1* protein expression is activated by cisplatin or hyperthermia. (A) *HO-1* expression increased following cisplatin treatment of OVCAR-3 cells. Hyperthermia-dependent activation of *HO-1* expression did not indicate any significant differences between 37 and 43°C. (B) In AGS cells, hyperthermia induced expression of *HO-1*, while treatment with cisplatin had no significant effect. *HO-1*-knockdown decreased expression levels of *HO-1* protein in (C) OVCAR-3 and (D) AGS cells compared with corresponding cells treated under normothermia and without cisplatin. * $P < 0.05$, compared with the control group (not treated with cisplatin, normothermia). Western blotting data is shown by selecting the most representative experiment. *HO-1*, heme oxygenase-1; siRNA, small interfering RNA.




Figure 3. Modulating *HO-1* expression differentially influences the viability of OVCAR-3 cells, however not AGS cells. (A) *HO-1*-silencing had no significant effect on the viability of OVCAR-3 cells treated with cisplatin in conditions of normothermia. *HO-1*-silencing enhanced the reduction of cell viability, following treatment with cisplatin and hyperthermia. (B) *HO-1*-silencing slightly decreased AGS cell viability, following treatment with cisplatin under normothermia, however there was no significant effect with hyperthermia. * $P < 0.05$, compared with the cells treated with cisplatin under normothermia. Data are equalized to untreated cells-100%. *HO-1*, heme oxygenase-1; siRNA, small interfering RNA.

Hyperthermia potentiated cisplatin cytotoxicity in AGS cells: viability dropped by 24% compared with 37°C. However, *HO-1*-silencing had no significant additional effect, whereas viability rates were similar in *HO-1*-silenced or unsilenced AGS cells following cisplatin treatment at 43°C (Fig. 3B).

HO-1-silencing prior to concomitant hyperthermia and cisplatin treatment increases ovarian and gastric cancer cell apoptosis. The exposure of OVCAR-3 and AGS cells to hyperthermia resulted in a better cell response to cisplatin with respect to apoptosis. Prior *HO-1*-silencing under normothermia increased cisplatin-induced apoptosis in OVCAR-3 and AGS cells by 2.07- and 2.63-fold, respectively. In addition, silencing of *HO-1* under hyperthermia enhanced the apoptosis of OVCAR-3 and AGS cells by 3.09- and 6.84-fold, respectively ($P < 0.05$; Fig. 4).

Hyperthermia enhances the effect of cisplatin on OVCAR-3, but not on AGS cells, following modulation of HO-1 expression. PCR analysis for 48 h following treatment indicated

that exposure to cisplatin resulted in a gradual decrease in the cell index of AGS and OVCAR-3 (*HO-1*-silenced) cells at 37°C. Hyperthermia at 43°C boosted this effect by inducing a gradual decrease of the OVCAR-3 (*HO-1*-silenced) cell index. However, the cell index of AGS (*HO-1*-silenced) cells following cisplatin treatment at 37 or 43°C remained similar (Fig. 5).

Discussion

In the present study, the *HO-1* protein was variably expressed at the basal level and variably induced following exposure to cisplatin and hyperthermia in OVCAR-3 and AGS cells. Cisplatin increased the expression levels of *HO-1* in OVCAR-3 cells, while hyperthermia at 43°C had no effect. In AGS cells, *HO-1* expression was slightly increased under hyperthermia, with no significant induction following exposure to cisplatin, indicating that the modulation of *HO-1* may serve a role in the response of cancer cells to cisplatin and hyperthermia and affect cancer treatment outcomes.




Figure 4. Prior *HO-1*-silencing and subsequent hyperthermia/cisplatin treatment enhances apoptosis in OVCAR-3 and AGS cells. (A) *HO-1*-silencing enhanced apoptosis in OVCAR-3 cells treated with cisplatin at 37°C. This effect was even more evident under conditions of hyperthermia. (B) *HO-1*-silencing significantly enhanced apoptosis in cisplatin-treated AGS cells under hyperthermia compared with normothermia. Dot plots represent the raw data of the corresponding bar. Dashed lines indicate apoptosis in control (untreated) cells. *P<0.05. The data were obtained from five independent experiments. *HO-1*, heme oxygenase-1; siRNA, small interfering RNA.




Figure 5. Hyperthermia enhances the effect of cisplatin only on OVCAR-3 cells. Exposure to cisplatin gradually decreased the cell index of OVCAR-3 and AGS (*HO-1* silenced) cells under normothermia. Hyperthermia enhanced this effect in OVCAR-3 cells, while the AGS cell index did not change. The data are shown by selecting the most representative experiment. *HO-1*, heme oxygenase-1; siRNA, small interfering RNA.

HIPEC is widely used in clinical settings, and promising results have been reported in the treatment of peritoneal dissemination of gastric and ovarian cancer (29,30). To the best of our knowledge, to date, there has been a lack of evidence regarding the synergy of chemotherapy and hyperthermia. In our previous studies, it was observed that gastrointestinal and ovarian cancer cells responded unpredictably following exposure to cisplatin and hyperthermia (16,31). One of the limits of this response may be the induction of cytoprotective enzymes associated with chemotherapy and/or hyperthermia, in particular *HO-1*. *HO-1* is known to be highly expressed in human gastric and ovarian cancer tissue (32). Anticancer treatment options, including chemotherapy and radiotherapy may increase *HO-1* expression (33). *HO-1* serves an important role in a number of pathophysiological conditions, including temperature rise and inflammation, and has been reported to be associated with cancer (34,35). *HO-1* expression is associated with cancer growth and progression by promoting angiogenesis in the tumor

itself and metastases and pro-proliferation in different types of tumors, including renal cell carcinoma, prostate and pancreatic cancer, melanoma, and hepatoma (36-40). Numerous studies have highlighted that cancer cells with high expression levels of *HO-1* are less sensitive to cisplatin treatment compared with cancer cells with low *HO-1* expression levels (41,42).

The mechanism underlying this cytoprotective effect relies on the ability of *HO-1* to catabolize free heme and prevent it from sensitizing cells to undergo programmed cell death (43). *HO-1* under normal conditions has various cellular functions, including catalyzing the heme molecule to form bile pigments (44). When stimulus (heat) is present, cellular *HO-1* synthesis is enhanced (45). Therefore, the present study suggests that *HO-1* is crucially important when dealing with intraperitoneally-spread cancer, particularly treating it with HIPEC. Following the administration of heated chemotherapy drugs into the abdominal cavity, tumor cells should start to defend themselves, by activating heat shock proteins, particu-

larly *HO-1*. The aim of the present study is to achieve better treatment results by downregulating *HO-1* expression.

To the best of our knowledge, there are no published data on the efficacy of HIPEC treatment while modulating the expression of *HO-1*. The results of our study demonstrate the impact of *HO-1* expression modulation in the combination treatment of hyperthermia at 43°C and cisplatin in OVCAR-3 cells.

Zhao *et al* (46) reported that the basal level of *HO-1* expression is higher in ovarian cancer cells compared with normal ovarian tissues. A high level of *HO-1* expression has also been associated with aggressive tumors and poor clinical outcomes (46). The ability of cisplatin to increase the expression of *HO-1* was also observed in different cancer types, including pancreatic and hepatic cancer (43,47,48). Was *et al* (37) reported the different abilities of tumor tissues to produce heat shock proteins. Nonetheless, a high level of *HO-1* is known to be associated with the reduced tumor growth observed in some types of cancers, including breast and prostate cancer and non-small-cell lung carcinoma (37).

The results of the present study indicate that the viability of *HO-1*-silenced OVCAR-3 cells decreased significantly following cisplatin treatment at 43°C. However, in AGS cells, the inhibition of *HO-1* did not improve the response to cisplatin treatment. These results are associated with the expression of *HO-1*. It is possible that the inhibition of *HO-1* only increases the effect of cisplatin in cancer cells, where *HO-1* is highly expressed. This is in accordance with the data reported by Lv *et al* (41), where cisplatin significantly induced the expression of *HO-1*. The study modulated *HO-1* expression using hemin (an inducer of *HO-1*) and ZnPPiX (an inhibitor of *HO-1*), and reported that hemin strongly inhibited cisplatin-induced cell death, while ZnPPiX significantly increased cell death (41,49). These effects following *HO-1* modulation can be explained by the cytoprotective ability of this protein. *HO-1* activates a cellular defense mechanism against oxidative stress through its catalytic products, including ferrous iron, carbon monoxide, and biliverdin (37). In addition, growing evidence has suggested that *HO-1* protects cells from chemotherapeutic agent-induced apoptosis, and the targeted knockdown of *HO-1* gene expression or suppression of *HO-1* activity *in vitro* significantly enhances the chemosensitivity of cancer cells (50). Furthermore, it has been reported that the inhibition of *HO-1* can increase cellular response to anticancer treatment (26).

Cisplatin can effectively induce and promote apoptosis in a wide range of solid tumors, including head and neck cancer, esophageal carcinoma, non-small cell lung carcinoma, and testicular, cervical, and ovarian cancer (51). Inhibition of *HO-1* may strengthen the pro-apoptotic effects of cisplatin (41). In the present study, the inhibition of *HO-1* increased the number of apoptotic cells in OVCAR-3 and AGS cell lines, however these results did not indicate any significant differences associated with *HO-1* expression and cell viability. This could be explained by the fact that the present study measured the number of cells in both early and late apoptosis, and early apoptosis can be reversible (52). Geske *et al* (52) reported that the early stages of apoptosis are reversible if the apoptotic stimulus is removed, which is the reason that PCR analysis was performed in the present study.

In this experimental model, hyperthermia alone did not induce the upregulation of *HO-1* expression in the tested cancer

cell lines. Nevertheless, *HO-1*-silencing resulted in the optimal response to cisplatin treatment in terms of cell viability in OVCAR-3 cells, and apoptosis in both OVCAR-3 and AGS cells, under conditions of hyperthermia. Therefore, a novel finding regarding the role of *HO-1* in HIPEC is presented in this study. In conclusion, the cytoprotective protein *HO-1* is induced in cancer cells by different stressors in a variable manner. In tumors with highly inducible *HO-1*, the present study indicated that prior silencing of this gene may significantly improve the cellular response to hyperthermia and cisplatin.

Acknowledgements

Not applicable.

Funding

This study was supported by the Research Council of Lithuania (grant no. SEN-01/2015).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

VC wrote the manuscript and analyzed the data. AS performed the western blot analysis and analyzed the data. GS and AJ performed semi-quantitative PCR and siRNA transfection. SP performed the PCR cell analysis and analyzed the data. ZD and AG revised the data and manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Shariff U, Seretis C and Youssef H: Management of colorectal cancer patients at high risk of peritoneal metastases. *J BUON* 20 (Suppl 1): S71-S79, 2015.
2. Zivanovic O, Chi DS, Filippova O, Randall LM, Bristow RE and O'Cearbhaill RE: It's time to warm up to hyperthermic intraperitoneal chemotherapy for patients with ovarian cancer. *Gynecol Oncol* 151: 555-561, 2018.
3. Yonemura Y, Endou Y, Sasaki T, Hirano M, Mizumoto A, Matsuda T, Takao N, Ichinose M, Miura M and Li Y: Surgical treatment for peritoneal carcinomatosis from gastric cancer. *Eur J Surg Oncol* 36: 1131-1138, 2010.
4. Stewart JH IV, Shen P and Levine EA: Intraperitoneal hyperthermic chemotherapy for peritoneal surface malignancy: Current status and future directions. *Ann Surg Oncol* 12: 765-777, 2005.
5. Stirrups R: HIPEC improves survival in stage III epithelial ovarian cancer. *Lancet Oncol* 19: e138, 2018.

6. Tentes AA, Pallas N, Karamveri C, Kyziridis D and Hristakis C: Cytoreduction and HIPEC for peritoneal carcinomatosis of pancreatic cancer. *J BUON* 23: 482-487, 2018.
7. Tonello M, Ortega-Perez G, Alonso-Casado O, Torres-Mesa P, Guíñez G and Gonzalez-Moreno S: Peritoneal carcinomatosis arising from rectal or colonic adenocarcinoma treated with cytoreductive surgery (CRS) hyperthermic intraperitoneal chemotherapy (HIPEC): Two different diseases. *Clin Transl Oncol* 20: 1268-1273, 2018.
8. Jacquet P and Sugarbaker PH: Peritoneal-plasma barrier. *Cancer Treat Res* 82: 53-63, 1996.
9. Markman M: Intraperitoneal chemotherapy in the management of malignant disease. *Expert Rev Anticancer Ther* 1: 142-148, 2001.
10. Helm CW: The role of hyperthermic intraperitoneal chemotherapy (HIPEC) in ovarian cancer. *Oncologist* 14: 683-694, 2009.
11. Dovern E, de Hingh IH, Verwaal VJ, van Driel WJ and Nienhuijs SW: Hyperthermic intraperitoneal chemotherapy added to the treatment of ovarian cancer: a review of achieved results and complications. *Eur J Gynaecol Oncol* 31: 256-261, 2010.
12. Zhu Y, Hanna N, Boutros C and Alexander HR Jr: Assessment of clinical benefit and quality of life in patients undergoing cytoreduction and Hyperthermic Intraperitoneal Chemotherapy (HIPEC) for management of peritoneal metastases. *J Gastrointest Oncol* 4: 62-71, 2013.
13. van de Vaart PJ, van der Vange N, Zoetmulder FA, van Goethem AR, van Tellingen O, ten Bokkel Huinink WW, Beijnen JH, Bartelink H and Begg AC: Intraperitoneal cisplatin with regional hyperthermia in advanced ovarian cancer: pharmacokinetics and cisplatin-DNA adduct formation in patients and ovarian cancer cell lines. *Eur J Cancer* 34: 148-154, 1998.
14. Yan TD, Cao CQ and Munkholm-Larsen S: A pharmacological review on intraperitoneal chemotherapy for peritoneal malignancy. *World J Gastrointest Oncol* 2: 109-116, 2010.
15. Kimura E and Howell SB: Analysis of the cytotoxic interaction between cisplatin and hyperthermia in a human ovarian carcinoma cell line. *Cancer Chemother Pharmacol* 32: 419-424, 1993.
16. Sukovas A, Cesna V, Jasukaitiene A, Barauskas G, Nadisauskiene RJ, Dambrauskas Z, Paskauskas S and Gulbinas A: Response of OVCAR-3 cells to cisplatin and hyperthermia: does hyperthermia really matter? *Anticancer Res* 37: 5011-5018, 2017.
17. Leung AW, Hung SS, Backstrom I, Ricaurte D, Kwok B, Poon S, McKinney S, Segovia R, Rawji J, Qadir MA, *et al*: Combined use of gene expression modeling and siRNA screening identifies genes and pathways which enhance the activity of cisplatin when added at no effect levels to non-small cell lung cancer cells in vitro. *PLoS One* 11: e0150675, 2016.
18. Skowron MA, Niegisch G, Albrecht P, van Koeveringe G, Romano A, Albers P, Schulz WA and Hoffmann MJ: Various mechanisms involve the nuclear factor (erythroid-derived 2)-like (NRF2) to achieve cytoprotection in long-term cisplatin-treated urothelial carcinoma cell lines. *Int J Mol Sci* 18: pii: E1680, 2017.
19. Kumar S, Stokes J III, Singh UP, Scisum Gunn K, Acharya A, Manne U and Mishra M: Targeting Hsp70: a possible therapy for cancer. *Cancer Lett* 374: 156-166, 2016.
20. Thul PJ, Åkesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A, Björk L, Breckels LM *et al*: A subcellular map of the human proteome. *Science* 356: pii: eaal3321, 2017.
21. Rushworth SA and O'Connell MA: Haem oxygenase-1 in inflammation. *Biochem Soc Trans* 32: 1093-1094, 2004.
22. Nemmiche S, Chabane-Sari D, Kadri M and Guiraud P: Cadmium-induced apoptosis in the BJAB human B cell line: involvement of PKC/ERK1/2/JNK signaling pathways in HO-1 expression. *Toxicology* 300: 103-111, 2012.
23. Shibahara S, Müller RM and Taguchi H: Transcriptional control of rat heme oxygenase by heat shock. *J Biol Chem* 262: 12889-12892, 1987.
24. Vile GF, Basu-Modak S, Waltner C and Tyrrell RM: Heme oxygenase 1 mediates an adaptive response to oxidative stress in human skin fibroblasts. *Proc Natl Acad Sci USA* 91: 2607-2610, 1994.
25. Yang RC, Chang CY, Lu TS and Chen SC: Effects of hyperthermia pretreatment on expression of heme oxygenase-1 and nitric oxide synthase in rats subjected to experimental anaphylactic shock. *Chin J Physiol* 48: 193-199, 2005.
26. Berberat PO, Dambrauskas Z, Gulbinas A, Giese T, Giese N, Künzli B, Autschbach F, Meuer S, Büchler MW and Friess H: Inhibition of heme oxygenase-1 increases responsiveness of pancreatic cancer cells to anticancer treatment. *Clin Cancer Res* 11: 3790-3798, 2005.
27. Schneider CA, Rasband WS and Eliceiri KW: NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9: 671-675, 2012.
28. Ke N, Wang X, Xu X and Abassi YA: The xCELLigence system for real-time and label-free monitoring of cell viability. *Methods Mol Biol* 740: 33-43, 2011.
29. Kang LY, Mok KT, Liu SI, Tsai CC, Wang BW, Chen IS, Chen YC, Chang BM and Chou NH: Intraoperative hyperthermic intraperitoneal chemotherapy as adjuvant chemotherapy for advanced gastric cancer patients with serosal invasion. *J Chin Med Assoc* 76: 425-431, 2013.
30. Halkia E and Spiliotis J: The role of cytoreductive surgery and HIPEC in epithelial ovarian cancer. *J BUON* 20 (Suppl 1): S12-S28, 2015.
31. Cesna V, Sukovas A, Jasukaitiene A, Naginiene R, Barauskas G, Dambrauskas Z, Paskauskas S and Gulbinas A: Narrow line between benefit and harm: additivity of hyperthermia to cisplatin cytotoxicity in different gastrointestinal cancer cells. *World J Gastroenterol* 24: 1072-1083, 2018.
32. Papatheodorou I, Fonseca NA, Keays M, Tang YA, Barrera E, Bazant W, Burke M, Füllgrabe A, Fuentes AM, George N, *et al*: Expression Atlas: gene and protein expression across multiple studies and organisms. *Nucleic Acids Res* 46 (D1): D246-D251, 2018.
33. Nitti M, Piras S, Marinari UM, Moretta L, Pronzato MA and Furfaro AL: HO-1 induction in cancer progression: a matter of cell adaptation. *Antioxidants (Basel)* 6: pii: E29, 2017.
34. Chau LY: Heme oxygenase-1: Emerging target of cancer therapy. *J Biomed Sci* 22: 22, 2015.
35. Wang TY, Liu CL, Chen MJ, Lee JJ, Pun PC and Cheng SP: Expression of haem oxygenase-1 correlates with tumour aggressiveness and BRAF V600E expression in thyroid cancer. *Histopathology* 66: 447-456, 2015.
36. Loboda A, Jozkowicz A and Dulak J: HO-1/CO system in tumor growth, angiogenesis and metabolism-Targeting HO-1 as an anti-tumor therapy. *Vascul Pharmacol* 74: 11-22, 2015.
37. Was H, Dulak J and Jozkowicz A: Heme oxygenase-1 in tumor biology and therapy. *Curr Drug Targets* 11: 1551-1570, 2010.
38. Becker JC, Fukui H, Imai Y, Sekikawa A, Kimura T, Yamagishi H, Yoshitake N, Pohle T, Domschke W and Fujimori T: Colonic expression of heme oxygenase-1 is associated with a better long-term survival in patients with colorectal cancer. *Scand J Gastroenterol* 42: 852-858, 2007.
39. Goodman AI, Choudhury M, da Silva JL, Schwartzman ML and Abraham NG: Overexpression of the heme oxygenase gene in renal cell carcinoma. *Proc Soc Exp Biol Med* 214: 54-61, 1997.
40. Maines MD and Abrahamsson PA: Expression of heme oxygenase-1 (HSP32) in human prostate: normal, hyperplastic, and tumor tissue distribution. *Urology* 47: 727-733, 1996.
41. Lv X, Song DM, Niu YH and Wang BS: Inhibition of heme oxygenase-1 enhances the chemosensitivity of laryngeal squamous cell cancer Hep-2 cells to cisplatin. *Apoptosis* 21: 489-501, 2016.
42. Jeon WK, Hong HY, Seo WC, Lim KH, Lee HY, Kim WJ, Song SY and Kim BC: Smad7 sensitizes A549 lung cancer cells to cisplatin-induced apoptosis through heme oxygenase-1 inhibition. *Biochem Biophys Res Commun* 420: 288-292, 2012.
43. Gozzelino R, Jeney V and Soares MP: Mechanisms of cell protection by heme oxygenase-1. *Annu Rev Pharmacol Toxicol* 50: 323-354, 2010.
44. Ewing JF, Haber SN and Maines MD: Normal and heat-induced patterns of expression of heme oxygenase-1 (HSP32) in rat brain: Hyperthermia causes rapid induction of mRNA and protein. *J Neurochem* 58: 1140-1149, 1992.
45. Choi AM and Alam J: Heme oxygenase-1: Function, regulation, and implication of a novel stress-inducible protein in oxidant-induced lung injury. *Am J Respir Cell Mol Biol* 15: 9-19, 1996.
46. Zhao Z, Xu Y, Lu J, Xue J and Liu P: High expression of HO-1 predicts poor prognosis of ovarian cancer patients and promotes proliferation and aggressiveness of ovarian cancer cells. *Clin Transl Oncol* 20: 491-499, 2018.
47. Doi K, Akaike T, Fujii S, Tanaka S, Ikebe N, Beppu T, Shibahara S, Ogawa M and Maeda H: Induction of haem oxygenase-1 nitric oxide and ischaemia in experimental solid tumours and implications for tumour growth. *Br J Cancer* 80: 1945-1954, 1999.

48. Nuhn P, Künzli BM, Hennig R, Mitkus T, Ramanauskas T, Nobiling R, Meuer SC, Friess H and Berberat PO: Heme oxygenase-1 and its metabolites affect pancreatic tumor growth in vivo. *Mol Cancer* 8: 37, 2009.
49. Kongpetch S, Kukongviriyapan V, Prawan A, Senggunprai L, Kukongviriyapan U and Buranrat B: Crucial role of heme oxygenase-1 on the sensitivity of cholangiocarcinoma cells to chemotherapeutic agents. *PLoS One* 7: e34994, 2012.
50. Gueron G, Giudice J, Valacco P, Paez A, Elguero B, Toscani M, Jaworski F, Leskow FC, Cotignola J, Marti M, *et al*: Heme-oxygenase-1 implications in cell morphology and the adhesive behavior of prostate cancer cells. *Oncotarget* 5: 4087-4102, 2014.
51. Barry MA, Behnke CA and Eastman A: Activation of programmed cell death (apoptosis) by cisplatin, other anticancer drugs, toxins and hyperthermia. *Biochem Pharmacol* 40: 2353-2362, 1990.
52. Geske FJ, Lieberman R, Strange R and Gerschenson LE: Early stages of p53-induced apoptosis are reversible. *Cell Death Differ* 8: 182-191, 2001.