

# ***NRP-1* and *KDR* polymorphisms are associated with survival time in patients with advanced gastric cancer**

YUE-JIAN ZHUO\*, YU SHI\* and TAO WU

Department of Medical Oncology, The Second Hospital Affiliated to Dalian Medical University,  
Dalian, Liaoning 116000, P.R. China

Received March 15, 2019; Accepted August 1, 2019

DOI: 10.3892/ol.2019.10842

**Abstract.** Neuropilin-1 (*NRP-1*), a member of the NRP-family, has been reported to be vital for tumor angiogenesis, growth and metastasis. As a co-receptor of vascular endothelial growth factor (*VEGF*), *NRP-1* can bind to *VEGF* and mediate vascular development through the *VEGF-VEGF* receptor 2 (*VEGFR2*) signaling pathway. Furthermore, *NRP-1* is capable of binding with platelet-derived growth factor (*PDGF*) to regulate the *PDGF-PDGF* receptor (*PDGR*) signaling pathway in tumor angiogenesis. In the present study, The DNA was obtained from the paraffin-embedded tissues of patients with advanced gastric cancer (AGC), amplified using PCR and subsequently sequenced to determine the polymorphisms within *NRP-1*, *VEGFR2* [kinase insert domain receptor (*KDR*)] and *PDGF*. The effect of the functional polymorphism of the aforementioned genes on the overall survival (OS) and progression-free survival (PFS) of 81 patients with advanced gastric cancer was examined. Three single nucleotide polymorphisms (SNPs) of *KDR* were significantly associated with clinical outcomes. The rs1870377 TT genotype was positively associated with longer OS and PFS times compared with the AA+AT genotype (PFS, P=0.012; OS, P=0.038), the rs7692791 wild-type TT genotype was positively associated with longer PFS time and the rs2034965 AA+GA genotype was associated with shorter OS time (P=0.034). With regards to the SNPs of *NRP-1*, the rs2065364 AA genotype was significantly associated with improved OS and PFS times (PFS, P=0.023; OS, P=0.045). Following multivariate analysis using Cox proportional hazards regression models, patients with the *KDR* rs7692791 TT genotype experienced a longer PFS time compared with those with the CT genotype (P=0.016), and patients with the

*NRP-1* rs2065364 variant-type AA genotype still experienced a longer PFS time compared with those patients with the AG+GG genotypes (P=0.006). Regarding OS, the results demonstrated that the *KDR* rs2034965 AG+GG genotypes presented with a significant reduction in OS time (P=0.029), and that the *KDR* rs1870377 AT+AA genotypes had worse OS times compared with the wild-type TT genotype (P=0.021). In addition, increased mortality risk and AGC progression were significantly associated with the number of adverse alleles for combinations of *NRP-1* rs2065364 and *KDR* rs1870377. In conclusion, the data from the present study demonstrated that the selected *KDR* and *NRP-1* gene polymorphisms may be potential prognostic biomarkers in AGC.

## **Introduction**

Gastric cancer is one of the most common cancer types worldwide and remains the third leading cause of cancer-associated mortality, accounting for >783,000 deaths worldwide in 2018 (1). The diagnostic rate of early gastric cancer is only 10% in China, and most patients are at an advanced stage when clinically diagnosed, which confers a poor prognosis (2).

Systematic chemotherapy is the major treatment for patients with advanced gastric cancer (AGC). Platinum-fluoropyrimidine- and paclitaxel-fluoropyrimidine-based chemotherapy regimens are recommended as the first-line treatments in line with the Chinese Society of Clinical Oncology guidelines (3).

However, patients with the same tumor stage and receiving similar treatment can exhibit different clinical outcomes, and gastric cancer is a complex disease and its prognosis and progression are significantly affected by genetic and environmental factors (4). Identifying predictive genetic biomarkers could therefore contribute to the development of individualized therapy and follow-up strategies (5).

Neuropilin-1 (*NRP-1*) is a type I transmembrane glycoprotein distributed at the surface of cells that has been reported to affect neuronal axon guidance and embryonic angiogenesis (6), and to serve as a co-receptor regulating tumorigenesis in the vascular endothelial growth factor (*VEGF*)-*VEGF* receptor 2 (*VEGFR2*) [kinase insert domain receptor (*KDR*)] or platelet-derived growth factor (*PDGF*)-*PDGF* receptor (*PDGFR*) signaling pathways (7,8). *VEGFRs* are a type of tyrosine kinase receptor, and include *VEGFR1* and *VEGFR2*

*Correspondence to:* Professor Tao Wu, Department of Medical Oncology, The Second Hospital Affiliated to Dalian Medical University, 467 Zhongshan Road, Dalian, Liaoning 116000, P.R. China

E-mail: wuyihong97@163.com

\*Contributed equally

**Key words:** gastric cancer, polymorphism, *NRP-1*, *KDR*, *PDGF*

(*KDR*), which can be activated by binding with *VEGF* ligands (9). The *VEGF-VEGFR2* signaling pathway is the leading pathway that activates the proliferation and migration of endothelial cells, therefore promoting angiogenesis and stimulating tumor growth and invasion (10,11). Furthermore, *PDGF* isoforms can transduce signals via binding to structurally similar  $\alpha$ - and  $\beta$ -tyrosine kinase receptors, known as *PDGFR $\alpha$*  and *PDGFR $\beta$* , respectively. The *PDGF-PDGFR* signaling pathway serves critical roles in regulating proliferation and survival of certain cell types (e.g. hematopoietic stem cell, vascular endothelial cell and vascular smooth muscle cell) during embryogenesis, and overexpression or mutation of the *PDGF-PDGFR* pathway can stimulate tumor cell proliferation (12,13). Previous studies have reported that polymorphisms within *VEGF* and *KDR* impacted their expression at the gene level (14,15). Thus, polymorphisms of these two signaling pathways may affect AGC prognosis by regulating the expression of the aforementioned genes and therefore affect the survival of patients with AGC. The present study investigated the association between polymorphisms of the *NRP-1*, *KDR*, *PDGF $\beta$* , *PDGFR $\beta$*  and *PDGFR $\alpha$*  genes and the prognosis of patients with AGC.

## Materials and methods

**Study population.** A total of 100 patients with AGC from the Second Affiliated Hospital of Dalian Medical University (Dalian, Liaoning, China) were recruited between January 2011 and June 2016. The inclusion criteria were as follows: i) Patients were histopathologically diagnosed with gastric adenocarcinoma; ii) patients had inoperable locally advanced, metastatic or recurrent gastric cancer (AGC); iii) patients had an Eastern Cooperative Oncology Group performance status (ECOG-PS)  $\leq 2$  (16); and iv) patients underwent at least 2 cycles of chemotherapy at the Second Affiliated Hospital when diagnosed with postoperative recurrence or inoperable advanced gastric cancer. The exclusion criteria were as follows: i) Patients who received chemotherapy, radiotherapy and/or biological treatment previously; ii) patients with an ECOG-PS  $>2$ ; and iii) patients with multiple primary malignant neoplasms. The 100 patients were followed up by clinic visits and phone calls every 2 months, and clinical outcomes were recorded until October 2018. Genotype information was not available for 8 patients, 5 cases were lost to follow-up and 6 patients failed to receive the protocol treatment. Therefore, 81 patients were analyzed in the present study. Unresectable patients were staged according to imaging and gastroscopy when histopathologically diagnosed by biopsy, and the postoperative recurrence patients were staged according to postoperative pathology. Tumors were staged using the 7th edition of the Tumor-Node-Metastasis (TNM) staging system of the International Union Against Cancer/American Joint Committee on Cancer (17). Chemotherapy was given prior to the present study, and regimens included platinum and fluoropyrimidine [cisplatin (D) 80 mg/m<sup>2</sup> on day 1 and fluorouracil (F) 750 mg/m<sup>2</sup> from day 1-4; D 80 mg/m<sup>2</sup> on day 1 and capecitabine (X) 1,000 mg/m<sup>2</sup> from day 1-14], and paclitaxel (P) and fluoropyrimidine [P, 150 mg/m<sup>2</sup> on day 1 and F 750 mg/m<sup>2</sup> from day 1-5; P 150 mg/m<sup>2</sup> on day 1 and X 1,000 mg/m<sup>2</sup> from day 1-14]. The regimens were repeated

every 21 days. Chemotherapy was stopped in case of disease progression, patient refusal or grade 3-4 toxicity according to National Cancer Institute Common Terminology Criteria for Adverse Events version 4.03 (18).

**SNP selection.** The SNP loci of the target genes were selected from the public SNP database of the 1,000 Genome Project in the National Center for Biotechnology Information (NCBI) using minor allele frequency (MAF)  $>0.1$  in the Chinese Han population and the Hardy-Weinberg equilibrium with a P-value of  $>0.1$ , then tagSNPs with a cut-off value of  $R^2 > 0.8$ , and covering the gene and flanking 3 kb either side of the gene regions were chosen by the Genome Variation Server (<https://gvs.gs.washington.edu>). In total, 66 tagSNPs (27 from *KDR* gene, 32 from *NRP-1* gene and 7 from *PDGF $\beta$* ) were selected, however due to financial constraints, 10 SNPs (rs7692791, rs6838752, rs2034965, rs1531290, rs13109660 from *KDR*, rs2070296, rs2804495, rs2065364 from *NRP-1*, and rs4821877, and rs9622978 from *PDGF $\beta$* ) were randomly selected from the tagSNPs. In addition, five disease-associated SNPs (rs1870377 and rs2305948 from *KDR*, rs6554162 and rs1800812 from *PDGFR $\alpha$* , and rs2302273 from *PDGFR $\beta$* ) were selected according to their use in previous literature (19-24). Finally, the 15 SNPs (Table SI) of *KDR* rs7692791, rs2305948, rs6838752, rs2034965, rs1531290, rs13109660 and rs1870377, of *NRP-1* rs2070296, rs2804495 and rs2065364, of *PDGF $\beta$*  rs4821877 and rs9622978, of *PDGFR $\alpha$*  rs6554162 and rs1800812, and of *PDGFR $\beta$*  rs2302273, were obtained from the SNP database of the NCBI (<http://www.ncbi.nlm.nih.gov/SNP>).

**SNP genotyping.** The tissues from patients with AGC were obtained via biopsy or surgery, fixed with 10% neutral buffer formalin for 24 h at room temperature, immersed in 60°C paraffin, embedded in a paraffin block and stored at 4°C. Genomic DNA was extracted from paraffin-embedded tissues of patients with AGC using the QIAamp DNA FFPE Tissue kit (Qiagen GmbH) according to the manufacturer's instructions. The polymerase chain reaction (PCR) primers for SNPs were designed using Sequenom Assay Design 3.1 software (Sequenom) and are listed in Table SII. A thermocycler (PTC-100PCR; MJ Research) and KAPA Taq HotStart DNA polymerase (Kapa Biosystems; Roche Diagnostics) were used for PCR amplification, the thermal cycling program employed was as follows: 94°C for 5 min, followed by 35 cycles of 30 sec at 94°C, then 30 sec of annealing at 60°C, 30 sec of extension at 72°C, and a final elongation step at 72°C for 10 min. The PCR products were sequenced using a 3730XL DNA Analyzer (Applied Biosystems; Thermo Fisher Scientific, Inc.).

**Statistical analysis.** In the present study, the genetic model was divided into 3 types, namely, general, dominant and recessive models, as follows: Dominant model, MW+MM vs. WW; recessive model, WW+WM vs. MM; and general model, MM vs. WM vs. WW, where W indicates the wild-type allele and M the mutant allele). Before analysis, the Hardy-Weinberg equation for the equilibrium of allele distributions was tested by the  $\chi^2$  test (Table I) and the SNPs with a P-value of  $<0.05$  were excluded. The progression-free survival (PFS) and overall survival (OS) probabilities were estimated using

Table I. Hardy-Weinberg equilibrium test results of selected SNPs.

Gene	SNP	$\chi^2$	P-value
<i>KDR</i>	rs7692791	0.11	0.739
	rs2305948	0.02	0.902
	rs6838752	0.65	0.418
	rs2034965	1.66	0.197
	rs13109660	0.03	0.860
	rs1870377	0.58	0.455
	rs1531290	3.37	0.067
<i>NRP-1</i>	rs2070296	3.46	0.062
	rs2804495	0.08	0.775
	rs2065364	1.00	0.317
<i>PDGF<math>\beta</math></i>	rs9622978	11.54	0.007 <sup>a</sup>
	rs4821877	0.00	0.998
<i>PDGFR<math>\alpha</math></i>	rs6554162	0.00	0.951
	rs1800812	30.3	<0.001 <sup>a</sup>
<i>PDGFR<math>\beta</math></i>	rs2302273	7.04	0.007 <sup>a</sup>

<sup>a</sup>P<0.05. *KDR*, kinase insert domain receptor; *NRP-1*, neuropilin-1; *PDGF*, platelet-derived growth factor; *PDGFR*, PDGF receptor; SNP, single nucleotide polymorphism.

the Kaplan-Meier method. The association between SNPs and PFS and OS were analyzed by log-rank tests and Cox regression analyses. Hazard ratios (HR) and 95% confidence intervals (CIs) were estimated for the univariate and multivariate analyses using Cox regression analyses. Bonferroni's correction was applied for multiple comparisons (with the significance level set at P<0.025). Statistical analyses were conducted using SPSS v.21.0 (IBM Corp.). All tests were two-sided, and P<0.05 was considered to indicate a statistically significant difference.

## Results

**Patient clinical characteristics.** A total of 81 patients were recruited in the present study, including 56 men (69.1%) and 25 women (30.9%). The age of the patients ranged from 30 to 83 years, and the mean age was 60.7±10.1 years. By October 2018, 79 patients were deceased, 2 had been lost during follow-up, and the median PFS and OS times were 5.5 and 11.0 months, respectively. The association between clinical pathological features and survival time are listed in Table II. The results demonstrated that TNM stage analyzed by Kaplan-Meier analysis was significantly associated with longer OS time (log-rank, P=0.047), and the platinum-based chemotherapy regimen was significantly associated with longer PFS time (log-rank, P=0.025). Associations between survival time and other clinical characteristics were not identified.

**Associations between genotype and survival time.** Associations between genotype and prognosis were estimated by Kaplan-Meier analysis, statistical significance was

determined by the log-rank test and the genotype information are listed in Table SIII. The associations between the three types of genetic models (general, dominant and recessive) and survival time were analyzed (Table III). The results demonstrated that of all the selected SNPs, five SNPs (*KDR* rs7692791, *KDR* rs1870377, *KDR* rs2034965, *NRP-1* rs2065364 and *NRP-1* rs2804495) were significantly associated with PFS or OS; however, SNPs from *PDGF* and *PDGFR* genes were not associated with clinical outcomes.

Following univariate analysis (Tables IV and V), the dominant model of *KDR* rs1870377 indicated that AA+AT carriers were associated with shorter PFS and OS times compared with TT carriers [PFS: HR, 2.618; 95% CI, 1.235-5.550; P=0.012; OS: HR, 2.041; 95% CI, 1.042-3.999; P=0.038] (Fig. 1). Furthermore, in a recessive model of *NRP-1* rs2065364, AA genotype carriers exhibited more favorable PFS and OS times compared with the GG+AG genotypes (PFS: HR, 2.896; 95% CI, 1.159-7.237; P=0.023; OS: HR, 2.367; 95% CI, 1.019-5.496; P=0.045) (Fig. 2). However, *KDR* rs1870377 variant AA and AT genotype were significantly associated with poor PFS times compared with wild-type TT (AA vs. TT: HR, 3.221; 95% CI, 1.356-7.651; P=0.008; AT vs. TT: HR, 2.545, 95% CI, 1.159-5.589; P=0.020; Table IV) (Fig. S1). Furthermore, the *KDR* rs7692791 CT genotype was associated with lower PFS times compared with the wild-type TT genotype (HR, 1.829, 95% CI, 1.091-3.066, P=0.022; Table IV) (Fig. 3). In addition, in the dominant model of *KDR* rs2034965, AA+GA genotypes were significantly associated with reduced OS times (HR, 1.687; 95% CI, 1.039-2.738; P=0.034; Table IV) (Fig. 3). Statistical significance between SNPs and survival time in other polymorphisms was not found.

For multivariate analysis (Tables IV and V), adjustments were performed for different variables in PFS and OS. Variables that were considered clinically relevant, such as age and TNM stage, or that presented an association with survival time following univariate analysis as listed in Table II were entered into a multivariate Cox proportional-hazards regression model. *KDR* rs7692791 remained significantly associated with PFS, and the TT genotype was associated with better prognosis compared with the CT genotype (HR, 1.969; 95% CI, 1.150-3.369; P=0.013). Furthermore, the association between *NRP-1* rs2065364 AG+GG genotypes and shorter PFS remained significant following adjustment (HR, 3.905; 95% CI, 1.485-10.268; P=0.006). Furthermore, *KDR* rs2034965 AA+GA genotypes remained significantly associated with worse OS following adjustment (HR, 1.978; 95% CI, 1.193-3.280, P=0.008), and the *KDR* rs1870377 AA+AT genotypes were also associated with shorter OS compared with the wild-type TT genotype following adjustment (HR, 2.264; 95% CI, 1.130-4.536; P=0.021).

**Effect of risk allele combinations on PFS and OS.** To study the combined effects of polymorphisms on survival time, risk alleles were selected according to the aforementioned results. The *NRP-1* rs2065364G allele and the *KDR* rs1870377 A allele were found to be unfavorable for PFS and OS. Subsequently, the *NRP-1* rs2065364/*KDR* rs1870377 combination was tested for its association with survival time and numbers of 'risk alleles' (Tables VI and VII). The results suggested that patients carrying >2 risk alleles were more likely to have shorter PFS

Table II. Association between characteristics and prognosis of patients with advanced gastric cancer.

Variables	n	mPFS (95% CI)	mOS (95% CI)	Log-rank P-value	
				PFS	OS
Sex				0.433	0.703
Male	56	5.0 (3.3-6.7)	11 (9.6-12.4)		
Female	25	6.0 (3.6-8.4)	12 (9.4-14.5)		
Age, years				0.773	0.898
>60	33	5.5 (2.8-8.2)	10.2 (6.0-14.4)		
≥60	48	6.0 (4.8-7.2)	11.0 (10.0-12.0)		
N stage				0.590	0.081
N1+N2	49	5.0 (3.6-6.4)	11.6 (10.4-12.8)		
N3	32	5.0 (2.8-7.2)	10.2 (8.6-11.8)		
TNM stage				0.080	0.047 <sup>a</sup>
I, II and III	26	6.8 (5.9-7.7)	12.0 (8.5-15.5)		
IV	55	4.5 (2.8-6.2)	10.5 (8.5-12.5)		
Tumor size, cm				0.803	0.916
>5	31	5.0 (2.9-7.1)	11.0 (9.0-13.0)		
≥5	50	6.0 (4.4-7.6)	11.0 (9.5-12.5)		
Differentiation				0.415	0.079
Well to moderate	27	6.0 (4.5-7.5)	14.8 (8.0-21.6)		
Poor	54	4.5 (3.5-5.5)	10.2 (8.2-12.2)		
Platinum chemotherapy regimen				0.025 <sup>a</sup>	0.359
Platinum included	38	6 (4.6-7.4)	11.6 (9.3-13.9)		
Non-platinum included	43	4.5 (3.3-5.7)	10.5 (9.0-12.0)		
Paclitaxel chemotherapy regimen				0.393	0.484
Paclitaxel included	39	4.4 (2.9-5.9)	11.0 (7.1-14.9)		
Non-paclitaxel included	42	6 (4.6-7.4)	11.0 (10.2-11.8)		

<sup>a</sup>P<0.05. CI, confidence interval; OS, overall survival (months); mOS, median overall survival; PFS, progression-free survival (months); mPFS, median progression-free survival; TNM, Tumor-Node-Metastasis.

Table III. Effect of SNPs in selected genes on the prognosis in patients with advanced gastric cancer.

Gene	SNP	Allelic change	Log-rank P-value for PFS			Log-rank P-value for OS		
			General	Dominant	Recessive	General	Dominant	Recessive
<i>KDR</i>	rs7692791	T/C	0.032 <sup>a</sup>	0.009 <sup>a</sup>	0.281	0.227	0.093	0.364
	rs2305948	C/T	0.619	0.329	0.871	0.277	0.109	0.748
	rs6838752	T/C	0.097	0.137	0.053	0.203	0.254	0.095
	rs2034965	G/A	0.155	0.065	0.240	0.065	0.031	0.883
	rs13109660	G/A	0.795	0.522	0.687	0.365	0.376	0.481
	rs1870377	T/A	0.030 <sup>a</sup>	0.008 <sup>a</sup>	0.256	0.091	0.032 <sup>a</sup>	0.250
	rs1531290	A/G	0.236	0.128	0.313	0.451	0.845	0.207
<i>NRP-1</i>	rs2070296	G/A	0.498	0.417	0.486	0.993	0.964	0.909
	rs2804495	G/T	0.064	0.150	0.028 <sup>a</sup>	0.085	0.308	0.029
	rs2065364	G/A	0.052	0.300	0.015 <sup>a</sup>	0.113	0.587	0.037 <sup>a</sup>
<i>PDGFβ</i>	rs4821877	C/T	0.712	0.490	0.862	0.949	0.933	0.747
<i>PDGFRα</i>	rs6554162	G/A	0.513	0.322	0.751	0.501	0.413	0.561

<sup>a</sup>P<0.05. *KDR*, kinase insert domain receptor; *NRP-1*, neuropilin-1; *PDGF*, platelet-derived growth factor; *PDGFR*, *PDGF* receptor; OS, overall survival; PFS, progression-free survival; SNP, single nucleotide polymorphism.

Table IV. Associations of SNPs in selected genes and PFS in patients with advanced gastric cancer.

SNP	Outcome	mPFS, months	Model	Log-rank P-value	Univariate analysis		Multivariate analysis	
					HR (95% CI)	P-value	HR (95% CI) <sup>a</sup>	P-value <sup>a</sup>
<i>KDR</i> rs7692791	PFS		General	0.032		0.020		0.012
		4.2	CC	0.099 <sup>b</sup>	1.926 (0.859-4.319)	0.112	2.053 (0.855-4.929)	0.107
		5.0	CT	0.018 <sup>b</sup>	1.829 (1.091-3.066)	0.022	1.969 (1.150-3.369)	0.013
		6.0	TT		Reference		Reference	
			Dominant	0.009		0.010		0.006
<i>KDR</i> rs1870377	PFS	6.0	TT		Reference		Reference	
		5.0	CC+CT		1.892 (1.156-3.098)	0.011	1.982 (1.196-3.284)	0.008
			General	0.030		0.017		0.127
		4.0	AA	0.005 <sup>b</sup>	3.221 (1.356-7.651)	0.008	2.892 (0.987-8.474)	0.053
		5.5	AT	0.015 <sup>b</sup>	2.545 (1.159-5.589)	0.020	1.778 (0.724-4.366)	0.209
<i>NRP-1</i> rs2065364	PFS	10.0	TT		Reference		Reference	
			Dominant	0.008		0.009		0.051
		10.0	TT		Reference		Reference	
		4.5	AA+AT		2.618 (1.235-5.550)	0.012	1.970 (0.861-4.503)	0.108
			Recessive	0.015		0.017		0.004
		8.0	AA		Reference		Reference	
		4.5	AG+GG		2.896 (1.159-7.237)	0.023	3.905 (1.485-10.268)	0.006

<sup>a</sup>Adjusted for age, sex, N stage, TNM stage, platinum included or not and differentiation. <sup>b</sup>Bonferroni-adjusted P-value=0.05/2, so P<0.025 was considered statistically significant. CI, confidence interval; HR, hazard ratio; *KDR*, kinase insert domain receptor; *NRP-1*, neuropilin-1; PFS, progression-free survival; mPFS, median progression-free survival; SNP, single nucleotide polymorphism.

Table V. Associations of SNPs in selected genes and OS in patients with advanced gastric cancer.

SNP	Outcome	mOS, months	Model	Log-rank P-value	Univariate analysis		Multivariate analysis	
					HR (95% CI)	P-value	HR (95% CI) <sup>a</sup>	P-value
<i>KDR</i> rs2034965	OS	11.6	Dominant	0.031	Reference	0.032	Reference	0.029
		10.3	GG		1.687 (1.039-2.738)	0.034	1.978 (1.193-3.280)	0.008
			AA+GA			0.039		0.105
<i>NRP1</i> rs2065364	OS	17.8	Recessive	0.037	Reference	0.045	Reference	0.112
		11.0	AA		2.367 (1.019-5.496)	0.034	2.048 (0.847-4.952)	0.035
			AG+GG					
<i>KDR</i> rs1870377	OS	16.0	Dominant	0.032	Reference	0.038	Reference	0.021
		10.8	TT		2.041 (1.042-3.999)	0.031	2.264 (1.130-4.536)	0.084
			AA+AT					
<i>NRP1</i> rs2804495	OS	8.8	Recessive	0.029	Reference	0.033	Reference	0.095
		12.0	TT		1.710 (1.046-2.796)		1.570 (0.924-2.667)	
			GT+GG					

<sup>a</sup> Adjusted for age, gender, N stage, TNM stage and differentiation. CI, confidence interval; HR, hazard ratio; *KDR*, kinase insert domain receptor; *NRP-1*, neuropilin-1; OS, overall survival; mOS, median overall survival; SNP, single nucleotide polymorphism.

and OS times compared with carriers with 1-2 risk alleles (PFS: HR, 0.427; 95% CI, 0.260-0.701; P=0.008; OS: HR, 0.523; 95% CI, 0.323-0.845; P=0.008; Tables VI and VII) (Fig. 4). Following adjustment, this association was also significant (PFS: HR, 0.427; 95% CI, 0.257-0.709; P=0.001; OS: HR, 0.511; 95% CI, 0.314-0.833; P=0.007).

## Discussion

The results from the present study demonstrated that polymorphisms of *NRP-1* and *KDR* genes were associated with clinical outcome in patients with AGC. Following univariate analysis, *KDR* rs1870377 AA+AT genotypes were found to be associated with shorter PFS and OS times compared with the wild-type TT genotype, and the *KDR* rs1870377 variant AA and AT genotypes were significantly associated with poor PFS time compared with wild-type TT genotype. Furthermore, the *NRP-1* rs2065364 homozygous mutant AA genotype was significantly associated with higher PFS and OS times compared with the GG+AG genotypes. The genotypes of *KDR* rs7692971 and *KDR* rs2034965 were also significantly associated with higher PFS and OS times, respectively. Following adjustment, the *KDR* rs7692971 TT genotype was associated with increased PFS time compared with the CT genotype, and the *NRP-1* rs2065364 AG+GG genotypes were associated with shorter PFS times compared with the AA genotype. The *KDR* rs2034965 AA+GA genotypes were associated with worse OS times compared with the GG genotype. The *KDR* rs1870377 AA+AT genotypes were associated with shorter OS times compared with the TT genotype. Additionally, increasing number of risk alleles with the *NRP-1* rs2065364/*KDR* rs1870377 combination was significantly associated with shorter OS and PFS times. These results demonstrated that *NRP-1* rs2065364, *KDR* rs7692971, *KDR* rs2034965 and *KDR* rs1870377 may be considered as independent indicators of prognosis in patients with AGC.

*NRP-1* was originally found to be crucial for neuronal axon guidance and embryonic angiogenesis, and was identified as a novel receptor involved in angiogenesis (6-8). Previous studies reported that the *NRP-1* gene is associated with tumorigenesis and progression. One study reported that *NRP-1* overexpression is associated with the promotion of gastric cancer migration, invasion and growth (25). Lin *et al* (26) demonstrated that *NRP-1* is a novel TEA domain transcription factor target that serves a crucial role in hepatocellular carcinoma tumorigenesis. A previous demonstrated that *NRP-1* is abnormally highly expressed in non-small cell lung tumor tissue, and is associated with patient prognosis (27). Another study reported that *NRP-1* affects the chemosensitivity of cancer cells (28), Wey *et al* (28) demonstrated that *NRP-1* overexpression in pancreatic cancer cell lines is associated with increased chemoresistance to gemcitabine *in vitro*. Yue *et al* (29) reported that *NRP-1* overexpression increases osteosarcoma cell survival following exposure to doxorubicin. To the best of our knowledge, no study has demonstrated the association between *NRP-1* SNPs and cancer. The present study confirmed that the *NRP-1* rs2065364 AA genotype was associated with increased PFS time compared with the AG+GG genotypes. Further molecular investigation is required to reveal the underlying mechanisms involved.






Figure 1. Effect of kinase insert domain receptor rs1870377 on survival time in patients carrying the AA+AT and TT genotypes. (A) PFS curve. (B) OS curve. P-values were obtained by log-rank tests. OS, overall survival; PFS, progression-free survival.




Figure 2. Effect of neuropilin-1 rs2065364 on survival time in patients carrying GG+AG and AA genotypes. (A) PFS curve. (B) OS curve. P-values were obtained by log-rank tests. OS, overall survival; PFS, progression-free survival.




Figure 3. Effect of kinase insert domain receptor rs7692791 and rs2034965 on survival time. (A) PFS curve of rs7692791 in patients with TT and CT genotypes. (B) OS curve of rs2034965 in patients with AA+AG and GG genotypes. P-values were obtained by log-rank tests. OS, overall survival; PFS, progression-free survival.

Table VI. Association between number of risk alleles and overall survival in patients with advanced gastric cancer.

		Univariate analysis		Multivariate analysis <sup>a</sup>	
Alleles combination	n	HR (95% CI)	P-value	HR (95% CI)	P-value
rs2065364/rs1870377					
1-2 risk alleles	39	0.523 (0.323-0.845)	0.008	0.511 (0.314-0.833)	0.007
3-4 risk alleles	42	Reference		Reference	

<sup>a</sup>Adjusted for age, sex, N stage, TNM stage, platinum included or not and differentiation. CI, confidence interval; HR, hazard ratio; OS, overall survival.

Table VII. Association between number of risk alleles and progression-free survival in patients with advanced gastric cancer.

		Univariate analysis		Multivariate analysis <sup>a</sup>	
Alleles combination	n	HR (95% CI)	P-value	HR (95% CI)	P-value
rs2065364/rs1870377					
1-2 risk alleles	39	0.427 (0.260-0.701)	0.008	0.427 (0.257-0.709)	0.001
3-4 risk alleles	42	Reference		Reference	

<sup>a</sup>Adjusted for age, sex, N stage, TNM stage, platinum included or not and differentiation. CI, confidence interval; HR, hazard ratio; PFS, progression-free survival.

*KDR* (*VEGFR-2*) is a tyrosine kinase receptor that can regulate signal transduction by binding to *VEGF* via its extracellular domain (9). *VEGF/VEGFR2* is an important signaling pathway that can promote proliferation, survival and migration of vascular endothelial cells and increase vascular permeability (9,10). The cellular processes mediated by the *VEGF-VEGFR2* signaling cascade can lead to angiogenesis and therefore regulate tumor growth and invasion, and therapeutic resistance (10,11). Previous studies reported that *KDR* gene polymorphisms are associated with clinical outcomes in various types of cancer, including colorectal cancer, glioma, hepatocellular carcinoma and gastric cancer. Torben *et al* (30) reported that *VEGFR2* 1192C>T and -604T>C polymorphisms were associated with increased microvessel density in colorectal cancer. A previous study of glioma in the Chinese population demonstrated that three SNPs of *VEGFR2* (rs7667298, rs2305948 and rs1870377) are correlated with an increased risk of a glioma when homozygous (31). Another study described that the *VEGFR-2* rs2305948 T polymorphism frequency is higher in patients with gastroenteropancreatic neuroendocrine neoplasms compared with that in the healthy population (19). In the present study, among the genetic variations of the *VEGFR2* gene, the *KDR* rs1870377 and *KDR* rs7692791 TT genotypes were found to be associated with a better prognosis, and the *KDR* rs2034965 GG genotype was associated with increased OS time. Zhu *et al* (20) demonstrated that the *VEGFR2* rs1870377 TT genotype confers a favorable prognosis in gastric cancer. Furthermore, Wang *et al* (21) investigated the correlation between polymorphisms of four genes from the epidermal growth factor receptor (*EGFR*) pathway and the clinical outcome of 363 patients with hepatocellular carcinoma, and reported that *EGFR* rs2034965 with the AA genotype is negatively correlated with disease-free survival.

These results were consistent with the results from the present study; however, inconsistent results were reported in other types of cancer, and Kim *et al* (22) reported that the *VEGFR2* rs1870377 TT genotype is associated with shorter OS time in patients with diffuse large B cell lymphoma. Furthermore, it was reported that rs7692791 C allele is significantly correlated with increased OS and DFS in hepatocellular carcinoma (21). These discordances may be partly attributed to the different types of cancer, the different clinical characteristics of the patients and the study sizes. The rs1870377 mutation is located in the coding region of *KDR* and is a missense mutation. The functional role of this gene polymorphism remains unclear.

Proteins from the *PDGF* family are crucial to stimulate the proliferation, survival and migration of mesenchymal cells (32). This family consists of 5 different isoforms, named disulphide-bonded homodimers of A-, B-, C- and D-polypeptide chains and the heterodimer *PDGF-AB*. *PDGFR* is classified as a receptor tyrosine kinase, and the 5 *PDGF* isoforms can activate cellular responses via *PDGFRα* and *PDGFRβ* (32,33). Overactivation of the *PDGF-PDGFR* signaling pathway has been reported to be associated with tumorigenesis (34). *PDGFR* gene mutations have been found in malignancies. Point mutations in *PDGFRα* were found in ~5% of gastrointestinal stroma tumors, which led to amino acid residue changes, therefore activating *PDGFR* kinase activity (35). In addition, a study reported that rs1800812 T allele and rs6554162 G allele in *PDGFRα* were related to decreased frequency in patients with papillary thyroid cancer compared with that in the healthy population (23). A previous study demonstrated that two SNPs in *PDGFRβ* (rs5757573 T>C and rs6001516 C>T) were associated with an increased risk of pancreatic cancer (36). Furthermore, Volz *et al* (24) found that the SNP (rs2302273 C>T) in *PDGFRβ* gene was associated






Figure 4. Association between number of risk alleles (rs1870377/rs2065364 combination) and survival time. (A) PFS curve. (B) OS curve. P-values were obtained by log-rank tests. OS, overall survival; PFS, progression-free survival.

with a significantly longer PFS time in patients with metastatic colorectal cancer. However, in the present study, no relevance was found between SNPs and prognosis.

The present study had some limitations. Firstly, only 81 patients with AGC were eligible for statistical analysis. Since the sample size was relatively small, the results from this study should be considered as preliminary data and for generation of a hypothesis for subsequent investigation. Secondly, since the patients studied had AGC, it is not known whether the results could be applicable to patients with other types of gastric cancer. Further investigation should therefore be conducted to validate the results.

In conclusion, the results from the present study demonstrated that *KDR* rs7692791 and *NRP-1* rs2065364 were positively associated with PFS. Furthermore, *KDR* rs2034965 and *KDR* rs1870377 significantly negatively correlated with OS time following multivariate analysis in patients with AGC. In addition, the numbers of 'risk alleles' of *NRP-1* rs2065364/*KDR* rs1870377 combination were significantly associated with survival time. These results suggested that genetic variants in *NRP-1* and *KDR* genes may affect the biological features and prognosis of patients with AGC. Due to limited funding, the underlying mechanisms were not explored, and further investigation is required to verify these results.

## Acknowledgements

The authors would like to thank Dr Dan Lv (Department of Medical Oncology, The Second Hospital Affiliated to Dalian Medical University) and Professor Na Gao (Department of Obstetrics and Gynecology, First Affiliated Hospital of Dalian Medical University) for their technical assistance.

## Funding

No funding was received.

## Availability of data and materials

All data and materials generated and/or used during the study are available from the corresponding author upon reasonable request.

## Authors' contributions

TW and YS designed the study. YS collected the patients' clinical data. YJZ and YS performed the experiments. YJZ and YS analyzed the data and wrote the manuscript. TW contributed to the revision of the manuscript. All the authors read and approved the final manuscript.

## Ethics approval and consent to participate

The present study was approved by the Clinical Research Ethics Committee of The Second Hospital Affiliated to Dalian Medical University and was conducted in accordance with The Declaration of Helsinki. Participants were fully informed of the procedures and provided written informed consent.

## Patient consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68: 394-424, 2018.
- Wang Y, Li Z, Shan F, Miao R, Xue K, Li Z, Gao C, Chen N, Gao X, Li S and Ji J: Current status of diagnosis and treatment of early gastric cancer in China-Data from China gastrointestinal cancer surgery union. *Zhonghua Wei Chang Wai Ke Za Zhi* 21: 168-174, 2018 (In Chinese).
- Wang FH, Shen L, Li J, Zhou ZW, Liang H, Zhang XT, Tang L, Xin Y, Jin J, Zhang YJ, *et al*: The Chinese Society of Clinical Oncology (CSCO): Clinical Guidelines for the Diagnosis and Treatment of Gastric Cancer. *Cancer Commun (Lond)*. 39: 10, 2019.
- Van Cutsem E, Sagaert X, Topal B, Haustermans K and Prenen H: Gastric cancer. *Lancet* 388: 2654-2664, 2016.
- Chen K, Yang D, Li X, Sun B, Song F, Cao W, Brat DJ, Gao Z, Li H, Liang H, *et al*: Mutational landscape of gastric adenocarcinoma in Chinese: Implications for prognosis and therapy. *Proc Natl Acad Sci USA* 112: 1107-1112, 2015.

6. Fujisawa H, Takagi S and Hirata T: Growth-associated expression of a membrane protein, neuropilin, in *Xenopus* optic nerve fibers. *Dev Neurosci* 17: 343-349, 1995.
7. Muhl L, Folestad EB, Gladh H, Wang Y, Moessinger C, Jakobsson L and Eriksson U: Neuropilin 1 binds PDGF-D and is a co-receptor in PDGF-D-PDGFR $\beta$  signaling. *J Cell Sci* 130: 1365-1378, 2017.
8. Raimondi C: Neuropilin-1 enforces extracellular matrix signaling via ABL1 to promote angiogenesis. *Biochem Soc Trans* 42: 1429-1434, 2014.
9. Ferrara N, Gerber HP and LeCouter J: The biology of VEGF and its receptors. *Nat Med* 9: 669-676, 2003.
10. Karaman S, Leppänen VM and Alitalo K: Vascular endothelial growth factor signaling in development and disease. *Development* 145: pii: dev151019, 2018.
11. Shibuya M and Claesson-Welsh L: Signal transduction by VEGF receptors in regulation of angiogenesis and lymphangiogenesis. *Exp Cell Res* 312: 549-560, 2006.
12. Farooqi AA and Siddik ZH: Platelet-derived growth factor (PDGF) signalling in cancer: Rapidly emerging signalling landscape. *Cell Biochem Funct* 33: 257-265, 2015.
13. Heldin CH, Lennartsson J and Westermark B: Involvement of platelet-derived growth factor ligands and receptors in tumorigenesis. *J Intern Med* 283: 16-44, 2018.
14. Sa-Nguanraksa D, Kooptiwut S, Chuangsuwanich T, Pongpruttipan T, Malasit P and O-Charoenrat P: Vascular endothelial growth factor polymorphisms affect gene expression and tumor aggressiveness in patients with breast cancer. *Mol Med Rep* 9: 1044-1048, 2014.
15. Glubb DM, Cerri E, Giese A, Zhang W, Mirza O, Thompson EE, Chen P, Das S, Jassem J, Rzyman W, *et al*: Novel functional germline variants in the VEGF receptor 2 gene and their effect on gene expression and microvessel density in lung cancer. *Clin Cancer Res* 17: 5257-5267, 2011.
16. Oken M, Creech R, Tormey D, Horton J, Davis TE, McFadden ET and Carbone PP: Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol* 5: 649-655, 1982.
17. Edge SB, Byrd DR, Compton CC, Fritz AG, Greene FL and Trotti A III: *AJCC Cancer Staging Manual*. 7th edition. Springer, New York, NY, 2010.
18. National Cancer Institute (NCI): Common Terminology Criteria for Adverse Events CTCAE. Version 4.03, 2010.
19. Berardi R, Torniai M, Partelli S, Rubini C, Pagliaretta S, Savini A, Polenta V, Santoni M, Giampieri R, Onorati S, *et al*: Impact of vascular endothelial growth factor (VEGF) and vascular endothelial growth factor receptor (VEGFR) single nucleotide polymorphisms on outcome in gastroenteropancreatic neuroendocrine neoplasms. *PLoS One* 13: e0197035, 2018.
20. Zhu X, Wang Y, Xue W, Wang R, Wang L, Zhu ML and Zheng L: The VEGFR-2 protein and the VEGFR-2 rs1870377 A>T genetic polymorphism are prognostic factors for gastric cancer. *Cancer Biol Ther* 20: 497-504, 2019.
21. Wang W, Ma XP, Shi Z, Zhang P, Ding DL, Huang HX, Saiyin HG, Chen TY, Lu PX, Wang NJ, *et al*: Epidermal growth factor receptor pathway polymorphisms and the prognosis of hepatocellular carcinoma. *Am J Cancer Res* 5: 396-410, 2015.
22. Kim MK, Suh C, Chi HS, Cho HS, Bae YK, Lee KH, Lee GW, Kim IS, Eom HS, Kong SY, *et al*: VEGFA and VEGFR2 genetic polymorphisms and survival in patients with diffuse large B cell lymphoma. *Cancer Sci* 103: 497-503, 2012.
23. Kim MJ, Kim SK, Park HJ, Chung DH, Park HK, Lee JS, Kwon KH and Chung JH: PDGFRA promoter polymorphisms are associated with the risk of papillary thyroid cancer. *Mol Med Rep* 5: 1267-1270, 2012.
24. Volz NB, Stintzing S, Zhang W, Yang D, Ning Y, Wakatsuki T, El-Khoueiry RE, Li JE, Kardosh A, Loupakis F, *et al*: Genes involved in pericyte-driven tumor maturation predict treatment benefit of first-line FOLFIRI plus bevacizumab in patients with metastatic colorectal cancer. *Pharmacogenomics J* 15: 69-76, 2015.
25. Peng Y, Liu YM, Li LC, Wang LL and Wu XL: MicroRNA-338 inhibits growth, invasion and metastasis of gastric cancer by targeting NRP1 expression. *PLoS One* 9: e94422, 2014.
26. Lin J, Zhang Y, Wu J, Li L, Chen N, Ni P, Song L and Liu X: Neuropilin 1 (NRP1) is a novel tumor marker in hepatocellular carcinoma. *Clin Chim Acta* 485: 158-165, 2018.
27. Ding M, Liu L, Hu C, Liu Y, Qiao Y and Jiang X: Expression of VEGFR2 and NRP-1 in non-small cell lung cancer and their clinical significance. *Chin J Cancer Res* 26: 669-677, 2014.
28. Wey JS, Gray MJ, Fan F, Belcheva A, McCarty MF, Stoeltzing O, Somcio R, Liu W, Evans DB, Klagsbrun M, *et al*: Overexpression of neuropilin-1 promotes constitutive MAPK signalling and chemoresistance in pancreatic cancer cells. *Br J Cancer* 93: 233-241, 2005.
29. Yue B, Ma JF, Yao G, Yang MD, Cheng H and Liu GY: Knockdown of neuropilin-1 suppresses invasion, angiogenesis and increases the chemosensitivity to doxorubicin in osteosarcoma cells-an in vitro study. *Eur Rev Med Pharmacol Sci* 18: 1735-1741, 2014.
30. Hansen TF, Sørensen FB, Spindler KL, Olsen DA, Andersen RF, Lindebjerg J, Brandslund I and Jakobsen A: Microvessel density and the association with single nucleotide polymorphisms of the vascular endothelial growth factor receptor 2 in patients with colorectal cancer. *Virchows Arch* 456: 251-260, 2010.
31. Zhang J, Yang J, Chen Y, Mao Q, Li S, Xiong W, Lin Y, Chen J and Ge J: Genetic variants of VEGF (rs201963 and rs3025039) and KDR (rs7667298, rs2305948, and rs1870377) are associated with glioma risk in a Han Chinese population: A case-control study. *Mol Neurobiol* 53: 2610-2618, 2016.
32. Andrae J, Gallini R and Betsholtz C: Role of platelet-derived growth factors in physiology and medicine. *Genes Dev* 22: 1276-1312, 2008.
33. Heldin CH and Westermark B: Mechanism of action and in vivo role of platelet-derived growth factor. *Physiol Rev* 79: 1283-1316, 1999.
34. Pietras K, Sjöblom T, Rubin K, Heldin CH and Ostman A: PDGF receptors as cancer drug targets. *Cancer Cell* 3: 439-443, 2003.
35. Heinrich MC, Corless CL, Duensing A, McGreevey L, Chen CJ, Joseph N, Singer S, Griffith DJ, Haley A, Town A, *et al*: PDGFRA activating mutations, in gastrointestinal stromal tumors. *Science* 299: 708-710, 2003.
36. Duan B, Hu J, Liu H, Wang Y, Li H, Liu S, Xie J, Owzar K, Abbruzzese J, Hurwitz H, *et al*: Genetic variants in the platelet-derived growth factor subunit B gene associated with pancreatic cancer risk. *Int J Cancer* 142: 1322-1331, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.