

Five extracellular matrix-associated genes upregulated in oral tongue squamous cell carcinoma: An integrated bioinformatics analysis

PINGPING ZHONG^{1-4*}, LIYA LIU^{5,6*}, ALING SHEN^{5,6}, ZHONGXIN CHEN⁷, XIAOYAN HU²⁻⁴,
YICHAO CAI²⁻⁴, JIE LIN²⁻⁴, BANGYAN WANG²⁻⁴, JIESEN LI²⁻⁴, YOUQIN CHEN⁸ and JUN PENG^{5,6}

¹Department of Orthodontics, School and Hospital of Stomatology, Fujian Medical University; ²Key Laboratory of Stomatology, Fujian Province University; ³Fujian Biological Materials Engineering and Technology Center of Stomatology; ⁴Institute of Stomatology, Fujian Medical University, Fuzhou, Fujian 350002; ⁵Academy of Integrative Medicine; ⁶Fujian Key Laboratory of Integrative Medicine on Geriatrics, Fujian University of Traditional Chinese Medicine, Fuzhou, Fujian 350122; ⁷West China School of Stomatology Sichuan University, Chengdu, Sichuan 610041, P.R. China; ⁸Department of Pediatrics, Case Western Reserve University School of Medicine, Rainbow Babies and Children's Hospital, Cleveland, OH 44106, USA

Received November 3, 2018; Accepted July 26, 2019

DOI: 10.3892/ol.2019.10982

Abstract. Despite advancements in treatment regimens, the mortality rate of patients with oral tongue squamous cell carcinoma (OTSCC) is high. In addition, the signaling pathways and oncoproteins involved in OTSCC progression remain largely unknown. Therefore, the aim of the present study was to identify specific prognostic marker for patients at a high risk of developing OTSCC. The present study used four original microarray datasets to identify the key candidate genes involved in OTSCC pathogenesis. Expression profiles of 93 OTSCC tissues and 76 normal tissues from GSE9844, GSE13601, GSE31056

and GSE75538 datasets were investigated. Differentially expressed genes (DEGs) were determined, and gene ontology enrichment and gene interactions were analyzed. The four GSE datasets reported five upregulated and six downregulated DEGs. Five upregulated genes (matrix metalloproteinase 1, 3, 10 and 12 and laminin subunit gamma 2) were localized in the extracellular region of cells and were associated with extracellular matrix disassembly. Furthermore, analysis for The Cancer Genome Atlas database revealed that the aforementioned five upregulated genes were also highly expressed in OTSCC and head and neck squamous cell carcinoma tissues. These results demonstrated that the five upregulated genes may be considered as potential prognostic biomarkers of OTSCC and may serve at understanding OTSCC progression. Upregulated DEGs may therefore represent valuable therapeutic targets to prevent or control OTSCC pathogenesis.

Correspondence to: Professor Youqin Chen, Department of Pediatrics, Case Western Reserve University School of Medicine, Rainbow Babies and Children's Hospital, 11100 Euclid Avenue, Cleveland, OH 44106, USA
E-mail: yxc571@case.edu

Professor Dr Jun Peng, Fujian Key Laboratory of Integrative Medicine on Geriatrics, Fujian University of Traditional Chinese Medicine, 1 Huatuo Road, Minhou Shangjie, Fuzhou, Fujian 350122, P.R. China
E-mail: pjunlab@hotmail.com

*Contributed equally

Abbreviations: HNSC, head and neck squamous cell carcinoma; OTSCC, oral tongue squamous cell carcinoma; DEG, differentially expressed gene; TCGA, The Cancer Genome Atlas; GEO, Gene Expression Omnibus; GO, Gene Ontology

Key words: head and neck squamous cell carcinoma, oral tongue squamous cell carcinoma, bioinformatics analysis, differentially expressed gene

Introduction

Head and neck squamous cell carcinoma (HNSC) is a common cancer worldwide and accounts for >600,000 new cases annually (1). Despite significant advancements in treatments, including reconstructive microvascular free tissue transfer, hyperfractionated radiotherapy and concurrent chemoradiation, the survival rate of patients with HNSC has not sufficiently improved over the last 50 years, with overall survival of ~50%, resulting in increased mortality rates worldwide every year (2). HNSC is a highly complex disease emerging from the oral cavity, tongue, pharynx or larynx (3). Each tumor harbors unique mutations, presents variable clinical outcomes and is associated with specific risk factors (4,5). For example, TP53 inactivation, either through somatic mutation or HPV infection, appears nearly universal in this malignancy (6). The present study particularly focused on one type of HNSC, the oral tongue squamous cell carcinoma (OTSCC), because of its poor diagnosis, high incidence

rate, aggressive clinical behavior and poor outcome (7-10). A recent study reported a five-year survival rate of 63% for patients with OTSCC in The Netherlands (11). In 2017, nearly 16,400 new cases of tongue cancer were diagnosed and 2,400 tongue cancer-associated mortality cases were recorded in the United States (12). Several prognostic factors for OTSCC exist, including occult node positivity, tumor depth, lymphovascular invasion and perineural invasion (13). Nevertheless, robust and reliable molecular prognostic biomarkers need to be determined in order to identify patients with advanced stages of OTSCC.

Several thousands of tumor biomarkers have been discovered and are associated with the prognosis of various types of cancer (14). In particular, markers for OTSCC, including microtubule associated scaffold protein 1 and beta-parvin, have attracted much attention due to their crucial role in OTSCC pathogenesis (15). Furthermore, it was demonstrated that downregulation of these markers markedly decreases cancer cell survival (15-19).

Thanks to the rapid development and extensive application of microarrays for gene identification in various types of cancer (20-22), a high number of differentially expressed genes (DEGs) in OTSCC have been identified (13,19). However, the results between studies are inconsistent, which might be due to the variability of tissue samples used. In addition, no reliable biomarkers for OTSCC have been established. Subsequently, the combination of bioinformatics methods and expression profiling techniques may represent a novel approach to resolve these problems. The present study used four microarray datasets [GSE9844 (3), GSE13601 (23), GSE31056 (24), and GSE75538 (25)] from the National Center for Biotechnology Information (NCBI)-Gene Expression Omnibus (GEO) database and an mRNA sequencing (mRNA-seq) dataset from The Cancer Genome Atlas (TCGA). DEGs were filtered using the GEO2R tool according to conventional data processing standards, and gene ontology (GO) enrichment analysis was performed to screen for DEGs using The Database for Annotation, Visualization and Integrated Discovery (DAVID). Expression levels of DEGs in samples from TCGA were assessed using Gene Expression Profiling Interactive Analysis (GEPIA), and corrections were applied for DEGs in samples from TCGA-HNSC using cBioPortal to identify potential oncogenes in OTSCC. The identified DEGs and their associated pathways may be considered as robust and reliable tumor biomarkers for OTSCC and serve as precise therapeutic targets for the prevention of OTSCC progression at early stages.

Materials and methods

GEO datasets. GEO (<https://www.ncbi.nlm.nih.gov/gds>) is a public repository at NCBI for storing high-throughput gene expression data (26). The gene expression profiles of GSE9844, GSE13601, GSE31056, and GSE75538 were selected from the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>). GSE9844 included 26 OTSCC and 12 normal tissues, GSE13601 included 31 OTSCC and 26 normal tissues, GSE31056 included 22 OTSCC and 24 normal tissues, and GSE75538 included 14 OTSCC and 14 normal tissues.

Identification of DEGs. GEO2R (<https://www.ncbi.nlm.nih.gov/geo/geo2r/>) was used to detect DEGs between OTSCC and

Table I. Characteristics of 147 patients with tongue tumor from The Cancer Genome Atlas database.

Characteristics	Patient number	Percentage
Sex		
Female	46	31.29
Male	101	68.71
Age at diagnosis, years		
Mean	58.60±12.71	
<40	10	6.80
40-49	20	13.61
50-59	38	25.85
60-69	53	36.05
70-79	21	14.29
>80	4	2.72
Tumor pathological stage		
T1-T2	70	47.62
T3-T4	72	48.98
Nodal pathological stage		
N0-N1	72	48.98
N2-N4	70	47.62
Smoking history		
Yes	98	66.67
No/Never	47	31.97
Unknown	2	1.36
Alcohol history		
Yes	99	67.35
No/Never	44	29.93
Unknown	4	2.72
Primary lymph node		
Yes	128	87.07
No	9	6.12

normal tissue samples (27). Adjusted P-values <0.01 and \log_2 fold change (FC) >2 were set as cutoffs. Co-expressed DEGs that were downregulated or upregulated in the two sets of gene expression profiles were identified using Venn diagrams (<http://bioinformatics.psb.ugent.be/webtools/Venn/>).

GO enrichment analysis of DEGs. Usually, genes and their products are annotated to identify characteristic biological function and processes of a high-throughput genome or transcriptome (28). DAVID (<https://david.ncifcrf.gov/>) is a web-based bioinformatics resource for gene annotation and visualization with an integrated discovery function. It is therefore useful for determining gene biological attributes (29). P<0.01 and the Benjamini corrected P<0.01 (30) were set as the cutoff. Sequential pathways, molecular and cellular components and biological functions of DEGs could be visualized by using DAVID (<https://david.ncifcrf.gov/>).

Comparison of gene expression in patients with OTSCC. The mRNA-seq data of the five OTSCC genes of interest from HNSC samples were obtained from the TCGA database

Table II. Gene ontology analysis of upregulated genes associated with oral tongue squamous cell carcinoma.

Category	Term	Count	Genes	P-value	Benjamini corrected P-value
GOTERM_BP_DIRECT	GO:0022617~extracellular matrix disassembly	5	MMP10, LAMC2, MMP3, MMP12, MMP1	3.9x10 ⁻¹⁰	7.7x10 ⁻⁹
GOTERM_BP_DIRECT	GO:0030574~collagen catabolic process	4	MMP10, MMP3, MMP12, MMP1	2.1x10 ⁻⁷	2.1x10 ⁻⁶
GOTERM_MF_DIRECT	GO:0004222~metalloendopeptidase activity	4	MMP10, MMP3, MMP12, MMP1	1.2x10 ⁻⁶	9.3x10 ⁻⁶
GOTERM_CC_DIRECT	GO:0005578~proteinaceous extracellular matrix	4	MMP10, MMP3, MMP12, MMP1	1.2x10 ⁻⁵	1.1x10 ⁻⁴
GOTERM_MF_DIRECT	GO:0004252~serine-type endopeptidase activity	4	MMP10, MMP3, MMP12, MMP1	1.3x10 ⁻⁵	5.4x10 ⁻⁵
GOTERM_MF_DIRECT	GO:0004175~endopeptidase activity	3	MMP3, MMP12, MMP1	6.0x10 ⁻⁵	1.6x10 ⁻⁴
GOTERM_CC_DIRECT	GO:0005576~extracellular region	5	MMP10, LAMC2, MMP3, MMP12, MMP1	6.1x10 ⁻⁵	2.7x10 ⁻⁴
GOTERM_BP_DIRECT	GO:0006508~proteolysis	4	MMP10, MMP3, MMP12, MMP1	1.0x10 ⁻⁴	6.8x10 ⁻⁴
GOTERM_MF_DIRECT	GO:0005509~calcium ion binding	4	MMP10, MMP3, MMP12, MMP1	3.0x10 ⁻⁴	5.9x10 ⁻⁴
GOTERM_MF_DIRECT	GO:0008270~zinc ion binding	4	MMP10, MMP3, MMP12, MMP1	1.3x10 ⁻³	2.0x10 ⁻³

(<https://cancergenome.nih.gov/>) (31). In addition, clinical data of OTSCC were downloaded using TCGA Assembler (Table I). mRNA-seq data of 147 OTSCC tissues and 15 adjacent normal tongue tissues were obtained on an Illumina HiSeq RNASeq platform (32). Since the TCGA dataset was developed as a community resource project, no additional approval was required for this study from Fujian Medical University. The present study complied with the TCGA publication guidelines and data access policies (33).

Analysis of gene expression patterns in all tumors from TCGA through GEPIA. The expression patterns of the five genes of interest in various types of cancers and normal tissues were determined using GEPIA (<http://gepia.cancer-pku.cn>) (34). The program was used to analyze RNA sequencing data of 33 types of cancer (including adrenocortical carcinoma, bladder urothelial carcinoma, breast invasive carcinoma (BRCA), cervical squamous cell carcinoma and endocervical adenocarcinoma (CESC), cholangiocarcinoma (CHOL), colon adenocarcinoma (COAD), lymphoid neoplasm diffuse large B-cell lymphoma, esophageal carcinoma (ESCA), glioblastoma multiforme, head and neck squamous cell carcinoma (HNSC), kidney chromophobe (KICH), kidney renal clear cell carcinoma, kidney renal papillary cell carcinoma, acute myeloid leukemia, brain lower grade glioma, liver hepatocellular carcinoma, lung adenocarcinoma (LUAD), lung squamous cell carcinoma (LUSC), mesothelioma, ovarian serous cystadenocarcinoma, pancreatic adenocarcinoma, pheochromocytoma and paraganglioma, prostate adenocarcinoma, rectum adenocarcinoma (READ), sarcoma, skin cutaneous melanoma, stomach adenocarcinoma (STAD), testicular germ cell tumor,

thyroid carcinoma, thymoma, uterine corpus endometrial carcinoma, uterine carcinosarcoma, uveal melanoma) and adjacent normal samples from TCGA following the standard processing pipeline. P-value <0.01 and $|\log_{10}FC| > 2$ were set as cutoffs. Box plots were generated to visualize the associations.

Correlations among genes expression in HNSC samples from TCGA through cBioPortal. The correlations between expression patterns of the genes of interest in cancer tissues from TCGA-HNSC dataset were analyzed using Pearson and Spearman correlation coefficient via cBioPortal (<http://www.cbioportal.org>) (35-36).

Statistical analysis. Statistical analyses were performed using Graphpad Prism 5.0 (GraphPad Software, Inc.). Mann-Whitney U test was performed to compare gene expression between tumor and adjacent normal tissues. A two-tailed P-value <0.01 was considered statistically significant.

Results

Some biomarkers distinguished OTSCC tissues from normal tissues. Four GEO datasets (GSE9844, GSE13601, GSE31056, and GSE75538) were imported into the GEO2R analysis tool. Based on DEGs between OTSCC and normal tissues, five upregulated genes [matrix metalloproteinases 1, 3, 10 and 12 (*MMP1*, *MMP3*, *MMP10* and *MMP12*) and laminin subunit gamma 2 (*LAMC2*)] (Fig. 1A) and six downregulated genes (dermatopontin, cartilage intermediate layer protein, keratin 4, ATP binding cassette subfamily A member 8, alcohol dehydrogenase 1B and protein phosphatase 1 regulatory




Figure 1. Venn diagram of (A) upregulated and (B) downregulated DEGs in four GEO datasets (GSE9844, GSE13601, GSE31056 and GSE75538). Each colored area represents the number of DEGs in one corresponding dataset. The cross-areas indicated altered DEGs. DEGs were identified using Mann-Whitney *U* test and statistically significant DEGs were defined for $P < 0.01$ and (A) $\log FC > 2$ (A) or (B) $\log FC < -2$ as cutoffs. DEGs, differentially expressed genes.




Figure 2. Expression levels of (A) *MMP1*, (B) *LAMC2*, (C) *MMP3*, (D) *MMP10* and (E) *MMP12* in oral tongue squamous cell carcinoma and normal tissues based on TCGA database. * $P < 0.01$. LAMC2, laminin subunit gamma 2; MMP1, 3, 10 and 12, matrix metalloproteinase 1, 3, 10 and 12; N, normal tissue; T, cancer tissue.




Figure 3. Comparison of *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* expressions between cancer (Red) and normal tissues (Green) among 23 various types of cancer from The Cancer Genome Atlas according to GEPIA program. (A) *MMP1*; (B) *LAMC2*; (C) *MMP3*; (D) *MMP10*; (E) *MMP12*. The Y axis indicates the \log_2 (TPM + 1) for gene expression. The red and the green bars represent the cancer and normal tissues, respectively. These figures were derived from GEPIA. * $P < 0.01$. BLCA, bladder urothelial carcinoma; BRCA, breast invasive carcinoma; CESC, cervical squamous cell carcinoma and endocervical adenocarcinoma; CHOL, cholangiocarcinoma; COAD, colon adenocarcinoma; ESCA, esophageal carcinoma; GEPIA, Gene Expression Profiling Interactive Analysis; HNSC, head and neck squamous cell carcinoma; KICH, kidney chromophobe; KIRC, kidney renal clear cell carcinoma; LAMC2, laminin subunit gamma 2; LUAD, lung adenocarcinoma; LUSC, lung squamous cell carcinoma; MMP1, 3, 10 and 12, matrix metalloproteinase 1, 3, 10 and 12; PAAD, pancreatic adenocarcinoma; READ, rectum adenocarcinoma; STAD, stomach adenocarcinoma; UCEC, uterine corpus endometrioid carcinoma; TPM, transcripts per million.

subunit 3C) (Fig. 1B) were highlighted in all four datasets. Whether the five upregulated genes could be considered as potential biomarkers for distinguishing OTSCC tissues from normal tissues was therefore assessed.

GO function enrichment analysis. GO enrichment analysis was performed using DAVID tool. The five specific upregulated genes *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* were uploaded into the DAVID software. GO results indicated that these genes were specifically involved in certain biological processes, including 'extracellular matrix disassembly', 'collagen catabolism' and 'proteolysis'. Furthermore, with regards to the molecular function, the products of these five genes mainly comprised metalloendopeptidases, serine-type endopeptidases, calcium ion binders and zinc ion binders. Similarly, GO cell component analysis revealed that the proteins encoded by these genes were significantly enriched for functions associated with the proteinaceous extracellular matrix and extracellular region ($P < 0.01$, Table II).

Gene expression of *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* in OTSCC samples from TCGA. TCGA database was used to analyze the expression levels of the five genes of interest in OTSCC and adjacent normal tissues. The results demonstrated that the expression levels of *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* were significantly upregulated in OTSCC tissues compared with adjacent normal tissues (Fig. 2).

Gene expression of *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* in 23 cancer samples from TCGA. Consistent results were obtained for bladder urothelial carcinoma, BRCA, COAD, ESCA, HNSC, LUAD, LUSC, READ and STAD. *MMP1* expression level was significantly increased in all cancer tissues compared with adjacent normal tissues (Fig. 3A). *LAMC2* was upregulated in CESC, CHOL, ESCA, HNSC, STAD and uterine corpus endometrial carcinoma, whereas it was significantly downregulated in BRCA, kidney chromophobe and kidney renal clear cell carcinoma (Fig. 3B). Furthermore, *MMP3* expression was upregulated in COAD,




Figure 4. Correlations between *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* expression levels in HNSC tissues. (A) Correlation between *MMP1* and *LAMC2* expression levels in HNSC tissues. (B) Correlation between *MMP1* and *MMP3* expression levels in HNSC tissues. (C) Correlation between *MMP1* and *MMP10* expression levels in HNSC tissues. (D) Correlation between *LAMC2* and *MMP10* expression levels in HNSC tissues. Analyses were performed using cBioPortal on the provisional datasets from The Cancer Genome Atlas. HNSC, head and neck squamous cell carcinoma; LAMC2, laminin subunit gamma 2; MMP1, 3, 10 and 12, matrix metalloproteinase 1, 3, 10 and 12.

ESCA, HNSC, and READ (Fig. 3C). *MMP10* expression was significantly upregulated in ESCA, HNSC and LUSC (Fig. 3D). In addition, *MMP12* was upregulated in CESC, ESCA, HNSC, LUAD, LUSC and STAD, whereas it was downregulated in pancreatic adenocarcinoma (Fig. 3E). Overall, the expression level of *MMP1*, *LAMC2*, *MMP3*, *MMP10* and *MMP12* was significantly upregulated in HNSC tissues.

Correlations among genes expression in samples from the TCGA-HNSC database. To select the five genes of interest for further investigation, the correlations between gene expression patterns in samples from TCGA were analyzed using cBioPortal (35-36). The results demonstrated that *MMP1* expression level was positively correlated with *LAMC2* (Fig. 4A), *MMP3* (Fig. 4B) and *MMP10* (Fig. 4C) expression levels. In addition, *LAMC2* expression level was positively correlated with *MMP10* expression level (Fig. 4D).

Pearson correlation coefficients ranged from 0.33 to 0.69, and Spearman correlation coefficients ranged from 0.55 to 0.83. All together, these results suggested that *MMP1*, *LAMC2*, *MMP3* and *MMP10* expression levels were positively correlated in HNSC tissues.

Discussion

The American Joint Committee on Cancer has updated the Tumor-Node-Metastasis system for cancer staging system according to diagnostic class, treatment choices and prognosis for OTSCC (37). However, with the development of novel modern therapeutic approaches for cancer, genetic analysis is an outstanding tool for early diagnosis of cancer that can prolong patient survival (38). Previous clinical studies on OTSCC development and progression reported increased incidence and mortality rates due to the lack of knowledge on

OTSCC (39,40). The present study identified 11 DEGs (five upregulated and six downregulated genes) from four mRNA expression profile datasets via GEO. These DEGs were significantly associated with OTSCC pathogenesis. The present study therefore focused on the five upregulated DEGs since they may be considered as potential reliable biomarkers. These five target genes were categorized into three classes according to their associated biological functions (molecular functions, biological processes and cellular components) thanks to the GO enrichment using multiple approaches. The results demonstrated that all genes were involved in extracellular matrix disassembly and were localized in the extracellular region during cancer development. Furthermore, expression levels of these genes were positively correlated with each other and upregulated in OTSCC and HNSC tissues compared with adjacent normal tissues, as confirmed using DAVID.

Expression levels of four MMP family members, including *MMP1*, *MMP3*, *MMP10* and *MMP12*, were upregulated in HNSC tissues compared with adjacent normal tissues. These genes serve important functions in numerous physiological and pathological processes involved in tumor progression and can promote tumor-induced angiogenesis and extracellular matrix disassembly, enhancing therefore tumor invasion and metastasis (41,42). *MMP1* is known to be significantly associated with poor clinical outcomes and is therefore a robust prognostic factor for various types of cancer, including colorectal cancer and breast cancer (43-45). Furthermore, *MMP3* has been proposed as a vital tumor oncogene in numerous cancers, including pancreatic, pulmonary, and mammary carcinoma (46). In addition, *MMP10* has been demonstrated to be a potential clinical marker for cancer stem-like cells/cancer-initiating cells in epithelial ovarian cancer (EOC) and may serve as a therapeutic target in chemotherapy-resistant EOCs (47). *MMP12* has been reported to be highly expressed in nasopharyngeal carcinoma both *in vitro* and *in vivo*, and is therefore a powerful tumor marker for cancer metastasis (48). Furthermore, *LAMC2*, which encodes the major component of laminin-5, was one of the upregulated genes in the present study. Highly expression levels of *LAMC2* have been previously reported in numerous invasive tumors (49,50), including carcinomas of the lung (51), the colorectum (52,53) and the pancreas (54,55). In addition, it has been reported that *LAMC2* overexpression in esophageal cancer is associated with poor survival (56,57). Subsequently, *MMP1*, *MMP3*, *MMP10*, *MMP12* and *LAMC2* may serve as novel promising prognostic factors in advanced metastasis of OTSCC. The protein expression of *MMP1*, *MMP3*, *MMP10*, *MMP12* and *LAMC2* should be further confirmed in OTSCC tissues. Furthermore, the correlation between *MMP1*, *MMP3*, *MMP10*, *MMP12* and *LAMC2* protein expression and the survival and clinic characteristics of patients with OTSCC should be further assessed. Tissue samples from patients with OTSCC are currently being collected for this purpose, and will be used for immunohistochemistry analysis to detect MMPs protein expression. Furthermore, the clinical significance, biological function and underlying mechanism of MMPs on the pathogenesis and progression of OTSCC will be further investigated.

In conclusion, the present study identified five upregulated genes that may be considered as robust and reliable biomarkers in OTSCC prognosis. The findings from this study may help discovering the role of oncogenes in cancer progression and

determining the underlying pathways in order to develop novel therapeutic strategies for OTSCC.

Acknowledgements

Not applicable.

Funding

The present study was funded by the National Natural Science Foundation of China (grant no. 81673721) and the Natural Science Foundation of Fujian Province (grant no. 2017I0007).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

PZ, LL, YC and JP conceived and designed the experiments. PZ, AS, ZC and XH performed GEO and GEO2R analyses. LL, XH, YC and JL conducted gene ontology analyses using DAVID. BW, JL, YC and JP analyzed TCGA database using GEPIA. LL, AS, ZC and YC analyzed expressions of genes in OTSCC from the TCGA database. XH, JL, BW and JL assessed the correlations among gene expressions using cBioPortal. PZ, LL, YC and JP drafted the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interest

The authors declare that they have no competing interests.

References

1. Jemal A, Bray F, Center MM, Ferlay J, Ward E and Forman D: Global cancer statistics. *CA Cancer J Clin* 61: 69-90, 2011.
2. Ledgerwood LG, Kumar D, Eterovic AK, Wick J, Chen K, Zhao H, Tazi L, Manna P, Kerley S, Joshi R, *et al*: The degree of intratumor mutational heterogeneity varies by primary tumor sub-site. *Oncotarget* 7: 27185-27198, 2016.
3. Ye H, Yu T, Temam S, Ziober BL, Wang J, Schwartz JL, Mao L, Wong DT and Zhou X: Transcriptomic dissection of tongue squamous cell carcinoma. *BMC Genomics* 9: 69, 2008.
4. Döbrossy L: Epidemiology of head and neck cancer: Magnitude of the problem. *Cancer Metastasis Rev* 24: 9-17, 2005.
5. Tímár J, Csuka O, Remenár E, Répássy G and Kásler M: Progression of head and neck squamous cell cancer. *Cancer Metastasis Rev* 24: 107-127, 2005.
6. Stransky N, Egloff AM, Tward AD, Kostic AD, Cibulskis K, Sivachenko A, Kryukov GV, Lawrence MS, Sougnez C, McKenna A, *et al*: The mutational landscape of head and neck squamous cell carcinoma. *Science* 333: 1157-1160, 2011.
7. Patel SC, Carpenter WR, Tyree S, Couch ME, Weissler M, Hackman T, Hayes DN, Shores C and Chera BS: Increasing incidence of oral tongue squamous cell carcinoma in young white women, age 18 to 44 years. *J Clin Oncol* 29: 1488-1494, 2011.

8. Ng JH, Iyer NG, Tan MH and Edgren G: Changing epidemiology of oral squamous cell carcinoma of the tongue: A global study. *Head Neck* 39: 297-304, 2017.
9. Bello IO, Soini Y and Salo T: Prognostic evaluation of oral tongue cancer: Means, markers and perspectives (I). *Oral Oncol* 46: 630-635, 2010.
10. Almangush A, Heikkinen I, Mäkitie AA, Coletta RD, Läärä E, Leivo I and Salo T: Prognostic biomarkers for oral tongue squamous cell carcinoma: A systematic review and meta-analysis. *Br J Cancer* 117: 856-866, 2017.
11. van Dijk BA, Brands MT, Geurts SM, Merks MA and Roodenburg JL: Trends in oral cavity cancer incidence, mortality, survival and treatment in the Netherlands. *Int J Cancer* 139: 574-583, 2016.
12. Siegel RL, Miller KD and Jemal A: Cancer statistics, 2017. *CA Cancer J Clin* 67: 7-30, 2017.
13. Thangaraj SV, Shyamsundar V, Krishnamurthy A, Ramani P, Ganesan K, Muthuswami M and Ramshankar V: Molecular portrait of oral tongue squamous cell carcinoma shown by integrative meta-analysis of expression profiles with validations. *PLoS One* 11: e0156582, 2016.
14. Ding X, Zhang N, Cai Y, Li S, Zheng C, Jin Y, Yu T, Wang A and Zhou X: Down-regulation of tumor suppressor MTUS1/ATIP is associated with enhanced proliferation, poor differentiation and poor prognosis in oral tongue squamous cell carcinoma. *Mol Oncol* 6: 73-80, 2012.
15. Eslami A, Miyaguchi K, Mogushi K, Watanabe H, Okada N, Shibuya H, Mizushima H, Miura M and Tanaka H: PARVB overexpression increases cell migration capability and defines high risk for endophytic growth and metastasis in tongue squamous cell carcinoma. *Br J Cancer* 112: 338-344, 2015.
16. Zhao T, Ding X, Chang B, Zhou X and Wang A: MTUS1/ATIP3a down-regulation is associated with enhanced migration, invasion and poor prognosis in salivary adenoid cystic carcinoma. *BMC Cancer* 15: 203, 2015.
17. Rogler A, Hoja S, Giedl J, Ekici AB, Wach S, Taubert H, Goebell PJ, Wullich B, Stöckle M, Lehmann J, *et al*: Loss of MTUS1/ATIP expression is associated with adverse outcome in advanced bladder carcinomas: Data from a retrospective study. *BMC Cancer* 14: 214, 2014.
18. Bravou V, Antonacopoulou A, Papanikolaou S, Nikou S, Lilis I, Giannopoulou E and Kalofonos HP: Focal adhesion proteins α - and β -parvin are overexpressed in human colorectal cancer and correlate with tumor progression. *Cancer Invest* 33: 387-397, 2015.
19. Qiu Z, Sun W, Gao S, Zhou H, Tan W, Cao M and Huang W: A 16-gene signature predicting prognosis of patients with oral tongue squamous cell carcinoma. *PeerJ* 5: e4062, 2017.
20. Bullinger L, Döhner K, Bair E, Fröhling S, Schlenk RF, Tibshirani R, Döhner H and Pollack JR: Use of gene-expression profiling to identify prognostic subclasses in adult acute myeloid leukemia. *N Engl J Med* 350: 1605-1616, 2004.
21. Valk PJ, Verhaak RG, Beijnen MA, Erpelinck CA, Barjesteh van Waalwijk van Doorn-Khosrovani S, Boer JM, Beverloo HB, Moorhouse MJ, van der Spek PJ, Löwenberg B and Delwel R: Prognostically useful gene-expression profiles in acute myeloid leukemia. *N Engl J Med* 350: 1617-1628, 2004.
22. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA Jr and Kinzler KW: Cancer genome landscapes. *Science* 339: 1546-1558, 2013.
23. Estilo CL, O-charoenrat P, Talbot S, Socci ND, Carlson DL, Ghossein R, Williams T, Yonekawa Y, Ramanathan Y, Boyle JO, *et al*: Oral tongue cancer gene expression profiling: Identification of novel potential prognosticators by oligonucleotide microarray analysis. *BMC Cancer* 9: 11, 2009.
24. Reis PP, Waldron L, Perez-Ordóñez B, Pintilie M, Galloni NN, Xuan Y, Cervigne NK, Warner GC, Makitie AA, Simpson C, *et al*: A gene signature in histologically normal surgical margins is predictive of oral carcinoma recurrence. *BMC Cancer* 11: 437, 2011.
25. Krishnan NM, Dhas K, Nair J, Palve V, Bagwan J, Siddappa G, Suresh A, Kekatpure VD, Kuriakose MA and Panda B: A minimal DNA methylation signature in oral tongue squamous cell carcinoma links altered methylation with tumor attributes. *Mol Cancer Res* 14: 805-819, 2016.
26. Clough E and Barrett T: The gene expression omnibus database. *Methods Mol Biol* 1418: 93-110, 2016.
27. Davis S and Meltzer PS: GEOquery: A bridge between the Gene Expression Omnibus (GEO) and BioConductor. *Bioinformatics* 23: 1846-1847, 2007.
28. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al*: Gene ontology: Tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25: 25-29, 2000.
29. Huang da W, Sherman BT and Lempicki RA: Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57, 2009.
30. Mandel-Brehm C, Retallack H, Knudsen GM, Yamana A, Hajj-Ali RA, Calabrese LH, Tihan T, Sample HA, Zorn KC, Gorman MP, *et al*: Exploratory proteomic analysis implicates the alternative complement cascade in primary CNS vasculitis. *Neurology* 93: e433-e444, 2019.
31. Zhu Y, Qiu P and Ji Y: TCGA-assembler: Open-source software for retrieving and processing TCGA data. *Nat Methods* 11: 599-600, 2014.
32. Doecke JD, Chekouo TT, Stingo F and Do KA: miRNA target gene identification: Sourcing miRNA target gene relationships for the analyses of TCGA illumina MiSeq and RNA-Seq HiSeq platform data. *Int J Hum Genet* 14: 17-22, 2017.
33. Chen WJ, Tang RX, He RQ, Li DY, Liang L, Zeng JH, Hu XH, Ma J, Li SK and Chen G: Clinical roles of the aberrantly expressed lncRNAs in lung squamous cell carcinoma: A study based on RNA-sequencing and microarray data mining. *Oncotarget* 8: 61282-61304, 2017.
34. Tang Z, Li C, Kang B, Gao G, Li C and Zhang Z: GEPIA: A web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res* 45: W98-W102, 2017.
35. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, *et al*: The cBio cancer genomics portal: An open platform for exploring multi-dimensional cancer genomics data. *Cancer Discov* 2: 401-404, 2012.
36. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, *et al*: Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6: p11, 2013.
37. Amit M, Tam S, Takahashi H, Choi KY, Zafereo M, Bell D and Weber RS: Prognostic performance of the American Joint Committee on Cancer 8th edition of the TNM staging system in patients with early oral tongue cancer. *Head Neck* 41: 1270-1276, 2019.
38. Cancer Genome Atlas Network: Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature* 517: 576-582, 2015.
39. Ashley EA: The precision medicine initiative: A new national effort. *JAMA* 313: 2119-2120, 2015.
40. Pitiyage G, Tilakaratne WM, Tavassoli M and Warnakulasuriya S: Molecular markers in oral epithelial dysplasia: Review. *J Oral Pathol Med* 38: 737-752, 2009.
41. Duffy MJ: Use of biomarkers in screening for cancer. *Adv Exp Med Biol* 867: 27-39, 2015.
42. Kerkelä E and Saarialho-Kere U: Matrix metalloproteinases in tumor progression: Focus on basal and squamous cell skin cancer. *Exp Dermatol* 12: 109-125, 2003.
43. Stamenkovic I: Matrix metalloproteinases in tumor invasion and metastasis. *Semin Cancer Biol* 10: 415-433, 2000.
44. Li H, Zhong A, Li S, Meng X, Wang X, Xu F and Lai M: The integrated pathway of TGF β /Snail with TNF α /NF κ B may facilitate the tumor-stroma interaction in the EMT process and colorectal cancer prognosis. *Sci Rep* 7: 4915, 2017.
45. Shen CJ, Kuo YL, Chen CC, Chen MJ and Cheng YM: MMP1 expression is activated by Slug and enhances multi-drug resistance (MDR) in breast cancer. *PLoS One* 12: e0174487, 2017.
46. Lu X, Wang Q, Hu G, Van Poznak C, Fleisher M, Reiss M, Massagué J and Kang Y: ADAMTS1 and MMP1 proteolytically engage EGF-like ligands in an osteolytic signaling cascade for bone metastasis. *Genes Dev* 23: 1882-1894, 2009.
47. Mehner C, Miller E, Nassar A, Bamlet WR, Radisky ES and Radisky DC: Tumor cell expression of MMP3 as a prognostic factor for poor survival in pancreatic, pulmonary, and mammary carcinoma. *Genes Cancer* 6: 480-489, 2015.
48. Mariya T, Hirohashi Y, Torigoe T, Tabuchi Y, Asano T, Saijo H, Kuroda T, Yasuda K, Mizuuchi M, Saito T and Sato N: Matrix metalloproteinase-10 regulates stemness of ovarian cancer stem-like cells by activation of canonical Wnt signaling and can be a target of chemotherapy-resistant ovarian cancer. *Oncotarget* 7: 26806-26822, 2016.

49. Chung IC, Chen LC, Chung AK, Chao M, Huang HY, Hsueh C, Tsang NM, Chang KP, Liang Y, Li HP and Chang YS: Matrix metalloproteinase 12 is induced by heterogeneous nuclear ribonucleoprotein K and promotes migration and invasion in nasopharyngeal carcinoma. *BMC Cancer* 14: 348, 2014.
50. Koshikawa N, Moriyama K, Takamura H, Mizushima H, Nagashima Y, Yanoma S and Miyazaki K: Overexpression of laminin gamma 2 chain monomer in invading gastric carcinoma cells. *Cancer Res* 59: 5596-5601, 1999.
51. Pyke C, Salo S, Ralfkiaer E, Rømer J, Danø K and Tryggvason K: Laminin-5 is a marker of invading cancer cells in some human carcinomas and is coexpressed with the receptor for urokinase plasminogen activator in budding cancer cells in colon adenocarcinomas. *Cancer Res* 55: 4132-4139, 1995.
52. Kagesato Y, Mizushima H, Koshikawa N, Kitamura H, Hayashi H, Ogawa N, Tsukuda M and Miyazaki K: Sole expression of laminin gamma 2 chain in invading tumor cells and its association with stromal fibrosis in lung adenocarcinomas. *Jpn J Cancer Res* 92: 184-192, 2001.
53. Hlubek F, Jung A, Kotzor N, Kirchner T and Brabletz T: Expression of the invasion factor laminin gamma2 in colorectal carcinomas is regulated by beta-catenin. *Cancer Res* 61: 8089-8093, 2001.
54. Sordat I, Bosman FT, Dorta G, Rousselle P, Aberdam D, Blum AL and Sordat B: Differential expression of laminin-5 subunits and integrin receptors in human colorectal neoplasia. *J Pathol* 185: 44-52, 1998.
55. Soini Y, Määttä M, Salo S, Tryggvason K and Autio-Harmainen H: Expression of the laminin gamma 2 chain in pancreatic adenocarcinoma. *J Pathol* 180: 290-294, 1996.
56. Takahashi S, Hasebe T, Oda T, Sasaki S, Kinoshita T, Konishi M, Ochiai T and Ochiai A: Cytoplasmic expression of laminin gamma2 chain correlates with postoperative hepatic metastasis and poor prognosis in patients with pancreatic ductal adenocarcinoma. *Cancer* 94: 1894-1901, 2002.
57. Yamamoto H, Itoh F, Iku S, Hosokawa M and Imai K: Expression of the gamma(2) chain of laminin-5 at the invasive front is associated with recurrence and poor prognosis in human esophageal squamous cell carcinoma. *Clin Cancer Res* 7: 896-900, 2001.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.