

Expression and function of long non-coding RNA LINC01420 in thyroid cancer

JIN-ZHU LUO^{1*}, LU QIN^{2*} and LING-JIE ZHANG³

Departments of ¹Clinical Laboratory and ²Thyroid Vascular Surgery, Jingzhou Central Hospital, The Second Clinical Medical College, Yangtze University, Jingzhou, Hubei 434020; ³Department of Anesthesiology, Hubei Provincial Hospital of Integrated Chinese and Western Medicine, Wuhan, Hubei 430015, P.R. China

Received February 1, 2019; Accepted September 20, 2019

DOI: 10.3892/ol.2019.11076

Abstract. The Human Genome Project revealed that >90% of the human genome was found to transcribe non-coding RNAs, including micro RNAs and long non-coding RNAs (lncRNAs). lncRNAs have been identified to play a crucial role in cancer progression. Thyroid cancer (TC) is a common type of endocrine cancer; however, the functional roles of lncRNAs in TC have yet to be fully elucidated. The present study investigated whether LINC01420 was upregulated in TC tissues, compared with normal thyroid tissues, and the results suggested that LINC01420 may play a regulatory role in TC. Bioinformatics analysis demonstrated that LINC01420 was associated with translation, rRNA processing, mRNA splicing, regulation of transcription, DNA repair and double-strand break repair. Furthermore, the exact role of LINC01420 in TC was explored by performing a loss-of-function assay, which revealed that the knockdown of LINC01420 inhibited TC cell proliferation and cell cycle progression. The findings of the present study provide a novel insight into the molecular mechanisms underlying TC development. Moreover, they suggest that LINC01420 may serve as a potential therapeutic target for the treatment of TC, and that increased LINC01420 expression levels show potential as a prognostic marker for the disease.

Introduction

The completion of the Human Genome Project illustrated that >90% of the human genome transcribes non-coding RNAs (ncRNAs). It has been demonstrated that ncRNAs [including microRNAs (miRNAs) and long non-coding RNAs (lncRNAs)] play a key role in cancer progression. lncRNAs (non-coding RNA molecules >200 nucleotides in length) are dysregulated and may function as potential biomarkers of various malignancies, such as breast (1), gastric (2), lung (3) and prostate cancer (4). Moreover, lncRNAs may act as oncogenes in certain types of cancer. For example, small nucleolar RNA host gene 15 promotes colon cancer progression by interacting with, and stabilizing snail family transcriptional repressor 2 (5) and differentiation antagonizing non-protein coding RNA (6), increasing the proliferative and invasive capacities of gastric cancer cells. However, lncRNAs can also act as tumor suppressors; for instance, lncRNA overexpressed in colon carcinoma-1 suppressed colorectal cancer-cell proliferation by destabilizing HuR protein (7), and maternally expressed 3 suppressed liver cancer-cell proliferation through the inhibition of β -catenin (8). Therefore, investigation into the functional roles of lncRNAs in cancer may provide new insights into the identification of novel diagnostic and therapeutic targets.

Thyroid cancer (TC) is one of the most common endocrine malignancies. In previous years, the functional roles of several lncRNAs in TC have been revealed. Pvt1 oncogene (PVT1) was the first lncRNA to be reported as having a functional role in TC (9). Zhou *et al* (9) found that PVT1 contributed to TC tumorigenesis through the recruitment of enhancer of zeste 2 polycomb repressive complex 2 subunit and the regulation of thyroid stimulating hormone receptor expression. Furthermore, lncRNA CDKN2B antisense RNA 1 was shown to promote TC metastasis through modulation of the transforming growth factor- β /Smad signaling pathway (10). lncRNAs were also found to be involved in the prognosis of TC; lncRNA papillary thyroid carcinoma susceptibility candidate 3 was identified as a tumor suppressor gene in TC (11). Moreover, low expression levels of growth arrest specific 5 were found to be associated with poor prognosis in patients with TC (12). High expression levels of LINC01420 have been associated with poorer overall survival (OS) in patients with nasopharyngeal carcinoma, and

Correspondence to: Professor Ling-Jie Zhang, Department of Anesthesiology, Hubei Provincial Hospital of Integrated Chinese and Western Medicine, 11 Lingjiaohu Road, Jiangnan, Wuhan, Hubei 430015, P.R. China
E-mail: 22457419@qq.com

*Contributed equally

Abbreviations: lncRNAs, long non-coding RNAs; TC, thyroid cancer; ncRNA, non-coding RNAs; miRNA, microRNAs; TCGA, The Cancer Genome Atlas; PTC, papillary thyroid carcinoma; GO, gene ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes

Key words: long non-coding RNA, LINC01420, proliferation, thyroid cancer, prognostic marker

LINC01420-knockdown inhibited nasopharyngeal carcinoma cell migration. However, the functions and underlying molecular mechanisms of LINC01420 in TC progression remain largely unknown.

The present study investigated whether LINC01420 was of value as a biomarker for TC. The expression of LINC01420 was evaluated by analyzing a dataset containing TC patient information retrieved from The Cancer Genome Atlas (TCGA). Additionally, bioinformatics analysis was performed to reveal the functional roles of LINC01420 in TC progression. Finally, the effect of LINC01420 on cell proliferation and cell cycle progression was investigated. Improved understanding of the role of LINC01420 in TC progression may indicate its use as either a biomarker, or a potential therapeutic target.

Materials and methods

TCGA dataset retrieval and analysis. The TCGA Thyroid carcinoma (THCA) dataset containing specimens from both TC patients and disease-free subjects was retrieved from TCGA (https://www.cbiportal.org/study/summary?id=thym_tcga), and the LINC01420 expression levels were determined (13). In total, 502 PTC tissue samples and 58 normal thyroid tissue samples were included in the TCGA-THCA dataset. Clinical information regarding LINC01420 was downloaded using cBioPortal (<http://www.cbiportal.org/>). The tumor-node-metastasis classification system (detailed in the American Joint committee on Cancer Manual) was used to determine disease stage. The inclusion criteria have been previously reported in TCGA.

Co-expression network, gene ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis. Correlation between the expression of LINC01420 in cancerous and disease-free tissues was calculated using the Pearson's correlation coefficient in cBioPortal (<http://www.cbiportal.org/>). The co-expressed LINC01420-mRNA pair with an absolute Pearson's correlation coefficient ≥ 0.3 was selected for analysis. GO and KEGG pathway analysis were used to predict the biological functions of LINC01420 using the MAS3.0 system (<http://bioinfo.capitalbio.com/mas3/>). $P < 0.05$ was considered to indicate a statistically significant difference. Furthermore, key positively- and negatively-related gene-mediated protein-protein interaction networks were identified using the Search Tool for the Retrieval of Interacting Genes/Proteins database (<http://string.embl.de/>) (14). Cytoscape software (version 3.6.1; <https://cytoscape.org/>) is a free visualization software (15). A confidence score > 0.4 was considered as the criterion of judgment.

Cell culture and transfection. All cell lines were obtained from the American Type Culture Collection. CAL62 and SW579 cells were cultured in L-15 medium supplemented with 10% FBS (Gibco; Thermo Fisher Scientific, Inc.) in a 37°C incubator with 5% CO₂. The short interfering (si)RNA for LINC01420 (5'-CAUCUCAGGUCUCUUGGCUUUGCCA-3') and an siRNA negative control were purchased from Guangzhou RiboBio Co., Ltd. Cells were transfected with siRNAs (10 nM) using Lipofectamine™ 3000 reagent (Thermo Fisher Scientific, Inc.).

Reverse transcription-quantitative (RT-q)PCR analysis. Total RNA was extracted from transfected CAL62 and SW579 cells using TRIzol® (Sigma-Aldrich; Merck KGaA) reagent, according to the manufacturer's protocol. Total RNA was then reverse transcribed into cDNA using the PrimeScript RT Master Mix (Takara Bio, Inc.), according to the manufacturer's protocol. qPCR was performed using the AceQ qPCR SYBR® Green Master Mix (Vazyme) on a Roche LightCycler 480 according to the manufacturer's protocol. The C value of β -actin was used as an internal control to calibrate the Cq values of the genes of interest, in order to determine the differential expression levels. Relative RNA expression was calculated using the $2^{-\Delta\Delta Cq}$ method (16), and each sample was run in triplicate.

Cell proliferation and cell cycle distribution. The Cell counting kit (CCK)-8 assay (Dojindo Molecular Technologies, Inc.) was used to detect cell proliferation following transfection according to the manufacturer's protocol. After 48 h transfection, cells were seeded in 96 well plate at the density of 4×10^3 cells/100 μ l per well, and proliferation was detected at 0, 1, 2, 3 and 4 days. CCK-8 reagent (10 μ l medium/well) was added prior to detection and after incubation for 1.5 h at 37°C, and the absorbance was measured at 450 nm using a microplate reader. Absorbance at 630 nm was used as the control.

For the cell cycle assays, 3×10^5 cells/well were harvested from a 6 well plate and fixed in ice-cold 70% (v/v) ethanol for 24 h at 4°C. The cell pellet was collected following centrifugation at 500 x g for 10 min at 4°C and resuspended in PBS. Cells were then stained with a mixture of RNase (10 μ g/ml) and propidium iodide (50 μ g/ml; Beyotime Institute of Biotechnology) in sodium citrate containing 0.5% Triton X-100 for 20 min, in the dark and at room temperature. The cells were subsequently analyzed using a flow cytometer (Gallios, Beckman Coulter, Inc.). The ModFit software version 4.0 (Verity Software House, Inc.) was used for data analysis.

Statistical analysis. Statistical analysis was performed using the SPSS software package, version 15.0 (SPSS, Inc.). Significant differences between two groups were compared using two-tailed Student's t-test. Statistical comparisons between two paired groups was performed using a paired t-test. For comparison of > 2 groups, one-way ANOVA was used, followed by the Newman-Keuls post hoc test. Kaplan-Meier and Cox regression analyses were used to evaluate the association between LINC01420 and disease-free survival (DFS), as well as the prognosis of patients with TC. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

LINC01420 is overexpressed in TC tissue samples. The TCGA-THCA dataset was analyzed to evaluate the expression levels of LINC01420 in TC. The data suggested that the expression of LINC01420 was significantly higher in TC tissues, compared with that in normal tissue samples (Fig. 1A). Furthermore, LINC01420 expression was analyzed using 50 paired TC samples from TCGA dataset. Of these paired samples, 80% exhibited a higher expression level




Figure 1. LINC01420 expression levels in TC tissues. (A) LINC01420 is upregulated in TC tissues compared with normal tissues. (B and C) In 50 paired TC samples from The Cancer Genome Atlas, LINC01420 was upregulated in the TC samples compared with the adjacent matched normal samples in 80% of cases. ***P<0.001. TC, thyroid cancer.

of LINC01420 in TC samples, compared with the adjacent normal tissues (Fig. 1B and C). The association of LINC01420 expression levels with clinicopathological data was also analyzed (Table SI). The results of these analyses suggest that LINC01420 may play a regulatory role in TC.

Co-expression analysis of LINC01420 in TC. Co-expression analysis is widely used to explore the potential roles of lncRNAs in human disease. Considering that LINC01420 is a novel lncRNA implicated in TC, a LINC01420-mediated co-expression network was constructed to identify its potential mechanism of action. A Pearson's correlation coefficient value of ≥ 0.50 was selected as the cut-off for the identification of reliable LINC01420-mRNA pairs. A total of 948 mRNAs were positively co-expressed and 568 mRNAs were negatively

co-expressed in this network. The two largest hub networks of positively and negatively co-expressed mRNAs are shown in Fig. 2A and C, respectively.

Furthermore, key positively- and negatively-related gene-mediated protein-protein interaction networks were identified using the Search Tool for the Retrieval of Interacting Genes/Proteins database. GO analysis revealed that genes positively related to LINC01420 were significantly associated with 'translation', 'rRNA processing', 'translational initiation', 'mRNA splicing', 'regulation of cellular amino acid metabolic processes', 'NIK/NF- κ B signaling', 'mitochondrial translation', 'regulation of mRNA stability', 'Wnt signaling pathway', 'stimulatory C-type lectin receptor signaling pathway', 'protein targeting to mitochondrion', 'spliceosomal snRNP assembly', 'protein




Figure 2. Co-expression analysis of LINC01420 in thyroid cancer. Top two hub networks were constructed for LINC01420 (A) positively co-expressed mRNAs and (C) negatively co-expressed mRNAs. Gene Ontology analysis for LINC01420 (B) positively-related and (D) negatively-related genes using the MAS3.0 system (<http://bioinfo.capitalbio.com/mas3/>).

polyubiquitination’, ‘TNF- α -mediated signaling pathway’ and ‘T-cell receptor signaling pathway’ (Fig. 2B). Genes negatively associated with LINC01420 were significantly involved in ‘regulation of transcription’, ‘peptidyl-serine phosphorylation’, ‘DNA repair’, ‘double-strand break repair’, ‘protein polyubiquitination’, ‘stem cell population maintenance’, ‘protein import into nucleus’, ‘protein K48-linked deubiquitination’, ‘telomere maintenance’ and ‘cytoplasmic microtubule organization’ (Fig. 2D).

Knockdown of LINC01420 inhibits cell proliferation in TC. The CCK-8 was used to evaluate the functional roles of LINC01420 in TC. Following transfection, the expression of

LINC01420 was found to be significantly downregulated in the LINC01420-knockdown group compared with that in the control group, in both CAL62 (Fig. 3A) and SW579 cells (Fig. 3C). The results of the CCK-8 assay indicated that compared with the control group, LINC01420 silencing significantly suppressed CAL62 (Fig. 3B) and SW579 (Fig. 3D) cell proliferation at 72 h.

LINC01420-knockdown inhibits cell cycle progression in TC. The effect of LINC01420 on the cell cycle was also investigated. Flow cytometry demonstrated that suppressing LINC01420 expression modulated the cell cycle by inducing G₀/G₁ arrest (compared with the control groups) in CAL62 (Fig. 4A-B) and SW579 cells (Fig. 4C-D).




Figure 3. LINC01420-knockdown inhibits the proliferation of thyroid cancer cells. Relative expression levels of LINC01420 in (A) CAL62 and (C) SW579 cells. LINC01420-knockdown significantly suppresses the proliferation of (B) CAL62 and (D) SW579 cells compared with the control-transfected group. Comparison between two groups was performed using two-tailed Student's t-tests. ***P<0.001 vs. control-transfected group. NC, negative control; si, small interfering RNA.




Figure 4. Knockdown of LINC01420 inhibits cell cycle progression in thyroid cancer. Silencing of LINC01420 in TC cells modulated the cell cycle by inducing G₀/G₁ arrest in (A and B) CAL62 and (C and D) SW579 cells. *P<0.05, **P<0.01 and ***P<0.001. TC, thyroid cancer; NC, negative control; si, small interfering RNA.

Discussion

lncRNAs have been identified as important regulators of cancer progression, binding to DNA, RNA and proteins to regulate epigenetic modification and protein and gene post-transcriptional regulation (17). lncRNAs act as both oncogenes and tumor suppressor genes; for example, epigenetically-induced lncRNA1 was reported to be oncogenic, promoting cell cycle progression by interacting with the MYC proto-oncogene (18). Conversely, testis associated oncogenic lncRNA promotes cancer progression and mRNA stabilization by interacting with insulin-like growth factor 2 mRNA binding protein 1 (19). Moreover, certain lncRNAs, such as X-inactive specific transcript, nuclear paraspeckle assembly transcript 1 and HOX transcript antisense RNA, were found to act as miRNA molecular sponges, which are implicated in human cancer progression (20-24). The current study investigated the role of LINC01420 in TC progression. LINC01420 expression was previously reported to be upregulated in nasopharyngeal carcinoma, and LINC01420-knockdown inhibited nasopharyngeal carcinoma-cell migration (25). In the present study, a loss-of-function assay was performed and revealed that LINC01420-knockdown inhibited TC cell proliferation by arresting cell cycle progression. Taken together, these findings provide evidence to support the oncogenic nature of LINC01420.

TC is a common endocrine malignancy; however no sensitive biomarkers are currently available for TC diagnosis. Previous studies have primarily focused on identifying diagnostic and therapeutic targets for TC. For example, Read *et al* (26) reported that higher PTTG1 regulator of sister chromatid separation, securin and zinc finger protein 395 expression predicted poor patient outcomes in TC. Downregulation of serum dickkopf WNT signaling pathway inhibitor 1 was also found to be associated with a poor prognosis in PTC patients (27). Furthermore, lncRNAs are often dysregulated in TC; NEAT1_2 (28), TNRC6C-AS1 (29), CNALPTC1 (30) and AFAP1-AS1 (31) were found to be overexpressed in TC. However, GAS8-AS1 (32), GAS5 (12) and BANC1 (33) were all found to be downregulated. To the best of our knowledge, the present study is the first to demonstrate the upregulation of LINC01420 in TC compared with normal tissues, and to determine a correlation between the aforementioned upregulation and poor prognosis. Furthermore, high LINC01420 expression was also associated with shorter DFS time. Thus the results of the present study suggest that LINC01420 may play a regulatory role in TC development, and consequently, may be a useful biomarker.

The functional roles and underlying molecular mechanisms of LINC01420 in TC progression remain largely unclear. Consequently, GO and KEGG pathway analysis were also performed in the present study. The findings demonstrated that LINC01420 was significantly associated with translation, rRNA processing, translational initiation, mRNA splicing, regulation of transcription, DNA repair and double-strand break repair.

In conclusion, the present study determined that LINC01420 is implicated in proliferation and cell cycle progression in TC. It was also observed that LINC01420 expression was upregulated in human TC tissues. Bioinformatics analyses demonstrated that LINC01420 was associated with translation, rRNA processing, translational initiation, mRNA splicing, regulation of transcription, DNA repair and double-strand

break repair. Therefore, the results of the present study indicate that LINC01420 may serve as a potential therapeutic target, and that increased LINC01420 levels may be used as a novel prognostic biomarker for TC.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets generated and/or analyzed during the current study are available in the TCGA-THCA repository (https://www.cbiportal.org/study/summary?id=thym_tcga).

Authors' contributions

Experimental conception and design were conducted by JZL and LJZ. JZL, LQ and LJZ developed the methodology, and analyzed and interpreted the data. JZL, LQ and LJZ wrote, reviewed and revised the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Lv M, Xu P, Wu Y, Huang L, Li W, Lv S, Wu X, Zeng X, Shen R, Jia X, *et al*: lncRNAs as new biomarkers to differentiate triple negative breast cancer from non-triple negative breast cancer. *Oncotarget* 7: 13047-13059, 2016.
2. Zhou X, Yin C, Dang Y, Ye F and Zhang G: Identification of the long non-coding RNA H19 in plasma as a novel biomarker for diagnosis of gastric cancer. *Sci Rep* 5: 11516, 2015.
3. Wang HM, Lu JH, Chen WY and Gu AQ: Upregulated lncRNA-UCA1 contributes to progression of lung cancer and is closely related to clinical diagnosis as a predictive biomarker in plasma. *Int J Clin Exp Med* 8: 11824-11830, 2015.
4. Crea F, Watahiki A, Quagliata L, Xue H, Pikor L, Parolia A, Wang Y, Lin D, Lam WL, Farrar WL, *et al*: Identification of a long non-coding RNA as a novel biomarker and potential therapeutic target for metastatic prostate cancer. *Oncotarget* 5: 764-774, 2014.
5. Jiang H, Li T, Qu Y, Wang X, Li B, Song J, Sun X, Tang Y, Wan J, Yu Y, *et al*: Long non-coding RNA SNHG15 interacts with and stabilizes transcription factor Slug and promotes colon cancer progression. *Cancer Lett* 425: 78-87, 2018.
6. Mao Z, Li H, Du B, Cui K, Xing Y, Zhao X and Zai S: lncRNA DANCR promotes migration and invasion through suppression of lncRNA-LET in gastric cancer cells. *Biosci Rep* 37: pii: BSR20171070, 2017.
7. Pibouin L, Villaudy J, Ferbus D, Muleris M, Prosperi MT, Remvikos Y and Goubin G: Cloning of the mRNA of overexpression in colon carcinoma-1: A sequence overexpressed in a subset of colon carcinomas. *Cancer Genet Cytogenet* 133: 55-60, 2002.

8. He Y, Wu YT, Huang C, Meng XM, Ma TT, Wu BM, Xu FY, Zhang L, Lv XW and Li J: Inhibitory effects of long noncoding RNA MEG3 on hepatic stellate cells activation and liver fibrogenesis. *Biochim Biophys Acta* 1842: 2204-2215, 2014.
9. Zhou Q, Chen J, Feng J and Wang J: Long noncoding RNA PVT1 modulates thyroid cancer cell proliferation by recruiting EZH2 and regulating thyroid-stimulating hormone receptor (TSHR). *Tumour Biol* 37: 3105-3113, 2016.
10. Zhao JJ, Hao S, Wang LL, Hu CY, Zhang S, Guo LJ, Zhang G, Gao B, Jiang Y, Tian WG and Luo DL: Long non-coding RNA ANRIL promotes the invasion and metastasis of thyroid cancer cells through TGF- β /Smad signaling pathway. *Oncotarget* 7: 57903-57918, 2016.
11. Fan M, Li X, Jiang W, Huang Y, Li J and Wang Z: A long non-coding RNA, PTCSC3, as a tumor suppressor and a target of miRNAs in thyroid cancer cells. *Exp Ther Med* 5: 1143-1146, 2013.
12. Guo LJ, Zhang S, Gao B, Jiang Y, Zhang XH, Tian WG, Hao S, Zhao JJ, Zhang G, Hu CY, *et al*: Low expression of long non-coding RNA GAS5 is associated with poor prognosis of patients with thyroid cancer. *Exp Mol Pathol* 102: 500-504, 2017.
13. Cancer Genome Atlas Research Network: Integrated genomic characterization of papillary thyroid carcinoma. *Cell* 159: 676-690, 2014.
14. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A, Santos A, Tsafou KP, *et al*: STRING v10: Protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 43: D447-D452, 2015.
15. Smoot ME, Ono K, Ruscheinski J, Wang PL and Ideker T: Cytoscape 2.8: New features for data integration and network visualization. *Bioinformatics* 27: 431-432, 2011.
16. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
17. Schmitt AM and Chang HY: Long noncoding RNAs in cancer pathways. *Cancer Cell* 29: 452-463, 2016.
18. Wang Z, Yang B, Zhang M, Guo W, Wu Z, Wang Y, Jia L, Li S, Cancer Genome Atlas Research Network, Xie W and Yang D: lncRNA epigenetic landscape analysis identifies EPIC1 as an oncogenic lncRNA that interacts with MYC and promotes cell-cycle progression in cancer. *Cancer Cell* 33: 706-720.e9, 2018.
19. Chen W, Chen M, Xu Y, Chen X, Zhou P, Zhao X, Pang F and Liang W: Long non-coding RNA THOR promotes human osteosarcoma cell growth in vitro and in vivo. *Biochem Biophys Res Commun* 499: 913-919, 2018.
20. Chen DL, Ju HQ, Lu YX, Chen LZ, Zeng ZL, Zhang DS, Luo HY, Wang F, Qiu MZ, Wang DS, *et al*: Long non-coding RNA XIST regulates gastric cancer progression by acting as a molecular sponge of miR-101 to modulate EZH2 expression. *J Exp Clin Cancer Res* 35: 142, 2016.
21. Kong Q, Zhang S, Liang C, Zhang Y, Kong Q, Chen S, Qin J and Jin Y: LncRNA XIST functions as a molecular sponge of miR-194-5p to regulate MAPK1 expression in hepatocellular carcinoma cell. *J Cell Biochem* 119: 4458-4468, 2018.
22. Zhang J, Li Y, Dong M and Wu D: Long non-coding RNA NEAT1 regulates E2F3 expression by competitively binding to miR-377 in non-small cell lung cancer. *Oncol Lett* 14: 4983-4988, 2017.
23. Liu XH, Sun M, Nie FQ, Ge YB, Zhang EB, Yin DD, Kong R, Xia R, Lu KH, Li JH, *et al*: Lnc RNA HOTAIR functions as a competing endogenous RNA to regulate HER2 expression by sponging miR-331-3p in gastric cancer. *Mol Cancer* 13: 92, 2014.
24. Ren K, Li Y, Lu H, Li Z, Li Z, Wu K, Li Z and Han X: Long noncoding RNA HOTAIR controls cell cycle by functioning as a competing endogenous RNA in esophageal squamous cell carcinoma. *Transl Oncol* 9: 489-497, 2016.
25. Yang L, Tang Y, He Y, Wang Y, Lian Y, Xiong F, Shi L, Zhang S, Gong Z, Zhou Y, *et al*: High expression of LINC01420 indicates an unfavorable prognosis and modulates cell migration and invasion in nasopharyngeal carcinoma. *J Cancer* 8: 97-103, 2017.
26. Read ML, Fong JC, Modasia B, Fletcher A, Imruetaicharoenchoke W, Thompson RJ, Nieto H, Reynolds JJ, Bacon A, Mallick U, *et al*: Elevated PTTG and PBF predicts poor patient outcome and modulates DNA damage response genes in thyroid cancer. *Oncogene* 36: 5296-5308, 2017.
27. Zhao YP, Wang W, Wang XH, Xu Y, Wang Y, Dong ZF and Zhang JJ: Downregulation of serum DKK-1 predicts poor prognosis in patients with papillary thyroid cancer. *Genet Mol Res* 14: 18886-18894, 2015.
28. Sun W, Lan X, Zhang H, Wang Z, Dong W, He L, Zhang T, Zhang P, Liu J and Qin Y: NEAT1_2 functions as a competing endogenous RNA to regulate ATAD2 expression by sponging microRNA-106b-5p in papillary thyroid cancer. *Cell Death Dis* 9: 380, 2018.
29. Hou S, Lin Q, Guan F and Lin C: LncRNA TNRC6C-AS1 regulates UNC5B in thyroid cancer to influence cell proliferation, migration and invasion as a competing endogenous RNA of miR-129-5p. *J Cell Biochem* 119: 8304-8316, 2018.
30. Chen C, Zhou L, Wang H, Chen J, Li W, Liu W, Shen M, Liu H and Fu X: Long noncoding RNA CNALPTC1 promotes cell proliferation and migration of papillary thyroid cancer via sponging miR-30 family. *Am J Cancer Res* 8: 192-206, 2018.
31. Dai W, Tian Y, Jiang B and Chen W: Down-regulation of long non-coding RNA AFAP1-AS1 inhibits tumor growth, promotes apoptosis and decreases metastasis in thyroid cancer. *Biomed Pharmacother* 99: 191-197, 2018.
32. Zhang D, Liu X, Wei B, Qiao G, Jiang T and Chen Z: Plasma lncRNA GAS8-AS1 as a potential biomarker of papillary thyroid carcinoma in Chinese patients. *Int J Endocrinol* 2017: 2645904, 2017.
33. Liao T, Qu N, Shi RL, Guo K, Ma B, Cao YM, Xiang J, Lu ZW, Zhu YX, Li DS and Ji QH: BRAF-activated lncRNA functions as a tumor suppressor in papillary thyroid cancer. *Oncotarget* 8: 238-247, 2017.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.