

# Biglycan as a potential diagnostic and prognostic biomarker in multiple human cancers

SHOU-FENG ZHAO<sup>1</sup>, XUE-JING YIN<sup>2</sup>, WEN-JU ZHAO<sup>3</sup>, LE-CUI LIU<sup>4</sup> and ZHI-PENG WANG<sup>5</sup>

<sup>1</sup>Central Laboratories, Qingdao Municipal Hospital, Qingdao, Shandong 266000; <sup>2</sup>Department of Pharmacy, Qingdao Mental Health Centre; <sup>3</sup>Operating Room, Qingdao Jinhua Gynecology Hospital, Qingdao, Shandong 266235; <sup>4</sup>Neonatal Department, Wulian County People's Hospital, Wulian, Shandong 262300; <sup>5</sup>Marine Science and Engineering College, Qingdao Agricultural University, Qingdao, Shandong 266109, P.R. China

Received February 16, 2019; Accepted August 16, 2019

DOI: 10.3892/ol.2020.11266

**Abstract.** Biglycan (BGN), a key member of the small leucine-rich proteoglycan family, is an important component of the extracellular matrix. Clinical studies have demonstrated that upregulation of BGN is associated with poor prognosis in patients with various types of solid cancer. The present study analyzed the mRNA expression levels of BGN in various types of solid cancer when compared with that in normal tissues via the Oncomine database. The UALCAN, OncoLnc and Kaplan-Meier Plotter databases were additionally used to evaluate the prognostic values of BGN in patients with solid cancer and co-expression gene analysis was conducted using the protein-protein interaction networks of BGN. The present study observed that the mRNA expression levels of BGN were increased in bladder, brain and central nervous system, breast, colorectal, esophageal, gastric, head and neck, lung, ovarian and 28 subtypes of cancer compared with normal tissues. The increased expression of BGN was identified to be associated with a poor outcome in ovarian and gastric cancer. Based on the co-expression network, BGN was identified as the key gene in a 43-gene network. The present findings of increased expression of BGN in solid tumors and its positive association with poor outcome on patient survival indicate that BGN may serve as a prognostic marker and as a target for novel therapeutics for multiple types of cancer.

## Introduction

Cancer is considered as one of the four major non-communicable diseases (1). Due to a delay in diagnosis, its poor prognosis and high recurrence rate, cancer is becoming one of the leading causes of mortality worldwide (2,3). Cancer incidence

and mortality rates have increased over the last decade. The global cancer burden is estimated to have risen to 18.1 million new cancer cases, and 9.6 million cancer-associated mortalities were reported in 2018 (4), compared with 12.7 million and 7.6 million, respectively, in 2008 (5). Therefore, there is an urgent need to explore novel potential cancer biomarkers that will have beneficial prognostic and therapeutic implications.

Biglycan (BGN; also known as proteoglycan-1 and dermatan sulfate PG-1) is a single-copy gene localized on the long arm of human X chromosome Xq13-qter (6). This gene contains at least two introns and it spans ~6 kb in length (7). BGN is a key member of the small leucine-rich proteoglycan family that resides at the cell surface or in the pericellular space of tissues (8). BGN is typically expressed in the nerve, bone, cartilage, skin and muscles, modulating the morphology, growth, adhesion, bone mineralization, inflammation, migration and differentiation of epithelial cells (9). The upregulation of BGN has been reported in multiple types of solid cancer, including ovarian carcinoma (10), prostate cancer (11), pancreatic cancer (12), gastric cancer (13) and colon cancer (14). Overexpressed BGN has been reported to be associated with the aggressive growth and metastasis of tumors (13,14), and with a worse prognosis for patients with gastric cancer (15) and pancreatic adenocarcinoma (16). These findings suggest that the *BGN* gene may act as either a potential therapeutic target or prognostic biomarker in multiple types of cancer. However, the transcriptional expression and prognostic value of the *BGN* gene in human cancers requires further investigation.

The present study investigated the mRNA expression levels of BGN in human normal and cancer tissues, using the Oncomine database. In addition, the prognostic value of BGN mRNA expression in patients with cancer was also assessed using the UALCAN, OncoLnc and the Kaplan-Meier Plotter databases. Finally, co-expression gene analysis was conducted using the protein-protein interaction (PPI) networks of BGN.

## Materials and methods

*Analysis of BGN expression in multiple cancers using Oncomine.* Oncomine is a cancer microarray database and web-based data-mining platform aimed at facilitating discovery from genome-wide expression analyses, as well as

*Correspondence to:* Dr Zhi-Peng Wang, Marine Science and Engineering College, Qingdao Agricultural University, 17 Wenhai Road, Jimo, Qingdao, Shandong 266109, P.R. China  
E-mail: wangzp@ysfri.ac.cn

*Key words:* biglycan, biomarker, cancer

comparing the transcriptome data in multiple types of cancer, respective to normal tissues (17). To date, the OncoPrint database contains 19 cancer types, 715 datasets and 86,733 samples, corresponding to ~48 million gene expression measurements. Differential mRNA level analyses of BGN were compared between normal tissues and malignant human tissues in different types of cancer, using the OncoPrint database. In the present study, the thresholds were set at 2-fold change,  $P < 1 \times 10^{-4}$  and the top 10% gene rank.

**Analysis using the Kaplan-Meier (KM) plotter, UALCAN and OncoPrint databases.** The prognostic significance of the mRNA expression levels of BGN in various types of cancer was evaluated using KM plotter (<http://www.kmplot.com>), UALCAN (<http://ualcan.path.uab.edu>) and OncoPrint (<http://oncoprint.org>). These online databases can be used to assess the effect of gene expression on cancer prognosis. The three databases of KM plotter, OncoPrint and UALCAN contain the same RNA-seq data (from TCGA) for 20 ([http://www.kmplot.com/analysis/index.php?p=service&cancer=pancancer\\_rnaseq](http://www.kmplot.com/analysis/index.php?p=service&cancer=pancancer_rnaseq)), 21 (18) and 35 (19) types and subtypes of cancer, respectively. KM plotter also contain the gene chip data [from GEO (breast cancer: GSE12276, GSE16391, GSE12093, GSE11121, GSE9195, GSE7390, GSE6532, GSE5327, GSE4922, GSE3494, GSE2990, GSE2034, GSE1456; ovarian cancer: GSE14764, GSE15622, GSE19829, GSE3149, GSE9891, GSE18520, GSE26712; lung cancer: GSE4573, GSE14814, GSE8894, GSE19188, GSE3141, GSE31210, GSE29013, GSE37745; gastric cancer: GSE44740, GSE51725, GSE13911, GSE43346, and GSE3526)] (20–23) for breast cancer (BC), lung cancer (LC), gastric cancer (GC) and ovarian cancer (OC). Therefore, the prognostic significance of the mRNA expression levels of BGN in various types of carcinomas, including BC, LC, GC and OC was evaluated using RNA-seq data and confirmed by gene chip data. The overall survival rate (OS) in patients with other types and subtypes of carcinomas was estimated using RNA-seq by KM plotter, OncoPrint or UALCAN databases.

The KM plotter is able to assess the effect of 54,675 genes on survival using 10,461 cancer samples. In this database, the types and subtypes of cancer samples were observed from RNA-sequencing (RNA-seq) data while the lung (22), ovarian (21), gastric (23), and breast (20) cancer samples were also analyzed from gene chip microarrays.

Patient samples were divided into two cohorts according to the median expression of the BGN gene (high vs. low expression). The present study analyzed the overall survival (OS) in patients using a Kaplan-Meier survival plot. Briefly, the BGN gene was uploaded into the respective databases to obtain the Kaplan-Meier survival plots, in which the number-at-risk was presented below the main plot. Affymetrix ID (or RNA-seq ID), log rank P-value and hazard ratio (HR) with 95% confidence intervals were calculated and displayed on the webpage.  $P < 0.05$  was considered to indicate a statistically significant difference.

UALCAN, is an interactive web resource for analyzing cancer transcriptome data, built on PERL-common gateway interface with high quality graphics using JavaScript and Cascading Style Sheets. UALCAN was used to construct an algorithm based on The Cancer Genome Atlas (TCGA) level 3 RNA-seq database (<https://portal.gdc.cancer.gov/>). UALCAN

can provide publication quality graphs and plots depicting gene expressions and patient survival information based on gene expression (19).  $P < 0.05$  was considered to indicate a statistically significant difference.

OncoPrint is a tool for interactively exploring survival correlations. OncoPrint contains survival data for 8,647 patients from 21 cancer studies performed by TCGA, along with RNA-seq expression for mRNAs and microRNAs from TCGA and long non-coding RNA expressions from MiTranscriptome ( $\beta$  release) (<http://www.mitranscriptome.com/>). OncoPrint stores precomputed survival analyses, allowing users to quickly explore survival correlations for up to 21 types of cancer in a single click (18). The BGN gene was uploaded into the database to obtain the patient survival information.  $P < 0.05$  was considered to indicate a statistically significant difference.

**Co-expression and PPI network construction.** The present study extracted the top 50 co-expressed genes that have similar expression pattern with BGN gene, based on Pearson correlation score across all tumor samples from the GEPIA database (<http://gepia.cancerpku.cn/index.html>). Then, the 11.0 Search Tool for the Retrieval of Interacting Genes/Proteins database (<http://string-db.org/>), was used to construct a PPI network with these co-expressed genes (24). The PPI pairs were extracted with a combined score of 0.4. Subsequently, the PPI network was visualized using the Cytoscape 3.7.0 software (<http://www.cytoscape.org/>).

## Results

**Increased expression of BGN in multiple types of solid cancer.** As illustrated in Fig. 1, OncoPrint contained a total of 421 research studies for the BGN gene. In 78 analyses, BGN exhibited statistically significant differences, 5 of which revealed lower mRNA expression levels in solid tumors compared with normal tissues, while 73 analyses indicated the opposite result (Fig. 1). BGN gene expression was the most upregulated in bladder, brain and central nervous system (CNS), breast, colorectal, gastric, head and neck, pancreatic cancer and other cancer, followed by esophageal, kidney, liver, ovarian cancer, and lung cancer (Fig. 1). Results in the investigation of BGN gene expression have been inconsistent in a number of studies (25–33), including kidney, liver, pancreatic, and prostate cancer. The expression levels of BGN in cervical cancer and melanoma were not significantly changed (Fig. 1). The BGN gene expression status in lymphoma, leukemia, sarcoma and myeloma was not analyzed in this study as they are not classified as solid cancers (34).

Solid cancers, in which BGN was reported to be upregulated consistently in different studies, were examined further in the various cancer subtypes using the OncoPrint datasets. The data summarized in Table I revealed that there was a significant increase in expression of BGN in BLCA, including subtype of IBLCA ( $n=53$ ), when compared with that in normal tissues ( $n=3$ ) in the study by Blaveri *et al.* (35). The BGN mRNA expression was significantly elevated in brain and CNS cancer, including subtype of GBM ( $n=130$ ) when compared with normal tissues ( $n=30$ ) in the study conducted by Bredel *et al.* (36). The mRNA expression of BGN was significantly elevated in BC, including subtypes BRCA, DBC,

Analysis Type by Cancer	Cancers vs. Normal	
Bladder Cancer	1	
Brain and CNS Cancer	3	
Breast Cancer	17	
Cervical Cancer		
Colorectal cancer	13	
Esophageal Cancer	3	
Gastric Cancer	10	
Head and Neck Cancer	1	
Kidney Cancer	2	1
Leukemia	1	
Liver Cancer	1	2
Lung Cancer	1	
Lymphoma	7	
Melanoma		
Myeloma		
Other Cancer	3	
Ovarian Cancer	3	
Pancreatic Cancer	3	1
Prostate Cancer		1
Sarcoma	4	
Significant Unique analyses	73	5
Total Unique analyses	421	



Figure 1. mRNA expression levels of BGN in different types of cancer from the Oncomine database. The schematic reveals the numbers of datasets with statistically significant mRNA overexpression (red) or underexpression (blue) of the target gene. Darker red indicates higher *BGN* expression, Darker blue indicates lower *BGN* expression. The color is determined by the best gene rank percentile for the analyses within the cell. The number in each cell represents the number of analyses that met the thresholds of: Gene, *BGN*; analysis type, cancer vs. normal; and data type, mRNA. The thresholds were set at 2-fold change,  $P < 1 \times 10^{-4}$  and the top 10% gene rank. For cervical cancer, melanoma, myeloma and prostate cancer, there was no analysis (white cell) that met the aforementioned thresholds. The gene rank was analyzed using the percentile of target gene in the top of all genes measured in each analysis. *BGN*, biglycan; CNS, central nervous system.

DCIS, IDBC, IDBCs, IDC, IDC-L, ILBC, ILC, LBC, MBC and TBC (n=2,540) compared with that in normal tissues (n=240), which was performed by Ma *et al* (37), Curtis *et al* (38), Karnoub *et al* (39), Perou *et al* (40), Zhao *et al* (41) and TCGA studies (42,43). The mRNA expression of *BGN* was also significantly upregulated in CC, including subtypes COAD, CMA, RSA, CeAC and READ (n=463), when compared with

that in normal tissues (n=150), which was reported in the studies conducted by Kaiser *et al* (44), Skrzypczak *et al* (45), Graudens *et al* (46), TCGA (47), Gaedcke *et al* (48), and Hong *et al* (49). *BGN* mRNA expression was also significantly upregulated in EC, including subtype ESCC (n=84), when compared with that in normal tissues (n=80), which was reported in the studies of Su *et al* (50), Hu *et al* (51) and Hao *et al* (52). *BGN* mRNA expression was also significantly higher in GC, including subtypes ITGA, DGAC and GMA (n=265), when compared with that in normal tissues (n=171), as reported by Chen *et al* (53), Cho *et al* (54), Wang *et al* (55), D'Errico *et al* (56) and Cui *et al* (57). The mRNA expression of *BGN* was also significantly increased in HNC, including subtype SGACC (n=16) compared with that in normal tissues (n=6) in the study by Frierson *et al* (58). The mRNA expression of *BGN* was significantly higher in LC, including subtype SCC (n=34) compared with that in normal tissues (n=28) in the study by Talbot *et al* (59). The mRNA expression of *BGN* was increased significantly in OC, including subtypes SSPC and OSC (n=799) compared with that in normal tissues (n=22), as reported by Welsh *et al* (60), and Bonome *et al* (61), and studies listed in TCGA (62). The mRNA expression of *BGN* was increased significantly in other types of cancer, including subtype SBCC (n=15) compared with that in normal tissues (n=4), as reported by Riker *et al* (63). Therefore, the expression of *BGN* was significantly increased in human solid cancers, including 10 types and 28 subtypes of carcinoma. These results indicate that the mRNA expression of *BGN* is elevated in a wide range of tumors, when compared with that in normal tissues.

*High BGN expression and survival outcome in multiple types of solid cancer.* The expression of *BGN* was significantly increased in certain types of human solid cancers, including bladder, brain and CNS, breast, colorectal, gastric, head and neck, esophageal, ovarian, lung and other cancers, however there was no data on the downregulation of the *BGN* from the database (Table I). The present study used the Kaplan-Meier Plotter, OncoLnc and UALCAN databases to identify the association between survival time and the mRNA levels of *BGN* in patients with different types and subtypes of solid cancer.

As shown in Table II, *BGN* gene with a significant association with patient survival can be identified in GC and OC ( $P < 0.05$ ). There is no significant association of *BGN* upregulation with patient survival in BC, EC ( $P > 0.05$ ), IBC, READ, COAD, and ESCC ( $P > 0.05$ ). The survival rate of patients with LC with  $P > 0.05$  in RNA-seq and  $P < 0.05$  in microarray analysis requires further investigation. Therefore, high *BGN* mRNA expression may potentially be associated with the prognosis in patients with BLC, LSCC, and OSC, as the present analyses provided RNA-seq analysis results ( $P < 0.05$ ) without microarray analysis confirmation (Table II). The association between *BGN* mRNA expression and prognosis in patients with other types of cancer and subtypes of cancer requires further investigation as there is no prognostic data in the KM plotter, OncoLnc and UALCAN database (Table III).

As presented in Fig. 2, high expression of *BGN* was significantly associated with shorter OS time in patients with GC [HR=1.9 (1.56-2.32),  $P = 1.3 \times 10^{-10}$  in microarray analysis; HR=1.59 (1.13-2.24),  $P = 6.8 \times 10^{-3}$  in RNA-seq analysis] and

Table I. Significant changes of biglycan mRNA expression between different subtypes of bladder, brain and CNS, breast, colorectal, esophageal, gastric, head and neck, lung, ovarian, other types of cancer and 28 subtypes cancer.

Type of cancer	Cancer Subtype (n)	Normal, n	Fold change	t-test	P-value	Rank, %	(Refs.)
Bladder	Infiltrating Bladder Urothelial Carcinoma (IBLCA) (53)	3	2.391	6.799	$7.02 \times 10^{-7}$	1	(35)
	Glioblastoma (GBM) (27)	4	3.055	13.066	$6.77 \times 10^{-14}$	1	(36)
Brain and CNS	Glioblastoma (GBM) (22)	3	13.32	7.773	$4.92 \times 10^{-5}$	4	
	Glioblastoma (GBM) (81)	23	2.163	7.599	$7.14 \times 10^{-11}$	7	
Breast	Ductal Breast Carcinoma in Situ Stroma (DCIS) (11)	14	13.191	7.155	$2.28 \times 10^{-11}$		(37)
	Invasive Ductal Breast Carcinoma Stroma (IDC) (9)	14	7.232	5.357	$1.27 \times 10^{-6}$	1	
	Invasive Lobular Breast Carcinoma (ILC) (36)	61	3.312	13.142	$1.58 \times 10^{-22}$	1	(42,43)
	Invasive Breast Carcinoma (BRCA) (76)	61	3.131	14.001	$1.22 \times 10^{-27}$	1	
	Male Breast Carcinoma (MBC) (3)	61	4.661	10.406	$3.76 \times 10^{-5}$	4	
	Mixed Lobular and Ductal Breast Carcinoma (IDC-L) (7)	61	2.780	5.576	$8.56 \times 10^{-5}$	4	
	Invasive Ductal Breast Carcinoma (IDC) (389)	61	2.714	14.829	$5.11 \times 10^{-25}$	5	
	Tubular Breast Carcinoma (TBC) (67)	144	3.014	18.174	$4.05 \times 10^{-41}$	1	(38)
	Invasive Lobular Breast Carcinoma (ILBC) (148)	144	3.087	19.715	$8.92 \times 10^{-56}$	1	
	Invasive Ductal and Invasive Lobular Breast Carcinoma (IDC and ILC) (90)	144	2.885	14.973	$4.81 \times 10^{-33}$	1	
Colorectal	Breast Carcinoma (BC) (14)	144	2.884	5.505	$3.00 \times 10^{-5}$	4	
	Invasive Ductal Breast Carcinoma (IDC) (1556)	144	2.988	26.199	$1.50 \times 10^{-62}$	5	
	Medullary Breast Carcinoma (MBC) (32)	144	2.256	6.302	$1.13 \times 10^{-7}$	9	
	Invasive Ductal Breast Carcinoma Stroma (IDBCs) (7)	15	3.546	6.337	$2.70 \times 10^{-6}$	1	(39)
	Ductal Breast Carcinoma (DBC) (36)	3	3.337	9.550	$7.02 \times 10^{-6}$	2	(40)
	Lobular Breast Carcinoma (LBC) (21)	3	3.337	10.925	$1.22 \times 10^{-7}$	1	(41)
	Invasive Ductal Breast Carcinoma (IDBC) (38)	3	6.756	12.230	$1.80 \times 10^{-6}$	5	
	Colon Adenocarcinoma (COAD) (41)	5	3.200	13.125	$1.74 \times 10^{-15}$	1	(44)
	Colon Mucinous Adenocarcinoma (CMA) (13)	5	2.942	9.781	$6.65 \times 10^{-8}$	1	
	Rectosigmoid Adenocarcinoma (RSA) (10)	5	3.799	7.377	$4.85 \times 10^{-6}$	2	
	Cecum Adenocarcinoma (CeAC) (17)	5	3.039	6.529	$1.17 \times 10^{-6}$	4	
	Colorectal Carcinoma (CC) (36)	24	4.795	8.671	$3.32 \times 10^{-12}$	1	(45)
	Colorectal Carcinoma (CC) (18)	12	2.563	5.442	$5.94 \times 10^{-6}$	4	(46)
	Colon Carcinoma (cc) (5)	10	7.740	15.965	$3.60 \times 10^{-10}$	2	(45)
	Colon Adenocarcinoma (COAD) (5)	10	4.099	14.219	$6.23 \times 10^{-9}$	2	
	Colon Mucinous Adenocarcinoma (CMA) (101)	22	2.324	11.956	$3.48 \times 10^{-19}$	3	(47)
	Rectal Adenocarcinoma (READ) (22)	22	3.772	10.097	$4.62 \times 10^{-11}$	3	
	Rectal Adenocarcinoma (READ) (60)	22	2.065	8.557	$3.90 \times 10^{-13}$	7	
	Rectal Adenocarcinoma (READ) (65)	65	2.068	11.751	$2.16 \times 10^{-19}$	6	(48)
	Colorectal Carcinoma (CC) (70)	12	3.508	9.208	$4.02 \times 10^{-9}$	7	(49)

Type of cancer	Cancer Subtype (n)	Normal, n	Fold change	t-test	P-value	Rank, %	(Refs.)
Esophageal	Esophageal Squamous Cell Carcinoma (ESCC) (53)	53	2.795	10.666	3.73x10 <sup>-17</sup>	2	(50)
	Esophageal Squamous Cell Carcinoma (ESCC) (17)	17	2.964	5.524	3.47x10 <sup>-6</sup>	5	(51)
	Esophageal Adenocarcinoma (EA) (14)	10	10.158	8.460	8.51x10 <sup>-5</sup>	5	(52)
Gastric	Gastric Intestinal Type Adenocarcinoma(ITGA) (62)	29	4.852	18.317	4.31x10 <sup>-32</sup>	1	(53)
	Diffuse Gastric Adenocarcinoma (DGAC) (13)	29	6.483	23.103	1.13x10 <sup>-17</sup>	1	
	Gastric mixed Adenocarcinoma (GMA) (8)	29	9.700	11.684	1.63x10 <sup>-6</sup>	1	
	Diffuse Gastric Adenocarcinoma (DGAC) (31)	19	3.287	8.446	2.38x10 <sup>-11</sup>	1	(54)
	Gastric Intestinal Type Adenocarcinoma (ITGA) (20)	19	3.038	6.251	4.01x10 <sup>-7</sup>	1	
	Gastric Cancer (GC) (15)	12	5.721	5.853	2.49x10 <sup>-6</sup>	1	(55)
	Diffuse Gastric Adenocarcinoma (DGAC) (6)	31	6.207	5.794	9.36x10 <sup>-5</sup>	2	(56)
	Gastric Mixed Adenocarcinoma (GMA) (4)	31	9.737	8.460	1.49x10 <sup>-5</sup>	3	
	Gastric Intestinal Type Adenocarcinoma (ITGA) (26)	31	4.413	7.512	2.59x10 <sup>-9</sup>	5	
	Gastric Cancer (GC) (80)	80	3.226	5.032	6.55x10 <sup>-7</sup>	2	(57)
	Salivary Gland Adenoid Cystic Carcinoma (SGACC) (16)	6	4.703	11.107	5.90x10 <sup>-10</sup>	1	(58)
	Squamous Cell Carcinoma (SCC) (34)	28	2.177	5.194	1.45x10 <sup>-6</sup>	10	(59)
	Ovarian Serous Surface Papillary Carcinoma (SSPC) (28)	4	36.947	7.443	2.62x10 <sup>-8</sup>	2	(60)
	Ovarian Carcinoma (OC) (185)	10	3.105	10.018	9.77x10 <sup>-9</sup>	7	(61)
Ovarian	Ovarian Serous Cystadenocarcinoma (OSC) (586)	8	2.706	8.141	2.09x10 <sup>-5</sup>	10	(62)
	Skin Basal Cell Carcinoma (SBCC) (15)	4	5.312	11.144	1.81 x10 <sup>-9</sup>	1	(63)
Other cancer							

Cancer samples were compared with normal tissues. CNS, central nervous system; TCGA, The Cancer Genome Atlas; IBLCA Infiltrating Bladder Urothelial Carcinoma.



Table II. Overall survival of patients with different types of cancer with overexpressed *BGN* gene.

Cancer types	RNA-seq P-value (database)	gene chip P-value (database)
Breast cancer	0.2556 <sup>a</sup>	0.7210 <sup>a</sup>
Esophageal adenocarcinoma	0.0965 <sup>a</sup>	N/A
Lung cancer	0.3010 <sup>a</sup>	0.0002 <sup>a</sup>
Gastric cancer	0.0068 <sup>a</sup>	1.3x10 <sup>-10a</sup>
Ovarian cancer	0.0093 <sup>a</sup>	0.0004 <sup>a</sup>
Bladder cancer	0.0025 <sup>a</sup>	N/A

Calculated using <sup>a</sup>Kaplan-Meier Plotter databases with RNA-seq data and gene chip data. *BGN*, biglycan; N/A, not applicable.

Table III. Overall survival of patients with different subtypes of cancer with overexpressed *BGN* gene.

Cancer subtype	RNA-seq P-value (database)	gene chip P-value (database)
Lung squamous cell carcinoma	0.0111 <sup>a</sup>	N/A
Ovarian serous cystadenocarcinoma	0.0290 <sup>c</sup>	N/A
Breast invasive carcinoma	0.9400 <sup>c</sup>	N/A
Colon adenocarcinoma	0.0963 <sup>b</sup>	N/A
Rectum adenocarcinoma	0.1776 <sup>a</sup>	N/A
Esophageal Squamous Cell Carcinoma	0.2707 <sup>a</sup>	N/A

Calculated using <sup>a</sup>Kaplan-Meier Plotter databases, <sup>b</sup>OncoLnc and <sup>c</sup>UALCAN with RNA-seq data. *BGN*, biglycan; N/A, not applicable.

OC [HR=1.28 (1.11-1.47), P=4.4x10<sup>-4</sup> in microarray analysis; HR=1.45 (1.09-1.93), P=9.3x10<sup>-3</sup> in RNA-seq analysis].

In summary, high *BGN* mRNA expression in gastric cancer and ovarian cancer was significantly associated with poor overall survival. High *BGN* mRNA expression was indicated to be associated with poor clinical outcome in the prognosis of patients with bladder cancer, lung squamous cell carcinoma, and ovarian serous cystadenocarcinoma. However, the association between *BGN* mRNA upregulation and prognosis in patients with other types and subtypes of cancer requires further examination.

**PPI network construction.** The GEPIA database was used to download the top 50 co-expressed genes, then the PPI was generated (Fig. 3). In the network, *BGN* directly interacted with 42 neighboring genes, including *ANTXR1*, *AEBP1*, *CDH11*, *CTHRC1*, *EFEMP2*, *FAP*, *LEPRE1*, *LRR15*, *LUM*, *MMP14*, *MRC2*, *MXRA5*, *OLFML2B*, *PCOLCE*, *PDGFRB*, *PXDN*, *SERPINH1*, *SFRP2*, *SPARC*, *SULF1*, *TGFβ3*, *THBS2*, *THY1*, genes of the disintegrin and metalloproteinase gene




Figure 2. OS curves of patients with different types of cancer divided by *BGN* expression. High expression levels of the *BGN* gene exhibited a significant association with lower OS time in patients with (A) gastric cancer (gene chip data), (B) gastric cancer (RNA-seq data), (C) ovarian cancer (gene chip data), (D) ovarian cancer (RNA-seq data). The plots were generated using the KM plotter database. The red lines indicate patients with *BGN* gene expression above the median value, and the black lines indicate patients with *BGN* gene expression below the median value. OS, overall survival; *BGN*, biglycan; HR, hazard ratio.




Figure 3. PPI network for *BGN* and its potential interacting proteins. Genes that were co-expressed in cancer together with *BGN* were extracted from the GEPIA database, and then a PPI network was established using the STRING database and visualized using the Cytoscape software. *TGFβ3* was highlighted in yellow because it may play an important role to regulate the expression of *BGN*. PPI, protein-protein interactions; *BGN*, biglycan; *TGFβ3*, transforming growth factor β3.

family, collagen family genes, the integrin subunit gene family and lastly the lysyl oxidase-like gene family, which jointly regulate the occurrence and development of human tumors. Jointly genes with similar expression patterns are likely to have related functions (64). For example, enrichment indicates that *BGN*-coexpressed genes are at least partially biologically connected in developing multiple cancers (65). The basic interaction between the neighboring genes is the 'functional association'. The two proteins that both contribute jointly to a specific biological function can interact specifically without touching at all, such as when a transcription factor helps to regulate the expression and production of another protein, or when two enzymes exchange a specific substrate via diffusion. The exact molecular mechanisms in cancer associated with *BGN* remain unclear (66). *BGN* upregulation has been implicated in the inflammatory response triggered by transforming growth factor  $\beta$  (TGF- $\beta$ ) (8,67,68). In the PPI network, TGF- $\beta$ 3 may play an important role to regulate the expression of *BGN*, however, the influence of TGF- $\beta$ 3 needs to be further investigated.

## Discussion

In the present study, the mRNA expression levels of *BGN* were systematically analyzed, and the results indicated that *BGN* was upregulated in various types of cancerous tissues, when compared with that in normal tissues. Previous studies have demonstrated that significantly increased levels of *BGN* are frequently detected in the clinical samples of patients with gastric (13), breast (69), colorectal (70), lung (71), ovarian (10) and pancreatic cancer (16). In addition, high expression of *BGN* in patients with solid cancer is significantly associated with poor outcome (15). Solid cancer include BLC, brain and CNS cancer, BC, cervical cancer, CC, EC, GC, HNC, kidney cancer, liver cancer, LC, melanoma, OC, pancreatic cancer, prostate cancer (72). Consistent with these studies, the present analyses demonstrated that *BGN* expression levels were increased in the majority of cancers, such as bladder, brain and central nervous system, breast, colorectal, esophageal, gastric, head and neck, lung, ovarian, and 28 subtype cancers, when compared with that in normal tissues. In addition, the current prognosis analyses revealed that high tissue *BGN* expression predicts worse survival in GC and OC. High *BGN* mRNA expression was associated with poor overall survival in patients with BLC, LSCC, and OSC. Therefore, *BGN* may be employed as either a novel prognostic biomarker or as a promising therapeutic target for human carcinomas, which is consistent with the findings of previous reports (15,73). The 43 genes with similar expression patterns are likely to have related functions in the aggressive growth and metastasis of cancers (64). In the PPI network, the genes of *AEBP1*, *MMP14*, *OLFML2B*, *PDGFRB*, *SERPINE1*, *SPARC*, *SFRP2*, *COL1A2*, *COL6A3*, *THBS2*, *COL5A2*, *COL11A1*, *FAP*, *MXRA5* and *THY1* were upregulated in solid cancer tissues, and significantly associated with the overall survival of patients with cancer (74-82). Some genes in the PPI network, including *AEBP1*, *OLFML2B*, *PDGFRB*, *SERPINE1*, *COL1A2*, *COL6A3*, and *THBS2* have been reported to be associated with metastasis, invasion and migration in cancer cells (74,76,80,83,84). The enrichment

of *BGN*-coexpressed genes indicates that the proteins are at least partially biologically connected as a group (64). However, a detailed understanding of the mechanism associated with the function of *BGN* is currently lacking; therefore, further functional studies are warranted in the future. In addition, the *BGN* protein expression levels or the signaling pathways potentially involved require further investigation. Finally, studies utilizing larger cohorts, specific cancers, or larger prospective studies also need to be conducted in order to validate the prognostic values of *BGN*.

In summary, the present study comprehensively analyzed the mRNA expression levels and prognostic value of *BGN* in the most common types of cancer, and the results indicated that *BGN* exhibited significantly high expression levels in cancer tissues compared with normal tissues in multiple types of cancer. The present findings indicated that *BGN* may serve as a promising prognostic biomarker and therapeutic target for patients with BLCA and STAD.

## Acknowledgements

Not applicable.

## Funding

No funding was received.

## Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Authors' contributions

SFZ conceived and designed the study. XJY, WJZ, LCL and ZPW made substantial contributions to the design of the current study, acquisition of data, interpretation of data and revising the manuscript. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

Not applicable.

## Patient consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

1. Chaker L, Falla A, van der Lee SJ, Muka T, Imo D, Jaspers L, Colpani V, Mendis S, Chowdhury R, Bramer WM, *et al*: The global impact of non-communicable diseases on macro-economic productivity: A systematic review. *Eur J Epidemiol* 30: 357-395, 2015.
2. Patel SA and DeMichele A: Adding adjuvant systemic treatment after neoadjuvant therapy in breast cancer: Review of the data. *Curr Oncol Rep* 19: 56, 2017.

3. Miller KD, Nogueira L, Mariotto AB, Rowland JH, Yabroff KR, Alfano CM, Jemal A, Kramer JL and Siegel RL: Cancer treatment and survivorship statistics, 2019. *CA Cancer J Clin* 69: 363-385, 2019.
4. Siegel RL, Miller KD and Jemal A: Cancer statistics, 2019. *CA Cancer J Clin* 69: 7-34, 2019.
5. Ferlay J, Shin HR, Bray F, Forman D, Mathers C and Parkin DM: Estimates of worldwide burden of cancer in 2008: *Int J Cancer* 127: 2893-2917, 2010.
6. Geerkens C, Vetter U, Just W, Fedarko NS, Fisher LW, Young MF, Termine JD, Robey PG, Wöhrle D and Vogel W: The X-chromosomal human biglycan gene BGN is subject to X inactivation but is transcribed like an X-Y homologous gene. *Hum Genet* 96: 44-52, 1995.
7. McBride OW, Fisher LW and Young MF: Localization of PGI (biglycan, BGN) and PGII (decorin, DCN, PG-40) genes on human chromosomes Xq13-qter and 12q, respectively. *Genomics* 6: 219-225, 1990.
8. Nastase MV, Young MF and Schaefer L: Biglycan: A multivalent proteoglycan providing structure and signals. *J Histochem Cytochem* 60: 963-975, 2012.
9. Fallon JR and McNally EM: Non-Glycanated biglycan and LTBP4: Leveraging the extracellular matrix for Duchenne muscular dystrophy therapeutics. *Matrix Biol* 68-69: 616-627, 2018.
10. Kocbek V, Hevir-Kene N, Bersinger NA, Mueller MD and Rižner TL: Increased levels of biglycan in endometriomas and peritoneal fluid samples from ovarian endometriosis patients. *Gynecol Endocrinol* 30: 520-524, 2014.
11. Jacobsen F, Kraft J, Schroeder C, Hube-Magg C, Kluth M, Lang DS, Simon R, Sauter G, Izbicki JR, Clauditz TS, *et al*: Up-regulation of biglycan is associated with poor prognosis and PTEN deletion in patients with prostate cancer. *Neoplasia* 19: 707-715, 2017.
12. Weber CK, Sommer G, Michl P, Fensterer H, Weimer M, Gansauge F, Leder G, Adler G and Gress TM: Biglycan is overexpressed in pancreatic cancer and induces G1-arrest in pancreatic cancer cell lines. *Gastroenterology* 121: 657-667, 2001.
13. Hu L, Duan YT, Li JF, Su LP, Yan M, Zhu ZG, Liu BY and Yang QM: Biglycan enhances gastric cancer invasion by activating FAK signaling pathway. *Oncotarget* 5: 1885-1896, 2014.
14. Xing X, Gu X, Ma T and Ye H: Biglycan up-regulated vascular endothelial growth factor (VEGF) expression and promoted angiogenesis in colon cancer. *Tumour Biol* 36: 1773-1780, 2015.
15. Wang B, Li GX, Zhang SG, Wang Q, Wen YG, Tang HM, Zhou CZ, Xing AY, Fan JW, Yan DW, *et al*: Biglycan expression correlates with aggressiveness and poor prognosis of gastric cancer. *Exp Biol Med* (Maywood) 236: 1247-1253, 2011.
16. Aprile G, Avellini C, Reni M, Mazzer M, Foltran L, Rossi D, Cereda S, Iaiza E, Fasola G and Piga A: Biglycan expression and clinical outcome in patients with pancreatic adenocarcinoma. *Tumor Biol* 34: 131-137, 2013.
17. Rhodes DR, Yu J, Shanker K, Deshpande N, Varambally R, Ghosh D, Barrette T, Pandey A and Chinnaiyan AM: ONCOMINE: A cancer microarray database and integrated data-mining platform. *Neoplasia* 6: 1-6, 2004.
18. Anaya J: OncoLnc: Linking TCGA survival data to mRNAs, miRNAs, and lncRNAs. *PeerJ Computer Sci* 2: e67, 2016.
19. Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Ponce-Rodriguez I, Chakravarthi BVSK and Varambally S: UALCAN: A portal for facilitating tumor subgroup gene expression and survival analyses. *Neoplasia* 19: 649-658, 2017.
20. Györfy B, Lanczky A, Eklund AC, Denkert C, Budczies J, Li Q and Szallasi Z: An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast Cancer Res Treat* 123: 725-731, 2010.
21. Györfy B, Lanczky A and Szállasi Z: Implementing an online tool for genome-wide validation of survival-associated biomarkers in ovarian-cancer using microarray data from 1287 patients. *Endocr Relat Cancer* 19: 197-208, 2012.
22. Györfy B, Surowiak P, Budczies J and Lanczky A: Online survival analysis software to assess the prognostic value of biomarkers using transcriptomic data in non-small-cell lung cancer. *PLoS One* 8: e82241, 2013.
23. Szász AM, Lanczky A, Nagy Á, Förster S, Hark K, Green JE, Boussiotas A, Busuttill R, Szabó A and Györfy B: Cross-validation of survival associated biomarkers in gastric cancer using transcriptomic data of 1,065 patients. *Oncotarget* 7: 49322-49333, 2016.
24. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth A, Santos A, Tsafou KP, *et al*: STRING v10: Protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 43: D447-D452, 2015.
25. Chen X, Cheung ST, So S, Fan ST, Barry C, Higgins J, Lai KM, Ji J, Dudoit S, Ng IO, *et al*: Gene expression patterns in human liver cancers. *Mol Biol Cell* 13: 1929-1939, 2002.
26. Higgins JP, Shinghal R, Gill H, Reese JH, Terris M, Cohen RJ, Fero M, Pollack JR, van de Rijn M and Brooks JD: Gene expression patterns in renal cell carcinoma assessed by complementary DNA microarray. *Am J Pathol* 162: 925-932, 2003.
27. Iacobuzio-Donahue CA, Maitra A, Olsen M, Lowe AW, van Heek NT, Rosty C, Walter K, Sato N, Parker A, Ashfaq R, *et al*: Exploration of global gene expression patterns in pancreatic adenocarcinoma using cDNA microarrays. *Am J Pathol* 162: 1151-1162, 2003.
28. Buchholz M, Braun M, Heidenblut A, Kestler HA, Klöppel G, Schmiegel W, Hahn SA, Lüttges J and Gress TM: Transcriptome analysis of microdissected pancreatic intraepithelial neoplastic lesions. *Oncogene* 24: 6626-6636, 2005.
29. Segara D, Biankin AV, Kench JG, Langusch CC, Dawson AC, Skalicky DA, Gotley DC, Coleman MJ, Sutherland RL and Henshall SM: Expression of HOXB2, a retinoic acid signaling target in pancreatic cancer and pancreatic intraepithelial neoplasia. *Clin Cancer Res* 11: 3587-3596, 2005.
30. Badea L, Herlea V, Dima SO, Dumitrascu T and Popescu I: Combined gene expression analysis of whole-tissue and microdissected pancreatic ductal adenocarcinoma identifies genes specifically overexpressed in tumor epithelia. *Hepatogastroenterology* 55: 2016-2027, 2008.
31. Mas VR, Maluf DG, Archer KJ, Yanek K, Kong X, Kulik L, Freise CE, Olthoff KM, Ghobrial RM, McIver P and Fisher R: Genes involved in viral carcinogenesis and tumor initiation in hepatitis C virus-induced hepatocellular carcinoma. *Mol Med* 15: 85-94, 2009.
32. Yusenko MV, Kuiper RP, Boethe T, Ljungberg B, van Kessel AG and Kovacs G: High-resolution DNA copy number and gene expression analyses distinguish chromophobe renal cell carcinomas and renal oncocytomas. *BMC Cancer* 9: 152, 2009.
33. Roessler S, Jia HL, Budhu A, Forgues M, Ye QH, Lee JS, Thorgeirsson SS, Sun Z, Tang ZY, Qin LX and Wang XW: A unique metastasis gene signature enables prediction of tumor relapse in early-stage hepatocellular carcinoma patients. *Cancer Res* 70: 10202-10212, 2010.
34. Franzmann E: Carcinoma. In: *Encyclopedia of Behavioral Medicine*. Gellman MD and Turner JR (eds.) Springer New York, New York, NY, pp329-330, 2013.
35. Blaveri E, Simko JP, Korkola JE, Brewer JL, Baehner F, Mehta K, Devries S, Koppie T, Pejavar S, Carroll P and Waldman FM: Bladder cancer outcome and subtype classification by gene expression. *Clin Cancer Res* 11: 4044-4055, 2005.
36. Bredel M, Bredel C, Juric D, Harsh GR, Vogel H, Recht LD and Sikic BI: Functional network analysis reveals extended gliomagenesis pathway maps and three novel MYC-interacting genes in human gliomas. *Cancer Res* 65: 8679-8689, 2005.
37. Ma XJ, Dahiya S, Richardson E, Erlander M and Sgroi DC: Gene expression profiling of the tumor microenvironment during breast cancer progression. *Breast Cancer Res* 11: R7, 2009.
38. Curtis C, Shah SP, Chin SF, Turashvili G, Rueda OM, Dunning MJ, Speed D, Lynch AG, Samarajiwa S, Yuan Y, *et al*: The genomic and transcriptomic architecture of 2,000 breast tumours reveals novel subgroups. *Nature* 486: 346-352, 2012.
39. Karnoub AE, Dash AB, Vo AP, Sullivan A, Brooks MW, Bell GW, Richardson AL, Polyak K, Tubo R and Weinberg RA: Mesenchymal stem cells within tumour stroma promote breast cancer metastasis. *Nature* 449: 557-563, 2007.
40. Perou CM, Sørlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, Pollack JR, Ross DT, Johnsen H, Akslen LA, *et al*: Molecular portraits of human breast tumours. *Nature* 406: 747-752, 2000.
41. Zhao H, Langerod A, Ji Y, Nowels KW, Nesland JM, Tibshirani R, Bukholm IK, Kåresen R, Botstein D, Børresen-Dale AL and Jeffrey SS: Different gene expression patterns in invasive lobular and ductal carcinomas of the breast. *Mol Biol Cell* 15: 2523-2536, 2004.



42. Cancer Genome Atlas Network: Comprehensive molecular portraits of human breast tumours. *Nature* 490: 61-70, 2012.
43. Ciriello G, Gatza ML, Beck AH, Wilkerson MD, Rhie SK, Pastore A, Zhang H, McLellan M, Yau C, Kandoth C, *et al*: Comprehensive molecular portraits of invasive lobular breast cancer. *Cell* 163: 506-519, 2015.
44. Kaiser S, Park YK, Franklin JL, Halberg RB, Yu M, Jessen WJ, Freudenberg J, Chen X, Haigis K, Jegga AG, *et al*: Transcriptional recapitulation and subversion of embryonic colon development by mouse colon tumor models and human colon cancer. *Genome Biol* 8: R131, 2007.
45. Skrzypczak M, Goryca K, Rubel T, Paziewska A, Mikula M, Jarosz D, Pachlewski J, Oledzki J and Ostrowski J: Modeling oncogenic signaling in colon tumors by multidirectional analyses of microarray data directed for maximization of analytical reliability. *PLoS One* 5: pii: e13091, 2010.
46. Graudens E, Boulanger V, Mollard C, Mariage-Samson R, Barlet X, Grémy G, Couillault C, Lajémi M, Piatier-Tonneau D, Zaborski P, *et al*: Deciphering cellular states of innate tumor drug responses. *Genome Biol* 7: R19, 2006.
47. Cancer Genome Atlas Network; Muzny DM, Bainbridge MN, Chang K, Dinh HH, Drummond JA, Fowler G, Kovar CL, Lewis LR, Morgan MB, *et al*: Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 487: 330-337, 2012.
48. Gaedcke J, Grade M, Jung K, Camps J, Jo P, Emons G, Gehoff A, Sax U, Schirmer M, Becker H, *et al*: Mutated KRAS results in overexpression of DUSP4, a MAP-kinase phosphatase, and SMYD3, a histone methyltransferase, in rectal carcinomas. *Genes Chromosomes Cancer* 49: 1024-1034, 2010.
49. Hong Y, Downey T, Eu KW, Koh PK and Cheah PY: A 'metastasis-prone' signature for early-stage mismatch-repair proficient sporadic colorectal cancer patients and its implications for possible therapeutics. *Clin Exp Metastasis* 27: 83-90, 2010.
50. Su H, Hu N, Yang HH, Wang C, Takikita M, Wang QH, Giffen C, Clifford R, Hewitt SM, Shou JZ, *et al*: Global gene expression profiling and validation in esophageal squamous cell carcinoma and its association with clinical phenotypes. *Clin Cancer Res* 17: 2955-2966, 2011.
51. Hu N, Clifford RJ, Yang HH, Wang C, Goldstein AM, Ding T, Taylor PR and Lee MP: Genome wide analysis of DNA copy number neutral loss of heterozygosity (CNNLOH) and its relation to gene expression in esophageal squamous cell carcinoma. *BMC Genomics* 11: 576, 2010.
52. Hao Y, Triadafilopoulos G, Sahbaie P, Young HS, Omary MB and Lowe AW: Gene expression profiling reveals stromal genes expressed in common between Barrett's esophagus and adenocarcinoma. *Gastroenterology* 131: 925-933, 2006.
53. Chen X, Leung SY, Yuen ST, Chu KM, Ji J, Li R, Chan AS, Law S, Troyanskaya OG, Wong J, *et al*: Variation in gene expression patterns in human gastric cancers. *Mol Biol Cell* 14: 3208-3215, 2003.
54. Cho JY, Lim JY, Cheong JH, Park YY, Yoon SL, Kim SM, Kim SB, Kim H, Hong SW, Park YN, *et al*: Gene expression signature-based prognostic risk score in gastric cancer. *Clin Cancer Res* 17: 1850-1857, 2011.
55. Wang Q, Wen YG, Li DP, Xia J, Zhou CZ, Yan DW, Tang HM and Peng ZH: Upregulated INHBA expression is associated with poor survival in gastric cancer. *Med Oncol* 29: 77-83, 2012.
56. D'Errico M, de Rinaldis E, Blasi MF, Viti V, Falchetti M, Calcagnile A, Sera F, Saieva C, Ottini L, Palli D, *et al*: Genome-wide expression profile of sporadic gastric cancers with microsatellite instability. *Eur J Cancer* 45: 461-469, 2009.
57. Cui J, Chen Y, Chou WC, Sun L, Chen L, Suo J, Ni Z, Zhang M, Kong X, Hoffman LL, *et al*: An integrated transcriptomic and computational analysis for biomarker identification in gastric cancer. *Nucleic Acids Res* 39: 1197-1207, 2011.
58. Frierson HF Jr, El-Naggar AK, Welsh JB, Sapinoso LM, Su AI, Cheng J, Saku T, Moskaluk CA and Hampton GM: Large scale molecular analysis identifies genes with altered expression in salivary adenoid cystic carcinoma. *Am J Pathol* 161: 1315-1323, 2002.
59. Talbot SG, Estilo C, Maghami E, Sarkaria IS, Pham DK, O-Charoenrat P, Socci ND, Ngai I, Carlson D, Ghossein R, *et al*: Gene expression profiling allows distinction between primary and metastatic squamous cell carcinomas in the lung. *Cancer Res* 65: 3063-3071, 2005.
60. Welsh JB, Zarrinkar PP, Sapinoso LM, Kern SG, Behling CA, Monk BJ, Lockhart DJ, Burger RA and Hampton GM: Analysis of gene expression profiles in normal and neoplastic ovarian tissue samples identifies candidate molecular markers of epithelial ovarian cancer. *Proc Natl Acad Sci USA* 98: 1176-1181, 2001.
61. Bonome T, Levine DA, Shih J, Randonovich M, Pise-Masison CA, Bogomolnii F, Ozbun L, Brady J, Barrett JC, Boyd J and Birrer MJ: A gene signature predicting for survival in suboptimally debulked patients with ovarian cancer. *Cancer Res* 68: 5478-5486, 2008.
62. Cancer Genome Atlas Research Network: Integrated genomic analyses of ovarian carcinoma. *Nature* 474: 609-615, 2011.
63. Riker AI, Enkemann SA, Fodstad O, Liu S, Ren S, Morris C, Xi Y, Howell P, Metge B, Samant RS, *et al*: The gene expression profiles of primary and metastatic melanoma yields a transition point of tumor progression and metastasis. *BMC Med Genomics* 1: 13, 2008.
64. Tang Z, Li C, Kang B, Gao G, Li C and Zhang Z: GEPIA: A web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res* 45: W98-W102, 2017.
65. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva NT, Morris JH, Bork P, *et al*: STRING v11: Protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* 47: D607-D613, 2019.
66. Appunni S, Anand V, Khandelwal M, Gupta N, Rubens M and Sharma A: Small leucine rich proteoglycans (decorin, biglycan and lumican) in cancer. *Clin Chim Acta* 491: 1-7, 2019.
67. Schaefer L, Babelova A, Kiss E, Hausser HJ, Baliova M, Krzyzankova M, Marsche G, Young MF, Mihalik D, Götte M, *et al*: The matrix component biglycan is proinflammatory and signals through Toll-like receptors 4 and 2 in macrophages. *J Clin Invest* 115: 2223-2233, 2005.
68. Hildebrand A, Romaris M, Rasmussen LM, Heinegård D, Twardzik DR, Border WA and Ruoslahti E: Interaction of the small interstitial proteoglycans biglycan, decorin and fibromodulin with transforming growth factor beta. *Biochem J* 302: 527-534, 1994.
69. Bischof AG, Yüksel D, Mammoto T, Mammoto A, Krause S and Ingber DE: Breast cancer normalization induced by embryonic mesenchyme is mediated by extracellular matrix biglycan. *Integr Biol (Camb)* 5: 1045-1056, 2013.
70. Liu B, Xu T, Xu X, Cui Y and Xing X: Biglycan promotes the chemotherapy resistance of colon cancer by activating NF-kappaB signal transduction. *Mol Cell Biochem* 449: 285-294, 2018.
71. Yang Z, Wen Y, Qian L and Hospital Z: Up-regulation of biglycan is associated with malignant phenotype of nonsmall cell lung cancer. *J Med Res* 45, 41-46, 2016.
72. Ferlay J, Colombet M, Soerjomataram I, Mathers C, Parkin DM, Piñeros M, Znaor A and Bray F: Estimating the global cancer incidence and mortality in 2018: GLOBOCAN sources and methods. *Int J Cancer* 144: 1941-1953, 2019.
73. Appunni S, Anand V, Khandelwal M, Seth A, Mathur S and Sharma A: Altered expression of small leucine-rich proteoglycans (Decorin, Biglycan and Lumican): Plausible diagnostic marker in urothelial carcinoma of bladder. *Tumour Biol* 39: 1010428317699112, 2017.
74. Liu JY, Jiang L, Liu JJ, He T, Cui YH, Qian F and Yu PW: AEBP1 promotes epithelial-mesenchymal transition of gastric cancer cells by activating the NF-κB pathway and predicts poor outcome of the patients. *Sci Rep* 8: 11955, 2018.
75. Kasurinen A, Gramolelli S, Hagström J, Laitinen A, Kokkola A, Miki Y, Lehti K, Yashiro M, Ojala PM, Böckelman C and Haglund C: High tissue MMP14 expression predicts worse survival in gastric cancer, particularly with a low PROX1. *Cancer Med* 8: 6995-7005, 2019.
76. Liu J, Liu Z, Zhang X, Gong T and Yao D: Bioinformatic exploration of OLFML2B overexpression in gastric cancer base on multiple analyzing tools. *BMC Cancer* 19: 227, 2019.
77. Wang G, Shi B, Fu Y, Zhao S, Qu K, Guo Q, Li K and She J: Hypomethylated gene NRPI is co-expressed with PDGFRB and associated with poor overall survival in gastric cancer patients. *Biomed Pharmacother* 111: 1334-1341, 2019.
78. Liao P, Li W, Liu R, Teer JK, Xu B, Zhang W, Li X, Mcleod HL and He Y: Genome-scale analysis identifies SERPINE1 and SPARC as diagnostic and prognostic biomarkers in gastric cancer. *Onco Targets Ther* 11: 6969-6980, 2018.

79. Wang H, Duan XL, Qi XL, Meng L, Xu YS, Wu T and Dai PG: Concurrent Hypermethylation of SFRP2 and DKK2 Activates the Wnt/ $\beta$ -catenin pathway and is associated with poor prognosis in patients with gastric cancer. *Mol Cells* 40: 45-53, 2017.
80. Ao R, Guan L, Wang Y and Wang JN: Silencing of COL1A2, COL6A3, and THBS2 inhibits gastric cancer cell proliferation, migration, and invasion while promoting apoptosis through the PI3k-Akt signaling pathway. *J Cell Biochem* 119: 4420-4434, 2018.
81. Zhuo C, Li X, Zhuang H, Tian S, Cui H, Jiang R, Liu C, Tao R and Lin X: Elevated THBS2, COL1A2, and SPP1 expression levels as predictors of gastric cancer prognosis. *Cell Physiol Biochem* 40: 1316-1324, 2016.
82. Hao S, Lv J, Yang Q, Wang A, Li Z, Guo Y and Zhang G: Identification of key genes and circular RNAs in human gastric cancer. *Med Sci Monit* 25: 2488-2504, 2019.
83. Suzuki S, Dobashi Y, Hatakeyama Y, Tajiri R, Fujimura T, Heldin CH and Ooi A: Clinicopathological significance of platelet-derived growth factor (PDGF)-B and vascular endothelial growth factor-A expression, PDGF receptor- $\beta$  phosphorylation, and microvessel density in gastric cancer. *BMC Cancer* 10: 659, 2010.
84. Mazzocchi G, Paziienza V, Panza A, Valvano MR, Benegiamo G, Vinciguerra M, Andriulli A and Piepoli A: ARNTL2 and SERPINE1: Potential biomarkers for tumor aggressiveness in colorectal cancer. *J Cancer Res Clin Oncol* 138: 501-511, 2012.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.