

Ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate exhibits anti-proliferative activity and induces apoptosis in promyelocytic leukemia HL-60 cells

AN-CHENG HUANG^{1*}, CHEN-SHENG LIN^{2*}, JIN-CHERNG LIEN^{3*},
HSUEH-CHOU LAI⁴, WEI-HUA LIN⁵ and CHENG-WEN LIN⁵⁻⁷

¹Department of Nursing, St. Mary's Junior College of Medicine, Nursing and Management, Yilan 26647;

²Division of Gastroenterology, Kuang Tien General Hospital, Taichung 43303; ³School of Pharmacy, China Medical University, Taichung 40402; ⁴Division of Hepato-Gastroenterology, Department of Internal Medicine, China Medical University Hospital, Taichung 40447; ⁵Department of Medical Laboratory Science and Biotechnology, China Medical University, Taichung 40402; ⁶Department of Biotechnology, Asia University, Wufeng, Taichung 41354; ⁷Chinese Medicine Research Center, China Medical University, Taichung 40402, Taiwan, R.O.C.

Received August 1, 2019; Accepted December 5, 2019

DOI: 10.3892/ol.2020.11342

Abstract. Furoquinolone and its derivatives exhibit anti-microbial, anti-allergic, anti-inflammatory and anticancer properties. The present study investigated the anti-tumor activity of synthesized intermediates of furoquinolone in human promyelocytic leukemia HL-60 cells. The biological effects of the active compound ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (compound 131) were examined in HL-60 cells. The following properties were analyzed: Cell survival, cell cycle profile, caspase-3 activity, Bax and Bcl-2 expression, the amount of intracellular Ca²⁺, the number of reactive oxygen species (ROS) and the mitochondrial membrane potential. Compound 131 (50% cytotoxic concentration, 23.5 μM) significantly reduced the proliferation of HL-60 cells and was revealed to induce apoptosis in HL-60 cells in a concentration-dependent manner. Moreover, this was associated with the activation of caspase-3, upregulation of Bax, an increase in intracellular Ca²⁺ and ROS production, and a decrease in mitochondrial membrane potential and Bcl-2 expression levels. Compound 131, a novel 4,5-dihydrofuran-3-carboxylate, induced apoptosis in HL-60 cells via the increase of intracellular Ca²⁺ and ROS to alter the mitochondrial membrane potential and the protein level of Bax and Bcl-2, as well as activating caspase-3. The results of

the current study indicate that compound 131 may represent a promising compound for the development of anti-leukemia therapeutics.

Introduction

Leukemia is a malignancy affecting leukocytes and can be subdivided into myeloid and lymphocytic leukemia. The mortality rate of chronic myeloid leukemia has reduced since imatinib methylate was approved as the treatment. However, total acute myeloid leukemia-associated mortalities have continued to rise over the past 20 years (1). Numerous furoquinoline alkaloids have been identified and studied, including dictamnine, confusameline, skimmianine and kokusaginine, all of which belong to the Rutaceae family (2). The majority of furoquinoline alkaloids influence multiple biological processes, such as antifungal activity (3), antiplatelet aggregation (4) and Ca²⁺ influx suppression (5). Acrophylline and acrophyllidine have structures that are similar to furoquinolone, and can be isolated from *Acronychia*, *Dictamnus*, *Ptelea*, *Glycosmis* and *Ruta* plants (6), which exhibit anti-microbial and anticancer activities (7,8). Acrophyllidine and its synthetic derivatives also exhibit significant anti-allergic activity via suppressing mast cell degranulation (9). Moreover, ethyl 2-(3-hydroxyanilino)-4-oxo-4,5-dihydrofuran-3-carboxylate, an intermediate in furoquinolone synthesis, exhibits anti-inflammatory activity (10) and ethyl 2-[N-p-chlorobenzyl-(2'-methyl)] anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (JOT01007) has also been revealed to induce apoptosis in mouse leukemia (WEHI-3) and human cervical cancer (CaSki) cell lines (11,12). Notably, JOT01006 activates BCL2 antagonist/killer 1, poly(ADP-ribose) polymerase 1 and caspase-3, resulting in apoptosis and inhibiting the migration of human cervical cancer HeLa cells (13). However, anticancer activity of the intermediates in furoquinolone synthesis is rarely reported in treating acute myeloid leukemia.

Correspondence to: Professor Cheng-Wen Lin, Department of Medical Laboratory Science and Biotechnology, China Medical University, 91 Hsueh-Shih Road, Taichung 40402, Taiwan, R.O.C.
E-mail: cwlin@mail.cmu.edu.tw

*Contributed equally

Key words: leukemia cell, anti-proliferation, apoptosis, intracellular Ca²⁺, reactive oxygen species

The present study aimed to characterize the anti-proliferative and apoptotic activity of ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (compound 131) in acute promyelocytic leukemia HL-60 cells. The current results indicate that compound 131 induces apoptosis in HL-60 cells, and this was associated with increased intracellular Ca^{2+} , increased reactive oxygen species (ROS), activation of caspase-3 and a decrease in mitochondrial membrane potential. Hence, compound 131 may represent a novel target for treating acute promyelocytic leukemia.

Materials and methods

Cells. Human promyelocytic leukemia HL-60 and plasma cell leukemia ARH-77 cells (Bioresource Collection and Research Centre) were cultured in RPMI-1640 medium plus 10% fetal bovine serum (FBS) (Thermo Fisher Scientific, Inc.), 100 U/ml penicillin and 100 μ g/ml streptomycin at 37°C and 5% CO_2 . Vero cells (a monkey kidney epithelial cell line) (Bioresource Collection and Research Centre) were cultivated in Eagle's Minimum Essential Medium (Thermo Fisher Scientific, Inc.) containing 10% FBS, 100 U/ml penicillin and 100 μ g/ml streptomycin at 37°C, 5% CO_2 .

Reagents. Ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (compound 131) was manufactured and refined using high-performance liquid chromatography as described in a previous study (10). Briefly, a mixture of diethyl malonate (32.0 g, 0.2 mol) in 50 ml tetrahydrofuran (THF) with chloroacetyl chloride (11.3 g, 0.1 mol) in 100 ml THF was incubated at 10–12°C for 1 h followed by 40–45°C for 1 h; ethyl 2-ethoxy-4-oxo-4,5-dihydrofuran-3-carboxylate was produced post cooling. Finally, the ethoxy group in the compound was substituted with aniline after stirring at room temperature for 1 h and heating on a water bath at 80°C for 3 h to yield ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (compound 131). After the product of the reaction had been confirmed by performing via thin layer chromatography on silica gel-protected aluminum sheets (Type 60 F254; Merck KGaA) in which the spots were detected using a UV-lamp, the reaction was further mixed with 100 cc of ice water to form a precipitate; white crystals of compound 131 (18.29 g; yield, 74%; melting point, 115–116°C) were generated after the precipitate was recrystallized from 90% ethanol at room temperature °C for 1–2 days. The structure of compound 131 (Fig. 1) was confirmed via mass spectrometry (m/z) as follows: 3267.87 (-NH-), 1695.26 (C4=O), 1672.59 (C3-CO-OEt); UV λ_{max} nm (MeOH) (log ϵ): 297 (4.523); ¹H-NMR (200 MHz, $CDCl_3$) δ : 1.24 (3H, t, $J=7$ Hz, H-2''), 4.20 (2H, q, $J=7$ Hz, H-1''), 4.67 (2H, s, H-5), 7.25 (5H, m, H-2', H-3', H-4', H-5', H-6'), 10.264 (1H, s, -NH-); ¹³C-NMR (200 MHz, DMSO- d_6) δ : 14.67 (C-2''), 59.37 (C-1''), 75.30 (C-5), 86.99 (C-3), 123.48 (C-2', C-6'), 126.35 (C-4'), 129.27 (C-3', C-5'), 135.24 (C-1'), 164.18 (C-2), 177.34 (C-3''), 188.84 (C-4).

MTT assay. HL-60, ARH-77 or Vero cells (1×10^4 /well) were cultured in 96-well plates in the presence or absence of compound 131 (0, 5, 25 and 50 μ M) at 37°C for 2 days followed by the addition of 20 μ l 0.5 mg/ml MTT (Sigma-Aldrich; Merck KGaA) per well for another 4-h incubation at 37°C.

After removing the cultured media, the reduced form of MTT in the cells was solubilized using 150 μ l dimethyl sulfoxide for 15 min. The survival rate of treated cells was measured as the ratio of the optical density (OD)_(570–630 nm) of treated cells to that of mock cells, similar to the calculation of the 50% cytotoxic concentration (CC₅₀).

Cell cycle analysis and caspase-3 fluorimetric assay. A total of 2×10^5 HL-60 cells were treated with compound 131 (0, 5, 25 or 50 μ M) for 48 h at 37°C. Subsequently, cells were collected after centrifugation at 100 x g (Kubota Corporation) for 5 min at room temperature and cell cycle profile analysis was performed using propidium iodide (PI) staining and caspase-3 activity analysis using BD ApoAlert Caspase Fluorescent assay kit (cat. no. 51-6632AK and 51-6632BK; BD Biosciences) as described in our previous report (14,15). PI-stained cells were analyzed at a wavelength of 488 nm by flow cytometry (BD Biosciences). The supernatants of treated cell lysates were added to the wells of the caspase-3 assay with BD ApoAlert Caspase Fluorescent assay kit for a 2-h incubation at 37°C according to the manufacturer's protocol. Subsequently, caspase-3 activity was examined using a fluorescent substrate and a fluorescent plate reader (BioTek China) using an excitation wavelength of 380 nm and an emission wavelength of 460 nm.

Western blotting assay. The cells treated with compound 131 were harvested and washed with ice-cold PBS by centrifuging at 100 x g (Kubota Corporation) for 5 min at 4°C, and then mixed with the RIPA lysis buffer (cat. no. R0278, Sigma-Aldrich; Merck KGaA) in microfuge tubes for 30 min at 4°C. The lysate of treated cells was collected after centrifugation at 12000 x g for 20 min at 4°C, and the protein concentration was determined using coomassie protein assay reagent (cat. no. 27813, Sigma-Aldrich; Merck KGaA) at an absorbance of 280 nm. A total of 20 μ g of lysate per lane from the cells treated with compound 131 was dissolved in 2X SDS-PAGE sample buffer (Sigma-Aldrich; Merck KGaA), boiled at 95°C for 5 min, loaded in 12% SDS-PAGE gels and then analyzed using a vertical electrophoresis system (Bio-Rad Laboratories, Inc.). The separated proteins in the gels were transferred to nitrocellulose papers, and were blocked with 5% skim milk at room temperature overnight. The immune reaction on the blots was performed using a 1:2,000 dilution of anti-caspase-3 (cat. no. 9662), anti-Bax (cat. no. 2772), anti-Bcl-2 (cat. no. 3498) and anti- β -actin (cat. no. 4970) antibodies in at room temperature overnight, and 1:3,000 dilution of horseradish peroxidase-conjugated goat anti-mouse IgG antibodies (cat. no. 7076) (Cell Signaling Technology, Inc.) at room temperature for 2 h. The immune-reactive bands were developed using enhanced chemiluminescence solution (Amersham Pharmacia Biotech Ltd.), as previously described (14,15).

Cytoplasmic-free Ca^{2+} , intracellular ROS and mitochondrial membrane potential ($\Delta\Psi_m$) assays using flow cytometry. To detect intracellular calcium levels, treated cells were collected and incubated with 10 μ M Fluo-3/AM for 30 min at 37°C in the dark and analyzed at 526 nm by flow cytometry (16). In the intracellular ROS assay, cells stained with




Figure 1. Chemical structure of compound 131 as ascertained by nuclear magnetic resonance spectroscopy. Compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate.

2,7-dichlorodihydrofluorescein diacetate (DCFH-DA) were examined using flow cytometry as described in our previous study (17). To measure the $\Delta\Psi_m$, compound 131-treated cells were analyzed with DiOC₆ staining at 37°C for 30 min and examined using flow cytometry (17).

Statistical analysis. Data from three independent experiments of mock cells and compound 131-treated cells were analyzed by one-way ANOVA followed by Scheffe's post hoc test using SPSS 12.0 (SPSS, Inc.). $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Treatment with compound 131 results in cell growth inhibition and apoptosis in HL-60 cells. Initially, 23 synthesized intermediates of furoquinolone at 50 μM were screened for anti-proliferative activity against HL-60 and Vero cells; only compound 131 (ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate) exhibited a significant inhibitory effect on the proliferation of HL-60 (but not Vero cells) (Fig. S1). Subsequently, the cells were treated with compound 131 at 0, 5, 25 and 50 μM for 48 h at 37°C; the cell viability was tested using the MTT assay to determine the CC_{50} values of compound 131 on the proliferation of HL-60, ARH-77 and Vero cells. The survival rates of HL-60 and ARH-77 cells treated with compound 131 were significantly lower compared with treated Vero cells (Fig. 2). The CC_{50} values of compound 131 were 23.5 μM for HL-60, 24.2 μM for ARH-77 and 87.0 μM for Vero cells. These results demonstrated that compound 131 exhibits significant anti-proliferative effects against HL60 and ARH-77 cells.

Activation of caspase-3 in HL-60 cells following treatment with compound 131. To determine whether compound 131 initiated apoptosis, the cell cycle and caspase-3 activity of treated cells were further analyzed using flow cytometry, western blotting and caspase-3 enzymatic activity assays (Figs. 3 and 4). Cell cycle analysis indicated that compound 131 increased the percentage of cells in the sub-G₁ phase (apoptotic fraction) in a concentration-dependent manner in HL-60 cells (Fig. 3A). The percentage of apoptotic cells (sub-G₁ fraction) was 0.2, 1.6, 75.4 and 80.0% in cells treated with compound 131 at 0, 5, 25 and 50 μM , respectively (Fig. 3B). Moreover, western blot




Figure 2. Survival rates of HL-60 cells and Vero cells in response to compound 131. HL-60, Vero and ARH-77 cells were treated with compound 131 at the indicated concentrations for 48 h. The cells were analyzed using an MTT assay; absorbance values at 570 nm were recorded. Survival data exhibited represent percentages of absorbance values of treated cells with compound 131 normalized to PBS-treated cells (Control). ** $P < 0.01$; *** $P < 0.001$ vs. PBS-treated cells. Compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate.

analysis of cell lysates indicated that compound 131 treatment increased the protein levels of pro- and active forms of caspase-3, and also upregulated Bax and downregulated Bcl-2 in a concentration-dependent manner (Fig. 4A and B). Fluorescence assays of caspase-3 enzymatic activity revealed that caspase-3 enzymatic activity in compound 131-treated cells was higher compared with mock cells by 6-, 43- and 140-fold for 5, 25 and 50 μM , respectively (Fig. 4C). The current results indicate that compound 131 significantly promotes caspase-3 and Bax activation, but suppresses Bcl-2 expression, in apoptotic cells.

Treatment with compound 131 results in increases of intracellular calcium and ROS in HL-60 cells. Intracellular Ca^{2+} accumulation and ROS generation serve a critical role in apoptosis (17-20). The effects of compound 131 on intracellular Ca^{2+} and ROS levels in HL-60 cells were examined (Figs. 5 and 6). Following treatment with compound 131 (0, 5, 25 or 50 μM) at 37°C for 48 h, cells were harvested, stained with Fluo-3/AM or DCFH-DA and analyzed using flow cytometry. Compound 131 treatment resulted in the increase of intracellular Ca^{2+} release from the ER in HL-60 cells. A 39.2% increase resulted from treatment with 5 μM , a 93.8% increase from treatment with 25 μM and an 80.4% increase following treatment with 50 μM compared with PBS-treated cells, respectively (Fig. 5). In addition, compound 131 significantly increased the production of intracellular ROS compared with mock cells (Fig. 6). The results indicate that compound 131 treatment significantly stimulates an increase in intracellular Ca^{2+} and ROS levels in HL-60 cells.

Treatment with compound 131 results in a decrease of mitochondrial membrane potential in HL-60 cells. The disruption of mitochondrial membrane potential (MMP) has been reported to correlate with ROS generation (21-23). Treated




Figure 3. Cell cycle profiles and apoptotic rates of HL-60 cells treated with compound 131. Cells were treated with compound 131 at 37°C for 48 h. (A) Cells stained with propidium iodide were analyzed using flow cytometry. (B) Proportions of apoptotic HL-60 cells after compound 131 treatment are presented as a bar graph. *** $P < 0.001$ vs. PBS-treated cells. Compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate.




Figure 4. Addition of compound 131 results in caspase-3 activation and Bax upregulation in HL-60 cells. Treated cells were harvested after a 2-day incubation with compound 131 at 37°C. (A) Lysates of treated HL-60 cells were analyzed using western blotting with anti-caspase-3, anti-Bax and anti-Bcl-2 antibodies. (B) Relative level of each immuno-reactive band was quantified and normalized to the protein level of β -actin. (C) Lysates reacted with fluorescent substrates at 37°C for 2 h; then, the fluorescence intensity of each well was analyzed using a fluorescent plate reader (BioTek China) (excitation at 380 nm and emission at 460 nm). $^{\circ}P < 0.05$ *** $P < 0.001$; ## $P < 0.01$; ### $P < 0.001$; +++ $P < 0.001$ vs. PBS-treated cells. Compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate.




Figure 5. Intracellular calcium production of HL-60 cells is influenced by compound 131. Treated cells were stained using Fluo-3/AM. (A) Intracellular calcium levels in HL-60 cells were analyzed by flow cytometry (absorbance at 526 nm). (B) Relative increases in the intracellular calcium levels were quantified and are presented as a bar graph. **P<0.01; ***P<0.001 vs. PBS-treated cells. Compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate.




Figure 6. Intracellular ROS production of HL-60 cells is influenced by compound 131. (A) Cells were stained with DCFH-DA and examined by flow cytometry. (B) Increment ratio of intracellular ROS in treated cells is shown by bar graph. ***P<0.001 vs. PBS-treated cells. ROS, reactive oxygen species; compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate; DCFH-DA, 2,7-dichlorodihydrofluorescein diacetate.




Figure 7. MMP detection in compound 131-treated HL-60 cells. (A) Treated HL-60 cells were stained with DiOC₆ and analyzed by flow cytometry. (B) Percentages of the MMP decrease in treated HL-60 cells are illustrated by bar graph. ***P<0.001 vs. PBS-treated cells. Compound 131, ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate; MMP, mitochondrial membrane potential.

cells were tested to determine the MMP levels by staining with DiOC₆ and analysis using flow cytometry (Fig. 7). Cells treated with compound 131 exhibited a significant decrease in DiOC₆ intensity by 19.6% for 5 μM, 32.5% for 25 μM and 27.45% for 50 μM. Thus, it was revealed that treatment with compound 131 also resulted in a decrease in MMP in HL-60 cells.

Discussion

To the best of our knowledge, this is the first report to demonstrate that ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (compound 131), an intermediate of furoquinoline synthesis, exerts anti-leukemic effects and induces apoptosis via the upregulation of caspase-3 and Bax in HL-60 cells. These effects are also associated with an increase in intracellular calcium, an increase in ROS levels and a decrease in MMP expression. The mechanism of compound 131-induced apoptosis was similar to previous reports of apoptosis induced by 4,5-dihydrofuran-3-carboxylate derivatives (11,12). In addition, a structure-function association study demonstrated that ethyl 2-(3-methoxyanilino)-4-oxo-4,5-dihydrofuran-3-carboxylate and ethyl-2-(3-oxyanilino)-4-oxo-4,5-dihydrofuran-3-carboxylate exhibited lower anti-proliferative properties than compound 131 (ethyl 2-anilino-4-oxo-4,5-dihydrofuran-3-carboxylate) (data not shown). The aforementioned result indicates that the aniline group in compound 131 serves a key role in its anti-proliferative activity against HL60 cells. Ethyl 2-[N-p-chlorobenzyl-(2'-methyl)] anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (JOT01007)

triggered the mitochondria-dependent pathway, which was significantly correlated with cytoplasmic Ca²⁺ levels in human cervical CaSki cancer cell apoptosis (11). Moreover, ethyl 2-[N-m-chlorobenzyl-(2'-methyl)] anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (JOT01006) induced the increase in ROS production that resulted in the caspase-dependent apoptosis of human cervical cancer cells (13). In addition to anticancer activity, the intermediates of furoquinoline synthesis and furoquinoline derivatives have been revealed to exhibit diverse pharmacological effects, such as 5-HT₂ receptor antagonist activity (24), vasorelaxation via the suppression of calcium influx (25), anti-allergic effects (9) and the blocking of outward K⁺ current and Na⁺ channels (5). A previous study revealed that the intermediates of furoquinoline synthesis exhibited moderate inhibitory effects on the growth of human ovarian cancer A2780 cells (26).

In summary, the present results indicated that compound 131 inhibits the proliferation and induces apoptosis in HL-60 cells; moreover, it was associated with the production of intracellular Ca²⁺ and ROS, and reduced the mitochondrial membrane potential. Compound 131, a novel 4,5-dihydrofuran-3-carboxylate, represents a promising compound that may inform the development of new anti-leukemia agents. The current results also revealed the biological properties of compound 131 and indicated the mechanisms underlying its anti-leukemia effects.

Acknowledgements

Not applicable.

Funding

Funding was received from the China Medical University (Taichung, Taiwan) under the Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education (grant nos. CHM106-6-2 and CMRC-CHM-2). The present study was also sponsored by grants from the Ministry of Science and Technology, Taiwan (grant nos. MOST107-2923-B-039-001-MY3 and MOST108-2320-B-039-039-MY3) and China Medical University (grant nos. CMU106-BC-1, CMU106-ASIA-06, CMU107-S-14 and CMU107-ASIA-12).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Authors' contributions

ACH, CSL, JCL, HCL, WHL and CWL designed and conducted the experiments. ACH, CSL, and JCL performed data analysis. ACH and CWL wrote the first draft of the manuscript. All authors reviewed and approved the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Hao T, Li-Talley M, Buck A and Chen W: An emerging trend of rapid increase of leukemia but not all cancers in the aging population in the United States. *Sci Rep* 9: 12070, 2019.
- Tarus PK, Coombes PH, Crouch NR, Mulholland DA and Moodley B: Furoquinoline alkaloids from the southern African Rutaceae *Tectlea natalensis*. *Phytochemistry* 66: 703-706, 2005.
- Zhao W, Wolfender JL, Hostettmann K, Xu R and Qin G: Antifungal alkaloids and limonoid derivatives from *Dictamnus dasycarpus*. *Phytochemistry* 47: 7-11, 1998.
- Chen KS, Chang YL, Teng CM, Chen CF and Wu YC: Furoquinolines with antiplatelet aggregation activity from leaves of *Melicope confusa*. *Planta Med* 66: 80-81, 2000.
- Su MJ, Chang GJ, Wu MH and Kuo SC: Electrophysiological basis for the antiarrhythmic action and positive inotropy of HA-7, a furoquinoline alkaloid derivative, in rat heart. *Br J Pharmacol* 122: 1285-1298, 1997.
- Lahey FN and McCamish M: Acrophylline and acrophyllidine. Two new alkaloids from *Acronychia haplophylla*. *Tetrahedron Lett* 12: 1525-1527, 1968.
- Zhao YL, Chen YL, Sheu JY, Chen IL, Wang TC and Tzeng CC: Synthesis and antimycobacterial evaluation of certain fluoroquinolone derivatives. *Bioorg Med Chem* 13: 3921-3926, 2005.
- Sheu JY, Chen YL, Tzeng CC, Hsu SL, Fang KC and Wang TC: Synthesis, and antimycobacterial and cytotoxic evaluation of certain fluoroquinolone derivatives. *Helv Chim Acta* 86: 2481-2489, 2003.
- Huang AC, Lin TP, Kuo SC and Wang JP: The anti-allergic activities of synthetic acrophylline and acrophyllidine. *J Nat Prod* 58: 117-120, 1995.
- Wang JP, Tsao LT, Raung SL, Hsu MF and Kuo SC: Inhibition by HAJ11 of respiratory burst in neutrophils and the involvement of protein tyrosine phosphorylation and phospholipase D activation. *Br J Pharmacol* 120: 79-87, 1997.
- Huang AC, Chung JG, Kuo SC, Lu HF and Lin TP: Synthesis and cytotoxic activity of certain 2,3,4,9-tetrahydrofuro [2,3-b] quinolin-3,4-dione and ethyl 2-(substituted aniline)-4-oxo-4,5-dihydrofuran-3-carboxylate derivatives in murine leukemia WEHI-3 cells. *In Vivo* 21: 227-236, 2007.
- Lin TP, Huang AC, Wei HC, Lin JG and Chung JG: Ethyl 2-[N-p-chlorobenzyl-(2'-methyl)] anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (JOT01007) induces apoptosis in human cervical cancer Ca Ski cells. *In Vivo* 21: 397-406, 2007.
- Huang AC, Lin TP, Weng YS, Ho YT, Lin HJ, Huang LJ, Kuo SC and Chung JG: Ethyl 2-[N-m-chlorobenzyl-(2'-methyl)] anilino-4-oxo-4,5-dihydrofuran-3-carboxylate (JOT01006) induces apoptosis in human cervical cancer HeLa cells. *Anticancer Res* 27: 2505-2514, 2007.
- Yang TC, Shiu SL, Chuang PH, Lin YJ, Wan L, Lan YC and Lin CW: Japanese encephalitis virus NS2B-NS3 protease induces caspase 3 activation and mitochondria-mediated apoptosis in human medulloblastoma cells. *Virus Res* 143: 77-85, 2009.
- Yang TC, Lai CC, Shiu SL, Chuang PH, Tzou BC, Lin YY, Tsai FJ and Lin CW: Japanese encephalitis virus down-regulates thioredoxin and induces ROS-mediated ASK1-ERK/p38 MAPK activation in human promonocyte cells. *Microbes Infect* 12: 643-651, 2010.
- Viarengo A, Cenesi L, Moore MN and Orunesu M: Effects of Hg²⁺ and Cu²⁺ on the cytosolic Ca²⁺ level in molluscan blood cells evaluated by confocal microscopy and spectrofluorimetry. *Marine Biol* 119: 557-564, 1994.
- Lin SS, Huang HP, Yang JS, Wu JY, Hsai TC, Lin CC, Lin CW, Kuo CL, Gibson Wood W and Chung JG: DNA damage and endoplasmic reticulum stress mediated curcumin-induced cell cycle arrest and apoptosis in human lung carcinoma A-549 cells through the activation caspases cascade- and mitochondrial-dependent pathway. *Cancer Lett* 272: 77-90, 2008.
- Mattson MP and Chan SL: Neuronal and glial calcium signaling in Alzheimer's disease. *Cell Calcium* 34: 385-397, 2003.
- Benali-Furet NL, Chami M, Houel L, De Giorgi F, Vernejoul F, Lagorce D, Buscaïl L, Bartenschlager R, Ichas F, Rizzuto R and Paterlini-Bréchet P: Hepatitis C virus core triggers apoptosis in liver cells by inducing ER stress and ER calcium depletion. *Oncogene* 24: 4921-4933, 2005.
- Haynes CM, Titus EA and Cooper AA: Degradation of misfolded proteins prevents ER-derived oxidative stress and cell death. *Mol Cell* 15: 767-776, 2004.
- Xue X, Piao JH, Nakajima A, Sakon-Komazawa S, Kojima Y, Mori K, Yagita H, Okumura K, Harding H and Nakano H: Tumor necrosis factor alpha (TNFalpha) induces the unfolded protein response (UPR) in a reactive oxygen species (ROS)-dependent fashion, and the UPR counteracts ROS accumulation by TNFalpha. *J Biol Chem* 280: 33917-33925, 2005.
- Kadenbach B: Intrinsic and extrinsic uncoupling of oxidative phosphorylation. *Biochim Biophys Acta* 1604: 77-94, 2003.
- Rego AC and Oliveira CR: Mitochondrial dysfunction and reactive oxygen species in excitotoxicity and apoptosis: Implications for the pathogenesis of neurodegenerative diseases. *Neurochem Res* 28: 1563-1574, 2003.
- Cheng JT, Chang TK and Chen IS: Skimmianine and related furoquinolines function as antagonists of 5-hydroxytryptamine receptors in animals. *J Auton Pharmacol* 14: 365-374, 1994.
- Yu SM, Ko FN, Su MJ, Wu TS, Wang ML, Huang TF and Teng CM: Vasorelaxing effect in rat thoracic aorta caused by fraxinellone and dictamine isolated from the Chinese herb *Dictamnus dasycarpus* Turcz: Comparison with cromakalim and Ca²⁺ channel blockers. *Naunyn Schmiedeberg's Arch Pharmacol* 345: 349-355, 1992.
- Cao S, Al-Rehaily AJ, Brodie P, Wisse JH, Moniz E, Malone S and Kingston DG: Furquinoline alkaloids of *Ertela (Monnieria) trifolia* (L.) Kuntze from the Suriname rainforest. *Phytochemistry* 69: 553-557, 2008.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.