

Wide-ranging analysis of survival-related alternative splicing events in invasive breast carcinoma

KEREN JIA¹, YINGCHENG WU¹, JING HUANG², JIANING CHEN¹, HUAGEN WEI¹ and HUIQUN WU³

¹School of Medicine, Nantong University, Nantong, Jiangsu 226001; ²School of Pharmacy, Nanjing University of Chinese Medicine, Nanjing, Jiangsu 210023; ³Department of Medical Informatics, Nantong University, Nantong, Jiangsu 226001, P.R. China

Received August 2, 2019; Accepted April 27, 2020

DOI: 10.3892/ol.2020.11695

Abstract. Invasive breast carcinoma (BRCA) is a serious disease that threatens the survival time of those affected. Alternative splicing (AS) involved in BRCA pathogenesis may be a potential therapeutic target. However, to the best of our knowledge, a systematic analysis of survival-related alternative splicing events (SREs) has not yet been reported. The aim of the present study was to identify SREs and analyze their potential biological functions as BRCA prognostic biomarkers. An UpSet plot demonstrated AS global characteristics. Cox's proportional hazards regression model quantitatively demonstrated the prognostic relevance of AS events. Functional enrichment analysis investigated the potential pathways through which AS events affect BRCA progression. The receiver operating characteristic curve model determined the clinical significance of AS events represented using percent-spliced-in (PSI) values. The regulatory network of splicing factors (SFs) and AS events laid the foundation for studying the role of SFs in BRCA. The present study identified 1,215 SREs and their distribution characteristics, suggesting that AS events in exon skipping (ES) primarily exerted normal physiological functions, while AS events in alternative terminator sites had the most significant prognostic effect. The present study demonstrated that survival-associated genes are involved primarily in certain biological processes of ribosomal proteins. In the diagnostic model, the alternative acceptor site, alternative donor site, alternative promoter site and ES performed well. ELAVL4 was the key gene associated with prognosis and SREs. In conclusion, a number of AS events affect BRCA initiation, progression and prognosis. The PSI value of AS events has the potential to diagnose BRCA and predict a prognosis; however, this must be confirmed in additional studies.

Introduction

Breast cancer is a global problem that primarily threatens the health of women, but can also affect men. In 2011, a study based on patients with breast cancer in Denmark, Finland, Geneva, Norway, Singapore and Sweden over the past 40 years reported that world standardized incidence rates of female breast cancer were 66.7 per 10⁵ individuals per year and those of male breast cancer were 0.40 per 10⁵ individuals per year (1). Epidemiological studies have demonstrated that breast cancer is closely associated with social factors, particularly quality of healthcare and ethnicity (2,3). In the United States, the mortality rate of non-Hispanic black women with breast cancer was 8.8% higher compared with that of non-Hispanic white women with breast cancer between 2010 and 2014 (4). The incidence rates of breast cancer have increased from about 100 per 10⁵ women in 1975 to about 125 per 10⁵ women in 2015 (5). As the breast is not an essential organ, cancer *in situ* does not usually lead directly to death. With invasive breast carcinoma (BRCA), some of the tumorigenic cells are in a poorly differentiated state and lose proper regulation ability. These cells can leave the lesion and spread with the blood to other tissues or lymph nodes and develop into new tumors, leading to organ dysfunction and patient death.

Alternative splicing (AS) is a tumorigenesis mechanism that has been studied in a number of tumors and is widely accepted as explaining the aforementioned phenomenon. Through AS, an mRNA precursor can produce a number of mRNA splicing isoforms, which generates protein diversity (6). Proteins created by AS exhibit a number of molecular properties and interactions that have a significant role in both normal and abnormal life activities (7,8).

The role of AS in BRCA has recently been investigated, with studies analyzing certain AS events in breast cancer; these studies have demonstrated the potential for specific AS events to classify and diagnose cancer (9,10). Tien *et al* (11) demonstrated that the mutation of cyclin-dependent kinase 12, which disrupts DNA repair, affected a *DNAJB6* isoform and the DNA damage response activator by regulating last-exon splicing; thereby causing tumorigenesis and invasion. The complexes of transactive response DNA binding protein 43 (TDP43) and serine/arginine-rich splicing factor 3 (SRSF3) can modulate the AS events of protease-activated receptor 3

Correspondence to: Professor Huiqun Wu, Department of Medical Informatics, Nantong University, 19 Qixiu Road, Nantong, Jiangsu 226001, P.R. China
E-mail: wuhuiqun@ntu.edu.cn

Key words: alternative splicing, splicing factor, invasive breast carcinoma, prognosis

(PAR3) and endocytic adaptor protein, which suggests that TDP43 or SRSF3 knockdown inhibits tumor progression, and the higher expression level of TDP43 in triple-negative breast cancer may suggest a poor patient prognosis (12).

As previous studies have demonstrated the potential of AS events as molecular markers and therapeutic targets (13,14), the investigation of full transcriptome AS events and survival-related alternative splicing events (SREs) in BRCA should be a priority in research. The aim of the present study was to reveal new features of BRCA by integrating and comparing AS events at the full transcriptome level and to validate their clinical values using a number of models.

Materials and methods

Acquisition of data on alternative splicing events. The Cancer Genome Atlas (TCGA), which comprises high-throughput sequencing data and clinical information on 33 types of tumor, including BRCA, was used in the present study (www.cancer.gov/tcga). TCGA SpliceSeq database calculates the percent-spliced-in (PSI) value for all AS events in each tumor, which can then be used for AS analyses (15). The PSI value is a ratio that presents the efficiency of splicing exons into transcripts (splicing isoforms). As presented in Fig. 1, AS events were further divided into seven types: Alternative promoter (AP), alternative donor site (AD), alternative terminator (AT), exon skipping (ES), mutually exclusive exons (ME), alternative acceptor site (AA) and retained intron (RI). The PSI values for the AS events in 1,207 tissue samples were downloaded from TCGA SpliceSeq comprising 1,094 samples from cancerous tissues and 113 samples from healthy tissues. As some volunteers donated both cancerous and healthy tissues, the present study contains 1,097 sets of clinical data, which means that each sample has the corresponding clinical information.

Identification of SREs. Some of the 1,094 cancerous tissue samples were removed by integrating clinical information to make the results more objective. Although males can also have breast cancer, the present study considered the large differences in sex hormones between men and women and selected 1,081 samples from only female patients. This slightly reduces the clinical significance; however, it notably avoids the bias caused by biological differences (16). From the 1,081 female samples, 1,019 patients with an overall survival (OS) from 31 days to >13 years were selected for univariate survival analysis. The R package survival (version 3.1-11) was used to conduct univariate survival analysis, which calculated the relationship between each AS event and OS. This step is performed in RStudio software (version 1.1.442) (17) using R language (v3.5.1) (18). OS is the time from the start of the random assignment to death. For patients who were lost to follow-up or survived until the end of the study, OS was considered to be the time from the start of the random assignment to the date of the last follow-up. Cox's proportional hazards regression analysis quantitatively demonstrated the association between the PSI values and OS using the following four key values: Hazard ratio (HR), coefficient (coef) value and maximum and minimum values for 95% confidence intervals. Generally, in cancer studies, a factor is considered to have a bad effect on a prognosis if its HR >1. The coef value is associated with

the direction and extent of the event and its influence on the outcome. Specifically, when an event decreases OS, the corresponding coef value is positive. Conversely, when an event increases OS, the corresponding coef value is negative. The stronger the influence of the event on the outcome, the greater the absolute coef value. The smaller P-value is, the more the reliable the result. Therefore, when the number of SREs of an AS type was >10, the present study chosen 10 SREs with the smallest P-value were used to calculate the weighted PSI value. As an aberrant biological process is caused by several AS events, the weighted PSI value of each AS type is theoretically more biologically significant compared with the PSI value of a single AS event. The weighted PSI value of AA is the sum of the PSI values of the top 10 survival-associated AS events multiplied by the corresponding coef values. In addition, the PSI values were calculated for the top 10 most significant AS events in all SREs and the weighted PSI value that was obtained was considered to represent all AS events. Samples were divided into two groups according to these weighted PSI values. The Kaplan-Meier estimator and the log-rank test were used to determine whether there was a significant difference in survival rates between the two groups. These calculations were performed by R packages: Survival (version 3.1-11) and survminer (version 0.4.6) and the result of $P < 0.05$ has statistical significance.

Distribution of AS events and the UpSet plot. For its greater efficiency, an UpSet plot, rather than a Venn plot, was used to display the intersections among multiple datasets. Using this plot, the present study could sort the data by gene frequency or by the number of AS types contained within the set to more clearly represent distribution features. Survival-related genes (SRGs) were those involved in SREs. The UpSet plot was used to visualize the distribution of AS event-related genes and SRGs within the different AS types. The plot was created using R package: UpSetR (v1.4.0) (19).

Protein-protein interaction (PPI) network and enrichment analysis. In order to identify the genes at the core of the pathological process and determine how they regulate each other, the present study submitted SRGs to the Search Tool for the Retrieval of Interacting Genes/Proteins (www.string-db.org/). The PPI network was constructed using a threshold of 0.4 to avoid missing key genes. The degree of connectivity was used to describe the association between nodes. A node with a high degree of connectivity may have a wide impact on other nodes and was considered a hub gene. Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG), and Reactome contain a large number of canonical descriptions of genes and pathways that can be used to study the functions and pathways with which target genes may be involved (20,21). The present study used ClueGO (v2.5.6) (22), a plugin for Cytoscape (v3.7.2) (23), to annotate the physiological functions of SRGs.

Diagnostic test and 5-year survival model. Diagnostics were conducted to test the ability of the weighted PSI values to distinguish between cancerous and healthy tissues. A receiver operating characteristic (ROC) curve was plotted to determine whether the weighted PSI values from the top 10 SREs with the most prognostic significance and those for each AS type




Figure 1. A total of seven alternative splicing types. Each rectangle represents a part of pre-mRNAs. The two rectangles connected by lines indicate that the corresponding two sequences are joined together after alternative splicing.

could be used to predict a prognosis. A total of two models in the present study consisted of some weighted PSI values, although it was unclear whether these values could be used as indicators for distinguishing the different groups; however, the area-under-curve (AUC) value could be used as an indicator. Therefore, the ROC curves were used to present the value of the indicators for distinguishing between cancerous and healthy tissues in the present study, and to assess the ability to predict whether patient OS could be >5 years. The distinction between the two models was that one reflected the impact of AS events on initiation of the disease, while the other was associated with its progression. After performing the diagnostic tests, a 5-year survival model was created using SPSS v19.0 (IBM Corp.).

SFs and regulation network. SFs comprise numerous types of proteins, such as serine/arginine-rich (SR) protein, which contains a protein domain with long repeats of serine and arginine amino acid residues (24). SpliceAid 2 (www.introni.it/spliceaid.html) is a database of SFs in cancerous and healthy tissues (25); 71 SFs that were identified in BRCA and their corresponding genes were obtained from this database. TCGA provided the third-level transcriptome data for these BRCA genes. In order to exclude interferences, such as gene length, sequencing amount and sample specificity, the original read counts were normalized to increase reliability. SR SFs were

identified using Cox's proportional hazards regression model. If the Pearson correlation coefficient >0.4, the corresponding SF was positively correlated with AS events. The regulatory network was visualized using Cytoscape (v3.7.2) (21).

Results

Characteristics of AS in invasive breast carcinoma. The present study identified 45,421 AS events in BRCA, which were associated with 10,480 genes (Fig. 2A). ES had the most AS events (17,702). A total of 233 AS events was found in ME, notably fewer compared with other AS types. ES had the largest number of associated genes (6,811). The number of genes associated with ME was 227, which was smaller than that of any other AS type. Some genes were found in only one type of AS event in BRCA, while others were demonstrated to be involved in several types (Fig. 3A). The largest group of genes (1,782) was contained in only the ES type, which accounted for 26.1% of the associated ES genes. The proportion of genes only associated with one AS type were 8.4, 9.7, 19.6, 28.3, 5.2 and 16.0% in AA, AD, AP, AT, ME and RI, respectively, which indicated that only a few genes were involved in only one type of AS. Therefore, the majority of event-associated genes were involved in more than one type of AS event. The group consisting of AT and ES contained most

Table I. Information of survival analysis based on the weighted PSI value.

AS type	HR	Coef	95% CI lower	95% CI upper	P-value
AA	0.487	-0.720	-0.916	-0.523	<0.001
AD	0.443	-0.815	-0.998	-0.632	<0.001
AP	1.511	0.413	0.292	0.533	<0.001
AT	0.697	-0.361	-0.449	-0.273	<0.001
ES	0.564	-0.572	-0.685	-0.459	<0.001
ME	2.079	0.732	0.347	1.116	<0.001
RI	0.701	-0.356	-0.482	-0.230	<0.001
TOP10	0.682	-0.382	-0.475	-0.290	<0.001

Coef is used to represent the quantitative relationship between variables and results. Absolute values represent correlation strength, positive numbers represent positive correlations, and negative numbers represent negative correlations. AS, alternative splicing; Coef, coefficient; CI, confidence interval; HR, hazard ratio; RI, retained intron; ME, mutually exclusive exons; ES, exon skipping; AT, alternative terminator; AP, alternative promoter; AD, alternative donor site; AA, alternative acceptor site; TOP10, 10 SREs with the smallest P-values.




Figure 2. Distribution of SREs and related genes in invasive breast carcinoma. The y-axis is AS type. The x-axis is the number of cases. (A) Green strips represent the number of AS events. Blue strips represent the number of AS event-associated genes. (B) Green strips represent the number of SREs. Blue strips represent the number of SRGs. AS, alternative splicing; SRE, survival-related alternative splicing event; SRG, survival-related gene; BRCA, breast carcinoma; RI, retained intron; ME, mutually exclusive exons; ES, exon skipping; AT, alternative terminator; AP, alternative promoter; AD, alternative donor site; AA, alternative acceptor site.

genes (680) compared with other groups with only two AS types. The group consisting of AP, AT and ES was the largest group (286 genes) among the groups that contained three types of AS events.

SREs and SRGs identified using survival analysis. A series of SREs and SRGs identified using Cox's proportional hazards regression model ($P < 0.05$) were investigated; the results of the survival analyses are presented in Table SI. In general, 1,215 events in 818 genes appeared to have potential links to OS. Although there were 17,702 AS events in ES, only 268 were associated with a prognosis. The AS type with the largest number of events and genes was not ES, but AT. Fig. 2B presents the number of SREs and SRGs in each type of AS.

The largest percentage of SRGs (778; 95.1%) was associated with only one type of AS event. The different distributions of AS event-associated genes and SRGs indicated that mRNA produced by SRGs appeared to be more specific. The group that included genes from only AT contained 298 SRGs and represented the group with the most genes. The intersections of the seven AS types are presented in Fig. 3.

Prognostic models based on the weighted PSI value of AS types. The PSI values of the 10 most significant splicing events in each AS type were weighted to obtain a weighted PSI value for each type. As ME had only four SREs, its weighted PSI value was calculated using only these four events (Table I). P-value was < 0.01 for all items, which indicated that the weighted calculation method was reliable. In the present study, AP and ME had higher hazard ratios (HRs) than the other AS types, which suggested that the AS events contained in AP and ME may increase both the risk of disease and a poor prognosis. All coef values for AA, AD, AT, ES, RI and the 10 most significant splicing events were negative, which indicated that their effects on prognosis were positive. Kaplan-Meier survival curves were plotted to display the differences in survival rate over time (Fig. 4). The ends of the curves for AA, AP and AT were very close and may have been affected by other factors, such as age (Fig. 4A, C and D). In addition, the weighted PSI values based on the 10 AS events in each AS type (a total of 70 events) were also calculated. The corresponding survival curve is presented in Fig. S1, which shows trends similar to those in Fig. 4G and indicates that the weighted PSI value




Figure 3. UpSet plot for all AS events. Blue strip shows the number of events included in each AS type. Dots and lines represent subsets of AS events. The AS types corresponding to the dots are contained in the subtype. The histogram represents the number of genes in each subset. (A) All genes. (B) Survival-related genes. AS, alternative splicing; RI, retained intron; ME, mutually exclusive exons; ES, exon skipping; AT, alternative terminator; AP, alternative promoter; AD, alternative donor site; AA, alternative acceptor site.

was a good measure for distinguishing between groups with a longer and shorter OS.

Fig. 5A presents the ROC curves that tested the ability of the weighted PSI value to determine the patients' 5-year OS rate; the quantitative results are presented in Table II. AD (AUC=0.723) was considered to be the best indicator of a prognostic model; AUC of all other values were <0.7.

PPI network and enrichment analysis based on SREs. The PPI network (Fig. 6) indicated that there were 13 hub genes connecting >20 nodes, the majority of which were ribosomal

protein genes; these were listed in Table III. A total of 22, 7 and 29 items were enriched in the GO, KEGG and Reactome databases, respectively. The results of the enrichment analysis are presented in Table IV and the P-values of all items were <0.05. The items in this table were sorted by the percentage of associated genes, which clearly showed the proportion of the submitted genes occupying the genes contained in each function or pathway. The items enriched in GO (biological processes), KEGG and Reactome databases with the highest percentage of associated genes were 'regulation of ribonuclease activity', 'ubiquinone and other terpenoid-quinone




Figure 4. Survival curve for the weighted percent-spliced-in value. Red and blue lines represent changes in survival probability in days in the group with lower and higher values, respectively. Red and blue areas represent the 95% confidence interval. The number of individuals in a group who remain alive at a certain point in time is the number at risk, which was exhibited in the lower part of each plot. (A) The AA type, (B) AD type, (C) AP type, (D) AT type, (E) ES type and (F) ME type.




Figure 4. Continued. (G) RI type, (H) based on the 10 most significant SREs. (I) Third-level transcriptome data for ELAVL4. AA, alternative acceptor; AD, alternative donor; AP, alternative promoter; AT, alternative promoter; ES, exon skipping; ME, mutually exclusive exon; RI, retained intron; SRE, alternative splicing event.

biosynthesis' and 'PKA activation in glucagon signaling', respectively.

Diagnostic tests. The diagnostic tests comprised eight indicators with weighted PSI values for the seven AS types and those calculated from the PSI values of the 10 most significant SREs. Fig. 5B shows the ROC curve for each indicator in the diagnostic test, which imply that certain AS types, such as AT and ME, are unsuitable for use as diagnostic indicators. The AUC values of AT and ME were 0.593 and 0.633, respectively, and, in general, an indicator with AUC <0.7 was not considered to be useful. AA appeared to be the most efficient indicator for distinguishing between the two groups (AUC=0.823). Quantitative results of each ROC curve are provided in Table V.

Regulatory network based on SFs and SREs. A total of 51 SFs were matched with read counts, which were used for survival

analysis using Cox's proportional hazards regression model; however, under the condition that P<0.05 was considered to be significant, only one gene, ELAVL4, was selected. The HR of ELAVL4 was 1.01, which suggested that it was a risk factor that may indicate a poor prognosis. The present study also demonstrated that OS changed with time (Fig. 4I). The group with a lower expression level of ELAVL4 had a higher OS rate than that with a higher expression level, which was more notable in patients with OS >7 years. By calculating the Pearson correlation coefficient, 11 SREs were selected, six of which were positively associated and five of which were negatively associated with ELAVL4 expression levels. In particular, the six that were positively associated were associated with a poor prognosis, while the five that were negatively associated were associated with an improved prognosis, which suggested the positive and negative effects of the association between SREs and ELAVL4 expression levels (Fig. 7).

Table II. Information of ROC curve on predicting 5-year survival.

AS type	Cut-off	Sensitivity	Specificity	AUC	95% CI lower	95% CI upper	P-value
AA	-76.343	0.633	0.670	0.675	0.616	0.733	<0.001
AD	-16.144	0.806	0.554	0.723	0.672	0.773	<0.001
AP	12.169	0.776	0.452	0.648	0.591	0.705	<0.001
AT	-0.797	0.480	0.762	0.655	0.598	0.713	<0.001
ES	-504.463	0.704	0.578	0.680	0.623	0.736	<0.001
ME	68.350	0.888	0.240	0.575	0.516	0.634	0.016
RI	-249.080	0.857	0.382	0.632	0.576	0.687	<0.001
TOP10	-332.922	0.592	0.666	0.660	0.603	0.716	<0.001

AUC, area under curve; CI, Confidence interval; RI, retained intron; ME, mutually exclusive exons; ES, exon skipping; AT, alternative terminator; AP, alternative promoter; AD, alternative donor site; AA, alternative acceptor site; TOP10, 10 SREs with the smallest P-values.

Table III. Degree value of hub genes in protein-protein interaction network.

Gene	Degree value
RPS15	27
RPL35A	26
RPL9	25
RPS15A	25
RPS29	24
RPS5	24
RPS3	23
RPL23	23
GNB2L1	22
RPL13	22
RPL7L1	22
RPS9	22
RPL18A	21

Degree value: The degree of connection was used to describe the connectivity of a particular node to other nodes.

Discussion

Biomarkers do not simply classify tumors into several types, but rather represent certain properties of the tumor, which are of significance for precision medical treatment. The traditional classification of a tumor is based primarily on its location, pathological morphology and distant metastasis; however, more molecular level-based classification methods have previously been proposed. For example, the presence of estrogen and/or progesterone receptors suggests that endocrine therapy has a favorable therapeutic effect (26). Gene expression level profiling is used to understand the differences in gene expression levels and pathogenesis of cancerous and healthy tissues in BRCA and could be used in BRCA classification for different prognoses (27-29). Although gene expression level profiling has been extensively studied, little is currently known about the AS profile of BRCA. Previous research has focused

on elucidating the pathological mechanisms of a single AS event or SF, which may lead to neglecting their population characteristics, regulatory relationships and clinical values. The present study identified a series of SREs through survival analysis and used a 5-year survival model and diagnostic tests to evaluate the ability of AS events to diagnose BRCA and predict a prognosis. Identification of SREs and the construction of the SF-AS event regulatory network has laid the foundation for subsequent classification.

Gene mutations are considered to be the major cause of tumorigenesis and >90% of coding genes are considered to have undergone AS events (25). Independently of gene mutations, AS events can also result in products expressed by deviant genes (8,26). For example, *TP53* is one of the first-discovered tumor-suppressor genes, and its isoform, produced by AS, modulates its tumor-suppressor function; whereas, the dysregulation of the *TP53* isoforms are found in a variety of tumors, which are closely associated with aberrant AS events (30). Aberrant AS produces cancer-specific mRNA that could further disrupt the normal function of tumor suppressors and activate oncogenic pathways (13). Furthermore, aberrant AS events are involved in the establishment of the tumor micro-environment, thus promoting tumor growth, invasion and metastasis (31). For example, RNA-binding proteins participate in the specific expression of isoforms of vascular endothelial growth factor through AS, resulting in a unique angiogenic profile of colorectal cancer (32). In the present study, AS events were represented by PSI values, which allowed them to be quantitatively analyzed. Based on the correlation between PSI values and prognosis, a series of SREs were identified, and their distribution characteristics in BRCA were exhibited using an UpSet plot.

The present study demonstrated that ES was the type with the most AS events, while AT had the most SREs. The distribution of AS-associated genes was consistent with AS events, which means that AT also has the most SRGs. The difference in the distribution between AS events and SREs suggested that AT had a notable effect on prognosis. The pathological role of ES is worthy of further investigation because the number of SREs in ES ranks second. In other tumor studies, ES is the most common type of AS, with ME events being the least common (33,34). SRE distribution exhibits different




Figure 5. ROC curves for 5-year survival and diagnostic models. The ROC curves present the sensitivity and (1-specificity) values for indicators at different cutoff values. (A) The 5-year survival model. (B) Diagnostic model. ROC, receiver operating characteristic; RI, retained intron; ME, mutually exclusive exons; ES, exon skipping; AT, alternative terminator; AP, alternative promoter; AD, alternative donor site; AA, alternative acceptor site. TOP10, 10 SREs with the smallest P-values.




Figure 6. Protein-protein interaction network for survival-associated genes. Color intensity represents the number of nodes connected to one node. Thick and thin lines represent strong and weak co-expression associations, respectively.

Table IV. Items with higher percent of associated genes in enrichment analysis.

A, Gene Ontology (Biological Processes)

Items	P-value	Associated genes, %	Gene number
1. Regulation of ribonuclease activity	<0.01	42.86	3
2. Ribosomal small subunit export from nucleus	<0.01	42.86	3
3. Ribosomal subunit export from nucleus	<0.01	35.71	5
4. Ribosomal RNA-containing ribonucleoprotein complex export from nucleus	<0.01	31.25	5
5. Cotranslational protein targeting to membrane	<0.01	17.14	18

B, Kyoto Encyclopedia of Genes and Genome

Items	P-value	Associated genes, %	Gene number
1. Ubiquinone and other terpenoid-quinone biosynthesis	0.01	27.27	3
2. Thiamine metabolism	0.01	25.00	4
3. Vasopressin-regulated water reabsorption	<0.01	18.18	8
4. Pyruvate metabolism	<0.01	17.95	7
5. Thyroid cancer	0.01	16.22	6

C, Reactome

Items	P-value	Associated genes, %	Gene number
1. PKA activation in glucagon signaling	<0.01	29.41	5
2. TP53 regulates transcription of genes involved in G ₂ cell cycle arrest	<0.01	27.78	5
3. Constitutive signaling by ligand-responsive EGFR cancer variants	<0.01	26.32	5
4. Signaling by EGFR in Cancer	<0.01	26.32	5
5. Signaling by ligand-responsive EGFR variants in cancer	<0.01	26.32	5

PKA, protein kinase A; EGFR, epidermal growth factor receptor.




Figure 7. Regulatory network for SFs and alternative splicing events. The green node represents the survival-related SF, named ELAVL4. Red and blue nodes represent SREs that increase and decrease risk, respectively. Purple and orange lines represent the positive and negative correlations of connected nodes, respectively. Thick lines indicate a strong correlation, thin lines indicate a weak correlation. SRE, alternative splicing events; SF, splicing factor; HR, hazard ratio.

characteristics. Overall, for BRCA and colon adenocarcinoma, the majority of SREs are found in AT, not in ES, while for esophageal, stomach and rectal adenocarcinomas, and diffuse

large B-cell lymphoma, the ES type contains more SREs than the AT type. This difference may correspond to the common properties of tumors from different tissues (35). In addition, SRGs exhibit characteristics different from those in AS-associated genes. Although the number of SRGs with only one AS event is highest, which may be due to the stability of the associated pre-mRNA, there are also a number of genes with two or more AS events; however, the pathways that influence a prognosis remain unclear.

AS events could be further regulated by a group of SFs. A previous study demonstrated that individual SR proteins can restore pre-mRNA splicing in cell extracts depleted of multiple SR family proteins (36). The SR protein family contains a number of proteins with phylogenetic conservation and structural relevance, with their characteristic domains containing multiple serine and arginine residues. Numerous human diseases, such as cancer and human immunodeficiency virus, are associated with the SR protein family (37,38). SpliceAid 2 integrates existing research and includes 71 SFs and their distributions in various tissues (25). SF gene mutations constitute early events that most likely play a role in initiation of the tumorigenesis of certain types of tumor (37). The results of a cohort study demonstrated that *SF3B1* mutations occur in >20% of patients

Table V. Information of receiver operating characteristic curve on diagnostic test.

AS type	Cut-off	Sensitivity	Specificity	AUC	95% CI lower	95% CI upper	P-value
AA	-77.313	0.788	0.735	0.823	0.786	0.861	<0.001
AD	-16.946	0.549	0.852	0.756	0.707	0.804	<0.001
AP	11.661	0.805	0.712	0.798	0.757	0.839	<0.001
AT	-1.520	0.796	0.422	0.593	0.548	0.638	0.001
ES	-505.318	0.823	0.711	0.819	0.785	0.853	<0.001
ME	68.415	0.460	0.739	0.633	0.582	0.684	<0.001
RI	-248.709	0.743	0.555	0.672	0.628	0.717	<0.001
TOP10	-333.612	0.832	0.478	0.692	0.652	0.733	<0.001

AUC, area under curve; CI, Confidence interval; RI, retained intron; ME, mutually exclusive exons; ES, exon skipping; AT, alternative terminator; AP, alternative promoter; AD, alternative donor site; AA, alternative acceptor site; TOP10, 10 SREs with the smallest P-values.

with uveal melanoma (39). SFs are also involved in the biological processes of BRCA, such as tumorigenesis, growth, infiltration and metastasis, and are thus involved in prognosis (9,12,40-42). SFs and AS have been found in the immune-evasion pathway of tumors (43). As one SF may regulate more than one AS event, each SF has multiple pathogenic pathways; this suggests that SF-based treatment may be a broad spectrum. AS events and SFs may be key targets for the treatment of cancer and merit further investigation. In addition, the present study incorporated SFs into the regulatory network to clarify the results. Enrichment analysis was used to determine the physiological functions and signaling pathways involved in SREs. The present study also examined whether SREs can be markers for diagnosis and prognosis, which will provide a reference for subsequent studies. Given the prognostic relevance of SRGs, their regulatory networks and biological functions in BRCA merit further study.

The PPI network examined the hub genes, which are primarily ribosomal protein genes. Previous studies reported that a number of ribosomal proteins were involved in the initiation and progress of BRCA. Knockdown of ribosomal protein S15A represses the proliferation of breast cancer cells *in vitro* by inducing apoptosis (44). Studies have demonstrated that ribosomal protein S6 kinase 4 had anti-invasive and anti-metastatic activities (45,46). Ribosomal protein S3 upregulates the X-linked inhibitor of apoptosis to confer the resistance of breast cancer cells to certain chemotherapeutic drugs (47). The PPI network suggests that AS may be one of the pathways by which ribosomal protein genes are involved in tumorigenesis. In previous studies, functional enrichment analyses were used to identify that the signaling pathways associated with ribosomes are found in esophageal, colon and rectal adenocarcinomas (33,34). The annotation of functions and pathways found in the GO, KEGG and Reactome databases were used to understand their pathological mechanisms. The results of enrichment analysis based on the GO database indicated that the proportion of SRGs within the gene group involved in the ribosome-associated biological processes was higher than that in other functional gene groups, such as the regulation of ribonuclease activity, ribosomal small-subunit export from the nucleus and ribosomal subunit export from the nucleus. To the best of our knowledge, few studies have previously noted changes in these biological processes and

how SRGs involved in these processes affect the prognosis for patients with BRCA.

In the present study, survival curves demonstrated that the combined effects of SREs were highly associated with patient OS; however, not all AS types represented by weighted PSI values were able to predict a prognosis. Some AS types with lower AUC values, such as AT and ME, did not appear to be suitable predictors for the 5-year survival outcome, which may have three possible explanations. First, the number of AS events used to calculate the weighted PSI values may have been too small to reflect the overall characteristics. Secondly, the calculation of SRGs participating in different functions may have masked some of the original attributes. The third cause may have been the special properties of BRCA as the indicators based on weighted PSI values were excellent in predicting a prognosis in gastrointestinal pan-adenocarcinomas (34). The diagnostic model implies that the weighted PSI values of AA, AD, AP and ES are reliable predictors. It is worth noting that although the sensitivity of AD is only 0.549, the specificity is 0.852, which suggests that there may be some AS events in AD that create significant changes in cancerous and healthy tissues; therefore, more research on SREs in AD is required.

ELAVL4 was associated with OS in patients with BRCA, which was consistent with the results of previous studies on non-small-cell lung cancer and meningioma (48,49). The present study indicates that ELAVL4 has a potential regulatory relationship with multiple SRGs, and that the AS is the potential mechanism by which they affect BRCA. Another uncertainty of the analyses in the present study is that it is difficult to infer the functional impact of AS and the altered protein structure. Certain AS events will totally drive structural changes in protein outputs; however, current algorithms may not precisely quantify those variations. New computational methods are necessary to replicate the present study to confirm the results. In addition, all results should be tested using another set of samples to determine the reliability of the results of the present study.

The present study systematically identified a number of SREs in BRCA, described the distribution characteristics of SREs and SRGs and mapped the regulatory networks based on the SRGs, as well as investigating potential pathological mechanisms. ELAVL4 has a potential regulatory relationship with multiple SRGs and is worth further investigation. Further

studies are needed to examine the potential of AS events as prognostic biomarkers and to provide insight on subsequent identification of therapeutic targets.

Acknowledgements

The authors would like to thank Mrs. Xuemei Guan (Suzhou Ninth People's Hospital) and Dr. Yongchun Gu (Suzhou Ninth People's Hospital) for their support and suggestions.

Funding

The present study was supported by a grant from Nantong '226 Project' (grant no. 2018436) and Excellent Key Teachers in the 'Qing Lan Project' of Jiangsu Colleges and Universities (grant no. 06190025).

Availability of data and materials

The datasets generated and/or analyzed during the present study are available in the figures, tables and supplementary files; TCGA database (<https://portal.gdc.cancer.gov/>); TCGA SpliceSeq database (<https://bioinformatics.mdanderson.org/public-software/tcgaspliceseq/>); SpliceAid 2 database (www.introni.it/spliceaid.html).

Authors' contributions

HQW, KRJ and YCW conceived and designed the current research. KRJ, JH, JNC and HGW collected and preprocessed the data. KRJ and YCW performed data analysis and wrote the manuscript. All authors had read and agreed with the final version of manuscript.

Ethics approval and consent for participation

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Miao H, Verkooijen HM, Chia KS, Bouchardy C, Pukkala E, Larønningen S, Mellemkjær L, Czene K and Hartman M: Incidence and outcome of male breast cancer: An international population-based study. *J Clin Oncol* 29: 4381-4386, 2011.
- Coughlin SS: Social determinants of breast cancer risk, stage, and survival. *Breast Cancer Res Treat* 177: 537-548, 2019.
- San Miguel Y, Gomez SL, Murphy JD, Schwab RB, McDaniels-Davidson C, Canchola AJ, Molinolo AA, Nodora JN and Martinez ME: Age-related differences in breast cancer mortality according to race/ethnicity, insurance, and socioeconomic status. *BMC Cancer* 20: 228, 2020.
- DeSantis CE, Ma J, Goding Sauer A, Newman LA and Jemal A: Breast cancer statistics, 2017, racial disparity in mortality by state. *CA Cancer J Clin* 67: 439-448, 2017.
- Siegel RL, Miller KD and Jemal A: Cancer statistics, 2019. *CA Cancer J Clin* 69: 7-34, 2019.
- Black DL: Protein diversity from alternative splicing: A challenge for bioinformatics and post-genome biology. *Cell* 103: 367-370, 2000.
- Yang X, Coulombe-Huntington J, Kang S, Sheynkman GM, Hao T, Richardson A, Sun S, Yang F, Shen YA, Murray RR, *et al*: Widespread expansion of protein interaction capabilities by alternative splicing. *Cell* 164: 805-817, 2016.
- Parikhshak NN, Swarup V, Belgard TG, Irimia M, Ramaswami G, Gandal MJ, Hartl C, Leppa V, Ubieta LT, Huang J, *et al*: Genome-wide changes in lncRNA, splicing, and regional gene expression patterns in autism. *Nature* 540: 423-427, 2016.
- Read A and Natrajan R: Splicing dysregulation as a driver of breast cancer. *Endocr Relat Cancer* 25: R467-R478, 2018.
- Tian N, Li J, Shi J and Sui G: From general aberrant alternative splicing in cancers and its therapeutic application to the discovery of an oncogenic DMTF1 isoform. *Int J Mol Sci* 18: E191, 2017.
- Tien JF, Mazloomian A, Cheng SG, Hughes CS, Chow CCT, Canapi LT, Oloumi A, Trigo-Gonzalez G, Bashashati A, Xu J, *et al*: CDK12 regulates alternative last exon mRNA splicing and promotes breast cancer cell invasion. *Nucleic Acids Res* 45: 6698-6716, 2017.
- Ke H, Zhao L, Zhang H, Feng X, Xu H, Hao J, Wang S, Yang Q, Zou L, Su X, *et al*: Loss of TDP43 inhibits progression of triple-negative breast cancer in coordination with SRSF3. *Proc Natl Acad Sci USA* 115: E3426-E3435, 2018.
- Climente-González H, Porta-Pardo E, Godzik A and Eyraes E: The functional impact of alternative splicing in cancer. *Cell Rep* 20: 2215-2226, 2017.
- Martinez-Montiel N, Rosas-Murrieta NH, Anaya Ruiz M, Monjaraz-Guzman E and Martinez-Contreras R: Alternative splicing as a target for cancer treatment. *Int J Mol Sci* 19: E545, 2018.
- Ryan M, Wong WC, Brown R, Akbani R, Su X, Broom B, Melott J and Weinstein J: TCGASpliceSeq a compendium of alternative mRNA splicing in cancer. *Nucleic Acids Res* 44: D1018-D1022, 2016.
- Gucalp A, Traina TA, Eisner JR, Parker JS, Selitsky SR, Park BH, Elias AD, Baskin-Bey ES and Cardoso F: Male breast cancer: A disease distinct from female breast cancer. *Breast Cancer Res Treat* 173: 37-48, 2019.
- RStudio Team (2015). RStudio: Integrated Development for R. RStudio, Inc., Boston, MA, 2015. <http://www.rstudio.com/>.
- R Core Team (2012). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, 2012. <http://www.R-project.org/>.
- Conway JR, Lex A and Gehlenborg N: UpSetR: An R package for the visualization of intersecting sets and their properties. *Bioinformatics* 33: 2938-2940, 2017.
- The Gene Ontology Consortium: Expansion of the gene ontology knowledgebase and resources. *Nucleic Acids Res* 45: D331-D338, 2017.
- Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, Haw R, Jassal B, Korninger F, May B, *et al*: The reactome pathway knowledgebase. *Nucleic Acids Res* 46: D649-D655, 2018.
- Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, Fridman WH, Pagès F, Trajanoski Z and Galon J: ClueGO: A cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* 25: 1091-1093, 2009.
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B and Ideker T: Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13: 2498-2504, 2003.
- Sahebi M, Hanafi MM, van Wijnen AJ, Azizi P, Abiri R, Ashkani S and Taheri S: Towards understanding pre-mRNA splicing mechanisms and the role of SR proteins. *Gene* 587: 107-119, 2016.
- Piva F, Giulietti M, Burini AB and Principato G: SpliceAid 2: A database of human splicing factors expression data and RNA target motifs. *Hum Mutat* 33: 81-85, 2012.
- Woolston C: Breast cancer. *Nature* 527: S101, 2015.
- Stricker TP, Brown CD, Bandalamudi C, McNeerney M, Kittler R, Montoya V, Peterson A, Grossman R and White KP: Robust stratification of breast cancer subtypes using differential patterns of transcript isoform expression. *PLoS Genet* 13: e1006589, 2017.
- Kim SY, Kawaguchi T, Yan L, Young J, Qi Q and Takabe K: Clinical relevance of microRNA expressions in breast cancer validated using the cancer genome atlas (TCGA). *Ann Surg Oncol* 24: 2943-2949, 2017.
- Heng YJ, Lester SC, Tse GM, Factor RE, Allison KH, Collins LC, Chen YY, Jensen KC, Johnson NB, Jeong JC, *et al*: The molecular basis of breast cancer pathological phenotypes. *J Pathol* 241: 375-391, 2017.

30. Wang ET, Sandberg R, Luo S, Khrebukova I, Zhang L, Mayr C, Kingsmore SF, Schroth GP and Burge CB: Alternative isoform regulation in human tissue transcriptomes. *Nature* 456: 470-476, 2008.
31. Biamonti G, Catillo M, Pignataro D, Montecucco A and Ghigna C: The alternative splicing side of cancer. *Semin Cell Dev Biol* 32: 30-36, 2014.
32. Hamdollah Zadeh MA, Amin EM, Hoareau-Aveilla C, Domingo E, Symonds KE, Ye X, Heesom KJ, Salmon A, D'Silva O, Betteridge KB, *et al*: Alternative splicing of TIA-1 in human colon cancer regulates VEGF isoform expression, angiogenesis, tumour growth and bevacizumab resistance. *Mol Oncol* 9: 167-178, 2015.
33. Xiong Y, Deng Y, Wang K, Zhou H, Zheng X, Si L and Fu Z: Profiles of alternative splicing in colorectal cancer and their clinical significance: A study based on large-scale sequencing data. *EBioMedicine* 36: 183-195, 2018.
34. Lin P, He RQ, Ma FC, Liang L, He Y, Yang H, Dang YW and Chen G: Systematic analysis of survival-associated alternative splicing signatures in gastrointestinal pan-adenocarcinomas. *EBioMedicine* 34: 46-60, 2018.
35. Zhang R, Lin P, Yang X, He RQ, Wu HY, Dang YW, Gu YY, Peng ZG, Feng ZB and Chen G: Survival associated alternative splicing events in diffuse large B-cell lymphoma. *Am J Transl Res* 10: 2636-2647, 2018.
36. Zahler AM, Lane WS, Stolk JA and Roth MB: SR proteins: A conserved family of pre-mRNA splicing factors. *Genes Dev* 6: 837-847, 1992.
37. Kedzierska H and Piekliko-Witkowska A: Splicing factors of SR and hnRNP families as regulators of apoptosis in cancer. *Cancer Lett* 396: 53-65, 2017.
38. Mahiet C and Swanson CM: Control of HIV-1 gene expression by SR proteins. *Biochem Soc Trans* 44: 1417-1425, 2016.
39. Decatur CL, Ong E, Garg N, Anbunathan H, Bowcock AM, Field MG and Harbour JW: Driver mutations in uveal melanoma: Associations with gene expression profile and patient outcomes. *JAMA Ophthalmol* 134: 728-733, 2016.
40. Hu Y, Sun Z, Deng J, Hu B, Yan W, Wei H and Jiang J: Splicing factor hnRNP2B1 contributes to tumorigenic potential of breast cancer cells through STAT3 and ERK1/2 signaling pathway. *Tumour Biol* 39: 1010428317694318, 2017.
41. Koedoot E, Fokkelman M, Rogkoti VM, Smid M, van de Sandt I, de Bont H, Pont C, Klip JE, Wink S, Timmermans MA, *et al*: Uncovering the signaling landscape controlling breast cancer cell migration identifies novel metastasis driver genes. *Nat Commun* 10: 2983, 2019.
42. Singh R, Gupta SC, Peng WX, Zhou N, Pochampally R, Atfi A, Watabe K, Lu Z and Mo YY: Regulation of alternative splicing of Bcl-x by BC200 contributes to breast cancer pathogenesis. *Cell Death Dis* 7: e2262, 2016.
43. Oltean S and Bates DO: Hallmarks of alternative splicing in cancer. *Oncogene* 33: 5311-5318, 2014.
44. Feng W, Liang C, Wang C, Yu X, Li Q and Yang H: Knockdown of ribosomal protein S15A inhibits proliferation of breast cancer cells through induction of apoptosis in vitro. *Cytotechnology* 70: 1315-1323, 2018.
45. Thakur A, Sun Y, Bollig A, Wu J, Biliran H, Banerjee S, Sarkar FH and Liao DJ: Anti-invasive and antimetastatic activities of ribosomal protein S6 kinase 4 in breast cancer cells. *Clin Cancer Res* 14: 4427-4436, 2008.
46. Zhu J, Li QY, Liu JL, Wei W, Yang HW and Tang W: RSK4 knock-down promotes proliferation, migration and metastasis of human breast adenocarcinoma cells. *Oncol Rep* 34: 3156-3162, 2015.
47. Ono H, Iizumi Y, Goi W, Sowa Y, Taguchi T and Sakai T: Ribosomal protein S3 regulates XIAP expression independently of the NF- κ B pathway in breast cancer cells. *Oncol Rep* 38: 3205-3210, 2017.
48. Wang F, Lu J, Li S, Huo X, Liu X, Du X, Li C, Wang J and Chen Z: Application of serum ELAVL4 (HuD) antigen assay for small cell lung cancer diagnosis. *Anticancer Res* 37: 4515-4522, 2017.
49. Wong KK, Rostomily R and Wong STC: Prognostic gene discovery in glioblastoma patients using deep learning. *Cancers (Basel)* 11: E53, 2019.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.