

Investigation of the association between C-X-C motif chemokine receptor subunits and tumor infiltration levels and prognosis in patients with early-stage pancreatic ductal adenocarcinoma

QIONG-YUAN WU¹, CHENG-KUN YANG², LIANG-JUN RONG¹, JUN-CHAN LI¹ and LONG-MING LEI¹

¹Department of Tuina, The First Affiliated Hospital of Guangxi University of Chinese Medicine;

²Department of Hepatobiliary Surgery, The First Affiliated Hospital of Guangxi Medical University, Nanning, Guangxi 530023, P.R.China

Received August 5, 2019; Accepted April 1, 2020

DOI: 10.3892/ol.2020.11877

Abstract. Pancreatic ductal adenocarcinoma (PDAC) is one of the malignancies with the highest mortality rate due to postoperative local invasion and distant metastasis. Although C-X-C motif chemokine receptor (CXCR) subunits have been reported as prognostic indicators in gastric cancer, the prognostic value of CXCR subunits in PDAC remains poorly understood. In the present study, the expression levels and biological functions of CXCR subunits were investigated using multiple publicly accessible bioinformatic platforms and databases. Survival analysis was used to evaluate the prognostic value of CXCR subunits in 112 early-stage PDAC cases by setting the median expression levels as the cut-off values. A nomogram was constructed to combine CXCR subunit expression levels and clinical data for prognosis prediction. Moreover, the association between CXCR subunit expression levels and tumor infiltration levels were detected in PDAC. The expression levels of CXCR subunits were elevated in PDAC tumor tissues. In the multivariate Cox proportional risk regression model, high CXCR2, CXCR4 and CXCR6 expression levels in early-stage PDAC were associated with a more favorable prognosis. Further, it was demonstrated that the differential

expression levels of CXCR subunits in PDAC for combined survival analysis could contribute to risk stratification. The nomogram model demonstrated the contribution of CXCR subunits and clinical features in the prognosis of PDAC. Gene Set Enrichment Analysis suggested that CXCR subunits serve a role in immunomodulatory functions. The expression levels and somatic copy number alterations of CXCR subunits were associated with tumor infiltration levels in PDAC. CXCR subunits were associated with prognosis in patients with early-stage PDAC and may be potential drug targets for the treatment of pancreatic cancer.

Introduction

Pancreatic cancer is known to be one of the most lethal malignancies globally. Currently, due to the aggressiveness and lack of effective treatment of pancreatic cancer, it ranks as the fourth leading cause of cancer-associated death in the United States in 2017, with a 5-year survival rate of 6% (1). In China, pancreatic cancer ranks ninth and sixth in cancer morbidity and mortality, respectively, and it is estimated that 91,000 cases were diagnosed in 2015, of which 79,400 died from this disease (2). The major histological type of pancreatic cancer (>95% of cases) is pancreatic ductal adenocarcinoma (PDAC) (3). Since there is a lack of early clinical symptoms and effective biomarkers, most patients are diagnosed at an advanced stage and only 10% of patients are eligible for surgery (4).

The tumor microenvironment serves a key role in tumor self-monitoring and control of malignant transformation. The function of tumor infiltrating immune cells can be mediated by chemokine heterocomplexes of chemokine agonists and specific receptors that can activate multiple chemokine receptors modulating cell recruitment, localization and conversion of cellular invasive components at different tumor stages (5). In inflammatory types of cancer, patients with established T cell tumor infiltration have an improved prognosis and response to immunotherapy compared with patients with non-inflammatory tumors (6-9). High immune cell scores, determined by scoring tumor samples based on the total number of immune cells at the center of the tumor and on the edge of invasion,

Correspondence to: Dr Long-Ming Lei, Department of Tuina, The First Affiliated Hospital of Guangxi University of Chinese Medicine, 89-9 Dongge Road, Nanning, Guangxi 530023, P.R. China
E-mail: leilongming@126.com

Abbreviations: PDAC, pancreatic ductal adenocarcinoma; CXCR, C-X-C motif chemokine receptor; GO, Gene Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; STITCH, Search Tool for Interactions of Chemicals; GTEX, Genotype-Tissue Expression; PAAD, pancreatic adenocarcinoma; TCGA, The Cancer Genome Atlas; OS, overall survival; DFS, disease-free survival; HR, hazard ratio; CI, confidence interval; ROC, receiver operating characteristic; AUC, area under curves; GSEA, Gene Set Enrichment Analysis; TIMER, Tumor Immune Estimation Resource

Key words: PDAC, CXCR, immune infiltration, survival analysis

were significantly associated with an improved prognosis in patients with early-stage PDAC after pancreaticoduodenal resection (10). Lianyuan *et al* (11) reported that low levels of stromal tumor-infiltrating lymphocytes are a poor indicator of prognosis and liver metastasis in patients with PDAC after surgery (11). Poschke *et al* (12) analyzed infiltration of T-cells in tumor biopsy of resectable PDAC by immunohistochemistry and reported that adoptive T-cell therapy has significant beneficial therapeutic impact.

Chemokines are a family of chemotactic cytokines that regulate immune cell activation and migration under normal and inflammatory conditions (13). The binding of chemokines to 7 transmembrane binding receptors and multi-level conduction facilitates intracellular delivery of activation signals for cell migration (14). A recent study reported that chemokines and their receptors are involved in tumor growth, tumor cell invasion and distant metastasis (5). C-X-C motif chemokine receptor (CXCR) subunits belong to an important subfamily of chemokines. High expression levels of CXCR subunits were detected in tumor tissues from diverse tumor types, including lung adenocarcinoma, gastric cancer, colorectal cancer and hepatocellular carcinoma (15-18). Another recent study reported the prognostic values of CXCR subunits in gastric cancer (19); however, the potential prognostic values of CXCR subunits in PDAC remains unclear. The present study aimed to examine the potential regulation pathway of CXCR subunits in network enrichment analysis and the prognostic value of CXCR subunits in patients with early-stage PDAC.

Materials and methods

Expression levels of CXCR subunits in tissues and enrichment analyses. The expression levels of CXCR subunits in normal human tissues were analyzed in Genotype-Tissue Expression projects (GTEx) (gtexportal.org/). The comparison of CXCR subunit expression levels in pancreatic adenocarcinoma (PAAD) tumor and non-tumor tissues were performed using Gene Expression Profiling Interactive Analysis (GEPIA; gepia.cancer-pku.cn/) (20), which provides comprehensive analysis of the expression profiling data from The Cancer Genome Atlas (TCGA) and GTEx projects. The enrichment analysis of Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways and annotation of Gene Ontology (GO) terms was performed using Database for Annotation, Visualization, and Integrated Discovery version 6.8 (david.ncifcrf.gov/home.jsp) (21).

In order to determine potential drug-target interaction profiles in CXCR subunits, the interactions between chemical compounds and proteins from the Search Tool for Interactions of Chemicals (STITCH) version 5.0 database (stitch.embl.de/) (22) were analyzed. To investigate the association between CXCR subunit mRNA expression levels, correlation analyses were performed using the *Corrplot* R package version 0.84 (cran.r-project.org/web/packages/corrplot/).

Survival analysis and prognostic model construction. The data on the expression levels of all CXCR subunits and the relevant clinical parameters were obtained from the UCSC Xena browser (xena.ucsc.edu/). Since the TCGA PAAD cohort contains a variety of histopathological types, inclusion and exclusion criteria of early-stage PDAC samples in TCGA were

processed as described in previous studies (23,24). Patients who underwent pancreaticoduodenectomy and whose clinical pathology was diagnosed as American Joint Committee on Cancer 7th stage (25) I or II PDAC were enrolled in the survival analysis. Then, a total of 112 PDAC cases with complete survival data were analyzed in the study. The median expression levels of the CXCR subunit was used as the cut-off for classifying patients into high- and low-expression groups. Estimation of survival distribution and overall survival (OS) and disease-free survival (DFS) times were performed using Kaplan-Meier method and analyzed using the log-rank test. A Cox proportional hazard model was established and used to calculate hazard ratios (HRs) and 95% confidence intervals (CIs) to identify independent prognostic predictors. The association between the prognosis of patients with early-stage PDAC and CXCR subunit expression levels was identified using combined survival analysis.

The *survivalROC* R package (cran.r-project.org/web/packages/survivalROC/) version 1.0.3 was used to construct time-dependent receiver operating characteristic (ROC) curves and further evaluate the predictive accuracy of CXCR subunits in the clinical outcomes of patients with PDAC. All area under the ROC curve (AUC) values were calculated. Further, a prognostic nomogram model was constructed using the *rms* (26) R package, according to the selection of prognosis-associated variables in a univariate Cox proportional hazards regression model.

Gene set enrichment analysis (GSEA). To investigate the potential mechanisms underlying the association of CXCR subunits between high- and low-expression groups and the prognosis of patients with PDAC, *c2* (*c2.all*, version 7.0, symbols) was used for KEGG pathway analysis and *c5* (*c5.all.v7.0*, symbols) was used for GO term analysis of the Molecular Signatures Database for GSEA (software.broadinstitute.org/gsea/index.jsp). $P < 0.05$ was considered to indicate a statistically significant difference.

Comprehensive analysis of tumor-infiltrating immune cells. The correlation between 6 immune infiltrates (B cells, CD4⁺ T cells, CD8⁺ T cells, neutrophils, macrophages and dendritic cells) and CXCR subunit expression levels in patients with PAAD was identified using the Tumor Immune Estimation Resource (TIMER) (27) and was analyzed using the purity-corrected partial Spearman method. The infiltration level for each somatic copy number alteration category was compared using the normal using two-sided Wilcoxon rank sum test, followed by Bonferroni's correction. The somatic copy number alteration categories were defined using GISTIC version 2.0 (28). Moreover, fractions of tumor-infiltrating immune cells in TCGA patients with PDAC were obtained from supplementary materials of a previously published literature (29). In addition, the sample IDs of all PDAC cases were matched with the sample IDs of the selected samples in the aforementioned steps for further analysis and other non-PDAC samples were excluded.

Statistical analysis. All statistical analyses were performed using SPSS version 24.0 (IBM Corp.) and R version 3.4.1 (R Foundation for Statistical Computing, Vienna, Austria).




Figure 1. Expression analysis of CXCR subunits. (A) Expression levels of CXCR subunits in The Cancer Genome Atlas pancreatic tumor and non-tumor tissues. Comparisons between groups were performed using unpaired Student's t-test. (B) Heatmap analysis of CXCR subunits expression in Genotype-Tissue Expression normal tissues. Red box indicates normal pancreatic tissue expression and the color of the cell indicates the level of expression. *P<0.05. CXCR, C-X-C motif chemokine receptor; T, tumor; N, normal; TPM, transcripts per kilobase million.

ISBN 3-900051-07-0, URL <http://www.R-project.org/>). The correlation of metric data was analyzed using Pearson's correlation coefficient. The comparison of CXCR subunit expression levels in PAAD tumor and non-tumor tissues were performed using a Student's t-test. P<0.05 was considered to indicate a statistically significant difference. All Kaplan-Meier survival curves were constructed using GraphPad Prism version 7.01 (GraphPad Software, Inc.).

Results

Expression levels of CXCR subunits in tumor and normal tissues and enrichment analysis. Public databases indicated that the expression levels of CXCR subunits (CXCR3, CXCR4, CXCR5 and CXCR6) were significantly elevated in patients with PAAD compared with healthy patients (Fig. 1A). The heatmap of the GTEx project showed that CXCR4 expression levels were slightly elevated in normal tissues, notably in whole blood and spleen (Fig. 1B). In normal pancreatic tissues, all CXCR subunits displayed a low expression levels. The enrichment analysis of CXCR subunits shows that the subunits may be involved in 'cytokine-cytokine receptor interaction', 'C-X-C chemokine receptor activity' and the 'G-protein coupled receptor signaling pathway' (Fig. 2A). Using the STITCH database, perlixafor was identified as a potential chemical

drug that acts on CXCR4, CXCL12 and CXCR6 (Fig. 2B). Furthermore, correlation analysis of each two CXCR subunits expression levels in PDAC presented a relation with a positive trend (Fig. 2C). The aforementioned results indicated that CXCR subunit expression levels are low in normal tissues and highly expressed in pancreatic cancer tissues. These CXCR expression levels were positively correlated with each other in PDAC cancer and para-cancer tissues and could be novel targets for pancreatic cancer treatment.

Survival analysis of CXCR subunit expression levels in patients with early-stage PDAC. The association between clinicopathological parameters and the prognosis of patients with early-stage PDAC in TCGA database is shown in Table SI. The survival curves of the association between CXCR subunit expression levels and prognosis of patients with PDAC are displayed in Fig. 3. In the univariate survival analysis, high expression level groups of CXCR3, CXCR4 and CXCR6 were associated with a more favorable OS time in patients with early-stage PDAC (P=0.033, 0.018 and 0.005, respectively; Fig. 3A-C). Compared with low expression levels of the CXCR subunits, PDAC patients with high CXCR3, CXCR4 and CXCR6 expression levels have longer OS times (HR=0.59, 0.56 and 0.49, respectively; Table I). Moreover, low CXCR2 and CXCR6 expression levels in patients with early-stage




Figure 3. Survival analysis of different CXCR subunit expression levels in patients with early-stage pancreatic ductal adenocarcinoma. (A-C) Survival curves of OS time for high and low CXCR3, CXCR4 and CXCR6 expression level groups, respectively. (D and E) Survival curves of DFS time for high and low CXCR2 and CXCR6 expression levels groups, respectively. (F and G) Survival curves of OS and DFS times for different combination of CXCR3, CXCR4 and CXCR6 expression level groups. (H and I) Survival curves of OS and DFS for combination of CXCR2 and CXCR6 expression groups. CXCR, C-X-C motif chemokine receptor; OS, overall survival; DFS, disease-free survival.

in patients with PDAC. The AUCs of time-independent ROC curves were 0.283-0.461 in different survival time (Fig. S1A and B). The time-dependent ROC curves are univariate analyses that require a consideration of clinical case information to improve accuracy. Therefore, a prognosis nomogram for OS and DFS time of patients with PDAC was constructed to predict the prognostic risk of different CXCR subunits mRNA levels and clinical characteristics and analyze its contribution to prognosis prediction. In the OS time nomogram model, low CXCR3 and CXCR6 expression levels were adverse factors for long-term OS time (Fig. 4A), which is consistent with the aforementioned results. Moreover, low CXCR2 expression levels, high pathological stage, high neoplasm histological grade and non-radical resection were the primary factors affecting DFS time in patients with PDAC (Fig. 4B). The aforementioned results showed that combined CXCR subunits expression levels and clinical data can predict postoperative clinical outcomes of patients with PDAC for further improvement of treatment plans.

GSEA report in patients with early-stage PDAC. The GSEA report suggested that CXCR subunits in the genome-wide expression profile dataset of the TCGA PDAC cohort serves a role in ‘immune response’ and ‘antigen binding’ (Fig. 5). Thus, CXCR subunits are associated with the cellular immune antigen presentation process for tumor cell recognition. Furthermore, CXCR subunits function in the regulation of

the immune cell activation and facilitate the adhesion and differentiation of immune cells (Fig. S2A-E), which may help immune cells to bind, recognize and destroy tumor cells to promote apoptosis.

Association of tumor infiltration levels with CXCR subunits expression levels in PDAC. The association between tumor infiltration levels with CXCR subunit expression levels were detected using TIMER and TCGA. Using heatmap clustering analysis, it was demonstrated that Th1, Th2 and TH17 cell fractions were associated with CXCR subunit expression levels, and high CXCR1 and CXCR2 expression levels were associated with less Th1 cell fractions and more Th2 and TH17 cell fractions. High CXCR3, CXCR4 and CXCR5 expression levels were associated with more Th1 cell fractions, less Th2 and TH17 cell fractions (Fig. 6A). Similarly, a number of immune cells in PDAC tumor tissues were associated with CXCR subunits expression levels: High CXCR1 and CXCR2 expression levels were associated with more neutrophils, eosinophils and dendritic cell fractions; high CXCR3, CXCR4 and CXCR5 expression levels were associated with more lymphocyte cell fractions and fewer macrophages cell fractions (Fig. 6B and C). In the purity-corrected partial Spearman's correlation analysis, CXCR subunits (CXCR2, CXCR3, CXCR4 and CXCR6) were positively correlated with tumor infiltration levels of 6 immune infiltrates, including B cells, CD4⁺ T cells, CD8⁺ T cells, neutrophils, macrophages

Table I. Survival analysis of the CXCR subunit expression levels in patients with early-stage PDAC.

Subunit	MST	OS						DFS						
		P-value	HR crude (95% CI)	P-value	Coefficient B ^a	HR adjusted (95% CI) ^b	P-value	MRT	P-value	HR crude (95%CI)	P-value	Coefficient B ^a	HR adjusted (95% CI) ^b	P-value
CXCR1		0.654												
Low	511		Ref.	0.654		Ref.	593	0.466	Ref.	0.467		Ref.		0.124
High	518		0.90 (0.55-1.45)		-0.110	0.75 (0.44-1.28)	716		0.78 (0.40-1.52)		-0.248	0.55 (0.25-1.18)		
CXCR2		0.056												
Low	476		Ref.	0.058		Ref.	581	0.006	Ref.	0.008		Ref.		0.008
High	596		0.63 (0.39-1.02)		-0.469	0.66 (0.38-1.16)	872		0.39 (0.20-0.78)		-0.939	0.32 (0.14-0.74)		
CXCR3		0.033												
Low	393		Ref.	0.035		Ref.	716	0.500	Ref.	0.501		Ref.		0.292
High	603		0.59 (0.37-0.97)		-0.522	0.61 (0.36-1.04)	620		0.78 (0.39-1.59)		-0.243	0.65 (0.30-1.45)		
CXCR4		0.018												
Low	473		Ref.	0.019		Ref.	593	0.543	Ref.	0.544		Ref.		0.692
High	592		0.56 (0.34-0.91)		-0.586	0.51 (0.28-0.91)	620		0.81 (0.41-1.59)		-0.208	0.84 (0.36-1.96)		
CXCR5		0.231												
Low	593		Ref.	0.233		Ref.	461	0.057	Ref.	0.062		Ref.		0.412
High	517		0.74 (0.46-1.21)		-0.296	0.79 (0.45-1.38)	872		0.50 (0.24-1.04)		-0.694	0.71 (0.31-1.62)		
CXCR6		0.005												
Low	458		Ref.	0.005		Ref.	443	0.027	Ref.	0.030		Ref.		0.237
High	603		0.49 (0.30-0.80)		-0.715	0.53 (0.31-0.92)	831		0.44 (0.21-0.93)		-0.819	0.60 (0.26-1.40)		

^aB, regression coefficient calculated using univariate Cox proportional hazards regression analysis. ^bAdjusted for pathological stage T and N stage, histological grade, radical resection and targeted molecular therapy. CXCR, C-X-C motif chemokine receptor; TCGA, The Cancer Genome Atlas; PDAC, pancreatic ductal adenocarcinoma; OS, overall survival time; DFS, disease-free survival time; MST, median survival time; MRT, median disease-free survival time; HR, hazard ratio; CI, confidence interval; Ref., reference.

Table II. Combined survival analysis of the CXCR subunits in patients with early-stage pancreatic ductal adenocarcinoma.

Variables	OS						DFS					
	MST	P-value	HR crude (95% CI)	P-value	HR adjusted (95% CI) ^a	P-value	MRT	P-value	HR crude (95% CI)	P-value	HR adjusted (95% CI) ^a	P-value
CXCR3+4+6		0.007						0.007				
All low	393		Ref.	0.011	Ref.	0.022	593		Ref.	0.013	Ref.	0.018
1 high	458		0.84 (0.44-1.62)	0.609	0.89 (0.43-1.83)	0.751	291		3.03 (1.07-8.57)	0.037	3.75 (1.15-12.18)	0.028
2 high	467		0.72 (0.35-1.46)	0.360	0.69 (0.32-1.49)	0.346	486		1.45 (0.46-4.54)	0.527	1.05 (0.30-3.63)	0.937
All high	913		0.29 (0.13-0.64)	0.002	0.25 (0.10-0.65)	0.004	872		0.75 (0.24-2.35)	0.623	0.76 (0.19-3.05)	0.696
CXCR2+6		0.013						0.020				
Low/low	381		Ref.	0.017	Ref.	0.087	439		Ref.	0.030	Ref.	0.057
Low/high	603		0.49 (0.24-0.99)	0.048	0.84 (0.37-1.92)	0.683	620		0.62 (0.24-1.61)	0.324	1.38 (0.43-4.37)	0.589
High/low	614		0.66 (0.34-1.28)	0.219	1.08 (0.51-2.27)	0.842	NA		0.52 (0.19-1.48)	0.222	0.46 (0.14-1.51)	0.199
High/high	596		0.36 (0.19-0.69)	0.002	0.44 (0.22-0.91)	0.026	872		0.25 (0.10-0.63)	0.003	0.29 (0.10-0.85)	0.024

^aAdjusted for pathological T and N stage, histological grade, radical resection and targeted molecular therapy. CXCR, C-X-C motif chemokine receptor; OS, overall survival time; DFS, disease-free survival time; MST, median survival time; MRT, median disease-free survival time; HR, hazard ratio; CI, confidence interval; NA, not available; Ref., reference.




Figure 4. Nomogram model constructed for CXCR subunits expression level for (A) overall survival and (B) disease-free survival in patients with early-stage pancreatic adenocarcinoma. To determine the value of each variable on the point axis, a vertical line is drawn on the total points axis according to the total value of variables and the predicted survival probability in the time-evaluated line. CXCR, C-X-C motif chemokine receptor.

and dendritic cells, in patients with PDAC, Spearman's correlation coefficient range from 0.234-0.54 ($P < 0.05$, Fig. 7). A comparison of tumor infiltration levels among tumors with different somatic copy number alterations for CXCR subunits also showed a significant correlation, different copy number alterations of CXCR subunits were mainly different in the infiltration levels of $CD4^+$ T cells ($P < 0.05$, Fig. S3).

Discussion

In the present study, the differential expression levels of CXCR subunits in patients with PDAC were identified using data from TCGA. High CXCR3, CXCR4 and CXCR6 expression levels were associated with a favorable OS in patients with early-stage PDAC. In addition, a significant association was found between high CXCR2 and CXCR6 expression levels and favorable DFS in patients with PDAC. The prognostic ability of CXCR subunits was evaluated using time-independent ROC

curves. A prediction model for the prognosis of patients with PDAC was then constructed using CXCR subunit expression levels and clinical features. Moreover, the results of GSEA analysis showed that CXCR subunit expression levels functioned in the regulation of immunity in PDAC. Heatmap clustering demonstrated that immune cell fragments in tumors were associated with CXCR expression levels. Expression level and somatic copy number alterations of CXCR subunits (CXCR2, CXCR3, CXCR4 and CXCR6) were positively correlated with tumor infiltration levels in pancreatic cancer. It was predicted that perixafor was associated with CXCR6, CXCR4 and its ligand CXCL12, using drug target network analysis.

In the present results of tumor infiltration levels, CXCR subunits expression were positively associated with infiltration levels of tumor-infiltrating immune cells in PDAC tissues, indicating that immune cells were mediated by chemokines. In the tumor microenvironment, a variety of cells in the tumor release different chemokines, leading to the recruitment and




Figure 5. Gene set enrichment analysis in patients with early-stage PDAC. Gene Set Enrichment Analysis reports for high C-X-C motif chemokine receptor subunit expression levels using high- vs. low-risk groups. Enrichment plots for: (A) Antigen binding, (B) adaptive immune response based on the somatic recombination of immune receptors built from immunoglobulin superfamily domains, (C) regulation of leukocyte-mediated immunity and (D) lymphocyte-mediated immunity. These panels provide the genome-wide potential pathways and molecular mechanisms in patients with PDAC based on gene expression. NES, normalized enrichment score; PDAC, pancreatic ductal adenocarcinoma; GO, Gene Ontology.

activation of different immune cells, modulating the balance between tumor-promoting and anti-tumor responses (30). CXCR subunits are expressed to coordinate leukocyte trafficking under both physiological and pathological conditions (31).

Tumor-associated neutrophils serve a role in the early stages of cancer, advanced progression and therapeutic drug resistance (32-34). The inhibition of CXCR2 expression in pancreatic tumors prevents neutrophil accumulation, leading to tumor growth under T cell-dependent suppression. Activated and functionalized T cells infiltrate pancreatic cancer in the absence of neutrophils (35). Ijichi *et al* (36) found that tumor-stromal interactions via a CXCR2-dependent chemokine can regulate the progression of PDAC. Idorn *et al* (37) identified that CXCR2, as a candidate for chemokine receptor transduction, can improve the recruitment of transduced tumor ascites lymphocytes toward the ovarian cancer microenvironment. However, the function of CXCR2 may be different in immune cells to regulate the balance between antitumor and

protumor responses. A recent study demonstrated that the CXC chemokine-receptor axis was associated with the invasion and migration of PDAC cells and that blockade of this axis prolonged survival and inhibited both tumor angiogenesis and PDAC microinvasion, following CXCR2 knockout in PKF mice (38). Moreover, high expression levels of CXCR2 were associated with a favorable DFS in patients with early-stage PDAC in the present study and the prognostic prediction model indicated that low CXCR2 expression level was a prognostic indicator for less favorable DFS.

A previous study reported that knockdown of CXCR2 diminished the DNA-damage response and CXCR2-binding chemokines reinforced growth arrest in senescent cells (39). In a further study, it was revealed that the overall effect of CXCR2 signaling was involved in the balance between tumor suppression and tumorigenicity, which induced senescence in benign lesions; however, this is reversed in more advanced tumors (40). It was suggested that the effect of CXCR2 signaling was determined by the stage and pathological status




Figure 6. Associations between CXCR subunit expression levels, clinical features and immune cell fractions in pancreatic ductal adenocarcinoma tumor tissues. (A) Helper cells. (B) Overall immune cells. (C) Immune cell subsets. Red to blue: High abundance of immune infiltration to low abundance. Various colored side boxes were used to characterize the age, sex, histological grade, targeted molecular therapy and CXCR subunit expression levels. Colors from light to dark indicate that the value of the continuity variable is from low to high. Multiple colors represent two or multiple categorical variables. CXCR, C-X-C motif chemokine receptor.

of the lesion, as well as its origin and genetic background. Studies performed in animal models require an understanding of the complex contribution of CXCR2 towards tumor progression in cell experiments. Intrinsic causes of the differences in results of CXCR2 towards tumor progression *in vitro* and *in vivo* experiments needs further verification to clarify the molecular mechanisms underpinning these differences.

In the present study, high expression levels of CXCR3 were associated with an improved OS in patients with early-stage PDAC. Studies have reported that agonists of CXCR3 promote

the recruitment of NK cells, CD4⁺ and CD8⁺ T lymphocytes to the tumor microenvironment, where they exert potent antitumor activity (30,41). The present study also identified that CXCR3 expression levels were positively correlated to tumor infiltration levels in PDAC and that CXCR3 expression levels were elevated in tumor tissues, enhancing the antitumor activity of immune cells.

In the previous studies, CXCR4 was upregulated in cancer tissues and extensively expressed in pancreatic cancer cell lines (42,43). In the SDF-1 α /CXCR4 axis, the migratory




Figure 7. Correlation of (A) CXCR2, (B) CXCR3, (C) CXCR4 and (D) CXCR6 expression levels with immune infiltration level in pancreatic cancer. The scatterplots show the purity-corrected partial Spearman's correlation and statistical significance. The abscissa and ordinate represent CXCR subunit expression levels of tumor tissues and tumor infiltration levels of immune cells, respectively. CXCR, C-X-C motif chemokine receptor; PAAD, pancreatic adenocarcinoma; RSEM, RNA-Seq by Expectation-Maximization; cor, correlation.

potential and invasion activity of pancreatic cancer cells were modulated by the induction of CXCR4 expression (42). Moreover, CXCR4 and its ligand CXCL12 are involved in the metastasis of different types of tumors, such as breast, prostate, lung and colorectal cancer (43). CXCR4 is important in the classical chemokine receptor response in adults, as well as neutrophil maturation (44). Maréchal *et al* (45) reported that patients with PAAD and high level of CXCR4 expression in tumor tissues had a shorter OS time compared with those with low CXCR4 expression levels, using immunohistochemistry. The group demonstrated that CXCR4 was expressed in 84.5% of PAAD tumor tissues, but CXCR4 expression levels were undetectable in 6 samples with a low proportion (<30%) of adenocarcinoma cells, indicating that CXCR4 expression levels are different in variable pathological types of PAAD. In contrast, Gebauer *et al* (46) reported that CXCR4 expression levels evaluated using immunohistochemistry in PAAD specimens were not associated with OS and DFS times. Gebauer *et al* (46) demonstrated that CXCR4 was expressed in 214 patients with PAAD (86.0%) and >90% patients received adjuvant gemcitabine chemotherapy, whereas Maréchal *et al* (45) reported only 17% of patients with PAAD to receive adjuvant chemotherapy. In the present study, high expression levels of CXCR4 were detected using RNA-sequencing of tumor tissues and were associated with longer OS in early-stage ductal adenocarcinoma type of PAAD. It was hypothesized that different types of histology

and adjuvant therapy may lead to differences in CXCR subunit expression levels. The present results demonstrated that CXCR4 expression levels were positively correlated with tumor infiltration levels of PDAC. Studies have reported that CXCR4 knock-out mice are embryonically lethal due to multiple organ failure, such as hematopoiesis damage (47,48). In addition, the high levels of immune cell infiltration in the microenvironment recruited by the CXCL12/CXCR4 axis is positively correlated with a favorable prognosis in breast, ovarian and cervical cancer (49-51). More studies are required to evaluate the prognostic value of CXCR4 in patients with PDAC.

A positive correlation was also identified in the present study between CXCR6 and immune cell-infiltrate levels in pancreatic cancer. High expression levels of CXCR6 were associated with a favorable OS and DFS in patients with early-stage PDAC. However, a number of studies have been reported where high CXCR6 expression levels indicated a less favorable clinical outcome for patients with gastric cancer and clear cell renal cell carcinoma (52,53). In the present study, it was hypothesized that the prognostic value of CXCR6 was associated with the type and stage of the tumor. Late-stage PDAC is characterized by the matrix portion that forms a thick layer of connective tissue across the epithelial portion of the tumor, functioning as a physical barrier mediating chemotherapy resistance and hindering T-cell migration (54,55). Using the nomogram model to predict the clinical outcomes of patients with PDAC, it was demonstrated that the prognostic

values of CXCR6 expression levels were different for OS and DFS. Based on the multivariate and combination survival analysis, high CXCR6 expression levels were associated with long-term OS and DFS of patients with PDAC in the present study. It was hypothesized that the prognostic value of CXCR may be associated with clinical information in the model, which needs further verification.

Using drug target network analysis, it was identified that plerixafor was associated with CXCR6, CXCR4 and its ligand CXCL12. Plerixafor is the only approved drug that targets CXCR4 and has been used in combination therapy clinical trials for other types of cancer, such as prostate and cervical cancer (56,57). Correlation analysis of the present study identified the association between tumor-associated immune cell-infiltrates levels, CXCR subunit expression levels and somatic copy number alterations, indicating that CXCR subunits are potential therapeutic targets for immunotherapy in PDAC. However, this needs further verification using *in vitro* and *in vivo* experiments to clarify the levels of tumor-associated immune cell infiltrates in PDAC tissues and associations between immunotherapy drugs and CXCR subunits in PDAC.

There are several limitations of the present study. Firstly, there was the lack of external and experimental validation of the results. Secondly, some important clinical baseline information and the postoperative adjuvant treatments in TCGA database were not available, such as chemotherapy and primary tumor location. Thirdly, the present study evaluated the association between CXCR subunits and tumor infiltration levels; however, no bioinformatic platforms were found for investigating the depth of PDAC cell infiltration. Moreover, an independent cohort is required to verify the constructed prognostic models and larger samples are needed for a more reliable prognostic assessment.

In conclusion, CXCR subunits are associated with infiltration levels of immune cells and the prognosis of patients with early-stage PDAC and these subunits may be potential drug targets for the treatment of pancreatic cancer.

Acknowledgements

Not available.

Funding

This work was supported by The National Nature Science Foundation of China (grant. nos. 81160457 and 81460747).

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

QW and LL designed the study. CY, QW, LR and JL performed data collection and integration. QW, CY, LR and JL interpreted the data. QW and CY wrote the manuscript. LL critically revised the manuscript and participated in the analysis and interpretation of the data. All authors read and approved the final manuscript.

Ethics approval and consent to participate

This study was approved by The Ethics Committee of the First Affiliated Hospital of Guangxi University of Chinese Medicine (Guangxi, China) (approval no. GXTCMU-KY-2-072).

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Siegel RL, Miller KD and Jemal A: Cancer statistics, 2017. *CA Cancer J Clin* 67: 7-30, 2017.
2. Chen W, Zheng R, Baade PD, Zhang S, Zeng H, Bray F, Jemal A, Yu XQ and He J: Cancer statistics in China, 2015. *CA Cancer J Clin* 66: 115-132, 2016.
3. Tanaka S: Molecular pathogenesis and targeted therapy of pancreatic cancer. *Ann Surg Oncol* 23 (Suppl 2): S197-S205, 2016.
4. Hidalgo M, Cascinu S, Kleeff J, Labianca R, Löhner JM, Neoptolemos J, Real FX, Van Laethem JL and Heinemann V: Addressing the challenges of pancreatic cancer: Future directions for improving outcomes. *Pancreatology* 15: 8-18, 2015.
5. D'Agostino G, Cecchinato V and Uguccioni M: Chemokine heterocomplexes and cancer: A novel chapter to be written in tumor immunity. *Front Immunol* 9: 2185, 2018.
6. Weiss SA, Han SW, Lui K, Tchack J, Shapiro R, Berman R, Zhong J, Krogsgaard M, Osman I and Darvishian F: Immunologic heterogeneity of tumor-infiltrating lymphocyte composition in primary melanoma. *Hum Pathol* 57: 116-125, 2016.
7. Tumeh PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, Chmielowski B, Spasic M, Henry G, Ciobanu V, *et al*: PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* 515: 568-571, 2014.
8. Fridman WH, Pages F, Sautès-Fridman C and Galon J: The immune contexture in human tumours: Impact on clinical outcome. *Nat Rev Cancer* 12: 298-306, 2012.
9. Becht E, Giraldo NA, Dieu-Nosjean MC, Sautès-Fridman C and Fridman WH: Cancer immune contexture and immunotherapy. *Curr Opin Immunol* 39: 7-13, 2016.
10. Tahkola K, Mecklin JP, Wirta EV, Ahtiaainen M, Helminen O, Böhm J and Kellokumpu I: High immune cell score predicts improved survival in pancreatic cancer. *Virchows Arch* 472: 653-665, 2018.
11. Lianyan T, Dianrong X, Chunhui Y, Zhaolai M and Bin J: The predictive value and role of stromal tumor-infiltrating lymphocytes in pancreatic ductal adenocarcinoma (PDAC). *Cancer Biol Ther* 19: 296-305, 2018.
12. Poschke I, Faryna M, Bergmann F, Flossdorf M, Lauenstein C, Hermes J, Hinz U, Hank T, Ehrenberg R, Volkmar M, *et al*: Identification of a tumor-reactive T-cell repertoire in the immune infiltrate of patients with resectable pancreatic ductal adenocarcinoma. *Oncoimmunology* 5: e1240859, 2016.
13. Martins-Green M, Petreaca M and Wang L: Chemokines and their receptors are key players in the orchestra that regulates wound healing. *Adv Wound Care (New Rochelle)* 2: 327-347, 2013.
14. Palomino DCT and Marti LC: Chemokines and immunity. *Einstein (Sao Paulo)* 13: 469-473, 2015 (In English, Portuguese).
15. Saintigny P, Massarelli E, Lin S, Ahn YH, Chen Y, Goswami S, Erez B, O'Reilly MS, Liu D, Lee JJ, *et al*: CXCR2 expression in tumor cells is a poor prognostic factor and promotes invasion and metastasis in lung adenocarcinoma. *Cancer Res* 73: 571-582, 2013.
16. Wang Z, Liu H, Shen Z, Wang X, Zhang H, Qin J, Xu J, Sun Y and Qin X: The prognostic value of CXCR2-chemokine receptor 2 (CXCR2) in gastric cancer patients. *BMC Cancer* 15: 766, 2015.
17. Xu C, Zheng L, Li D, Chen G, Gu J, Chen J and Yao Q: CXCR4 overexpression is correlated with poor prognosis in colorectal cancer. *Life Sci* 208: 333-340, 2018.

18. Gao Q, Zhao YJ, Wang XY, Qiu SJ, Shi YH, Sun J, Yi Y, Shi JY, Shi GM, Ding ZB, *et al*: CXCR6 upregulation contributes to a proinflammatory tumor microenvironment that drives metastasis and poor patient outcomes in hepatocellular carcinoma. *Cancer Res* 72: 3546-3556, 2012.
19. Yu C and Zhang Y: Characterization of the prognostic values of CXCR family in gastric cancer. *Cytokine* 123: 154785, 2019.
20. Tang Z, Li C, Kang B, Gao G, Li C and Zhang Z: GEPIA: A web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res* 45: W98-W102, 2017.
21. Dennis G Jr, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC and Lempicki RA: DAVID: Database for annotation, visualization, and integrated discovery. *Genome Biol* 4: P3, 2003.
22. Szklarczyk D, Santos A, Von Mering C, Jensen LJ, Bork P and Kuhn M: STITCH 5: Augmenting protein-chemical interaction networks with tissue and affinity data. *Nucleic Acids Res* 44: D380-D384, 2016.
23. Liao X, Huang K, Huang R, Liu X, Han C, Yu L, Yu T, Yang C, Wang X and Peng T: Genome-scale analysis to identify prognostic markers in patients with early-stage pancreatic ductal adenocarcinoma after pancreaticoduodenectomy. *Oncotargets Ther* 10: 4493-4506, 2017.
24. Yang C, Yu T, Liu Z, Ye X, Liao X, Wang X, Han C, Zhu G, Qin W and Peng T: Cystatin F as a key family 2 cystatin subunit and prognostic biomarker for early-stage pancreatic ductal adenocarcinoma. *Oncol Rep* 42: 79-90, 2019.
25. Edge SB and Compton CC: The American joint committee on cancer: The 7th edition of the AJCC cancer staging manual and the future of TNM. *Ann Surg Oncol* 17: 1471-1474, 2010.
26. Harrell FE Jr: Package 'rms': Regression modeling strategies. Package Version 5.1-4, 2016. <http://biostat.mc.vanderbilt.edu/wiki/Main/RmS>.
27. Li T, Fan J, Wang B, Traugh N, Chen Q, Liu JS, Li B and Liu XS: TIMER: A Web server for comprehensive analysis of tumor-infiltrating immune cells. *Cancer Res* 77: e108-e110, 2017.
28. Mermel CH, Schumacher SE, Hill B, Meyerson ML, Beroukhi R and Getz G: GISTIC2.0 facilitates sensitive and confident localization of the targets of focal somatic copy-number alteration in human cancers. *Genome Biol* 12: R41, 2011.
29. Thorsson V, Gibbs DL, Brown SD, Wolf D, Bortone DS, Ou Yang TH, Porta-Pardo E, Gao GF, Plaisier CL, Eddy JA, *et al*: The immune landscape of cancer. *Immunity* 48: 812-830.e14, 2018.
30. Chow MT and Luster AD: Chemokines in cancer. *Cancer Immunol Res* 2: 1125-1131, 2014.
31. Zlotnik A and Yoshie O: Chemokines: A new classification system and their role in immunity. *Immunity* 12: 121-127, 2000.
32. Coffelt SB, Wellenstein MD and de Visser KE: Neutrophils in cancer: Neutral no more. *Nat Rev Cancer* 16: 431-446, 2016.
33. Bronte V, Brandau S, Chen SH, Colombo MP, Frey AB, Greten TF, Mandruzzato S, Murray PJ, Ochoa A, Ostrand-Rosenberg S, *et al*: Recommendations for myeloid-derived suppressor cell nomenclature and characterization standards. *Nat Commun* 7: 12150, 2016.
34. Fridlender ZG and Albelda SM: Tumor-associated neutrophils: Friend or foe? *Carcinogenesis* 33: 949-955, 2012.
35. Chao T, Furth EE and Vonderheide RH: CXCR2-dependent accumulation of tumor-associated neutrophils regulates T-cell immunity in pancreatic ductal adenocarcinoma. *Cancer Immunol Res* 4: 968-982, 2016.
36. Ijichi H, Chytil A, Gorska AE, Aakre ME, Bierie B, Tada M, Mohri D, Miyabayashi K, Asaoka Y, Maeda S, *et al*: Inhibiting Cxcr2 disrupts tumor-stromal interactions and improves survival in a mouse model of pancreatic ductal adenocarcinoma. *J Clin Invest* 121: 4106-4117, 2011.
37. Idorn M, Olsen M, Halldórsdóttir HR, Skadborg SK, Pedersen M, Høgdaal C, Høgdaal E, Met Ó and Thor Straten P: Improved migration of tumor ascites lymphocytes to ovarian cancer microenvironment by CXCR2 transduction. *Oncoimmunology* 7: e141209, 2017.
38. Sano M, Ijichi H, Takahashi R, Miyabayashi K, Fujiwara H, Yamada T, Kato H, Nakatsuka T, Tanaka Y, Tateishi K, *et al*: Blocking CXCLs-CXCR2 axis in tumor-stromal interactions contributes to survival in a mouse model of pancreatic ductal adenocarcinoma through reduced cell invasion/migration and a shift of immune-inflammatory microenvironment. *Oncogenesis* 8: 8, 2019.
39. Acosta JC, O'Loughlin A, Banito A, Guijarro MV, Augert A, Raguz S, Fumagalli M, Da Costa M, Brown C, Popov N, *et al*: Chemokine signaling via the CXCR2 receptor reinforces senescence. *Cell* 133: 1006-1018, 2008.
40. Acosta JC and Jesús G: A role for CXCR2 in senescence, but what about in cancer? *Cancer Res* 69: 2167-2170, 2009.
41. Tosolini M, Kirilovsky A, Mlecnik B, Fredriksen T, Mauger S, Bindea G, Berger A, Bruneval P, Fridman WH, Pagès F and Galon J: Clinical impact of different classes of infiltrating T cytotoxic and helper cells (Th1, th2, treg, th17) in patients with colorectal cancer. *Cancer Res* 71: 1263-1271, 2011.
42. Wang J, Wang H, Cai J, Du S, Xin B, Wei W, Zhang T and Shen X: Artemin regulates CXCR4 expression to induce migration and invasion in pancreatic cancer cells through activation of NF- κ B signaling. *Exp Cell Res* 365: 12-23, 2018.
43. Zlotnik A, Burkhardt AM and Homey B: Homeostatic chemokine receptors and organ-specific metastasis. *Nat Rev Immunol* 11: 597-606, 2011.
44. Machado ID, Spatti M, Hastreiter A, Santin JR, Fock RA, Gil CD, Oliani SM, Perretti M and Farsky SH: Annexin A1 is a physiological modulator of neutrophil maturation and recirculation acting on the CXCR4/CXCL12 pathway. *J Cell Physiol* 231: 2418-2427, 2016.
45. Maréchal R, Demetter P, Nagy N, Berton A, Decaestecker C, Polus M, Closset J, Devière J, Salmon I and Van Laethem JL: High expression of CXCR4 may predict poor survival in resected pancreatic adenocarcinoma. *Br J Cancer* 100: 1444-1451, 2009.
46. Gebauer F, Tachezy M, Effenberger K, von Loga K, Zander H, Marx A, Kaifi JT, Sauter G, Izbicki JR and Bockhorn M: Prognostic impact of CXCR4 and CXCR7 expression in pancreatic adenocarcinoma. *J Surg Oncol* 104: 140-145, 2011.
47. Ma Q, Jones D, Borghesani PR, Segal RA, Nagasawa T, Kishimoto T, Bronson RT and Springer TA: Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc Natl Acad Sci USA* 95: 9448-9453, 1998.
48. Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, Kitamura Y, Matsushima K, Yoshida N, Nishikawa S, *et al*: The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature* 393: 591-594, 1998.
49. Schmidt M, Böhm D, Von Törne C, Steiner E, Puhl H, Pilch H, Lehr HA, Hengstler JG, Kölbl H and Gehrmann M: The humoral immune system has a key prognostic impact in node-negative breast cancer. *Cancer Res* 68: 5405-5413, 2008.
50. Nedergaard BS, Ladekar M, Nyengaard JR and Nielsen K: A comparative study of the cellular immune response in patients with stage IB cervical squamous cell carcinoma. Low numbers of several immune cell subtypes are strongly associated with relapse of disease within 5 years. *Gynecol Oncol* 108: 106-111, 2008.
51. Milne K, Köbel M, Kalloger SE, Barnes RO, Gao D, Gilks CB, Watson PH and Nelson BH: Systematic analysis of immune infiltrates in high-grade serous ovarian cancer reveals CD20, FoxP3 and TIA-1 as positive prognostic factors. *PLoS One* 4: e6412, 2009.
52. Jin JJ, Dai FX, Long ZW, Cai H, Liu XW, Zhou Y, Hong Q, Dong QZ, Wang YN and Huang H: CXCR6 predicts poor prognosis in gastric cancer and promotes tumor metastasis through epithelial-mesenchymal transition. *Oncol Rep* 37: 3279-3286, 2017.
53. Chang Y, Zhou L, Xu L, Fu Q, Yang Y, Lin Z and Xu J: High expression of CXCR6 chemokine receptor 6 associates with poor prognosis in patients with clear cell renal cell carcinoma. *Urol Oncol* 35: 675.e17-675.e24, 2017.
54. Neesse A, Michl P, Frese KK, Feig C, Cook N, Jacobetz MA, Lolkema MP, Buchholz M, Olive KP, Gress TM and Tuveson DA: Stromal biology and therapy in pancreatic cancer. *Gut* 60: 861-868, 2011.
55. Hartmann N, Giese NA, Giese T, Poschke I, Offringa R, Werner J and Ryschich E: Prevailing role of contact guidance in intrastromal T-cell trapping in human pancreatic cancer. *Clin Cancer Res* 20: 3422-3433, 2014.
56. Chaudary N, Pintilie M, Jelveh S, Lindsay P, Hill RP and Milosevic M: Plerixafor improves primary tumor response and reduces metastases in cervical cancer treated with radio-chemotherapy. *Clin Cancer Res* 23: 1242-1249, 2017.
57. Conley-LaComb MK, Semaan L, Singareddy R, Li Y, Heath EI, Kim S, Cher ML and Chinni SR: Pharmacological targeting of CXCL12/CXCR4 signaling in prostate cancer bone metastasis. *Mol Cancer* 15: 68, 2016.

