

Delphinidin induces cell cycle arrest and apoptosis in HER-2 positive breast cancer cell lines by regulating the NF- κ B and MAPK signaling pathways

AILIN WU^{1,2*}, YANFENG ZHU^{3*}, BIN HAN^{4*}, JIAYUAN PENG¹, XIAOMIN DENG¹,
WEI CHEN¹, JINGCHANG DU⁴, YU OU⁵, XIAOLI PENG⁴ and XIAOPING YU³

¹Basic Medical School, Chengdu Medical College, Chengdu, Sichuan 610500; ²Ministry of Science and Technology, The Second Affiliated Hospital of Chengdu Medical College, China National Nuclear Corporation 416 Hospital, Chengdu, Sichuan 610051; ³Graduate School; ⁴School of Public Health; ⁵Division of Planning and Finance, Chengdu Medical College, Chengdu, Sichuan 610500, P.R. China

Received May 10, 2020; Accepted June 25, 2021

DOI: 10.3892/ol.2021.13093

Abstract. Delphinidin is an anthocyanidin monomer, commonly found in vegetables and fruits, and has demonstrated antitumor effects in the HER-2-positive MDA-MB-453 breast cancer cell line, with low cytotoxicity on normal breast cells. However, the direct functional mechanisms underlying the effect of delphinidin on HER-2-positive breast cancer cells has not been fully characterized. In the present study, it was found that delphinidin could induce G₂/M phase cell cycle arrest by inhibiting the protein expression level of cyclin B1 and Cdk1 in HER-2-positive breast cancer cell lines. In addition, delphinidin promoted the mitochondrial apoptosis pathway by inhibiting the ERK and NF- κ B signaling pathway and activating the JNK signaling pathway. Therefore, delphinidin markedly suppressed the viability of the HER-2-positive breast cancer cell lines by modulating the cell cycle and inducing apoptosis. Overall, the findings from the present study demonstrated that delphinidin treatment could induce the mitochondrial apoptosis pathway in human HER-2-positive breast cancer cell lines, providing an experimental basis for the prevention and treatment of HER-2-positive breast cancer by flavonoids.

Introduction

Breast cancer is among the most common type among women worldwide (1). Human epidermal growth factor receptor (HER)-2 is overexpressed in 20-30% of all breast cancer types (2) and is a preferred target for treating HER-2-positive breast cancer. Trastuzumab and pertuzumab, two HER2-targeted monoclonal antibodies, are used as adjuvant therapy to treat metastatic HER-2-positive breast cancer when combined with docetaxel. However, only a small percentage of patients benefit from this therapeutic method (3). Thus, identifying novel agents or drug combination and developing new therapeutic strategies is urgently required.

Delphinidin is an anthocyanidin monomer commonly found in vegetables and fruits (4). The benefits of delphinidin have been associated with its antioxidant, anti-inflammatory, antiangiogenic and anticancer effects (5). Delphinidin contains 3 hydroxyl groups on its B ring, making its oxidative biological effect stronger compared with that in other flavonoid compounds, such as cyanidin and malvidin (5-7). Delphinidin has the strongest antioxidant effect among all anthocyanidins. It is considered to be the most beneficial anthocyanidin, which has become the focus of anti-cancer research in anthocyanin monomers. Our previous studies confirmed the anticancer activities of anthocyanidin and delphinidin-3-glucoside in breast cancer cell lines. Luo *et al* (8) used structural biology methods to demonstrate that delphinidin could dock with the HER-2 tyrosine kinase region in the HER-2-positive MDA-MB-453 breast cancer cell line using hydrophobic forces and hydrogen bonding. This inhibited competitive ATP binding to the HER-2 receptor, thus preventing the autophosphorylation of HER-2 and exerting its anti-cancer effect (8-11). In addition, the therapeutic effects of delphinidin in cancer have been associated with the inhibition of MAPK and NF- κ B signaling, and the activation of the AP-1 signaling pathway (12,13). Therefore, the purpose of the current study was to determine the *in vitro* biological effects of delphinidin in 2 HER-2-positive breast cancer cell lines.

We hypothesized that delphinidin could induce mitochondrial pathway apoptosis by modulating NF- κ B and MAPK

Correspondence to: Dr Xiaoli Peng, School of Public Health, Chengdu Medical College, 783 Xindu Avenue, Chengdu, Sichuan 610500, P.R. China
E-mail: xiaopeng967@sohu.com

Dr Xiaoping Yu, Graduate School, Chengdu Medical College, 783 Xindu Avenue, Chengdu, Sichuan 610500, P.R. China
E-mail: yuxp@cmc.edu.cn

*Contributed equally

Key words: delphinidin, HER-2-positive breast cancer, cell cycle arrest, apoptosis, NF- κ B signaling pathway, MAPK signaling pathway

signaling in the HER-2-positive MDA-MB-453 and BT-474 breast cancer cell lines. The results of the present study may indicate that delphinidin is a valuable therapeutic agent for patients with breast cancer. This may provide new insight for the prevention and treatment of HER-2-positive breast cancer using delphinidin.

Materials and methods

Cell culture and treatment. The HER-2-positive breast cancer cell lines (MDA-MB-453 and BT-474) were purchased from the Chinese Academy of Sciences. The MDA-MB-453 cell line was cultured in L-15 medium (cat. no. 11415-064; Gibco; Thermo Fisher Scientific, Inc.), while the BT-474 cell line was cultured in RPMI-1640 medium (cat. no. A1049101; Gibco; Thermo Fisher Scientific, Inc.) and both supplemented with 10% fetal bovine serum (cat. no. ES-009-B; EMD Millipore), and at 37°C in a humidified incubator with 5% CO₂. Delphinidin (cat. no. 43725; purity ≥95%) was dissolved in dimethyl sulfoxide (DMSO; cat. no. D2650) (both from Sigma-Aldrich; Merck KGaA) to a stock concentration of 80 mM. The DMSO was added to the medium at a final concentration of 0.1% as a control.

Antibodies and reagents. The anti-rabbit IgG, HRP-linked (cat. no. 7074), anti-mouse IgG, HRP-linked (cat. no. 7076), p21^{Waf1/Cip1} (cat. no. 2947), c-Raf (cat. no. 53745), phosphorylated(p)-c-Raf (cat. no. 9431), MEK1/2 (cat. no. 9126), p-MEK1/2 (cat. no. 9154), p44/42 MAPK (Erk1/2) (cat. no. 4695), p-p44/42 MAPK (Erk1/2) (cat. no. 4370), SAPK/JNK (cat. no. 252), p-SAPK/JNK (cat. no. 4668), Bax (cat. no. 5023S), Bcl-2 (cat. no. 4223S), PKCα (cat. no. 59754), IKKα (cat. no. 11930), IKKβ (cat. no. 8943), p-IKKα/β (cat. no. 2697), IκBα (cat. no. 4814), NF-κB/p65 (cat. no. 8242), GAPDH (cat. no. 5174) antibodies, and Alexa Fluor 594 conjugate (cat. no. 8889) and Alexa Fluor 488 phalloidin (cat. no. 8878) were purchased from Cell Signaling Technology, Inc. The cyclin B1 (cat. no. ab32053), Cdk1 (cat. no. ab133327), p-PKCα (cat. no. ab23513), p-IκBα (cat. no. ab12134), p-NF-κB/p65 (cat. no. ab131109) antibodies were purchased from Abcam. The β-actin (cat. no. TA-09) antibody was purchased from OriGene Technologies, Inc.

Cytotoxicity assay. The MDA-MB-453 and BT-474 breast cancer cell lines were seeded in 96-well microtiter plates (1x10⁴ cells per well), then treated with delphinidin at different concentrations (10, 20, 40, 80 and 160 μM) or treated with the vehicle (DMSO) for 48 h. Cell Counting Kit-8 (CCK-8; cat. no. CK04; Dojindo Molecular Technologies, Inc.) was used to evaluate the viability of the cells. The absorbance was read at 450 nm using a microplate reader (Powerwave XS; Bio Tek China). Then SPSS v21.0 (IBM Corp.) was used to calculate IC₅₀.

Cellular morphological observation. Approximately 2x10⁵ cells were seeded in 6-well plates, containing glass coverslips. The cells were treated with delphinidin at different concentrations (20, 40 and 80 μM) or with DMSO for 48 h. The cells were then fixed using 4% paraformaldehyde for 20 min and stained (for 3-5 min with hematoxylin and 1 min with

eosin), both at room temperature, according to the instructions of the H&E staining kit (cat. no. D006; Nanjing Jiancheng Bioengineering Institute). Finally, the cells were observed with a light microscope at x200 magnification.

Cell cycle analysis. Approximately 2x10⁵ cells were seeded in 6-well plates, containing glass coverslips. The cells were treated with the delphinidin at different concentrations (20, 40 and 80 μM) or treated with DMSO for 48 h. The cells were then harvested in a culture dish, washed with ice-cold PBS and fixed with 80% ice-cold ethanol in PBS at 4°C for 4 h. The cells were then pelleted in a cool centrifuge (120 x g at 4°C for 5 min), resuspended in cold PBS and stained according to the instructions of the cell cycle detection kit (cat. no. KGA512; KeyGen Biotech Co., Ltd.). Following incubation at 37°C for 30 min, bovine pancreatic RNase was added at a final concentration of 2 mg/ml with 20 mg/ml propidium iodide for 20 min at room temperature. A total of 2x10⁴ cells were analyzed using a BD FACS Canto flow cytometer (Becton, Dickinson and Company). The data was analyzed using GraphPad Prism 7 (GraphPad Software, Inc.).

Protein expression analysis. The adherent cell lines were collected in radio immunoprecipitation assay (RIPA; cat. no. P0013B; Beyotime Institute of Biotechnology) buffer using a cell scraper and protein expression was analyzed using western blot analysis. A protein BCA assay kit (cat. no. 23227; Thermo Fisher Scientific, Inc.) was used to quantify the protein concentration. The samples (50 μg) were separated using 10% SDS-PAGE and the proteins were transferred to PVDF membranes, then blocked with 5% Blotting-Grade Blocker solution (cat. no. 180-6404; Bio-Rad Laboratories, Inc.). Subsequently, the membranes were incubated with the primary antibodies (diluted 1:1,000) overnight at 4°C, washed 3 times with TBS containing 1% Tween-20, then incubated with the secondary antibodies (diluted 1:1,000) for 2 h at 37°C. Finally, chemiluminescence reagents (cat. no. WBKLS0500; EMD Millipore) were used to visualize the target proteins and densitometry was analyzed using Image J v1 (National Institutes of Health.). GAPDH or β-actin served as the loading control.

TUNEL assay. The cells were seeded in 6-well plates, containing glass coverslips and treated with delphinidin, then fixed with 4% paraformaldehyde for 20 min and permeabilized with 1X PBS + 0.1% Triton for 2 min both at 4°C. TUNEL staining was performed using the Dead End Fluorometric TUNEL system in accordance with the manufacturer's instructions (cat. no. G3250; Promega Corporation). A total of 50 μl TUNEL detection solution was added to the sample, then incubated for 60 min at 37°C in the dark. Subsequently, 10 μl DAPI was added to the samples and incubated at room temperature for 5 min. The samples were observed and recorded using a fluorescence microscope (IX71; Olympus Corporation) at a x200 magnification, from at least 5 fields of views for each sample. Then, ImageJ v1 (National Institutes of Health) to record TUNEL-positive cells. The nuclei stained with bright green fluorescence were considered to be TUNEL-positive cells.

Mitochondrial membrane potential assay. Approximately 2x10⁵ cells were seeded in 6-well plates and treated with




Figure 1. Delphinidin inhibits the viability of the epidermal growth factor receptor-2-positive breast cancer cell lines. (A) MDA-MB-453 and BT-474 cells were treated with 10, 20, 40 or 80 μM delphinidin or DMSO. Relative cell numbers were determined using a Cell Counting Kit-8 assay after 48 h. # $P < 0.05$ vs. control; ^ $P < 0.05$ vs. 20 μM ; + $P < 0.05$ vs. 40 μM ; * $P < 0.05$ vs. 80 μM . (B) The MDA-MB-453 and BT-474 cell lines were treated with 20, 40 and 80 μM delphinidin or DMSO. After 48 h, the cells were fixed, stained with H&E, then analyzed using a microscope (magnification, x200). Experiments were repeated at least 3 times. DMSO, dimethyl sulfoxide.

delphinidin at different concentrations (20, 40 and 80 μM) or treated with DMSO for 8 h. The cells were washed twice with PBS and incubated with L-15 or RPMI-1640 medium containing 10 μg fluorescent lipophilic cationic JC-10 (cat. no. KGA608; Nanjing KeyGen Biotech Co., Ltd.) dye for 30 min at 37°C in the dark. The stained cells were harvested, washed and resuspended with PBS, then analyzed using a BD FACS Canto II flow cytometer and BD FACS Canto clinical software v2.4 (Becton, Dickinson and Company, Inc.) immediately. The lower-right quadrants represent cells with low membrane potential, while the upper-right quadrants represent cells with high membrane potential.

Detection of apoptosis using flow cytometry. An Annexin V-FITC Apoptosis Detection kit (cat. no. KGA108; Nanjing KeyGen Biotech Co., Ltd.) was used to detect apoptosis, as previously described (14). Annexin V⁺/PI⁻ cells were considered early apoptotic and Annexin V⁺/PI⁺ cells were considered late apoptotic. Double-negative cells (lower-left quadrant) were considered to be live cells. Annexin V⁻/PI⁺ cells (upper-left quadrant) were considered to be necrotic cells.

Immunofluorescence. Approximately 2×10^5 cells were seeded in 6-well plates, containing glass coverslips and treated with delphinidin at different concentrations (20, 40 and 80 μM) or treated with DMSO for 48 h, then fixed with 4% paraformaldehyde for 20 min at room temperature and permeabilized with 1X PBS + 0.5% Triton for 30 min at room temperature. The slides were blocked with PBS + 10% FBS (cat. no. ES-009-B; EMD Millipore) and incubated with anti-NF- κB /p65 (diluted

1:200) in PBS for 12 h at 4°C, then washed twice with PBS, and incubated with the Alexa Fluor 594-conjugated secondary antibody (diluted 1:500) for 2 h at room temperature. Following which, the slides were washed twice with PBS and mounted with Prolong Gold with Alexa Fluor 488 Phalloidin (diluted 1:20) and DAPI (cat. no. C0065; Beijing Solarbio Science and Technology Co., Ltd.). Finally, the images were captured using a confocal microscope (magnification, x600; FV1200; Olympus, Corporation). The results were determined using ImageJ software v.1 (National Institutes of Health) (15).

Statistical analysis. All the data are presented as the mean \pm standard error of the mean, and each experiment was performed at least three times. The statistical analyses were performed using one-way ANOVA and the Tukey's post hoc test using SPSS version 21.0 (IBM Corp.). $P < 0.05$ was considered to indicate a statistically significant difference. GraphPad Prism 7 (GraphPad software, Inc.) was used to generate the figures.

Results

Delphinidin inhibits the viability of the MDA-MB-453 and BT-474 breast cancer cell lines. To understand the effects of delphinidin on the breast cancer cell lines, the inhibitory effect of delphinidin on the viability of the HER-2-positive MDA-MB-453 and BT-474 breast cancer cell lines was analyzed (Fig. 1A). Delphinidin inhibited the viability of the HER-2-positive MDA-MB-453 and BT-474 breast cancer cell lines in a dose-dependent manner, with IC₅₀ values of




Figure 2. Effect of delphinidin on the cell cycle in the epidermal growth factor receptor-2-positive breast cancer cell lines. The MDA-MB-453 and BT-474 cells were treated with 20, 40 and 80 μ M delphinidin or dimethyl sulfoxide for 48 h. (A) Analysis of the cell cycle using flow cytometry following staining with PI. $n=3$. (B) Cyclin B1, Cdk2 and p21^{WAF1/Cip1} protein expression levels in the MDA-MB-453 and BT-474 cell lines were analyzed using western blot analysis and analyzed using ImageJ. GAPDH served as the loading control. The data are presented as the mean \pm standard error of the mean from at least 3 independent experiments. # $P<0.05$ vs. control; * $P<0.05$ vs. 20 μ M; $\dagger P<0.05$ vs. 40 μ M.

41.42 and 60.92 μ M, respectively (Fig. 1A). Therefore, 20, 40 and 80 μ M concentrations of delphinidin were used in subsequent experiments.

Following treatment of the cells with different concentrations of delphinidin and staining of the cells with H&E, some of the cells detached and floated, prior to fixation, and the cell volume gradually decreased as the concentration of delphinidin increased (Fig. 1B).

Delphinidin promotes cell cycle progression in the MDA-MB-453 and BT-474 breast cancer cell lines. To determine whether delphinidin could affect cell cycle regulation in the MDA-MB-453 and BT-474 breast cancer cell lines, the effect of delphinidin on G₂/M phase distribution was analyzed using flow cytometry, following staining with PI. As shown in Fig. 2A, delphinidin treatment significantly inhibited G₂/M phase arrest in the HER-2-positive breast cancer cell lines.

Further analysis revealed that the distribution of the cells in G₂/M phase was 7.5, 10.58 and 13.58% in the MDA-MB-453 breast cancer cell line with 20, 40 and 80 μ M delphinidin, respectively. The distribution of cells in the G₂/M phase was 13.43, 17.43 and 23.53% in the BT-474 breast cancer cell line with 20, 40 and 80 μ M delphinidin, respectively.

As delphinidin inhibited G₂/M phase progression, the protein expression levels of key cell cycle regulators were analyzed in the MDA-MB-453 and BT-474 breast cancer cell lines using western blot analysis (Fig. 2B). The results demonstrated that in the MDA-MB-453 and BT-474 breast cancer cell lines, 20, 40 and 80 μ M delphinidin significantly reduced the protein expression level of cyclin B1 and cdk2, and increased the protein expression level of p21^{WAF1/Cip1} following treatment for 48 h.

Delphinidin induces apoptosis in the MDA-MB-453 and BT-474 breast cancer cell lines. The aforementioned experiments confirmed that delphinidin could induce G₂/M phase cell cycle arrest in the MDA-MB-453 and BT-474 breast cancer cell lines. To determine whether cell cycle changes could lead to apoptosis and whether delphinidin inhibited the proliferation of the breast cancer cell lines, TUNEL staining was performed. The DNA fragment levels in the MDA-MB-453 and BT-474 breast cancer cell lines notably increased following delphinidin treatment, due to increase in fluorescent staining, and the number of dead cells increased with increasing concentrations of delphinidin, compared with that in the control group (P<0.05; Fig. 3A). As shown in Fig. 3A and B, concentrations of 20, 40 and 80 μ M delphinidin resulted in apoptosis rates of 3.3, 5.4 and 20.2% in the MDA-MB-453 cell line, respectively. Apoptosis rates reached and 1.4, 4.4 and 7.9% in the BT-474 breast cancer cell line following treatment with 20, 40 and 80 μ M delphinidin, respectively.

Mitochondria are producers of ATP, and their transmembrane potential is formed by the displacement of the transmembrane protons in the process of ATP generation (16). A previous study has shown that irreversible apoptosis occurs in breast cancer cell lines with very low mitochondrial membrane potential (17). Flow cytometry results showed that the mitochondrial membrane potential of the MDA-MB-453 and BT-474 breast cancer cell lines decreased with increasing concentrations of delphinidin compared with that in the control group (UR and UL quadrants; Fig. 3C). At concentrations of 40 and 80 μ M delphinidin, the apoptosis rate of the MDA-MB-453 cells was 49.58 and 53.87%, and those of the BT-474 cells was 17.55 and 55.18%, respectively (Fig. 3D). These results demonstrated that delphinidin could induce apoptosis in MDA-MB-453 and BT-474 breast cancer cell lines.

Delphinidin induces mitochondrial pathway apoptosis by inhibiting the ERK signaling pathway and activating the JNK signaling pathway. To determine whether the MAPK pathway played a role in mediating the effects of delphinidin, the expression levels of the Bcl-2 protein family were analyzed, as well as proteins in the MAPK signaling pathway, such as c-Raf, MEK1/2, ERK1/2, JNK, p-c-Raf, p-MEK1/2, p-ERK1/2 and p-JNK in the BT-474 and MDA-MB-453 breast cancer cell lines. The protein expression levels of Bax and Bcl-2 was

increased and decreased, respectively with increasing concentrations of delphinidin in the MDA-MB-453 and BT-474 cell lines (P<0.05; Fig. 4A and B). In addition, compared with that in the control group, the ratio of Bcl-2/Bax decreased significantly in the 40 and 80 μ M delphinidin treatment group (P<0.05). Furthermore, the protein expression levels of p-c-Raf, p-MEK1/2 and p-ERK1/2 decreased, and the protein expression level of p-JNK increased, in the MDA-MB-453 and BT-474 cell lines, indicating that delphinidin induced mitochondrial pathway apoptosis by inhibiting ERK signaling and activating JNK signaling in the HER-2-positive breast cancer cell lines (Fig. 4C and D).

Delphinidin induces apoptosis by inhibiting the NF- κ B signaling pathway in the HER-2-positive breast cancer cell lines. A previous study suggested that anthocyanins could induce apoptosis in various cancer cell lines, including stomach, prostate, intestine and breast cancer, by blocking the NF- κ B signaling pathway (18). To investigate whether delphinidin reduced the viability of the HER-2-positive breast cancer cell lines by inhibiting NF- κ B signaling, the protein expression levels of NF- κ B signaling-related molecules in the MDA-MB-453 and BT-474 breast cancer cell lines were analyzed. Compared with that in the control group, the protein expression levels of p-NF- κ B/p65, p-IKK α / β and p-PKC α decreased significantly in the MDA-MB-453 and BT-474 high-concentration (40 and 80 μ M) delphinidin treatment groups. In addition, the protein expression level of I κ B α , IKK α , IKK β and PKC α increased following delphinidin treatment (Fig. 5). The results suggested that delphinidin could block NF- κ B signaling in the MDA-MB-453 and BT-474 breast cancer cell lines by inhibiting NF- κ B/p65, I κ B α , and IKK α / β activity.

Immunofluorescence was used to detect the effect of delphinidin on nuclear translocation of NF- κ B/p65 in the MDA-MB-453 and BT-474 breast cancer cell lines. The results showed that the levels of NF- κ B/p65 nuclear translocation decreased in the delphinidin-treated groups. Thus, delphinidin reduced NF- κ B/p65 entry into the nucleus and inhibited transcription of nuclear genes (Fig. 6).

Discussion

Cancer is characterized by abnormal cell proliferation, cell cycle regulation and apoptosis (19). Lim *et al* (20,21) confirmed that delphinidin could reduce the survival rate of ovarian and liver cancer cells in a dose-dependent manner, and could also significantly inhibit proliferation, migration and invasion in a SKOV3 ovarian cancer cell line. Venancio *et al* (22) demonstrated that anthocyanins played an anti-inflammatory and anti-proliferating role by inhibiting the protein expression level of inflammatory cytokines, TNF- α , IL-1 β and IL-6 in colonic inflammatory fibroblasts and colon cancer cells. In the present study, it was found that delphinidin inhibited the viability of HER-2-positive MDA-MB-453 and BT-474 breast cancer cell lines and this inhibitory effect increased with delphinidin concentration.

Previous studies have demonstrated that anthocyanidin could induce G₂/M cell cycle arrest by increasing the phosphorylation of Cdk1 at Tyr15 and increasing the protein expression level of Cdk inhibitors, p27^{kip1} and p21^{WAF1/Cip1} in




Figure 3. Delphinidin induces apoptosis in the MDA-MB-453 and BT-474 cell lines. The cells were treated with 0, 20, 40 or 80 μ M delphinidin or dimethyl sulfoxide for 48 h. Representative images of TUNEL-positive (A) MDA-MB-453 and (B) BT-474 cell lines and the data was statistically analyzed. The cells were treated with 0, 20, 40 or 80 μ M delphinidin for 8 h. (C) Mitochondrial membrane potential was analyzed using flow cytometry. (D) Apoptosis was analyzed using flow cytometry. Magnification, $\times 200$. n=3. *P<0.05 vs. control; #P<0.05 vs. 20 μ M; ^P<0.05 vs. 40 μ M.

leukemia cell lines (23,24). Clemente-Soto *et al* (25) found that quercetin induced the protein expression level of p21^{WAF1/Cip1} by activating p53 and induced apoptosis in the HeLa and SiHa cell lines. In the present study, it was demonstrated

that delphinidin could induce G₂/M cell cycle arrest in the HER-2-positive MDA-MB-453 and BT-474 breast cancer cell lines and that the G₂/M cell cycle arrest was associated with the inhibition of multiple key cell cycle regulators, such as Cdk1




Figure 4. Effect of delphinidin on the cell cycle in the epidermal growth factor receptor-2-positive breast cancer cell lines. The MDA-MB-453 and BT-474 cells were treated with 20, 40 and 80 μM delphinidin or dimethyl sulfoxide for 48 h. Bcl-2 protein expression level and proteins in the MAPK signaling pathway in the (A and C) MDA-MB-453 and (B and D) BT-474 cell lines, respectively were analyzed using western blot analysis and densitometry. The data are presented as the mean and SD from at least 3 independent experiments. * $P<0.05$ vs. control; $^{\#}P<0.05$ vs. 20 μM ; $^{\#^{\wedge}}P<0.05$ vs. 40 μM . p, phosphorylated.




Figure 5. Delphinidin induces epidermal growth factor receptor-2-positive breast cancer cell apoptosis by blocking the NF- κB signaling pathway. After treatment with delphinidin for 48 h, total protein expression and phosphorylation of NF- κB /p65, I $\kappa\text{B}\alpha$, IKK α/β and PKC α were detected using western blot analysis. Phosphorylation of proteins in the NF- κB pathway in (A) MDA-MB-453 and (B) BT-474 cell lines. p, phosphorylated. The data are presented as the mean and SD from at least 3 independent experiments. #P<0.05 vs. control; #^P<0.05 vs. 20 μM ; #^+P<0.05 vs. 40 μM .

and cyclin B1. It can be hypothesized that delphinidin induced G₂/M phase arrest by increasing the protein expression level of p21^{WAF1/Cip1}, and may be involved in the induction of apoptosis in MDA-MB-453 and BT-474 breast cancer cell lines.

Loss of normal regulation of apoptosis is an important reason for the dysregulated proliferation of cancer cells (19,25,26). Induction of apoptosis may represent an effective way to treat malignant tumors. A previous study suggested that cytarabine, nitrogen mustard, cisplatin and topological isoenzyme inhibitors can induce apoptosis (27). There are 3 types of apoptotic pathways: The death receptor pathway, the mitochondrial apoptosis pathway and the endoplasmic reticulum apoptosis pathway. During the process of apoptosis, these 3 pathways may be initiated simultaneously or separately.

Kang *et al* (28) found that delphinidin inhibited the ERK/p38 MAPK signaling pathway by decreasing the phosphorylation levels of ERK and p38, and exerted anti-proliferative and pro-apoptotic effects in human osteosarcoma cells. Delphinidin inhibited the proliferation of ES2 and SKOV3 ovarian cancer cells by inhibiting the PI3K/AKT and ERK signaling pathways (29,30), which is consistent with the results of the present study. Several studies have also suggested that JNK could play an important role in cell cycle regulation, while JNK inhibited Cdk1 activation and caused G₂/M phase

arrest (31,32). Activated JNK stabilized the activity of p21^{WAF1/Cip1} by post-translational protein modification (32,33). In the present study, it was found that different concentrations of delphinidin significantly decreased the protein expression levels of p-c-Raf, p-MEK1/2 and p-ERK1/2, and increased the protein expression levels of p-JNK in the MDA-MB-453 and BT-474 breast cancer cell lines. It could be hypothesized that delphinidin-induced apoptosis may be associated with the ERK and JNK signaling pathway in the HER-2-positive breast cancer cell lines. The present study also found that delphinidin increased the protein expression level of p21^{WAF1/Cip1}, while inducing G₂/M phase arrest in the HER-2-positive breast cancer cells, which may be associated with the activation of the JNK signaling pathway.

The NF- κB signaling pathway plays an important role in the survival and proliferation of tumor cells, as well as inflammation, apoptosis. In the steady state, NF- κB is bound to its inhibitor (I κB) and remains inactive (32,33). Numerous extracellular stimulating factors, such as TNF- α , IL-1, lipopolysaccharide (LPS) and other chemicals or physical stressors (such as ultraviolet light), can phosphorylate I κB causing its degradation (32,33). The dissociated NF- κB then translocates to the nucleus and exposes nuclear recognition sites, thereby participating in target gene transcription (34,35). Previous studies have shown that




Figure 6. Delphinidin inhibits NF-κB/p65 nuclear translocation in epidermal growth factor receptor-2-positive breast cancer cell lines. MDA-MB-453 and BT-474 cells were treated with different concentrations of delphinidin for 48 h and intracellular NF-κB nuclear translocation was detected using immunofluorescence and confocal microscopy. NF-κB/p65 nuclear localization in the (A) MDA-MB-453 and (B) BT-474 cell lines. Magnification x600.




Figure 7. Schematic diagram of the molecular mechanism underlying inhibition of HER-2-positive breast cancer cell viability by delphinidin. HER, epidermal growth factor receptor.

NF-κB was abnormally activated in breast cancer cell lines. Luo *et al* (34) demonstrated that ursolic acid inhibited breast cancer growth by inhibiting proliferation, inducing apoptosis and suppressing inflammatory responses via NF-κB signaling *in vitro*. In addition, NF-κB could inhibit JNK-mediated apoptosis (36-38). Papa *et al* (38) showed that NF-κB could be inhibited during TNF-α-induced apoptosis. It caused activation

of JNK, which ultimately lead to JNK signaling pathway-mediated apoptosis. Lee *et al* (39) demonstrated that delphinidin inhibited LPS-induced macrophage NF-κB/p65 nuclear translocation. Haseeb *et al* (40) reported that delphinidin could inhibit IL-1β-induced activation of NF-κB/p65 in chondrocytes and blocked NF-κB/p65 nuclear translocation. The present study found that the phosphorylation levels of NF-κB/p65, IKKα/β,

PKC α significantly decreased in the MDA-MB-453 and BT-474 breast cancer cell lines following treatment with 40 and 80 μ M delphinidin. In addition, the nuclear translocation of NF- κ B/p65 decreased as the concentration of delphinium increased. These findings are consistent with a previous study (40) demonstrating that delphinidin inhibited the activation and nuclear translocation of NF- κ B/p65. Based on these results, it may be hypothesized that the antitumor activity of delphinidin was associated with the inhibition of the NF- κ B signaling pathway and the activation of the JNK signaling pathway.

The BT-474 cells are a HER-2-positive cell line, therefore the conclusions made in the present study could be applied to all HER-2-positive breast cancer cell lines. However, the effect of delphinidin was only analyzed in HER-2-positive breast cancer cell lines; therefore, further investigation is required in different breast cancer cell lines.

In conclusion, the present study identified the molecular mechanism in which delphinidin inhibited the viability of the HER-2-positive breast cancer cell lines. Delphinidin inhibited the ERK signaling pathway by decreasing the protein expression level of p-c-Raf, p-MEK1/2 and p-ERK1/2, regulating the protein expression level of Bax and Bcl-2 and reducing the proportion of Bax/Bcl-2 heterodimers, thereby mediating apoptosis in the mitochondrial pathway. Furthermore, delphinidin inhibited the activation of NF- κ B and nuclear translocation of NF- κ B/p65 by inhibiting the degradation of I κ B- α , blocking the NF- κ B signaling pathway and promoting the JNK signaling pathway, thereby inducing G₂/M phase arrest and apoptosis (Fig. 7).

Acknowledgements

Not applicable.

Funding

This study was supported by the National Natural Science Foundation of China (grant nos. 81573154, 82073539 and 81773432).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

AW, XY, YZ and BH confirm the authenticity of all the raw data. AW, YZ, WC, XP and XY designed the study. AW, JP, XD, JD, YO and BH conducted the research and analyzed the data. AW, XP, YZ and BH wrote the manuscript with input from the other authors. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Chan CW, Law BM, So WK, Chow KM and Waye MM: Novel strategies on personalized medicine for breast cancer treatment: An update. *Int J Mol Sci* 18: 2423, 2017.
- Akram M, Iqbal M, Daniyal M and Khan AU: Awareness and current knowledge of breast cancer. *Biol Res* 50: 33, 2017.
- Nami B, Maadi H and Wang Z: Mechanisms underlying the action and synergism of trastuzumab and pertuzumab in targeting HER2-positive breast cancer. *Cancers (Basel)* 10: 342, 2018.
- Hafeez BB, Siddiqui IA, Asim M, Malik A, Afaq F, Adhami VM, Saleem M, Din M and Mukhtar H: A dietary anthocyanidin Delphinidin induces apoptosis of human prostate cancer PC3 cells in vitro and in vivo: Involvement of nuclear factor-kappaB signaling. *Cancer Res* 68: 8564-8572, 2008.
- Jing P, Qian B, Zhao S, Qi X, Ye L, Mónica Giusti M and Wang X: Effect of glycosylation patterns of Chinese eggplant anthocyanins and other derivatives on antioxidant effectiveness in human colon cell lines. *Food Chem* 172: 183-189, 2015.
- Eghbaliferiz S and Iranshahi M: Prooxidant activity of polyphenols, flavonoids, anthocyanins and carotenoids: Updated review of mechanisms and catalyzing metals. *Phytother Res* 30: 1379-1391, 2016.
- Fang J: Classification of fruits based on anthocyanin types and relevance to their health effects. *Nutrition* 31: 1301-1306, 2015.
- Luo LP, Han B, Yu XP, Chen XY, Zhou J, Chen W, Zhu YF, Peng XL, Zou Q and Li SY: Anti-metastasis activity of black rice anthocyanins against breast cancer: Analyses using an ErbB2 positive breast cancer cell line and tumoral xenograft model. *Asian Pac J Cancer Prev* 15: 6219-6225, 2014.
- Chen XY, Zhou J, Luo LP, Han B, Li F, Chen JY, Zhu YF, Chen W and Yu XP: Black rice anthocyanins suppress metastasis of breast cancer cells by targeting RAS/RAF/MAPK pathway. *Biomed Res Int* 2015: 414250, 2015.
- Yang X, Luo E, Liu X, Han B, Yu X and Peng X: Delphinidin-3-glucoside suppresses breast carcinogenesis by inactivating the Akt/HOTAIR signaling pathway. *BMC Cancer* 16: 423, 2016.
- Chen J, Zhu Y, Zhang W, Peng X, Zhou J, Li F, Han B, Liu X, Ou Y and Yu X: Delphinidin induced protective autophagy via mTOR pathway suppression and AMPK pathway activation in HER-2 positive breast cancer cells. *BMC Cancer* 18: 342, 2018.
- Kang NJ, Lee KW, Kwon JY, Hwang MK, Rogozin EA, Heo YS, Bode AM, Lee HJ and Dong Z: Delphinidin attenuates neoplastic transformation in JB6 C141 mouse epidermal cells by blocking Raf/mitogen-activated protein kinase/extracellular signal-regulated kinase signaling. *Cancer Prev Res (Phila)* 1: 522-531, 2008.
- Yun JM, Afaq F, Khan N and Mukhtar H: Delphinidin, an anthocyanidin in pigmented fruits and vegetables, induces apoptosis and cell cycle arrest in human colon cancer HCT116 cells. *Mol Carcinog* 48: 260-270, 2009.
- Chen S, Cheng AC, Wang MS and Peng X: Detection of apoptosis induced by new type gosling viral enteritis virus in vitro through fluorescein Annexin V-FITC/PI double labeling. *World J Gastroenterol* 14: 2174-2178, 2008.
- Miao Y, Wang X, Lai Y, Lin W, Huang Y, Yin H, Hou R and Zhang F: Mitochondrial calcium uniporter promotes cell proliferation and migration in esophageal cancer. *Oncol Lett* 22, 2021.
- Zorova LD, Popkov VA, Plotnikov EY, Silachev DN, Pevzner IB, Jankauskas SS, Babenko VA, Zorov SD, Balakireva AV, Juhaszova M, *et al*: Mitochondrial membrane potential. *Anal Biochem* 552: 50-59, 2018.
- Cui L, Bu W, Song J, Feng L, Xu T, Liu D, Ding W, Wang J, Li C, Ma B, *et al*: Apoptosis induction by alantolactone in breast cancer MDA-MB-231 cells through reactive oxygen species mediated mitochondrial dependent pathway. *Arch Pharm Res* 41: 299-313, 2018.
- Boivin D, Blanchette M, Barrette S, Moghrabi A and Béliveau R: Inhibition of cancer cell proliferation and suppression of TNF induced activation of NFkappaB by edible berry juice. *Anticancer Res* 27: 937-948, 2007.
- Ouyang L, Shi Z, Zhao S, Wang FT, Zhou TT, Liu B and Bao JK: Programmed cell death pathways in cancer: A review of apoptosis, autophagy and programmed necrosis. *Cell Prolif* 45: 487-498, 2012.

20. Lim WC, Kim H and Ko H: Delphinidin inhibits epidermal growth factor-induced epithelial-to-mesenchymal transition in hepatocellular carcinoma cells. *J Cell Biochem* 120: 9887-9899, 2019.
21. Lim WC, Kim H, Kim YJ, Park SH, Song JH, Lee KH, Lee IH, Lee YK, So KA, Choi KC and Ko H: Delphinidin inhibits BDNF-induced migration and invasion in SKOV3 ovarian cancer cells. *Bioorg Med Chem Lett* 27: 5337-5343, 2017.
22. Venancio VP, Cipriano PA, Kim H, Antunes LM, Talcott ST and Mertens-Talcott SU: Cocoplum (*Chrysobalanus icaco* L.) anthocyanins exert anti-inflammatory activity in human colon cancer and non-malignant colon cells. *Food Funct* 8: 307-314, 2017.
23. Tsai TC, Huang HP, Chang KT, Wang CJ and Chang YC: Anthocyanins from roselle extract arrest cell cycle G2/M phase transition via ATM/Chk pathway in p53-deficient leukemia HL-60 cells. *Environ Toxicol* 32: 1290-1304, 2017.
24. El-Deiry WS: p21(WAF1) mediates cell-cycle inhibition, relevant to cancer suppression and therapy. *Cancer Res* 76: 5189-5191, 2016.
25. Clemente-Soto AF, Salas-Vidal E, Milan-Pacheco C, Sánchez-Carranza JN, Peralta-Zaragoza O and González-Maya L: Quercetin induces G2 phase arrest and apoptosis with the activation of p53 in an E6 expression in dependent manner in HPV positive human cervical cancer derived cells. *Mol Med Rep* 19: 2097-2106, 2019.
26. Jiao Y, Wu Y and Du D: Polydatin inhibits cell proliferation, invasion and migration, and induces cell apoptosis in hepatocellular carcinoma. *Braz J Med Biol Res* 51: e6867, 2018.
27. Liu L, Fan J, Ai G, Liu J, Luo N, Li C and Cheng Z: Berberine in combination with cisplatin induces necroptosis and ovarian cancer cells. *Biol Res* 52: 37, 2019.
28. Kang HM, Park BS, Kang HK, Park HR, Yu SB and Kim IR: Delphinidin induces apoptosis and inhibits epithelial-to-mesenchymal transition via the ERK/p38 MAPK-signaling pathway in human osteosarcoma cell lines. *Environ Toxicol* 33: 640-649, 2018.
29. Lim W and Song G: Inhibitory effects of Delphinidin on the proliferation of ovarian cancer cells via PI3K/AKT and ERK 1/2 MAPK signal transduction. *Oncol Lett* 14: 810-818, 2017.
30. Lim W, Jeong W and Song G: Delphinidin suppresses proliferation and migration of human ovarian clear cell carcinoma cells through blocking AKT and ERK1/2 MAPK signaling pathways. *Mol Cell Endocrinol* 422: 172-181, 2016.
31. Wu J, Sun J and Xue Y: Involvement of JNK and P53 activation in G2/M cell cycle arrest and apoptosis induced by titanium dioxide nanoparticles in neuron cells. *Toxicol Lett* 199: 269-276, 2010.
32. Zhu N, Shao Y, Xu L, Yu L and Sun L: Gadd45-alpha and Gadd45-gamma utilize p38 and JNK signaling pathways to induce cell cycle G2/M arrest in Hep-G2 hepatoma cells. *Mol Biol Rep* 36: 2075-2085, 2009.
33. Wang CC, Chiang YM, Sung SC, Hsu YL, Chang JK and Kuo PL: Plumbagin induces cell cycle arrest and apoptosis through reactive oxygen species/c-Jun N-terminal kinase pathways in human melanoma A375.S2 cells. *Cancer Lett* 259: 82-98, 2008.
34. Luo J, Hu YL and Wang H: Ursolic acid inhibits breast cancer growth by inhibiting proliferation, inducing autophagy and apoptosis, and suppressing inflammatory responses via the PI3K/AKT and NF-κB signaling pathways in vitro. *Exp Ther Med* 14: 3623-3631, 2017.
35. Wang H, Huang W, Liang M, Shi Y, Zhang C, Li Q, Liu M, Shou Y, Yin H, Zhu X, *et al*: (+)-JQ1 attenuated LPS-induced microglial inflammation via MAPK/NFκB signaling. *Cell Biosci* 8: 60, 2018.
36. Bubici C, Papa S, Pham CG, Zazzeroni F and Franzoso G: NF-kappaB and JNK: An intricate affair. *Cell Cycle* 3: 1524-1529, 2004.
37. Sakon S, Xue X, Takekawa M, Sasazuki T, Okazaki T, Kojima Y, Piao JH, Yagita H, Okumura K, Doi T and Nakano H: NF-kappaB inhibits TNF-induced accumulation of ROS that mediate prolonged MAPK activation and necrotic cell death. *EMBO J* 22: 3898-3909, 2003.
38. Papa S, Zazzeroni F, Pham CG, Bubici C and Franzoso G: Linking JNK signaling to NF-kappaB: A key to survival. *J Cell Sci* 117: 5197-5208, 2004.
39. Lee SG, Kim B, Yang Y, Pham TX, Park YK, Manatou J, Koo SI, Chun OK and Lee JY: Berry anthocyanins suppress the expression and secretion of proinflammatory mediators in macrophages by inhibiting nuclear translocation of NF-κB independent of NRF2-mediated mechanism. *J Nutr Biochem* 25: 404-411, 2014.
40. Haseeb A, Chen D and Haqqi TM: Delphinidin inhibits IL-1β-induced activation of NF-κB by modulating the phosphorylation of IRAK-1(Ser376) in human articular chondrocytes. *Rheumatology (Oxford)* 52: 998-1008, 2013.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.