Application of robotic surgery and traditional laparoscopic surgery in lymph node dissection for gynecological cancer: A meta-analysis

YUANYUAN LU, JINGPING CHEN, RENJI WEI, WENTING LIN, YUDONG CHEN, YICHENG SU, LIJUAN LIU, YUKUN LIANG and MULAN WEI

Department of Gynecology, Guigang People's Hospital, Guigang, Guangxi 537100, P.R. China

Received October 7, 2022; Accepted January 30, 2023

DOI: 10.3892/ol.2023.13761

Abstract. Since the advantages of robotic surgery and laparoscopic surgery in the number of lymph node resections are not well understood, this meta-analysis used evidence-based medicine to assess the difference in the number of lymph nodes retrieved in gynecological cancer between the two surgical methods to guide clinical treatment. In the present meta-analysis, the Pubmed, Embase, Cochrane, China National Knowledge Infrastructure and Wanfang libraries were searched for articles that were published from the time of the database's inception to January 2021, including cohort studies and randomized controlled trials, where the observation group underwent robotic surgery to treat gynecological cancers and the control group underwent laparoscopic surgery to treat gynecological cancers, including cervical and ovarian cancers and endometrial cancers. Duplicate publications, studies with no full text, incomplete information or where the authors were unable to perform data extraction, animal experiments, reviews and systematic reviews were excluded. STATA 15.1 was used to analyze the data. Robotic surgery resulted in a significant increase in the number of lymph nodes retrieved from the pelvis [standard mean difference (SMD)=0.24; 95% CI, 0.04-0.45; P=0.007] and para-aortic (SMD=0.41; 95%) CI, 0.13-0.69; P=0.004) regions compared with the number retrieved by laparoscopic surgery. Furthermore, there was no significant difference in operating time between robotic and laparoscopic surgery, despite the use of different instruments (SMD=0.12; 95% CI, -0.35-0.58; P=0.616). The amount of blood lost during robotic surgery was significantly less compared with that lost during laparoscopic surgery [SMD=-0.40; 95% CI, -0.58-(-0.22); P<0.001]. The present study evaluated cancer recurrence and death in further detail, and no statistically significant difference was demonstrated between robotic surgery and laparoscopic surgery in terms of recurrence rate [odds ratio (OR)=0.59; 95% CI, 0.21-1.65; P=0.318] and mortality rate (OR=0.31; 95% CI, 0.08-1.30; P=0.109). The present study demonstrated that robotic surgery was able to retrieve more pelvic and para-aortic lymph nodes than traditional laparoscopic surgery, which was consistent with previous reports. With regards to blood loss, The difference in operation time between the two surgical methods was not statistically significant, whereas the estimated blood loss of robotic surgery was significantly lower than that of traditional laparoscopic surgery. There was no statistically significant difference in the recurrence rate and mortality rate of the two surgical modality.

Introduction

Gynecological malignancies are the leading cause of female mortality in the United States (1). Despite advancements in the identification and treatment of gynecological malignancies, the survival rate has remained constant for decades (2). Gynecological cancers predominately comprise cervical, endometrial and ovarian cancers (3). Surgical care for these malignancies has transformed during the past decade as a result of the development of techniques that are less invasive such as laparoscopic and robotic surgeries.

As a result of the US Federal Drug Administration's approval of the da Vinci Surgical System for gynecological surgery in 2005, the use of robot-assisted surgery in the treatment of gynecological malignancies has become more common (4). There are several benefits to using robotic technology, including 3D and high-definition optics, endoscopic devices that may be used in numerous ways, the removal of human tremors, and greater accuracy and autonomy for the surgeon (5-7). The advantages of robot-assisted surgery over conventional laparoscopic surgery are mainly due to these aforementioned factors (8).

Women with gynecological malignancies who have solitary pelvic lymph node or para-aortic lymph node metastases at the time of their first diagnosis have a poor prognosis (9,10). It is possible to compare the number of clear

Correspondence to: Dr Yuanyuan Lu, Department of Gynecology, Guigang People's Hospital, 1 Zhongshan Road, Gangbei Square, Guigang, Guangxi 537100, P.R. China E-mail: luyuanyuan-009@163.com

Key words: robotic surgery, laparoscopic surgery, lymph node dissection, gynecological cancer

lymph nodes in gynecological cancers treated with robotic surgery and traditional laparoscopic surgery, to assess the relative benefits and drawbacks of the two surgical methods in terms of prognosis. Lymph node dissection is a controversial procedure, and the outcomes of the two surgical approaches are still debated (11,12). Robot-assisted surgery and laparoscopic surgery may not vary significantly in the number of lymph node dissections, according to certain previous reports (13,14). However, numerous studies reported that robotic surgery has a higher lymph node clearance rate than laparoscopic surgery (15,16). Therefore, this meta-analysis used evidence-based medicine to explore the difference in the number of lymph nodes retrieved in gynecological cancer between robotic surgery and laparoscopic surgery by pooling the results of relevant previous studies.

Materials and methods

Protocol registration. The protocol used for the present study was registered in the International Platform of Registered Systematic Review and Meta-analysis Protocols (https://inplasy. com/), the registration number was INPLASY2022120046 (DOI, 10.37766/inplasy2022.12.0046).

Literature inclusion and exclusion criteria. The following inclusion criteria were used for reports in the literature: The study was a cohort study or a randomized controlled trial (RCT) study; the study was reported in either English or Chinese; and studies where the observation group was treated by robotic surgery for gynecological cancer (taken to include cervical cancer, ovarian cancer and endometrial cancer) and the control group was treated by laparoscopic surgery for gynecological cancer.

The following exclusion criteria were used for reports in the literature: Repeated publication of the same study; studies without full text, incomplete information or for which data extraction could not be performed; animal experiments; and reviews and systematic reviews.

Search strategy. The Pubmed (https://pubmed.ncbi.nlm. nih.gov/), Embase (https://www.embase.com), Cochrane library (https://www.cochranelibrary.com/), China National Knowledge Infrastructure (https://www.cnki.net/) and Wanfang (https://www.wanfangdata.com.cn/index.html) literature databases were used in the present meta-analysis, and were searched between their formation and January 1st 2021. The mesh terms used were as follows: 'robotic-assisted', 'robotic surgery' and 'laparoscopic', 'laparoscopy' and 'gynecological cancer', 'Uterine Cervical Neoplasms', 'Endometrial Neoplasms', 'Ovarian Neoplasms'. Detailed search strategies were presented in Data S1.

Literature screening and data extraction. The literature search, screening and extraction of relevant material was completed by two researchers, separately. When there were questions or conflicts, a third person was consulted before making a decision. Basic features (author, year, research type, field of research on gynecological cancers, number of patients, age of patients and place of diagnosis) and outcome indicators (number of retrieved pelvic lymph nodes and para-aortic

lymph nodes, operative time, estimated blood loss, recurrence rate and mortality rate) were included in the data extracted.

Literature quality assessment. The Newcastle-Ottawa Scale (NOS) for evaluating the quality of published literature was performed separately by two academics (17)], in the present study it was used to evaluate the quality of the 16 cohort studies. The NOS includes 4 items (4 points) for 'Research Subject Selection', 1 item (2 points) for 'Comparability between Groups' and 3 items (3 points) for 'Result Measurement', for a maximum score of 9 points. A score of \geq 7 is classified as high-quality literature, <7 is classified into lower-quality literature. Following assessment using the NOS, the quality of the remaining two RCT studies was assessed against the Cochrane Risk Assessment Scale using the ReviewManager 5.3 (Cochrane) software risk assessment tool.

Data synthesis and statistical analysis. The data was analyzed using the STATA (version 15.1, StataCorp LP) (18). The present study used the combined effect size of standard mean difference (SMD) (with a 95% confidence interval) as a continuous variable to assess the number of retrieved pelvic lymph nodes and para-aortic lymph nodes, as well as the amount of time spent operating and estimated blood loss, and the odds ratio (OR; with a 95% confidence interval) as a binary variable to measure the rates of recurrence and mortality. The I² statistic was used to determine the degree of heterogeneity. If the results of the test for heterogeneity were P \ge 0.1 and I² \le 50%, this indicated that the studies were homogeneous, and the fixed effects model (I-V) was used for combined analysis; however, if the results were P<0.1 and I^2 >50%, this indicated that the studies were heterogeneous. Since heterogeneity is always expected for the intervention effects among multiple studies from different groups and geographical locations, only the random effects model was used to combine the results. Sensitivity analysis or subgroup analysis was required to identify the source of heterogeneity. If the heterogeneity was still large, the random effects model (D+L) was used or the combination of results was stopped and descriptive analysis was used. Egger's bias test was used to analyze the publication prejudice.

Results

The results of literature search. In total, the database search yielded 1735 studies. A total of 746 studies remained after duplicates were removed. After browsing titles and abstracts, 307 studies were evaluated. Finally, 18 papers were finally meta-analyzed after being read in their entirety (Fig. 1).

Baseline characteristics and quality assessment of the included studies

Baseline characteristics. The present meta-analysis included 18 publications, 16 of which were cohort studies and two of which were RCTs. The sample sizes ranged from 42 to 933, and the present meta-analysis included a total of 2,381 patients (all women; age, >44.1 years). Among the included studies, 11 reported cervical cancer, three reported ovarian cancer, three reported endometrial cancer and one reported gyne-cological cancers, including cervical cancer, ovarian cancer



Figure 1. Flow diagram for selection of studies. CNKI, China National Knowledge Infrastructure.

and endometrial cancer. A total of 16 cohort studies had NOS ratings of at \geq 7 and satisfied all other conditions (Table I). The quality evaluation of the randomized controlled trials was presented in Fig. 2.

Results of the meta-analysis

Number of retrieved lymph nodes. i) Number of retrieved pelvic lymph nodes. A total of >2,000 patients participated in 15 studies, which reported on the number of pelvic lymph nodes that were retrieved. The meta-analysis was performed using a random-effects model because of the large amount of variation in the data (I²=68.6%, P<0.001). The pooled results demonstrated that the number of pelvic lymph nodes retrieved by robotic surgery was significantly higher than that of laparoscopic surgery (SMD=0.24; 95% CI, 0.04-0.45; P=0.007; Fig. 3A).

ii) Number of retrieved para-aortic lymph nodes. A total of eight studies including 751 patients reported on the number of para-aortic lymph nodes that were removed. Meta-analysis was performed using a random-effects model due to significant heterogeneity ($I^2=65.2\%$, P=0.004). The combined data demonstrated that the number of para-aortic lymph nodes



Figure 2. The quality assessment of randomized controlled studies. Green circles indicated low risk and yellow circles indicated undescribed.

First author/s, year	Study type	Study area	No. patients		Age, years (mean ± SD or median with range)				
			Robotic surgery	Laparos- copic surgery	Robotic surgery	Laparos- copic surgery	Site of gynecological cancer	NOS score	(Refs.)
Magrina <i>et al</i> , 2011	Cohort	USA	25	27	62.0±15.0	61.0±16.0	Ovarian cancer	7	(27)
Sert and Abeler, 2011	Cohort	Norway	35	7	44.1±10.5	45.0±12.9	Cervical Cancer	8	(28)
Chong <i>et al</i> , 2013	Cohort	South Korea	50	50	48.0±9.9	47.1±11.0	Cervical Cancer	7	(29)
Chen <i>et al</i> , 2014 [14]	Cohort	China	24	32	53.7±15.3	51.2±11.9	Cervical Cancer	7	(14)
Coronado <i>et al</i> , 2014	Cohort	Spain	32	30	57.5 (47.5-66.7)	59.0 (38.6-66.3)	Gynecological cancer	8	(30)
Díaz-Feijoo <i>et al</i> , 2014	Cohort	Spain	83	17	51.0 (24.0-74.0)	49.0 (29.0-66.0)	Cervical Cancer	7	(31)
Yim <i>et al</i> , 2014 [11]	Cohort	South Korea	60	42	46.3±9.9	49.8±11.4	Cervical Cancer	7	(11)
Chen <i>et al</i> , 2016	Cohort	China	44	21	44.3±12.3	43.8±10.3	Ovarian cancer	7	(32)
Manchana <i>et al</i> , 2015	Cohort	Thailand	28	47	55.5 (48.2-61.5)	54.0 (49.0-62.0)	Endometrial cancer	7	(33)
Gallotta <i>et al</i> , 2017	Cohort	Italy	32	64	49.0 (32.0-76.0)	49.0 (27.0-73.0)	Ovarian cancer	8	(15)
Mäenpää <i>et al</i> , 2016	RCT	Finland	50	49	67.0 (43.0-84.0)	70.0 (48.0-83.0)	Endometrial	/	(34)
Mendivil <i>et al</i> , 2016	Cohort	USA	58	49	47.3±11.2	47.8±12.0	Cervical Cancer	7	(35)
Lee <i>et al</i> , 2018	Cohort	South Korea	26	16	56.7±6.9	51.1±7.8	Endometrial cancer	8	(4)
Pellegrino <i>et al</i> , 2017	Cohort	Italy	34	18	46.9±9.5	48.2±13.1	Cervical Cancer	8	(36)
Nie <i>et al</i> , 2017	Cohort	China	100	833	47.1±9.5	45.9±8.9	Cervical	7	(37)
Luo <i>et al</i> , 2018	RCT	China	30	30	65.0 (62.0-67.0)	64.0 (62 0-66 0)	Cervical	/	(38)
Oyama <i>et al</i> , 2019	Cohort	Japan	64	57	43.4±12.2	44.9±12.3	Cervical	7	(16)
Loverix <i>et al</i> , 2020	Cohort	Belgium	55	162	49.0 (22.0-79.0)	48.0 (22.0-78.0)	Cervical Cancer	8	(12)

Table I. The baseline characteristics and quality assessment of the cohort studies.

recovered by robotic surgery was significantly greater than the number retrieved by laparoscopic surgery, which was a considerable improvement (SMD=0.41; 95% CI, 0.13-0.69; P=0.004; Fig. 3B).

Operation time. Operation time was recorded in 14 studies, which included 2112 patients. To account for the substantial variability in the data, a random effects model

was used ($I^2=94.0\%$, P<0.001). The data indicated that, robotic surgery and laparoscopic surgery took the same amount of time to perform (SMD=0.12; 95% CI, -0.35-0.58; P=0.616; Fig. 3C).

Estimated blood loss. A total of 13 studies including 1102 patients, estimated blood loss. Because of the substantial heterogeneity, meta-analysis was performed using a



Figure 3. Comparison in the number of (A) pelvic and (B) para-aortic lymph nodes retrieved by robotic surgery and laparoscopic surgery; comparison in (C) operation time and (D) estimated blood loss between robotic surgery and laparoscopic surgery; comparison in (E) recurrence rate and (F) mortality rate between robotic surgery and laparoscopic surgery. SMD, standard mean difference.

random-effects model (I^2 =44.3%, P=0.043). There was a significant difference in blood loss between robotic and laparoscopic surgery, demonstrated by the pooled findings [SMD=-0.40; 95% CI, -0.58-(-0.22); P<0.001; Fig. 3D].

robotic and laparoscopic surgery, according to the pooled statistics (OR=0.59; 95% CI, 0.21-1.65; P=0.318; Fig. 3E).

Disease prognosis. i) Recurrence rate. A total of four studies, including 1170 patients, reported recurrence rate. Meta-analysis was performed using a random-effects model. Recurrence rates were not significantly different between

ii) Mortality rate. The death rate was recorded in 4 studies, which included 1147 patients. Meta-analysis was performed using a random-effects model. Mortality rate was not significantly different between robotic and laparoscopic surgery, according to the pooled statistics (OR=0.31; 95% CI, 0.08-1.30; P=0.109; Fig. 3F)



Figure 4. Subgroup analysis of comparison in the number of (A) pelvic and (B) para-aortic lymph nodes retrieved by robotic surgery and laparoscopic surgery; comparison of (C) operation time and (D) estimated blood loss between robotic surgery and laparoscopic surgery. SMD, standard mean difference; D + L, random effects model; I-V, fixed effects model.

Subgroup analysis. Subgroup analysis of the number of retrieved lymph nodes, operation time and estimated blood loss according to different gynecological cancers was performed.

Number of retrieved lymph nodes. i) Number of retrieved pelvic lymph nodes. The pooled results showed that the number of pelvic lymph nodes retrieved by robotic surgery was significantly larger than that obtained by laparoscopic surgery in cervical cancer (SMD=0.38; 95% CI, 0.08-0.67; P=0.012; I²=77.6%; P<0.001) (Fig. 4A). However, in ovarian cancer (SMD=0.11; 95% CI, -0.33-0.55; P=0.634; I²=57.3%; P=0.096) and endometrial cancer (SMD=0.02; 95% CI, -0.26-0.29; P=0.904; I²=0.0%; P=0.420), the difference between robotic surgery and laparoscopic surgery in the number of retrieved pelvic lymph nodes was not statistically significant (Fig. 4A).

ii) Number of retrieved para-aortic lymph nodes. The pooled results demonstrated that in cervical cancer (SMD=0.49; 95% CI, 0.06-0.91; P=0.024; I²=70.3%, P=0.034) and endometrial cancer (SMD=0.75; 95% CI, 0.36-1.14; P<0.001; I²=0.0%; P=0.562), the

number of para-aortic lymph nodes retrieved by robotic surgery were significantly increased compared with laparoscopic surgery (Fig. 4B). For ovarian cancer therapy, the number of para-aortic lymph nodes removed by robotic surgery and laparoscopic surgery did not differ significantly between the two techniques, according to the results of the present meta-analysis (SMD=0.21; 95% CI, -0.47-0.88; P=0.554; I²=73.7%; P=0.051; Fig. 4B).

Operation time. The pooled results demonstrated that in cervical cancer (SMD=0.11; 95% CI, -0.44-0.66; P=0.701; I²=94.0%; P<0.001), ovarian cancer (SMD=0.121; 95% CI, -0.92-1.15; P=0.827; I²=88.4%; P=0.003) and endometrial cancer (SMD=0.50; 95% CI, -2.48-3.49; P=0.616; I²=98.6%; P<0.001), the difference in operation time between robotic surgery and laparoscopic surgery was not statistically significant (Fig. 4C).

Estimated blood loss. The pooled results demonstrated that in cervical cancer [SMD=-0.52; 95% CI, -0.80-(-0.25); P<0.001; I^2 =57.5%; P=0.029] and ovarian cancer [SMD=-0.54; 95%



Figure 5. Publication bias of the analysis of the number of (A) pelvic and (B) para-aortic lymph nodes retrieved by robotic surgery and laparoscopic surgery; comparison of (C) operation time and (D) estimated blood loss between robotic surgery and laparoscopic surgery. SMD, standard mean difference; se, standard error.

CI, -0.88-(-0.20); P=0.002; I²=0.0%; P=0.670], the estimated blood loss of robotic surgery was significantly lower than that of laparoscopic surgery (Fig. 4D). There was no statistically significant difference in estimated blood loss when compared between robotic surgery and laparoscopic surgery for endometrial cancer (SMD=-0.05; 95% CI, -0.32-0.22; P=0.725; I²=0.0%; P=0.766; Fig. 4D).

Sensitivity analysis. The remaining investigations were subjected to a pooled analysis to assess whether any of the included studies had a disproportionate influence on the meta-analysis's overall results, which was accomplished using sensitivity analyses that eliminated each included publication one at a time. The results of this meta-analysis indicated that no research had a disproportionate influence on its results, which suggested that the findings were steady and credible (Figs. S1-S6).

Publication bias. Egger's test (P=0.305; P=0.684; P=0.547; and P=0.366) was used to assess the four funnel plots created in the present study and found no significant publication bias was demonstrated (Fig. 5).

Discussion

The development of minimally invasive technologies is important in surgical systems. Minimally invasive surgery can result in better treatment outcomes, faster recovery times, shorter hospital stays, and reduced physical and psychological trauma (19). Recently, a growing number of studies have reported the efficiency and safety of da Vinci robots for the treatment of benign and malignant gynecological tumors (20-22). Due to subjective factors such as the patient's subjective choice of surgical method, difficultly in ensuring the informed consent of patients and their families with randomized surgical methods and difficulty in achieving double-blinding of the wound type after surgery, it is difficult to randomize surgical methods. Therefore, clinical randomized controlled trials exploring different surgical techniques are currently lacking. Since lymph node metastasis has an impact on the prognosis of tumors, including recurrence rate and mortality, it is very important to evaluate the number of lymph nodes removed. This meta-analysis included 18 articles with a total of 2,381 patients who underwent either robotic surgery or conventional laparoscopic surgery for gynecological cancers. The clinical outcomes of da Vinci robot-assisted laparoscopic surgery for gynecological cancers were compared with those of conventional laparoscopic surgery. The number of lymph nodes removed (pelvic and abdominal aortic lymph nodes), length of surgery, expected blood loss, recurrence rate and mortality rate were analyzed.

The number of lymph nodes removed was the first factor assessed. In 15 of the studies, a total of 2002 patients

underwent pelvic lymph node removal. The pooled results in the present study demonstrated that the number of pelvic lymph nodes retrieved by robotic surgery was significantly higher than that of laparoscopic surgery (SMD=0.24; 95% CI, 0.04-0.45; P=0.007). Furthermore, 751 patients in eight studies had their para-aortic lymph nodes removed. The pooled results demonstrated that significantly more para-aortic lymph nodes were removed using robotic surgery compared with laparoscopic surgery (SMD=0.41; 95% CI, 0.13-0.69; P=0.004). This result may be due to the ability of the da Vinci robot's simulated wrist endoscopy operating instrument to complete a 7-direction degrees of freedom operation, which means it can reach positions that cannot be reached by human hands (23). Consequently, the robotic system can operate more precisely, allowing for the removal of more lymph nodes. For gynecological cancers, lymph node metastases are an essential channel of spread, and the positive identification rate of lymph nodes is directly linked to postoperative therapy and prognosis (24). The present study performed subgroup analyses for different forms of gynecological cancer. In cervical cancer, the number of pelvic lymph nodes retrieved via robotic surgery was significantly greater than that retrieved via laparoscopic surgery (SMD=0.38; 95% CI, 0.08-0.67; P=0.012). The results of the present study also demonstrated that only in cervical (SMD=0.49; 95% CI, 0.06-0.91; P=0.024) and endometrial (SMD=0.75; 95% CI, 0.36-1.14; P<0.001) cancers were the number of para-aortic lymph nodes retrieved by robotic surgery significantly increased compared with laparoscopic surgery. This may have been due to the differences in the anatomical morphology of gynecological cancer at different locations, which suggested that the combination of the number of lymph nodes removed with the location of the gynecological cancer should be used to choose the most appropriate surgical method for each patient.

A total of 14 publications, which included 2,112 patients, were pooled to examine the difference in operating time between laparoscopic and robotic surgery. All available data demonstrated that there was no significant difference in the operating time between the two types of surgery (SMD=0.12; 95% CI, -0.35-0.58; P=0.616). This result may be due to the da Vinci robot-assisted laparoscopic system being an emerging technology and surgical operators in different countries and regions have different proficiencies in robotic and laparoscopic surgery. As the proficiency of laparoscopic operators is improved, the operation time will be reduced in the future. The difference in estimated blood loss between robotic and laparoscopic surgery was also assessed in the present study. Compared with laparoscopy, the robot-assisted technique has been previously reported to minimize blood loss in gynecological malignancies (25). The pooled results of the present study also demonstrated that the estimated blood loss in robotic surgery was significantly lower compared with that in laparoscopic surgery [SMD=-0.40; 95% CI, -0.58-(-0.22), P<0.001]. This has been reported to be because the robotic surgical system can provide a high-definition, 10-15x magnification view of the surgical field of view, which has a magnifying effect on blood vessels and can avoid the problem of missing small blood vessels in the surgical area (26). When the subsets of patients were assessed, there was no statistically significant difference in estimated blood loss between robotic and laparoscopic surgery for endometrial cancer (SMD=-0.05; 95% CI, -0.32-0.22; P=0.725), which indicated that the blood loss in certain sections of the malignancy may vary.

The cancer recurrence and death rates were further evaluated. Robotic surgery demonstrated a significantly lower death rate compared with laparoscopic surgery (OR=0.2; 95% CI, 0.08-0.95; P=0.041); however, there was no significant difference in recurrence rate and mortality rate between the two procedures. However, few studies have reported recurrence and mortality rates, and other reasons, such as failure to complete postoperative adjuvant treatment during the follow-up period, may affect the postoperative recurrence rate and mortality rate for certain patients. Therefore, to evaluate the long-term efficacy of robotic surgery the sample size should expanded in further research.

This meta-analysis has several limitations. Firstly, the majority of the studies included in this research were retrospective cohort studies, with only two randomized controlled studies; therefore, the quality of the literature was uneven, which could lead to selection bias. Secondly, the literature included in this study did not describe the surgeon's proficiency in the use of the da Vinci Robotic Surgery System, laparoscopic surgery or open surgery, and differences between surgeons' skills may also be a reason for the heterogeneity in this study. Therefore, it will be necessary to include additional studies in the future and expand the sample size to further verify the findings of the present study. Thirdly, as the patients included in each study were not all at a specific cancer stage, it was not possible to perform further subgroup analyses by cancer stage. Fourth, as the complication rates analyzed in the included studies were all robotic surgery and conventional laparoscopic surgery for gynecological cancers, it was not possible to analyze complications after lymph node dissection. If a relevant study is performed, a correlation analysis should be performed.

The results of the present study have demonstrated that lymph nodes from the pelvic and para-aortic areas may be retrieved more effectively using robotic surgery than with laparoscopy. The difference in operation time between the two surgical methods w not statistically significant; however, the estimated blood loss of robotic surgery is significantly lower than that of traditional laparoscopic surgery. Furthermore, robotic surgery has a significantly lower death rate than laparoscopic surgery; however, the recurrence rates were not statistically different.

Acknowledgements

Not applicable.

Funding

The present study was supported by the Guigang Self-funded Scientific Research Project (grant no. 2100031).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

YLu conceived the study and wrote the manuscript; WL, YS, LL, RW and MW performed the acquisition of the data. YLi, YC and YLu performed the data analyses; JC helped to perform the analysis. YLu and YC confirm the authenticity of all the raw data. All authors have read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Keyvani V, Kheradmand N, Navaei ZN, Mollazadeh S and Esmaeili SA: Epidemiological trends and risk factors of gynecological cancers: An update. Med Oncol 40: 93, 2023.
- Wu Z, Tang H, Xiong Q, Liu D, Xia T, Liang H and Ye Q: Prognostic role of microRNA-205 in human gynecological cancer: A meta-analysis of fourteen studies. DNA Cell Biol 39: 875-889, 2020.
- Lu J, Gu Y, Li Q, Zhong H, Wang X, Zheng Z, Hu W and Wen L: Wilms' tumor 1 (WT1) as a prognosis factor in gynecological cancers: A meta-analysis. Medicine (Baltimore) 97: e11485, 2018.
- 4. Lee HJ, Lee YH, Chong GO, Hong DG and Lee YS: Comparison of robotic-assisted versus laparoscopy for transperitoneal infrarenal para-aortic lymphadenectomy in patients with endometrial cancer. J Obstet Gynaecol Res 44: 547-555, 2018.
- Seamon LG, Cohn DE, Richardson DL, Valmadre S, Carlson MJ, Phillips GS and Fowler JM: Robotic hysterectomy and pelvic-aortic lymphadenectomy for endometrial cancer. Obstet Gynecol 112: 1207-1213, 2008.
- Lowe MP, Johnson PR, Kamelle SA, Kumar S, Chamberlain DH and Tillmanns TD: A multiinstitutional experience with robotic-assisted hysterectomy with staging for endometrial cancer. Obstet Gynecol 114: 236-243, 2009.
- Lim PC, Kang É and Park DH: A comparative detail analysis of the learning curve and surgical outcome for robotic hysterectomy with lymphadenectomy versus laparoscopic hysterectomy with lymphadenectomy in treatment of endometrial cancer: A case-matched controlled study of the first one hundred twenty two patients. Gynecol Oncol 120: 413-418, 2011.
 Holloway RW, Ahmad S, DeNardis SA, Peterson LB, Sultana N,
- Holloway RW, Ahmad S, DeNardis SA, Peterson LB, Sultana N, Bigsby GE IV, Pikaart DP and Finkler NJ: Robotic-assisted laparoscopic hysterectomy and lymphadenectomy for endometrial cancer: Analysis of surgical performance. Gynecol Oncol 115: 447-452, 2009.
- Diaz-Feijoo B, Luna-Guibourg R, Cabrera S, Manrique S and Gil-Moreno A: Laparoscopic extraperitoneal pelvic lymph node debulking in locally advanced cervical cancer. J Minim Invasive Gynecol 26: 366, 2019.
- Fastrez M, Goffin F, Vergote I, Vandromme J, Petit P, Leunen K and Degueldre M: Multi-center experience of robot-assisted laparoscopic para-aortic lymphadenectomy for staging of locally advanced cervical carcinoma. Acta Obstet Gynecol Scand 92: 895-901, 2013.
- Yim GW, Kim SW, Nam EJ, Kim S, Kim HJ and Kim YT: Surgical outcomes of robotic radical hysterectomy using three robotic arms versus conventional multiport laparoscopy in patients with cervical cancer. Yonsei Med J 55: 1222-1230, 2014.
 Loverix L, Salihi RR, Van Nieuwenhuysen E, Concin N, Han S,
- Loverix L, Salihi RR, Van Nieuwenhuysen E, Concin N, Han S, van Gorp T and Vergote I: Para-aortic lymph node surgical staging in locally-advanced cervical cancer: Comparison between robotic versus conventional laparoscopy. Int J Gynecol Cancer 30: 466-472, 2020.

- Magrina JF, Cetta RL, Chang YH, Guevara G and Magtibay PM: Analysis of secondary cytoreduction for recurrent ovarian cancer by robotics, laparoscopy and laparotomy. Gynecol Oncol 129: 336-340, 2013.
- 14. Chen CH, Chiu LH, Chang CW, Yen YK, Huang YH and Liu WM: Comparing robotic surgery with conventional laparoscopy and laparotomy for cervical cancer management. Int J Gynecol Cancer 24: 1105-1111, 2014.
- 15. Gallotta V, Cicero C, Conte C, Vizzielli G, Petrillo M, Fagotti A, Chiantera V, Costantini B, Scambia G and Ferrandina G: Robotic versus laparoscopic staging for early ovarian cancer: A case-matched control study. J Minim Invasive Gynecol 24: 293-298, 2017.
- 16. Oyama K, Kanno K, Kojima R, Shirane A, Yanai S, Ota Y and Andou M: Short-term outcomes of robotic-assisted versus conventional laparoscopic radical hysterectomy for early-stage cervical cancer: A single-center study. J Obstet Gynaecol Res 45: 405-411, 2019.
- Cook DA and Reed DA: Appraising the quality of medical education research methods: The medical education research study quality instrument and the Newcastle-Ottawa scale-education. Acad Med 90: 1067-1076, 2015.
 Holzman SA, Grant C, Zee RS, Rana MS, Sprague B, Blum E
- Holzman SA, Grant C, Zee RS, Rana MS, Sprague B, Blum E and Rushton HG: High incidence of abnormal imaging findings in circumcised boys diagnosed with urinary tract infections. J Pediatr Urol 16: 560-565, 2020.
- Nezhat FR, Datta MS, Liu C, Chuang L and Zakashansky K: Robotic radical hysterectomy versus total laparoscopic radical hysterectomy with pelvic lymphadenectomy for treatment of early cervical cancer. JSLS 12: 227-237, 2008.
- 20. Shi C, Gao Y, Yang Y, Zhang L, Yu J and Zhang T: Comparison of efficacy of robotic surgery, laparoscopy, and laparotomy in the treatment of ovarian cancer: A meta-analysis. World J Surg Oncol 17: 162, 2019.
- 21. Su MC, Zheng Y, Yang F and Liu YY: Placement of robotic single-site surgery using the da Vinci Xi platform and the tumor-free technique for early cervical cancer. Asian J Surg: S1015-9584(22)01307-0, 2022 (Epub ahead of print).
- 22. Yotsumoto F, Sanui A, Ito T, Miyahara D, Yoshikawa K, Shigekawa K, Noguchi Y, Yasunaga S and Miyamoto S: Cumulative summation analysis of learning curve for robotic-assisted hysterectomy in patients with gynecologic tumors. Anticancer Res 42: 4111-4117, 2022.
- 23. Lanfranco AR, Castellanos AE, Desai JP and Meyers WC: Robotic surgery: A current perspective. Ann Surg 239: 14-21, 2004.
- 24. Ji Y and Wang H: Prognostic prediction of systemic immune-inflammation index for patients with gynecological and breast cancers: A meta-analysis. World J Surg Oncol 18: 197, 2020.
- Reza M, Maeso S, Blasco JA and Andradas E: Meta-analysis of observational studies on the safety and effectiveness of robotic gynaecological surgery. Br J Surg 97: 1772-1783, 2010.
- Camarillo DB, Krummel TM and Salisbury JK Jr: Robotic technology in surgery: Past, present, and future. Am J Surg 188 (4A Suppl): 2S-15S, 2004.
- 27. Magrina JF, Zanagnolo V, Noble BN, Kho RM and Magtibay P: Robotic approach for ovarian cancer: Perioperative and survival results and comparison with laparoscopy and laparotomy. Gynecol Oncol 121: 100-105, 2011.
- Sert MB and Abeler V: Robot-assisted laparoscopic radical hysterectomy: Comparison with total laparoscopic hysterectomy and abdominal radical hysterectomy; one surgeon's experience at the Norwegian Radium Hospital. Gynecol Oncol 121: 600-604, 2011.
- 29. Chong GO, Lee YH, Hong DG, Cho YL, Park IS and Lee YS: Robot versus laparoscopic nerve-sparing radical hysterectomy for cervical cancer: A comparison of the intraoperative and perioperative results of a single surgeon's initial experience. Int J Gynecol Cancer 23: 1145-1149, 2013.
- 30. Coronado PJ, Fasero M, Magrina JF, Herraiz MA and Vidart JA: Comparison of perioperative outcomes and cost between robotic-assisted and conventional laparoscopy for transperitoneal infrarenal para-aortic lymphadenectomy (TIPAL). J Minim Invasive Gynecol 21: 674-681, 2014.
- 31. Díaz-Feijoo B, Gil-Ibáñez B, Pérez-Benavente A, Martínez-Gómez X, Colás E, Sánchez-Iglesias JL, Cabrera-Díaz S, Puig-Puig O, Magrina JF and Gil-Moreno A: Comparison of robotic-assisted vs conventional laparoscopy for extraperitoneal paraaortic lymphadenectomy. Gynecol Oncol 132: 98-101, 2014.

- 32. Chen CH, Chiu LH, Chen HH, Chan C and Liu WM: Comparison of robotic approach, laparoscopic approach and laparotomy in treating epithelial ovarian cancer. Int J Med Robot 12: 268-275, 2016.
- 33. Manchana T, Puangsricharoen P, Sirisabya N, Worasethsin P, Vasuratna A, Termrungruanglert W and Tresukosol D: Comparison of perioperative and oncologic outcomes with laparotomy, and laparoscopic or robotic surgery for women with endometrial cancer. Asian Pac J Cancer Prev 16: 5483-5488, 2015. 34. Mäenpää MM, Nieminen K, Tomás EI, Laurila M, Luukkaala TH
- and Mäenpää JU: Robotic-assisted vs traditional laparoscopic surgery for endometrial cancer: A randomized controlled trial. Am J Obstet Gynecol 215: 588.e1-588.e7, 2016.
- Mendivil AA, Rettenmaier MA, Abaid LN, Brown JV III, Micha JP, Lopez KL and Goldstein BH: Survival rate comparisons amongst cervical cancer patients treated with an open, robotic-assisted or laparoscopic radical hysterectomy: A five year experience. Surg Oncol 25: 66-71, 2016.
- 36. Pellegrino A, Damiani GR, Loverro M, Pirovano C, Fachechi G, Corso S and Trojano G: Comparison of robotic and laparoscopic radical type-B and C hysterectomy for cervical cancer: Long term-outcomes. Acta Biomed 88: 289-296, 2017.
- 37. Nie JC, Yan AQ and Liu XS: Robotic-assisted radical hysterectomy results in better surgical outcomes compared with the traditional laparoscopic radical hysterectomy for the treatment of cervical cancer. Int J Gynecol Cancer 27: 1990-1999, 2017.
- 38. Luo C, Liu M and Li X: Efficacy and safety outcomes of robotic radical hysterectomy in Chinese older women with cervical cancer compared with laparoscopic radical hysterectomy. BMC Womens Health 18: 61, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.