

Combined use of a 980-nm diode laser and preoperative intravesical instillation of pirarubicin for the prevention of short-term recurrence of non-muscle invasive bladder cancer: A pilot study

ZHEN TANG^{1*}, JIAO PENG^{2*}, SHANGXIN WU¹, FENFEN ZHAO¹ and MINGUI WU¹

Departments of ¹Urology and ²Medical Imaging, The Fourth Affiliated Hospital of Guangxi Medical University, Liuzhou, Guangxi Zhuang Autonomous Region 545005, P.R. China

Received February 18, 2023; Accepted May 31, 2023

DOI: 10.3892/ol.2023.13908

Abstract. At present, transurethral resection of bladder tumors (TURBT) is the main surgical method for treating non-muscle invasive bladder cancer (NMIBC), but its postoperative recurrence needs to be prevented. The aim of the present study was to investigate the efficacy of a 980-nm diode laser combined with preoperative intravesical instillation of pirarubicin (THP) for the prevention of NMIBC recurrence. The data of 120 patients with NMIBC who underwent transurethral resection between May 2021 and July 2022 were retrospectively collected, and these patients were followed up. The patients were divided into four groups based on the surgical method used and preoperative intravesical instillation of THP as follows: i) 980-nm diode laser with THP (LaT); ii) 980-nm diode laser alone (La); iii) TURBT with THP (TUT); and iv) TURBT alone (TU). Clinicopathological variables, postoperative complications and short-term outcomes among the aforementioned groups were analyzed. The blood loss volume and the incidence of perforation and delayed bleeding were significantly lower in the LaT and La groups compared with those in the TUT and

TU groups. The days of bladder irrigation, catheter extubation and postoperative hospitalization were significantly shorter in the LaT and La groups compared with the TUT and TU groups. The detection rate of suspicious lesions was significantly higher in the THP irrigation groups (LaT and TUT) compared with that in the saline irrigation groups (La and TU). Tumor diameter and number, 980-nm laser and THP irrigation were shown to be independent risk factors in the Cox regression analysis. In addition, the recurrence-free survival (RFS) rate of the LaT group was significantly higher than that of the other three groups. In conclusion, a 980-nm diode laser can effectively reduce intraoperative blood loss and the incidence of perforation, and accelerate postoperative recovery. Preoperative intravesical instillation of THP is conducive to identifying suspicious lesions. The combination of a 980-nm laser with preoperative THP intravesical instillation can significantly prolong RFS time.

Introduction

Bladder cancer is one of the most common malignancies worldwide (1). In total, ~70% of patients with bladder cancer are initially diagnosed with non-muscle invasive bladder cancer (NMIBC), which is confined to the muscular layer and submucosa (2,3). Transurethral resection of bladder tumors (TURBT) is the primary surgical procedure for NMIBC (4,5), but >50% of patients experience recurrence within 1 to 2 years. Therefore, there is an urgent need to find novel treatments to reduce the rate of postoperative recurrence of NMIBC.

After recurrence of bladder cancer, progression to MIBC is likely (6), and radical resection of bladder cancer is needed at a later stage, which is associated with increased pain and a worse prognosis. Currently, TURBT combined with adjuvant therapy is considered to be effective in reducing the incidence of postoperative recurrence (7,8). Adjuvant therapy drugs, such as mitomycin C (MMC) (9,10), epirubicin (EPI) (11) and pirarubicin (THP) (12), have increased benefits due to their ability to be used in intravesical instillation. Due to its safety, low number of toxic side effects and price, THP has been widely used in the clinic. However, it is mainly used for postoperative adjuvant therapy, and there is still a lack of relevant studies

Correspondence to: Dr Zhen Tang or Dr Mingui Wu, Department of Urology, The Fourth Affiliated Hospital of Guangxi Medical University, 156 Heping Road, Liuzhou, Guangxi Zhuang Autonomous Region 545005, P.R. China
E-mail: tzykd104@163.com
E-mail: wumingui1968@163.com

*Contributed equally

Abbreviations: NMIBC, non-muscle invasive bladder cancer; TURBT, transurethral resection of bladder tumors; THP, pirarubicin; LaT, laser with THP; La, laser alone; TUT, TURBT with THP; TU, TURBT alone; RFS, recurrence-free survival; MMC, mitomycin C; EPI, epirubicin; ASA, American Society of Anesthesiologists; ONR, obturator nerve reflex; BI, bladder irrigation; CE, catheter extubation; PH, postoperative hospitalization

Key words: NMIBC, THP, 980-nm diode laser

proving the efficacy of preoperative intravesical instillation for the prevention of bladder cancer recurrence.

In recent years, laser en bloc resection with a 980-nm diode laser has been shown to exhibit a notably reduced incidence of complications and recurrence compared with conventional TURBT surgery (13). For diode lasers at a 980-nm wavelength, a new generation of laser technology, their energy can be absorbed by both water and hemoglobin. Compared with conventional transurethral electric resection, due to the tissue vaporization and cutting ability, coagulation characteristics and controllable penetration depth of 980-nm diode laser, this technique has shown increased efficacy and safety, and can markedly reduce the amount of bleeding and the risk of bladder perforation (13,14). However, the efficacy of the combination of 980-nm diode laser en bloc resection and preoperative THP intravesical instillation for the treatment of patients with NMIBC remains unclear. In the present study, the aim was to investigate the effect of combining 980-nm diode laser en bloc resection with preoperative intravesical instillation of THP on the rates of postoperative recurrence and complications of NMIBC.

Materials and methods

Patients. A total of 120 patients who were diagnosed with NMIBC at the Department of Urology, The Fourth Affiliated Hospital of Guangxi Medical University (Liuzhou, China) between May 2021 and July 2022 were retrospectively included in the present study. Patients with Ta-T1N0M0 (8th American Joint Committee on Cancer TNM staging system) (15) bladder tumors that were diagnosed by preoperative magnetic resonance imaging examination and who underwent surgery were included in the current study. Exclusion criteria included: i) Patients with recurrent or muscle-invasive bladder tumors; ii) patients with other tumors; and iii) patients with distant metastasis. The patients were treated using either a 980-nm diode laser combined with preoperative intravesical instillation of THP (LaT), a 980-nm laser alone (La), TURBT combined with preoperative intravesical instillation of THP (TUT) or TURBT alone (TU). In the laser surgery group, the low-power 980-nm diode laser was used to free the tumor in the submucosa, and then the tumor was completely removed. Subsequently, the high-power 980-nm diode laser was used to vaporize the edge and base of the tumor to further ensure negative margins.

The patients in the LaT and TUT groups received preoperative intravesical instillation of 30 mg THP/30 ml saline for 15 min. The patients in the La and TU groups received preoperative intravesical instillation of 30 ml saline for 15 min. During the operation, a biopsy was performed at the THP staining sites (orange staining) in the LaT and TUT groups. For those patients with ≤ 3 stained sites, a biopsy was taken within 2 cm of the tumor to supplement the remaining number. In the La and TU groups, a biopsy was performed at three random sites within 2 cm of the tumor. Intravesical instillation chemotherapy with 50 ml gemcitabine (20 mg/ml) was performed for 1 year after surgery. The first chemotherapy session started within 24 h after surgery, once a week for the first 2 months and once a month after 2 months. Each patient was followed up and underwent cystoscopy every 3 months.

Data collection. Clinicopathological data were also collected and evaluated, including sex, age, American Society of Anesthesiologists (ASA) grade (16), tumor location, maximum tumor diameter, tumor multiplicity, grade (17), tumor (T) stage (15), operation time, blood loss volume, perforation, obturator nerve reflex (ONR), days of bladder irrigation (BI), days of catheter extubation (CE), days of postoperative hospitalization (PH) and delayed bleeding.

Statistical analysis. All statistical analyses were performed using SPSS (version 23.0; IBM Corp.). One-way analysis of variance was performed to analyze the difference in continuous variables among the four groups, followed by the Bonferroni post hoc test. Continuous variables are presented as the mean \pm SD. χ^2 or Fisher's exact test was employed to evaluate categorical variables. Univariate and multivariate Cox regression analyses were applied to identify independent risk factors. Kaplan-Meier analysis was used to compare the survival differences among the four groups, with statistical analysis using the log-rank test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Baseline characteristics of patients. The clinicopathological characteristics of patients with NMIBC in the four different groups, namely, the LaT group ($n=32$), the La group ($n=30$), the TUT group ($n=29$) and the TU group ($n=29$), are shown in Table I. There was no significant difference with regard to sex, age, ASA, tumor location, tumor diameter, multiplicity of tumor, T stage and grade among the four groups.

Perioperative surgical outcomes and detection of lesions. The perioperative surgical outcomes among the four different groups are listed in Table II. No significant differences in operation time or ONR were observed among the four groups. However, ONR did not occur in the two 980-nm laser groups but it did occur in the two TURBT groups, with a single case in the TUT group (3.4%) and two cases in the TU group (6.9%). There were significant differences among the four groups in terms of blood loss, perforation, days of BI, CE and PH, and delayed bleeding ($P < 0.05$). Among the four groups, the blood loss in the two 980-nm laser groups (LaT group, 8.8 ± 2.6 ml; La group, 8.8 ± 2.3 ml) was significantly lower than that in the two TURBT groups (TUT group, 17.1 ± 5.3 ml; TU group, 15.7 ± 6.2 ml; $P < 0.001$). For perforation, the incidence was similar to ONR, but statistically significant, with no perforation events occurring in the two laser groups but four cases in the TUT group (13.8%) and three cases in the TU group (10.3%). Similarly, the days of BI, CE and PH were observed to be significantly shorter in the two laser groups than in the two TURBT groups, and the time was shortened by ~ 2 days ($P < 0.001$). In addition, there was a significant difference in delayed bleeding among the four groups, with no delayed bleeding in the two laser groups but seven cases of delayed bleeding in the TURBT groups; three in the TUT group (10.3%) and four in the TU group (13.8%). As illustrated in Table III, the detection rate of suspicious lesions in the preoperative intravesical instillation of THP (LaT and TUT) groups was significantly higher than that in the other two (La and TU) groups without THP ($P < 0.001$).

Table I. Clinicopathological characteristics of patients with non-muscle invasive bladder cancer in the LaT (n=32), La (n=30), TUT (n=29) and TU (n=29) groups.

Clinicopathological characteristics	LaT group	La group	TUT group	TU group	P-value
Sex, n (%)					0.936
Male	26 (81.3)	24 (80.0)	22 (75.9)	22 (75.9)	
Female	6 (18.7)	6 (20.0)	7 (24.1)	7 (24.1)	
Age, years ^a	64.7±7.5	64.9±5.7	65.1±8.1	65.2±9.3	0.995
ASA, n (%)					0.985
I	17 (53.1)	14 (46.7)	14 (48.3)	12 (41.4)	
II	14 (43.8)	15 (50.0)	14 (48.3)	16 (55.2)	
III	1 (3.1)	1 (3.3)	1 (3.4)	1 (3.4)	
Tumor location, n (%)					0.274
Lateral	19 (59.4)	14 (46.7)	14 (48.3)	20 (69.0)	
Others	13 (40.6)	16 (53.3)	15 (51.7)	9 (31.0)	
Tumor diameter, cm ^a	2.6±0.8	2.4±0.9	2.8±1.0	2.3±0.6	0.090
Multiplicity of tumor, n (%)					0.800
Single	28 (87.5)	24 (80.0)	23 (79.3)	23 (79.3)	
Multiple	4 (12.5)	6 (20.0)	6 (20.7)	6 (20.7)	
T stage, n (%)					0.688
Ta	24 (75.0)	19 (63.3)	20 (69.0)	18 (62.1)	
T1	8 (25.0)	11 (36.7)	9 (31.0)	11 (37.9)	
WHO 2004 grade, n (%)					0.206
PUNLMP	17 (53.1)	10 (33.3)	9 (31.0)	6 (20.7)	
Low	10 (31.3)	10 (33.3)	12 (41.4)	14 (48.3)	
High	5 (15.6)	10 (33.3)	8 (27.6)	9 (31.0)	

^aData are presented as the mean ± SD. PUNLMP, papillary urothelial neoplasm of low malignant potential; ASA, American Society of Anesthesiologists; T, tumor; LaT, 980-nm diode laser with pirarubicin; La, 980-nm diode laser alone; TUT, transurethral resection of bladder tumors with pirarubicin; TU, transurethral resection of bladder tumors alone.

Table II. Comparison of perioperative surgical outcomes in patients with non-muscle invasive bladder cancer in the LaT (n=32), La (n=30), TUT (n=29) and TU (n=29) groups.

Perioperative surgical outcomes	LaT group	La group	TUT group	TU group	P-value
Operation time, min ^a	45.6±8.2	43.8±9.4	44.2±8.4	47.4±8.5	0.377
Blood loss, ml ^a	8.8±2.6	8.8±2.3	17.1±5.3	15.7±6.2	<0.001
Perforation, n (%)	0 (0.0)	0 (0.0)	4 (13.8)	3 (10.3)	<0.05
ONR, n (%)	0 (0.0)	0 (0.0)	1 (3.4)	2 (6.9)	0.142
Days of BI ^a	1.4±0.5	1.6±0.5	3.3±1.0	3.4±1.2	<0.001
Days of CE ^a	2.4±0.5	2.6±0.5	4.3±1.0	4.4±1.2	<0.001
Days of PH ^a	3.7±0.7	3.9±0.7	5.7±1.0	5.7±1.0	<0.001
Delayed bleeding, n (%)	0 (0.0)	0 (0.0)	3 (10.3)	4 (13.8)	<0.05

^aData are presented as the mean ± SD. BI, bladder irrigation; CE, catheter extubation; PH, postoperative hospitalization; LaT, 980-nm diode laser with pirarubicin; La, 980-nm diode laser alone; TUT, transurethral resection of bladder tumors with pirarubicin; TU, transurethral resection of bladder tumors alone; ONR, obturator nerve reflex.

Cox regression analysis and survival analysis. To further identify independent risk factors for NMIBC recurrence, both univariate and multivariate Cox regression analyses were employed. As shown in Table IV, the univariate and

multivariate Cox regression analyses revealed that maximum tumor diameter ≥3 cm, tumor multiplicity, 980-nm laser surgery and preoperative intravesical instillation of THP were independent risk factors for recurrence of NMIBC. Among

Table III. Difference in the detection rate of suspected lesions by preoperative intravesical instillation of THP in groups with no irrigation with THP (La and TU) (n=177) and irrigation with THP (LaT and TUT) (n=183)^a.

Groups	No irrigation with THP	Irrigation with THP	P-value
Lesions, n (%)	6 (3.4)	24 (13.1)	<0.001

^aThree biopsy samples were taken per patient. THP, pirarubicin.

Table IV. Results of univariate and multivariate Cox regression analyses.

Variable	Univariate			Multivariate		
	HR	95% CI	P-value	HR	95% CI	P-value
Sex	1.679	0.487-5.781	0.412			
Age, years	1.034	0.972-1.100	0.283			
ASA	1.334	0.565-3.147	0.511			
Lateral	1.778	0.713-4.429	0.217			
Diameter, >3 cm	6.140	2.032-18.554	<0.01	6.155	1.716-22.083	<0.01
Multiplicity	5.134	2.073-12.715	<0.001	4.819	1.778-13.059	<0.01
Operation time	1.017	0.960-1.078	0.558			
Blood loss, ml	1.023	0.955-1.097	0.515			
Grade	1.822	1.030-3.224	<0.05	1.180	0.558-2.494	0.665
T stage	2.656	1.077-6.552	<0.05	1.652	0.560-4.872	0.363
980-nm laser	0.364	0.138-0.959	<0.05	0.266	0.088-0.803	<0.05
Irrigation with THP	0.394	0.149-1.036	0.059	0.195	0.065-0.584	<0.01

HR, hazard ratio; CI, confidence interval; T, tumor; ASA, American Society of Anesthesiologists; THP, pirarubicin.




Figure 1. Comparison of RFS in patients with non-muscle invasive bladder cancer divided into the four treatment groups. The RFS rate of the LaT group was higher than that of the other three groups, including the La, TUT and TU groups. RFS, recurrence-free survival; LaT, 980-nm diode laser combined with preoperative intravesical instillation of pirarubicin; La, 980-nm diode laser alone; TUT, transurethral resection of bladder tumors combined with preoperative intravesical instillation of pirarubicin; TU, transurethral resection of bladder tumors alone.

tumor multiplicity (HR, 4.819; 95% CI, 1.778-13.059; $P<0.01$) were adverse risk factors, while 980-nm laser surgery (HR, 0.266; 95% CI, 0.088-0.803; $P<0.05$) and irrigation with THP (HR, 0.195; 95% CI, 0.065-0.584; $P<0.01$) were protective risk factors. Moreover, a survival analysis was conducted to further investigate the effect of combining the 980-nm laser with preoperative intravesical instillation of THP on the recurrence-free survival (RFS) rate of patients with NMIBC. The Kaplan-Meier survival curves are plotted in Fig. 1. The results indicated that the RFS rate of the LaT group was significantly higher than that of the other three groups ($P=0.033$).

Discussion

To date, traditional TURBT is the standard surgical treatment for NMIBC, but it has some intraoperative complications, including bleeding, bladder perforation and ONR (18), which are closely associated with the surgical experience and skill of surgeons. Meanwhile, due to the segmented resection method of TURBT surgery, tumor cells may be shed during the operation to some extent, thereby increasing the risks of recurrence and metastasis (19,20). To reduce the occurrence of these complications, surgeons have conducted additional research on new approaches. In 1978, Staehler *et al* (21) first explored the use of laser surgery for bladder neoplasms. Studies have shown that compared with TURBT surgery, laser procedures

them, the maximum tumor diameter ≥ 3 cm [hazard ratio (HR), 6.155; 95% confidence interval (CI), 1.716-22.083; $P<0.01$] and

such as holmium (22-24), 2- μm (25-27), 1.9- μm Vela (28) and potassium-titanyl-phosphate (29,30) lasers used to treat NMIBC can not only reduce the incidence of complications, but can also reduce the postoperative recurrence rate when used with en bloc resection. However, studies on en bloc resection of NMIBC with a 980-nm diode laser have rarely been reported (13,31).

In the present study, the intraoperative and postoperative characteristics, and the recurrence status of patients with NMIBC were retrospectively compared in the LaT, La, TUT and TU groups. It was shown that 980-nm diode laser en bloc resection was more effective at reducing intraoperative blood loss than TURBT surgery. Furthermore, intraoperative perforation and ONR did not occur in the two 980-nm diode laser surgery groups, while four cases of perforation and a single case of ONR occurred in the TUT group, and three cases of perforation and two cases of ONR occurred in the TU group. Due to its improved hemostatic effect, the number of days of PI, CE and PH was significantly decreased compared with that in the TURBT group. In addition, no delayed bleeding was observed in the two 980-nm laser groups. According to the aforementioned results, compared with TURBT surgery, the 980-nm diode laser has similar advantages to other lasers (27,28,32-34), and the treatment of NMIBC with a 980-nm diode laser can not only decrease the incidence of intraoperative and postoperative complications, but can also accelerate postoperative recovery. More importantly, the laser groups had a lower 1-year recurrence rate, which significantly improved RFS time in the patients with NMIBC.

To prevent recurrence in patients with NMIBC, transurethral resection is usually followed by adjuvant therapy. Postoperative intravesical instillation of chemotherapeutic agents, such as MMC, EPI and THP, can effectively improve prognosis and prolong RFS time in patients with NMIBC (7). On the one hand, it has been reported that THP can be rapidly absorbed by tumor cells, and enter the nucleus to inhibit DNA synthesis (35). A meta-analysis study showed that intravesical instillation of THP significantly reduced the risk of tumor recurrence and was superior to MMC and EPI (8). On the other hand, according to the characteristics of the THP solution, preoperative intravesical instillation of THP can stain tumors orange, which may be conducive to the detection of small lesions and reduce the risk of missed detection. However, the effect of preoperative intravesical instillation of THP on postoperative recurrence in patients with NMIBC remains unclear. Considering the advantages of both the 980-nm diode laser and the intravesical instillation of THP solution, the combined effect of preoperative intravesical instillation of THP and the 980-nm diode laser was investigated on the postoperative recurrence of NMIBC in the present study.

In the present study, the proportion of suspicious lesions detected in the two THP groups, LaT and TUT, was significantly higher than that in the random sampling groups, La and TU, suggesting that preoperative intravesical instillation of THP was beneficial for detecting suspicious and unobtrusive lesions. Furthermore, Cox regression analysis showed that both the preoperative intravesical instillation of THP and 980-nm diode laser surgery were protective factors for the recurrence of NMIBC, which significantly reduced the risk of tumor recurrence. In addition, the survival analysis showed that

among the four groups, the LaT group exhibited a significantly higher 1-year RFS rate for patients with NMIBC compared with that in the other three groups, followed by the La, TUT and TU groups. The results indicated that preoperative THP intravesical instillation or en bloc resection with a 980-nm diode laser could effectively prolong RFS time in patients with NMIBC and reduce the postoperative recurrence rate.

However, there are some limitations to the current study. The study is retrospective rather than a double-blind prospective randomized controlled study, which may have led to selection bias to some extent. Additionally, the small sample size of this study may also have led to the deviation of results. In the future, a randomized controlled study will be performed to draw more rigorous conclusions.

In summary, compared with TURBT surgery alone, 980-nm diode laser en bloc resection combined with preoperative intravesical instillation of THP not only assists in reducing the incidence of intraoperative and postoperative complications, but also accelerates postoperative recovery in patients and reduces the risk of recurrence of NMIBC.

Acknowledgements

Not applicable.

Funding

The present study was funded by the Guangxi Zhuang Autonomous Region Health and Family Planning Commission (grant no. Z20210478).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

ZT prepared the manuscript and performed the statistical analysis. JP performed the data analysis. SW and FZ contributed to the data acquisition, analysis and interpretation. MW presented the concept of the study and designed the study. ZT and PJ confirm the authenticity of all the raw data. All authors read and approved the final manuscript.

Ethics approval and consent to participate

This study was approved by the Ethics Committee of The Fourth Affiliated Hospital of Guangxi Medical University (Liuzhou, China) (approval no. KY2021053) and written informed consent was obtained from all patients.

Patient consent for publication

Written informed consent was obtained from the patients to publish this paper.

Competing interests

The authors declare that they have no competing interests.

References

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A and Bray F: Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 71: 209-249, 2021.
- Kirkali Z, Chan T, Manoharan M, Algaba F, Busch C, Cheng L, Kiemeny L, Kriegmair M, Montironi R, Murphy WM, *et al*: Bladder cancer: Epidemiology, staging and grading, and diagnosis. *Urology* 66: 4-34, 2005.
- Kamat AM, Hahn NM, Efstathiou JA, Lerner SP, Malmström PU, Choi W, Guo CC, Lotan Y and Kassouf W: Bladder cancer. *Lancet* 388: 2796-2810, 2016.
- Babjuk M, Böhle A, Burger M, Tobisawa Y, Yoneyama T, Yamamoto H, Imai H, Yoneyama T, Hashimoto Y, *et al*: EAU Guidelines on Non-Muscle-invasive Urothelial Carcinoma of the Bladder: Update 2016. *Eur Urol* 71: 447-461, 2017.
- Nishiyama H: Asia Consensus Statement on NCCN Clinical Practice Guideline for bladder cancer. *Jpn J Clin Oncol* 48: 3-6, 2018.
- Sylvester RJ: Natural history, recurrence, and progression in superficial bladder cancer. *ScientificWorldJournal* 6: 2617-2625, 2006.
- Sylvester RJ, Oosterlinck W, Holmang S, Sydes MR, Birtle A, Gudjonsson S, De Nunzio C, Okamura K, Kaasinen E, Solsona E, *et al*: Systematic review and individual patient data meta-analysis of randomized trials comparing a single immediate instillation of chemotherapy after transurethral resection with transurethral resection alone in patients with stage pTa-pT1 urothelial carcinoma of the bladder: Which patients benefit from the instillation? *Eur Urol* 69: 231-244, 2016.
- Kang M, Jeong CW, Kwak C, Kim HH and Ku JH: Single, immediate postoperative instillation of chemotherapy in non-muscle invasive bladder cancer: A systematic review and network meta-analysis of randomized clinical trials using different drugs. *Oncotarget* 7: 45479-45488, 2016.
- Solsona E, Iborra I, Ricos JV, Monros JL, Casanova J and Dumont R: Effectiveness of a single immediate mitomycin C instillation in patients with low risk superficial bladder cancer: Short and long-term followup. *J Urol* 161: 1120-1123, 1999.
- Tolley DA, Parmar MK, Grigor KM, Lallemand G, Benyon LL, Fellows J, Freedman LS, Grigor KM, Hall RR, Hargreave TB, *et al*: The effect of intravesical mitomycin C on recurrence of newly diagnosed superficial bladder cancer: A further report with 7 years of follow up. *J Urol* 155: 1233-1238, 1996.
- Oosterlinck W, Kurth KH, Schroder F, Bultinck J, Hammond B and Sylvester R: A prospective European Organization for Research and Treatment of Cancer Genitourinary Group randomized trial comparing transurethral resection followed by a single intravesical instillation of epirubicin or water in single stage Ta, T1 papillary carcinoma of the bladder. *J Urol* 149: 749-752, 1993.
- Okamura K, Ono Y, Kinukawa T, Matsuura O, Yamada S, Ando T, Fukatsu T, Ohno Y and Ohshima S: Nagoya University Urological Oncology Group: Randomized study of single early instillation of (2'R)-4'-O-tetrahydropyranyl-doxorubicin for a single superficial bladder carcinoma. *Cancer* 94: 2363-2368, 2002.
- Mao T, Zhang H, Cui J, Zhao Z, Jiao D and Zhang W: The 980 nm diode laser treatment for non-muscle-invasive bladder tumor with en bloc technique: Single-center experience. *World J Surg Oncol* 20: 324, 2022.
- Wendt-Nordahl G, Huckele S, Honeck P, Alken P, Knoll T, Michel MS and Häcker A: 980-nm Diode laser: A novel laser technology for vaporization of the prostate. *Eur Urol* 52: 1723-1728, 2007.
- Magers MJ, Lopez-Beltran A, Montironi R, Williamson SR, Kaimakliotis HZ and Cheng L: Staging of bladder cancer. *Histopathology* 74: 112-134, 2019.
- De Cassai A, Boscolo A, Tonetti T, Ban I and Ori C: Assignment of ASA-physical status relates to anesthesiologists' experience: A survey-based national study. *Korean J Anesthesiol* 72: 53-59, 2019.
- Varma M, Delahunt B and van der Kwast T: Grading Noninvasive Bladder Cancer: World Health Organisation 1973 or 2004 May Be the Wrong Question. *Eur Urol* 76: 413-415, 2019.
- Xu J, Wang C, Ouyang J, Sun J and Hu C: Efficacy and Safety of transurethral laser surgery versus transurethral resection for Non-Muscle-invasive bladder cancer: A Meta-Analysis and systematic review. *Urol Int* 104: 810-823, 2020.
- Engilbertsson H, Aaltonen KE, Björnsson S, Kristmundsson T, Patschan O, Rydén L and Gudjonsson S: Transurethral bladder tumor resection can cause seeding of cancer cells into the bloodstream. *J Urol* 193: 53-57, 2015.
- Liang H, Yang T, Wu K, He D and Fan J: En bloc resection improves the identification of muscularis mucosae in non-muscle invasive bladder cancer. *World J Urol* 37: 2677-2682, 2019.
- Stähler G, Schmiedt E and Hofstetter A: Destruction of bladder neoplasms by means of transurethral neodym-YAG-laser coagulation. *Helv Chir Acta* 45: 307-311, 1978 (In German).
- D'Souza N and Verma A: Holmium laser transurethral resection of bladder tumor: Our experience. *Urol Ann* 8: 439-443, 2016.
- Kramer MW, Rassweiler JJ, Klein J, Martov A, Baykov N, Lusuardi L, Janetschek G, Hürle R, Wolters M, Abbas M, *et al*: En bloc resection of urothelium carcinoma of the bladder (EBRUC): A European multicenter study to compare safety, efficacy, and outcome of laser and electrical en bloc transurethral resection of bladder tumor. *World J Urol* 33: 1937-1943, 2015.
- Xishuang S, Deyong Y, Xiangyu C, Tao J, Quanlin L, Hongwei G, Jibin Y, Dongjun W, Zhongzhou H, Jianbo W, *et al*: Comparing the safety and efficiency of conventional monopolar, plasmakinetic, and holmium laser transurethral resection of primary non-muscle invasive bladder cancer. *J Endourol* 24: 69-73, 2010.
- Chen X, Liao J, Chen L, Qiu S, Mo C, Mao X, Yang Y, Zhou S and Chen J: En bloc transurethral resection with 2-micron continuous-wave laser for primary non-muscle-invasive bladder cancer: A randomized controlled trial. *World J Urol* 33: 989-995, 2015.
- Liu H, Wu J, Xue S, Zhang Q, Ruan Y, Sun X and Xia S: Comparison of the safety and efficacy of conventional monopolar and 2-micron laser transurethral resection in the management of multiple nonmuscle-invasive bladder cancer. *J Int Med Res* 41: 984-992, 2013.
- Zhong C, Guo S, Tang Y and Xia S: Clinical observation on 2 micron laser for non-muscle-invasive bladder tumor treatment: Single-center experience. *World J Urol* 28: 157-161, 2010.
- Xu H, Ma J, Chen Z, Yang J, Yuan H, Wang T, Liu J, Yang W and Ye Z: Safety and efficacy of en bloc transurethral resection With 1.9 microm vella laser for treatment of Non-Muscle-invasive bladder cancer. *Urology* 113: 246-250, 2018.
- Cheng B, Qiu X, Li H and Yang G: The safety and efficacy of front-firing green-light laser endoscopic en bloc photoselective vapo-enucleation of non-muscle-invasive bladder cancer. *Ther Clin Risk Manag* 13: 983-988, 2017.
- Yang D, Xue B, Zang Y, Liu X, Zhu J, Zhou Y and Shan Y: Efficacy and safety of potassium-titanyl-phosphate laser vaporization for clinically non-muscle invasive bladder cancer. *Urol J* 11: 1258-1263, 2014.
- Tao W, Sun C, Yao Q, Fu K, Shan Y, Zhang Y, Xue B and Yang D: The clinical study of en bloc transurethral resection with 980 nm laser for treatment of primary non-muscle invasive bladder cancer. *J Xray Sci Technol* 28: 563-571, 2020.
- Chen J, Zhao Y, Wang S, Jin X, Sun P, Zhang L and Wang M: Green-light laser en bloc resection for primary non-muscle-invasive bladder tumor versus transurethral electro-resection: A prospective, nonrandomized two-center trial with 36-month follow-up. *Lasers Surg Med* 48: 859-865, 2016.
- Migliari R, Buffardi A and Ghabin H: Thulium laser endoscopic en bloc Enucleation of Nonmuscle-Invasive bladder cancer. *J Endourol* 29: 1258-1262, 2015.
- Xu Y, Guan W, Chen W, Xie C, Ouyang Y, Wu Y and Liu C: Comparing the treatment outcomes of potassium-titanyl-phosphate laser vaporization and transurethral electroresection for primary nonmuscle-invasive bladder cancer: A prospective, randomized study. *Lasers Surg Med* 47: 306-311, 2015.
- Kunimoto S, Miura K, Takahashi Y, Takeuchi T and Umezawa H: Rapid uptake by cultured tumor cells and intracellular behavior of 4'-O-tetrahydropyranyladriamycin. *J Antibiot (Tokyo)* 36: 312-317, 1983.



Copyright © 2023 Tang et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.