

Pioglitazone, a peroxisome proliferator-activated receptor γ agonist, induces cell death and inhibits the proliferation of hypoxic HepG2 cells by promoting excessive production of reactive oxygen species

GUOHAO HUANG^{*}, MENG FAN ZHANG^{*}, MANZHOU WANG, WENZE XU,
XUHUA DUAN, XINWEI HAN and JIANZHUANG REN

Department of Interventional Radiology, The First Affiliated Hospital of Zhengzhou University,
Zhengzhou, Henan 450052 P.R. China

Received August 31, 2023; Accepted January 19, 2024

DOI: 10.3892/ol.2024.14294

Abstract. Hypoxia is a hallmark of solid tumors. Hypoxic cancer cells adjust their metabolic characteristics to regulate the production of cellular reactive oxygen species (ROS) and facilitate ROS-mediated metastasis. Peroxisome proliferator-activated receptor γ (PPAR γ) is a nuclear receptor that regulates the transcription of fatty acid metabolism-related genes that have a key role in the survival and proliferation function of hypoxic cancer cells. In the present study, mRNA expression in HepG2 cells under chemically induced hypoxia was assessed. The protein expression levels of hypoxia-inducible factor 1 α (HIF-1 α) were measured using western blotting. Following treatment with the PPAR γ agonist pioglitazone, cell viability was assessed using a Cell Counting Kit-8 assay, whilst cell proliferation and death were determined using 5-ethynyl-2'-deoxyuridine incorporation staining, and calcein-acetoxymethyl ester and propidium iodide staining, respectively. Cellular ROS production was assessed using dihydroethidium staining. Cobalt chloride was used to induce hypoxia in HepG2 cells, which was evaluated using HIF-1 α expression. The results revealed that the mRNA expression of PPAR γ , CD36, acetyl-co-enzyme A dehydrogenase (ACAD) medium chain (ACADM) and ACAD short-chain (ACADS) was downregulated in hypoxic HepG2 cells. The PPAR γ agonist pioglitazone decreased

the cell viability of hypoxic HepG2 cells by inhibiting cell proliferation and inducing cell death. Following treatment with the PPAR γ agonist pioglitazone, hypoxic HepG2 cells produced excessive ROS. ROS-mediated cell death induced by the PPAR γ agonist pioglitazone was rescued with the antioxidant N-acetyl-L-cysteine. The downregulated mRNA expression of PPAR γ , CD36, ACADM and ACADS was not reverted by a PPAR γ agonist in hypoxic HepG2 cells. By contrast, the PPAR γ agonist suppressed the mRNA expression of BCL2, which was upregulated in hypoxic HepG2 cells. In summary, the PPAR γ agonist stimulated excessive ROS production to inhibit cell proliferation and increase the death of hypoxic HepG2 cells by decreasing BCL2 mRNA expression, suggesting a negative association between PPAR γ and BCL2 in the regulation of ROS production in hypoxic HepG2 cells.

Introduction

Oxidative stress is a condition that involves the excessive production of reactive oxygen species (ROS) to higher levels than antioxidants, and is closely associated with carcinogenesis and cancer progression (1). Excessive ROS production is toxic for cell survival; however, cancer cells exhibit metabolic characteristics enabling them to adjust the antioxidant status and promote ROS-mediated metastasis (2,3). Peroxisome proliferator-activated receptors (PPARs) are a type of nuclear receptor with three members: PPAR α , PPAR γ and PPAR δ . Among them, PPAR γ acts as a pivotal regulator of fatty acid degradation. PPAR γ activation facilitates fatty acid uptake (4,5), and it has been reported that PPAR γ agonists promote the expression of the fatty acid transport protein and acetyl-co-enzyme A (CoA) dehydrogenase (ACAD) medium chain (ACADM) in rat liver (6). PPAR γ has also been reported to be downregulated in cells treated with hydrogen peroxide (H₂O₂), which indicates an association between oxidative stress and PPAR γ signaling (7). In addition, PPAR γ interacts with the Wnt/ β -catenin pathway to modulate oxidative stress and promote carcinogenesis (8).

Correspondence to: Dr Jianzhuang Ren, Department of Interventional Radiology, The First Affiliated Hospital of Zhengzhou University, 1 Jianshe Road, Zhengzhou, Henan 450052, P.R. China
E-mail: fccrenjz@zzu.edu.cn

^{*}Contributed equally

Key words: hypoxia, peroxisome proliferator-activated receptor γ , reactive oxygen species, HepG2, oxidative stress

Hypoxia is a hallmark of solid tumors. Cancer cells reprogram their metabolic characteristics to adapt to hypoxic conditions. Apart from the well-known Warburg effect, fatty acid metabolism reprogramming provides energy and macromolecules required for the proliferation, division and survival of cancer cells (9). Fatty acid degradation mainly occurs via β -oxidation, is processed in mitochondria, and is involved in mitochondrial ROS and ATP production and acetyl-CoA recycling (10). β -oxidation is the main form of fatty acid degradation. ACADs consist of ACAD short-chain (ACADS), ACADM, ACAD long-chain and ACAD very-long chain, which degrade short, medium, long and very long-chain fatty acids, respectively (11). Overactive fatty acid β -oxidation is reported to produce ROS and promote metastasis (12). Inhibition of fatty acid oxidation reduces tumor growth and metastasis (13,14); however, ACADM activity suppresses cancer progression (11,15). This conflicting evidence indicates that the role of fatty acid β -oxidation in cancer cells remains elusive.

Oxidative stress and hypoxia are promoters of carcinogenesis, and involve PPAR γ . However, the real effect of PPAR γ signaling in hypoxic cancer cells remains unclear. Therefore, the present study aimed to assess the effect of the modulation of PPAR γ signaling with its agonist, pioglitazone, on hypoxic HepG2 cells.

Materials and methods

Cell culture. HepG2 cells were purchased from the China Center for Type Culture Collection and were cultured in DMEM (Gibco; Thermo Fisher Scientific, Inc.) supplemented with 10% FBS (Gibco; Thermo Fisher Scientific, Inc.), 100 U/ml penicillin and 10 μ g/ml streptomycin in an incubator containing 5% CO $_2$ at 37°C. The potential presence of mycoplasma in the cell line was detected regularly with a PCR kit (cat. no. C0301S, Beyotime Biotech. Inc., China) followed by the manufacturer's protocol, and no mycoplasma was detected.

Drug administration protocols. The concentration and duration of drug administration depend on the sensitivity and resistance of the cells used. Therefore, the concentration of cobalt chloride (CoCl $_2$) and pioglitazone (Sigma-Aldrich) used in the experiments ranged from 100 to 300 μ M and from 10 to 100 μ M, respectively (16-21). Based on the literature (16-21), a concentration gradient of CoCl $_2$ and pioglitazone was tested by Cell Counting Kit-8 (CCK-8) assay *in vitro* (data not shown). Cells were then treated with CoCl $_2$ (200 μ M) and pioglitazone (40 μ M) for 24, 48 or 72 h.

CCK-8 assay. HepG2 cells were seeded into a 96-well plate and cultured overnight. Before treatment, cells were replenished with medium and treated with different concentrations of CoCl $_2$ ranging from 100 to 300 μ M. When the treatment ended at 24 h or 48 h, 10 μ l CCK-8 reagent (Vazyme Biotech Co., Ltd.) was added to each well, and the plate was incubated at 37°C for 1 h. Subsequently, the absorbance at 450 nm was measured using a microplate reader (SpectraMax i3x; Molecular Devices, LLC). The optical density of each well was normalized to the control group and calculated.

Reverse transcription (RT)-quantitative (q)PCR. Gene expression levels were quantified by RT-qPCR. Total RNA was isolated from cells using TRI Reagent[®] (Sigma-Aldrich; Merck KGaA) according to the manufacturer's protocol. The concentration of RNA was determined using NanoDrop[™] 2000 (Thermo Fisher Scientific, Inc.). cDNA was synthesized from 0.5-2.5 μ g RNA by using the HiScript[®] II 1st Strand cDNA Synthesis Kit (Vazyme Biotech Co., Ltd.) at 42°C for 5 min, 37°C for 15 min and 85°C for 5 sec, performed on a T100 thermocycler (Bio-Rad Laboratories, Inc.). Gene expression was determined by qPCR using specific primers and SYBR Green (Vazyme Biotech Co., Ltd.) in a QuantStudio[™] 3 system (Thermo Fisher Scientific, Inc.) with the following thermocycling conditions: Denaturing at 95°C for 10 min, followed by 40 cycles of denaturing at 95°C for 15 sec and annealing and extension at 60°C for 1 min. Relative gene expression was calculated using the 2^{- $\Delta\Delta$ C_q} method (22). The primers used for qPCR are listed in Table I. Gene expression was measured in duplicate and was normalized using ribosomal protein S18 as the housekeeping gene.

Western blotting. Protein samples were prepared in lysis buffer (25 mmol/l HEPES, 150 mmol/l potassium acetate, 2 mmol/l EDTA pH 8.0, 0.1% NP-40, 10 mmol/l sodium fluoride, 50 mmol/l PMSF, 1 μ g/ μ l aprotinin, 1 μ g/ μ l pepstatin, 1 μ g/ μ l leupeptin and 1 mmol/l dithiothreitol). The protein determination method was the BCA protein assay (Beyotime Institute of Biotechnology), which was performed according to the manufacturer's protocol using bovine serum albumin (BSA; Servicebio, Ltd.) to prepare a standard curve. SDS-PAGE was performed using 10-20 μ g protein/lane and 4-15% gels (Beyotime Institute of Biotechnology), followed by transblotting to a 0.2 μ m nitrocellulose membrane (Amersham; Cytiva). The membranes were blocked with 5% skimmed milk (in buffer containing 10 mM Tris pH 8.0, 150 mM NaCl and 0.05% Tween 20) for 1 h at room temperature. The membranes were incubated overnight with primary antibody at 4°C and then incubated with HRP-conjugated secondary antibody for 1 h at room temperature. Protein band intensities were determined and detected with BeyoECL Star (Beyotime Institute of Biotechnology) using the Amersham[™] Imager 680 system (Amersham; Cytiva). The primary antibodies included anti-hypoxia-inducible factor (HIF)-1 α (rabbit monoclonal antibody (mAb); cat. no. 36169S; Cell Signaling Technology, Inc.) and β -actin (rabbit mAb; cat. no. 4970S; Cell Signaling Technology, Inc.), diluted 1:1,000 in 1% BSA. The secondary antibody was HRP-conjugated goat anti-rabbit immunoglobulin G (cat. no. 5127S; Cell Signaling Technology, Inc.) diluted 1:1,000.

Fluorescence microscopy

Live and dead cell staining. Live and dead cell staining was performed using the Live & Dead Kit (BioScience) following the manufacturer's instructions. HepG2 cells treated in the presence or without 200 μ M CoCl $_2$ (Co), 40 μ M pioglitazone (P) and both (Co + P) for 72 h, or HepG2 cells treated with or without 200 μ M CoCl $_2$ (Co), 40 μ M pioglitazone (P) and both (Co + P) for 48 h. At the end of the treatment, cells were replenished with serum-free DMEM containing

Table I. Primers for reverse transcription-quantitative PCR.

Gene	Forward primer, 5'→3'	Reverse primer (5'-3')
RPS18	TGCGAGTACTCAACACCAACA	CTTCGGCCCCACACCCTTAAT
PPAR γ	AGAGCCTTCCAACCTCCCTCA	TCTCCGGAAGAAACCCCTTGC
CD36	TGTGCAAAATCCACAGGAAGTG	GGCTAGAAAACGAACTCTGTACG
ACADM	GGGTTCTGGGCGATGCTG	CTGCTGTTTCGGTGAACCTCAAA
ACADS	TGAATGGAACCAAAGCCTGGA	AGGCACTGATGCCCTTGTTTT
HMOX1	ACCTTCCCCAACATTGCCAG	CAACTCCTCAAAGAGCTGGATG
BCL2	AGATTGATGGGATCGTTGCCT	AGTCTACTTCCTCTGTGATGTTGT

RPS18, ribosomal protein S18; PPAR γ , peroxisome proliferator-activated receptor γ ; ACAD, acetyl-co-enzyme A dehydrogenase; ACADM, ACAD medium-chain; ACADS, ACAD short-chain; HMOX1, heme oxygenase 1; BCL2, B cell lymphoma-2.

2 μ M calcein-acetoxymethyl ester (calcein-AM) and 4.5 μ M propidium iodide (PI), and then incubated for another 30 min at 37°C. The plate was observed under a fluorescence microscope (Leica DFC450; Leica Microsystems, Inc.). The integrated density of calcein-AM and PI was analyzed using ImageJ software (National Institutes of Health) version 1.54.

5-Ethynyl-2'-deoxyuridine (EdU) staining. EdU staining was performed using the BeyoClick EdU-488 kit (Beyotime Institute of Biotechnology) following the manufacturer's instructions. Cells were seeded into coverslips, and then cultured and treated as aforementioned. A total of 10 μ M EdU solution was added to the culture medium 6 h before the end of treatment. After incorporation of EdU, coverslips were washed with PBS and fixed with 4% paraformaldehyde for 30 min at room temperature. Coverslips were then washed again with PBS and incubated with 0.3% Triton (Beyotime Institute of Biotechnology) for 10 min at room temperature. After washing with PBS, coverslips were incubated with BeyoClick-reactive solution containing labeled azide for 30 min at room temperature. Coverslips were washed with PBS, sealed with mounting medium containing DAPI and then observed under a fluorescence microscope (Olympus ix73; Olympus Corporation). Cell count was carried out with ImageJ software (National Institutes of Health) version 1.54 using manual and automatic methods.

ROS staining. ROS staining was performed using a DHE (Dihydroethidium) assay (Abcam) following the manufacturer's instructions. Cells were cultured and treated as aforementioned. At the end of treatment, cells were replenished with serum-free DMEM containing 5 μ M DHE. Cells were cultured for another 30 min at 37°C and then observed under a fluorescence microscope (Leica DFC450; Leica Microsystems, Inc.). Integrated fluorescence density was analyzed using ImageJ.

Statistical analysis. Data are presented as the mean \pm standard deviation from ≥ 3 independent experiments. Data analysis was performed using GraphPad Prism (version 7; GraphPad; Dotmatics). Multiple group comparisons were first performed with analysis of variance (one-way ANOVA), followed by Bonferroni post hoc test if the results of ANOVA achieved statistical significance. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

PPAR γ -associated fatty acid metabolism genes are downregulated in hypoxic HepG2 cells. CoCl₂ has been widely used to induce the hypoxia of cultured cells *in vitro* (23). In the present study, CoCl₂ was used to imitate the hypoxic conditions in which HepG2 cells could survive. The maximum dose (400 μ M) of CoCl₂ that maintained $>80\%$ cell viability was determined using a CCK-8 assay (Fig. 1A). The expression of the HIF-1 α protein was notably increased in a dose-dependent manner in CoCl₂-treated HepG2 cells, which demonstrated that HepG2 cells were under hypoxic conditions (Fig. 1B). The expression of the antioxidant gene heme oxygenase 1 (HMOX1) was significantly increased ~ 10 -fold higher in hypoxic HepG2 cells treated with 400 μ M CoCl₂ than control HepG2 cells (Fig. 1C). By contrast, the mRNA expression of PPAR γ was significantly downregulated about 50% in HepG2 cells treated with 400 μ M CoCl₂ (Fig. 1C). PPAR γ is the gene encoding PPAR γ (24). In-line with the downregulation of PPAR γ , several key regulator genes associated with fatty acid metabolism, including CD36, ACADM and ACADS (11), were significantly downregulated in HepG2 cells treated with 400 μ M CoCl₂ than control HepG2 cells (Fig. 1C). These results indicated that downregulation of PPAR γ was accompanied by an increased severity of hypoxia in HepG2 cells.

PPAR γ agonist pioglitazone decreases cell proliferation and induces cell death in hypoxic HepG2 cells. To determine whether the activation of PPAR γ promoted the death of hypoxic HepG2 cells, pioglitazone, an established PPAR γ agonist (25), was added to activate PPAR γ signaling in hypoxic HepG2 cells. Pioglitazone (20-60 μ M) in combination with 200 μ M CoCl₂ significantly reduced the viability of HepG2 cells treated with 200 μ M CoCl₂ alone (Fig. 2A). Following treatment with pioglitazone, the proliferation of hypoxic HepG2 cells was further inhibited, and markedly few EdU⁺ cells were observed following treatment with a combination of CoCl₂ and pioglitazone compared with cells treated with CoCl₂ alone (Fig. 2B). The EdU⁺ cell ratio of hypoxic HepG2 cells was calculated, and the results revealed that pioglitazone significantly inhibited the proliferation of hypoxic HepG2 cells treated with CoCl₂ (Fig. 2C). In the combination group of hypoxia and pioglitazone, the number of calcein-AM⁺ cells and




Figure 1. mRNA expression of PPAR γ is downregulated in CoCl₂-treated HepG2 cells. (A) Viability of HepG2 cells treated with a gradient dose of CoCl₂ for 24 h. (B) Protein expression of HIF-1 α in HepG2 cells treated with different doses of CoCl₂ for 24 h. β -actin was used as a loading control. (C) mRNA expression of PPAR γ , CD36, ACADM, ACADS and HMOX1 in HepG2 cells treated with different doses of CoCl₂ for 24 h (n=3-4). *P<0.05; **P<0.01; ***P<0.001. PPAR γ , peroxisome proliferator-activated receptor γ ; HIF-1 α , hypoxia-inducible factor-1 α ; ACAD, acetyl-co-enzyme A dehydrogenase; ACADM, ACAD medium-chain; ACADS, ACAD short-chain; HMOX1, heme oxygenase 1; ns, not significant; CoCl₂, cobalt chloride; Cq, quantification cycle.




Figure 2. A PPAR γ agonist inhibits the proliferation and promotes the death of hypoxic HepG2 cells. (A) HepG2 cells were treated with 20, 40 and 60 μ M P with or without 200 μ M Co for 48 h. P was added 24 h before Co. Cell viability was measured using a Cell Counting Kit-8 assay. (B) EdU staining of HepG2 cells treated with or without 200 μ M Co, 40 μ M P, and both Co + P for 48 h (magnification, x40). P was added 24 h before Co. (C) EdU⁺ cells were counted using ImageJ software version 1.54, and the positive ratio was compared between different groups. (D) Calcein-AM and PI double staining of HepG2 cells treated with or without 200 μ M Co, 40 μ M P and both Co + P for 72 h (scale bar, 500 μ m). P was added 24 h before Co. (E) Integrated density of calcein-AM and PI was calculated using ImageJ (n=3-4). ***P<0.001. PPAR γ , peroxisome proliferator-activated receptor γ ; P, pioglitazone; Co, cobalt chloride; EdU, 5-ethynyl-2'-deoxyuridine; Ctrl, control; calcein-AM, calcein-acetoxymethyl ester; PI, propidium iodide; ns, not significant; BF, bright field.

their fluorescence intensity were notably reduced compared with those in HepG2 cells treated with hypoxia only (Fig. 2D). The integrated density of calcein-AM and PI was calculated in repeated staining experiments. The cell confluence was low in the CO+P. PI staining cannot stain the detached dead cells. With optimization of treatment time, the CO + P showed the most dead cells, and the results suggested that pioglitazone in combination with CoCl₂ promoted the death of HepG2 cells treated with CoCl₂ alone (Fig. 2E). These results demonstrated that the activation of PPAR γ abolished the survival and proliferation abilities of hypoxic HepG2 cells.

PPAR γ agonist pioglitazone induces the death of hypoxic HepG2 cells via aggravation of oxidative stress. In the present study, pioglitazone was demonstrated to induce the death of hypoxic HepG2 cells treated with CoCl₂. Given the

PPAR γ agonist pioglitazone increased intracellular ROS production and then exacerbated the oxidative stress of hypoxic HepG2 cells, it was hypothesized that supplementation of an antioxidant could rescue hypoxic HepG2 cells. To assess this hypothesis, N-acetyl-L cysteine (NAC) was added to scavenge intracellular ROS in hypoxic HepG2 cells treated with pioglitazone. Following addition of NAC, the number of calcein-AM positive cells was markedly increased, whilst the number of PI positive cells was notably decreased compared to cells treated with Co + P (Fig. 3A). Furthermore, following NAC treatment, the integrated fluorescence intensity of PI of cells was significantly diminished compared with cells treated with Co + P (Fig. 3C). DHE staining was then performed to directly assess intracellular ROS production in HepG2 cells. The staining results demonstrated that increased red fluorescence in the Co + P group




Figure 3. A PPAR γ agonist increases the levels of intracellular reactive oxygen species to induce cell death. (A) Calcein-AM and PI double staining of HepG2 cells treated with or without 200 μ M Co, 40 μ M P and both Co + P for 48 h (magnification, x50). (B) DHE staining of HepG2 cells treated with or without 200 μ M Co, 40 μ M P and both Co + P for 48 h (magnification, x50). Integrated density of fluorescence was measured and calculated in samples of panels (C) A and (D) B. (n=3-4). **P<0.01; ***P<0.001. PPAR γ , peroxisome proliferator-activated receptor γ ; calcein-AM, calcein-acetoxymethyl ester; PI, propidium iodide; P, pioglitazone; Co, cobalt chloride; NAC, N-acetyl-L cysteine; DHE, dihydroethidium; Ctrl, control; ns, not significant; BF, bright field.

vs. Co group, indicated the PPAR γ agonist notably increased intracellular ROS in hypoxic HepG2 cells, which could be scavenged by NAC (Fig. 3B). The comparison of quantified integrated fluorescence density revealed that the PPAR γ agonist significantly increased integrated fluorescence in the Co + P group vs. Co group, whilst NAC significantly reduced integrated fluorescence in the Co + P + Nac group vs. Co + P group (Fig. 3D). The results demonstrated that the PPAR γ agonist pioglitazone induced cell death through the

production of excessive ROS and the induction of oxidative stress in hypoxic HepG2 cells.

PPAR γ agonist-mediated induction of ROS production may not occur through fatty acid oxidation but through impairment of B-cell lymphoma-2 (BCL2) expression. Fatty acid oxidation has been reported to be a source of ROS in cancer cells (17). Therefore, whether PPAR γ increased intracellular ROS production in hypoxic HepG2 cells was assessed by enhancing




Figure 4. A PPAR γ agonist downregulates anti-apoptotic gene expression but does not revert downregulated expression of fatty acid metabolism-related genes. HepG2 cells were treated with or without 200 μ M Co, 40 μ M P or both Co + P for 48 h. The mRNA expression of (A) PPAR γ , (B) CD36, (C) ACADM, (D) ACADS, (E) HMOX1 and (F) BCL2 was measured and analyzed (n=3-4). *P<0.05; **P<0.01; ***P<0.001. PPAR γ , peroxisome proliferator-activated receptor γ ; Co, cobalt chloride; P, pioglitazone; ACAD, acetyl-co-enzyme A dehydrogenase; ACADM, ACAD medium-chain; ACADS, ACAD short-chain; HMOX1, heme oxygenase 1; BCL2, B-cell lymphoma-2; Ctrl, control; ns, not significant; Cq, quantification cycle.

fatty acid oxidation. The mRNA expression of PPARG, CD36, ACADM and ACADS was measured. The expression of genes including PPARG, CD36, ACADM and ACADS significantly reduced about 50% in HepG2 cells treated with CoCl₂. However, the downregulated expression of these genes was not reverted by the PPAR γ agonist (Fig. 4A-D). In addition, the expression of the antioxidant gene HMOX1 was also measured to evaluate the antioxidant reactivity of hypoxic HepG2 cells treated with the PPAR γ agonist. However, the results revealed

that the PPAR γ agonist did not significantly stimulate the expression of HMOX1 in Co + P group compare to Co group (Fig. 4E).

BCL2 has been reported to be a key regulator of mitochondrial ROS production, and is able to confer anti-apoptotic effects to cancer cells with oxidative stress (18). Therefore, whether the aforementioned PPAR γ agonist regulated intracellular ROS production by modulating BCL2 expression was assessed. The mRNA expression of BCL2 was measured, and

the data indicated that hypoxic HepG2 cells had a significantly higher expression of BCL2 than control cells, and the PPAR γ agonist significantly decreased the expression of BCL2 in Co + P group cells compared to Co group cells (Fig. 4F). These results indicated that the PPAR γ agonist pioglitazone increased intracellular ROS production in hypoxic HepG2 cells, not via upregulation of the expression of fatty acid degrading genes, but via downregulation of the expression of the anti-apoptotic gene BCL2.

Discussion

The generation of a hypoxic environment and the activation of HIF-1 are common features of advanced cancer (26). The response to hypoxia is mainly attributed to HIFs. CoCl₂-induced chemical hypoxia is one of the most commonly used models of cell hypoxia *in vitro*, and the use of CoCl₂ *in vitro* has been reported to increase HIF-1 α /2 α in a dose- and time-dependent manner (23). CoCl₂ strongly stabilizes HIFs, thereby mimicking hypoxia and inducing the upregulation of a range of hypoxic adaptive responses, many of which have potential carcinogenic effects (27). As alteration of the cellular adaptation to hypoxia is also fundamental in cancer treatment (26), it was hypothesized that it would be meaningful to assess the mechanism and to identify an approach to abolish the resistance of cells to CoCl₂. Nevertheless, the hypoxic response can also be detrimental for tumorigenesis. Cobalt is cytotoxic and induces apoptosis and necrosis at high concentrations (27). Horev-Azaria *et al* (28) reported that CoCl₂ decreased the viability of NCI-H441 cells by 30-40% at 0.4 mM after 48 and 72 h, and of HepG2 cells by 70-100% at 0.4 mM after 48 and 72 h via MTT assay.

The experiments carried out in the present study revealed that the downregulation of the expression of PPAR γ and fatty acid oxidation genes was accompanied by increased expression of HIF-1 α in hypoxic HepG2 cells. The PPAR γ agonist pioglitazone specifically stimulated excessive ROS production in hypoxic HepG2 cells to induce cell death and inhibit cell proliferation. Notably, the effect of this PPAR γ agonist did not depend on fatty acid oxidation, but on the downregulation of BCL2 expression in hypoxic HepG2 cells. Thus, the results of the current study reveal a negative association between PPAR γ and BCL2 in regulating ROS production in hypoxic HepG2 cells.

Under hypoxic conditions, cancer cells dynamically modulate the intake and degradation of fatty acids to balance ROS production (11,14,29). Fatty acid degradation mainly occurs via β -oxidation and is catalyzed by ACADs. Fatty acid oxidation can be regulated in numerous tissues by activating the PPAR signaling pathway. PPARs are nuclear receptors that regulate lipid metabolism by promoting gene transcription (30,31). It has been reported that PPAR γ serves an important role in tumor cell proliferation and death, as well as in angiogenesis, invasion and metastasis (32). Once activated by a ligand, PPAR γ binds to DNA-specific PPAR response elements and modulates the transcription of its target genes, such as CD36, ACADM and ACADS, which may be important regulatory targets of fatty acid metabolism in tumor cells. CD36 stimulates tumor development and metastasis by allowing cells to absorb lipids from the extracellular environment,

and promotes fatty acid synthetase oxidation to produce ATP (33,34). CD36 is expressed at a high level in glioblastoma, and the reduction in CD36 leads to a loss of self-renewal and tumor initiation ability (35). ACADM reflects fatty acid metabolism and gemcitabine sensitivity in pancreatic cancer, potentially providing a reliable way to measure the efficacy of chemotherapy (36). ACADS is a potential methylation biomarker associated with the proliferation and metastasis of hepatocellular carcinoma (37). PPAR γ is the key regulator of ACADM transcription, which first catalyzes the reaction of the β -oxidation cycle for 4-10-carbon fatty acids (6). PPAR γ signaling is dysregulated in cancer cells under hypoxia or oxidative stress (7,38). In the present study, it was hypothesized that the association between PPAR γ and fatty acid oxidation may be essential for ROS regulation in hypoxic cancer cells. PPAR γ and its downstream genes, including CD36, ACADM and ACADS, were all downregulated in hypoxic HepG2 cells. HMOX1 is a key enzymes to antagonize harmful ROS of cytoplasm (39). The marked increase in HMOX1 gene expression in hypoxic HepG2 cells observed in the current study suggests a cellular adjustment of the redox status to regulate ROS production.

Pioglitazone is an established PPAR γ agonist, and is approved for the clinical treatment of diabetes. Despite its effect on glucose metabolism, pioglitazone has been reported to enhance fatty acid β -oxidation (25). Given that impaired PPAR γ activity confers cancer cells tolerance to hypoxia, activation of PPAR γ signaling should disrupt the intracellular metabolic balance of hypoxic HepG2 cells. The results of the present study revealed that pioglitazone decreased the viability of hypoxic HepG2 cells by inhibiting cell proliferation and inducing cell death. However, the effect of PPAR γ in hypoxia is controversial. It has been reported that the PPAR agonist pioglitazone protects against hypoxia-induced fetal growth inhibition (40). Similarly, another PPAR γ agonist has been reported to have a protective effect against hypoxia in cardiac myocytes (41). However, inhibition of PPAR γ and HIF-1 α potentiated the sensitivity of HepG2 cells to a tyrosine kinase inhibitor (42). It could be assumed that the conflicting PPAR γ effect in hypoxia may be due to differences in metabolic characteristics between cancer and benign cells. In the present study, it was demonstrated that the PPAR γ agonist pioglitazone abolished the hypoxia tolerance of HepG2 cells.

Oxidative stress is a status of excessive ROS compared with antioxidants. Cancer cells have aberrant redox homeostasis to tolerate high ROS levels, and adjust their antioxidant status to facilitate ROS-driven proliferation and to avoid ROS-induced senescence, apoptosis or ferroptosis (1). Glutathione (GSH) is a fast reactive endogenous antioxidant that scavenges intracellular ROS (43). Cysteine is the necessary source of GSH biosynthesis, and NAC is widely used as the source of GSH production (44). It can be hypothesized that the PPAR γ agonist pioglitazone induces the death of hypoxic HepG2 cells by increasing the production of excessive ROS, and the effect of pioglitazone may be neutralized by NAC-stimulated GSH production. The results of the present study demonstrated that the above PPAR γ agonist specifically increased intracellular ROS production in hypoxic HepG2 cells. By contrast, NAC protected hypoxic HepG2 cells against this PPAR γ agonist by scavenging intracellular ROS and

reducing PPAR γ agonist-mediated cell death. An association between the PPAR γ agonist pioglitazone and excessive ROS production was also demonstrated in hypoxic HepG2 cells. Similarly, this PPAR γ agonist has been reported to increase intracellular ROS production in lung cancer cells exposed to γ -radiation (45). Nevertheless, pioglitazone has been reported to inhibit the ROS production of cardiac fibroblasts treated under anoxia-reoxygenation conditions (46). Therefore, it can be hypothesized that the diverse effects of this PPAR γ agonist result from the dysregulated redox status of hypoxic cancer cells.

Mitochondria are the main producers of cellular ROS, including superoxide and/or H₂O₂, via aerobic metabolism (47). ROS production is tightly controlled, and HIFs alter the structure and activity of the electron transport chain to regulate ROS production under hypoxia (48). Stimulating fatty acid oxidation has been reported to disturb the redox balance of hypoxic cancer cells. Furthermore, increased expression of ACADM has been reported to decrease cell proliferation and invasion in solid tumors, whilst impaired ACADM activity promotes cancer progression (11,15). In the present study, it was demonstrated that the mRNA expression of fatty acid-degrading genes, including PPARG, ACADM and ACADS, was downregulated, and the PPAR γ agonist pioglitazone promoted excessive ROS production in hypoxic HepG2 cells. However, the results of the present study do not suggest that the PPAR γ agonist upregulated fatty acid oxidation in hypoxic HepG2 cells. PPAR γ expression has been reported to be dysregulated in esophageal cancer, and a PPAR γ agonist inhibited cancer cell proliferation *in vitro* and cancer progression *in vivo* via the Akt-P21^{CIP1} signaling pathway (49). Moreover, the PPAR γ agonist lobeglitazone has been reported to inhibit thyroid cancer cell metastasis by suppressing the MAPK signaling pathway (50). The evidence implies that the effect of PPAR γ agonists is not confined to modulating fatty acid metabolism. Moreover, the PPAR α activator fenofibrate has been reported to induce apoptosis in human hepatocellular carcinoma cells by increasing ROS production (51). In view of PPAR α -dependent apoptosis, PPAR α can be considered an E3 ubiquitin ligase able of inducing the ubiquitination and degradation of BCL2, thus leading to apoptosis (52). BCL2 is a mitochondrial membrane protein that regulates ROS production and serves as an anti-apoptotic effector in the endogenous apoptosis pathway (53). Therefore, downregulation of BCL2 by PPAR γ agonists stimulates excessive ROS production in hypoxic HepG2 cells. The results of the present study indicate a negative association between BCL2 expression and PPAR γ activity in hypoxic HepG2 cells. This mechanism appears to be essential for regulating ROS production in hypoxic HepG2 cells.

In conclusion, PPAR γ downregulation appears to be required for redox homeostasis in hypoxic HepG2 cells. In the present study, the PPAR γ agonist pioglitazone stimulated excessive ROS production in hypoxic HepG2 cells to induce cell death and inhibit cell proliferation via downregulation of BCL2 expression. Notably, differences between lack of oxygen-induced hypoxia and CoCl₂-induced chemical hypoxia may lead to proapoptotic responses, and the genes induced by the two modes of hypoxia may not overlap. However, other anti-apoptotic genes potentially modulated by pioglitazone

were not included in the present study, and therefore further experiments are needed to fully assess the role of pioglitazone in CoCl₂-induced hypoxic HepG2 cells.

Acknowledgements

The authors would like to thank the Translational Medical Center at The First Affiliated Hospital of Zhengzhou University for providing access to analytical instruments.

Funding

The present study was supported by the Health Commission of Henan Province (grant nos. LHGJ20190155 and LHGJ20210287).

Availability of data and materials

The data generated in the present study are included in the figures and/or tables of this article.

Authors' contributions

GH conceived the study, performed the experiments, and analyzed and interpreted the data. MZ conceived the study and analyzed and interpreted the data. MW and WX performed the experiments. XD and XH interpreted the data and revised the manuscript. XD and MZ confirm the authenticity of all the raw data. JR and MZ funded the study, and interpreted the data. All authors have read and approved the final version of the manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Hayes JD, Dinkova-Kostova AT and Tew KD: Oxidative Stress in Cancer. *Cancer Cell* 38: 167-197, 2020.
2. Kaminski MM, Sauer SW, Kaminski M, Opp S, Ruppert T, Grigaravicius P, Grudnik P, Grone HJ, Krammer PH and Gülow K: T cell activation is driven by an ADP-dependent glucokinase linking enhanced glycolysis with mitochondrial reactive oxygen species generation. *Cell Rep* 2: 1300-1315, 2012.
3. Ishikawa K, Takenaga K, Akimoto M, Koshikawa N, Yamaguchi A, Imanishi H, Nakada K, Honma Y and Hayashi J: ROS-Generating Mitochondrial DNA mutations can regulate tumor cell metastasis. *Science* 320: 661-664, 2008.
4. Zong X, Cao X, Wang H, Xiao X, Wang Y and Lu Z: Cathelicidin-WA Facilitated intestinal fatty acid absorption through enhancing PPAR- γ dependent barrier function. *Front Immunol* 10: 1674, 2019.
5. Angela M, Endo Y, Asou HK, Yamamoto T, Tumes DJ, Tokuyama H, Yokote K and Nakayama T: Fatty acid metabolic reprogramming via mTOR-mediated inductions of PPAR γ directs early activation of T cells. *Nat Commun* 7: 13683, 2016.

6. Seo YS, Kim JH, Jo NY, Choi KM, Baik SH, Park JJ, Kim JS, Byun KS, Bak YT, Lee CH, *et al*: PPAR agonists treatment is effective in a nonalcoholic fatty liver disease animal model by modulating fatty-acid metabolic enzymes. *J Gastroenterol Hepatol* 23: 102-109, 2008.
7. Small DM, Morais C, Coombes JS, Bennett NC, Johnson DW and Gobe GC: Oxidative stress-induced alterations in PPAR- γ and associated mitochondrial destabilization contribute to kidney cell apoptosis. *Am J Physiol Renal Physiol* 307: F814-F822, 2014.
8. Vallee A and Lecarpentier Y: Crosstalk between peroxisome proliferator-activated receptor gamma and the canonical WNT/ β -Catenin pathway in chronic inflammation and oxidative stress during carcinogenesis. *Front Immunol* 9: 745, 2018.
9. Koundouros N and Pouligiannis G: Reprogramming of fatty acid metabolism in cancer. *Br J Cancer* 122: 4-22, 2020.
10. Carracedo A, Cantley LC and Pandolfi PP: Cancer metabolism: fatty acid oxidation in the limelight. *Nat Rev Cancer* 13: 227-232, 2013.
11. Ma APY, Yeung CLS, Tey SK, Mao X, Wong SWK, Ng TH, Ko FCF, Kwong EML, Tang AHN, Ng IO, *et al*: Suppression of ACADM-Mediated fatty acid oxidation promotes hepatocellular carcinoma via aberrant CAV1/SREBP1 Signaling. *Cancer Res* 81: 3679-3692, 2021.
12. Wang C, Shao L, Pan C, Ye J, Ding Z, Wu J, Du Q, Ren Y and Zhu C: Elevated level of mitochondrial reactive oxygen species via fatty acid beta-oxidation in cancer stem cells promotes cancer metastasis by inducing epithelial-mesenchymal transition. *Stem Cell Res Ther* 10: 175, 2019.
13. Amoedo ND, Sarlak S, Obre E, Esteves P, Begueret H, Kieffer Y, Rousseau B, Dupis A, Izotte J, Bellance N, *et al*: Targeting the mitochondrial trifunctional protein restrains tumor growth in oxidative lung carcinomas. *J Clin Invest* 131: e133081, 2021.
14. Li XX, Wang ZJ, Zheng Y, Guan YF, Yang PB, Chen X, Peng C, He JP, Ai YL, Wu SF, *et al*: Nuclear Receptor Nur77 facilitates melanoma cell survival under metabolic stress by protecting fatty acid oxidation. *Mol Cell* 69: 480-492 e7, 2018.
15. Hsieh CH, Cheung CHY, Liu YL, Hou CL, Hsu CL, Huang CT, Yang TS, Chen SF, Chen CN, Hsu WM, *et al*: Quantitative Proteomics of Th-MYCN transgenic mice reveals aurora kinase inhibitor altered metabolic pathways and enhanced ACADM To suppress neuroblastoma progression. *J Proteome Res* 18: 3850-3866, 2019.
16. Pacary E, Tixier E, Coulet F, Roussel S, Petit E and Bernaudin M: Crosstalk between HIF-1 and ROCK pathways in neuronal differentiation of mesenchymal stem cells, neurospheres and in PC12 neurite outgrowth. *Mol Cell Neurosci* 35: 409-423, 2007.
17. Befani C, Mylonis I, Gkoutinakou IM, Georgoulis P, Hu CJ, Simos G and Liakos P: Cobalt stimulates HIF-1-dependent but inhibits HIF-2-dependent gene expression in liver cancer cells. *Int J Biochem Cell Biol* 45: 2359-2368, 2013.
18. Zhigalova N, Artemov A, Mazur A and Prokhortchouk E: Transcriptome sequencing revealed differences in the response of renal cancer cells to hypoxia and CoCl₂ treatment. *F1000Res* 4: 1518, 2015.
19. Zhao Y, Lützen U, Gohlke P, Jiang P, Herdegen T and Culman J: Neuroprotective and antioxidative effects of pioglitazone in brain tissue adjacent to the ischemic core are mediated by PI3K/Akt and Nrf2/ARE pathways. *J Mol Med (Berl)* 99: 1073-1083, 2021.
20. Jiao XX, Lin SY, Lian SX, Qiu YR, Li ZH, Chen ZH, Lu WQ, Zhang Y, Deng L, Jiang Y and Hu GH: Inhibition of the breast cancer by PPAR γ agonist pioglitazone through JAK2/STAT3 pathway. *Neoplasma* 67: 834-842, 2020.
21. Tsubaki M, Takeda T, Tomonari Y, Kawashima K, Itoh T, Imano M, Satou T and Nishida S: Pioglitazone inhibits cancer cell growth through STAT3 inhibition and enhanced AIF expression via a PPAR γ -independent pathway. *J Cell Physiol* 233: 3638-3647, 2017.
22. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408, 2001.
23. Munoz-Sanchez J and Chanez-Cardenas ME: The use of cobalt chloride as a chemical hypoxia model. *J Appl Toxicol* 39: 556-570, 2019.
24. Ran Y, Hu C, Wan J, Kang Q, Zhou R, Liu P, Ma D, Wang J and Tang L: Integrated investigation and experimental validation of PPARG as an oncogenic driver: Implications for prognostic assessment and therapeutic targeting in hepatocellular carcinoma. *Front Pharmacol* 14: 1298341, 2023.
25. Hsiao PJ, Chiou HC, Jiang HJ, Lee MY, Hsieh TJ and Kuo KK: Pioglitazone enhances cytosolic lipolysis, β -oxidation and autophagy to ameliorate hepatic steatosis. *Sci Rep* 7: 9030, 2017.
26. Petrova V, Annicchiarico-Petruzzelli M, Melino G and Amelio I: The hypoxic tumour microenvironment. *Oncogenesis* 7: 10, 2018.
27. Simonsen LO, Harbak H and Bennekou P: Cobalt metabolism and toxicology-A brief update. *Sci Total Environ* 432: 210-215, 2012.
28. Horev-Azaria L, Kirkpatrick CJ, Korenstein R, Marche PN, Maimon O, Ponti J, Romano R, Rossi F, Golla-Schindler U, Sommer D, *et al*: Predictive toxicology of cobalt nanoparticles and ions: Comparative in vitro study of different cellular models using methods of knowledge discovery from data. *Toxicol Sci* 122: 489-501, 2011.
29. Choi HJ, Jhe YL, Kim J, Lim JY, Lee JE, Shin MK and Cheong JH: FoxM1-dependent and fatty acid oxidation-mediated ROS modulation is a cell-intrinsic drug resistance mechanism in cancer stem-like cells. *Redox Biol* 36: 101589, 2020.
30. Sprecher DL, Massien C, Pearce G, Billin AN, Perlstein I, Willson TM, Hassall DG, Ancellin N, Patterson SD, Lobe DC and Johnson TG: Triglyceride:High-Density Lipoprotein cholesterol effects in healthy subjects administered a peroxisome proliferator activated receptor δ agonist. *Arterioscler Thromb Vasc Biol* 27: 359-365, 2007.
31. Froment P, Gizard F, Defever D, Staels B, Dupont J and Monget P: Peroxisome proliferator-activated receptors in reproductive tissues: From gametogenesis to parturition. *J Endocrinol* 189: 199-209, 2006.
32. Wagner N and Wagner KD: Peroxisome proliferator-activated receptors and the hallmarks of cancer. *Cells* 11: 2432, 2022.
33. Kuemmerle NB, Rysman E, Lombardo PS, Flanagan AJ, Lipe BC, Wells WA, Pettus JR, Froehlich HM, Memoli VA, Morganeli PM, *et al*: Lipoprotein lipase links dietary fat to solid tumor cell proliferation. *Mol Cancer Ther* 10: 427-436, 2011.
34. Pepino MY, Kuda O, Samovski D and Abumrad NA: Structure-Function of CD36 and importance of fatty acid signal transduction in fat metabolism. *Annu Rev Nutr* 34: 281-303, 2014.
35. Hale JS, Otvos B, Sinyuk M, Alvarado AG, Hitomi M, Stoltz K, Wu Q, Flavahan W, Levison B, Johansen ML, *et al*: Cancer stem cell-specific scavenger receptor CD36 drives glioblastoma progression. *Stem Cells* 32: 1746-1758, 2014.
36. Yang Y, Gu H, Zhang K, Guo Z, Wang X, Wei Q, Weng L, Han X, Lv Y, Cao M, *et al*: Exosomal ACADM sensitizes gemcitabine-resistance through modulating fatty acid metabolism and ferroptosis in pancreatic cancer. *BMC Cancer* 23: 789, 2023.
37. Chen D, Feng X, Lv Z, Xu X, Lu Y, Wu W, Wu H, Liu H, Cao L, Ye S, *et al*: ACADS acts as a potential methylation biomarker associated with the proliferation and metastasis of hepatocellular carcinomas. *Aging (Albany NY)* 11: 8825-8844, 2019.
38. Xu R, Luo X, Ye X, Li H, Liu H, Du Q and Zhai Q: SIRT1/PGC-1 α /PPAR- γ correlate with hypoxia-induced chemoresistance in non-small cell lung cancer. *Front Oncol* 11: 682762, 2021.
39. Abe K, Ikeda S, Nara M, Kitadate A, Tagawa H and Takahashi N: Hypoxia-induced oxidative stress promotes therapy resistance via upregulation of heme oxygenase-1 in multiple myeloma. *Cancer Med* 12: 9709-9722, 2023.
40. Lane SL, Dodson RB, Doyle AS, Park H, Rathi H, Matarrazo CJ, Moore LG, Lorca RA, Wolfson GH and Julian CG: Pharmacological activation of peroxisome proliferator-activated receptor γ (PPAR- γ) protects against hypoxia-associated fetal growth restriction. *FASEB J* 33: 8999-9007, 2019.
41. Kilter H, Werner M, Roggia C, Reil JC, Schafers HJ, Kintscher U and Bohm M: The PPAR-gamma agonist rosiglitazone facilitates Akt rephosphorylation and inhibits apoptosis in cardiomyocytes during hypoxia/reoxygenation. *Diabetes Obes Metab* 11: 1060-1067, 2009.
42. Feng J, Dai W, Mao Y, Wu L, Li J, Chen K, Yu Q, Kong R, Li S, Zhang J, *et al*: Simvastatin re-sensitizes hepatocellular carcinoma cells to sorafenib by inhibiting HIF-1 α /PPAR- γ /PKM2-mediated glycolysis. *J Exp Clin Cancer Res* 39: 24, 2020.
43. Liu T, Sun L, Zhang Y, Wang Y and Zheng J: Imbalanced GSH/ROS and sequential cell death. *J Biochem Mol Toxicol* 36: e22942, 2021.
44. Aldini G, Altomare A, Baron G, Vistoli G, Carini M, Borsani L and Sergio F: N-Acetylcysteine as an antioxidant and disulphide breaking agent: The reasons why. *Free Radic Res* 52: 751-762, 2018.

45. Han EJ, Im CN, Park SH, Moon EY and Hong SH: Combined treatment with peroxisome proliferator-activated receptor (PPAR) γ ligands and gamma radiation induces apoptosis by PPAR γ -independent up-regulation of reactive oxygen species-induced deoxyribonucleic acid damage signals in non-small cell lung cancer cells. *Int J Radiat Oncol Biol Phys* 85: e239-248, 2013.
46. Chen K, Li D, Zhang X, Hermonat PL and Mehta JL: Anoxia-reoxygenation stimulates collagen type-I and MMP-1 expression in cardiac fibroblasts: Modulation by the PPAR-gamma ligand pioglitazone. *J Cardiovasc Pharmacol* 44: 682, 2004.
47. Cadenas S: Mitochondrial uncoupling, ROS generation and cardioprotection. *Biochim Biophys Acta Bioenerg* 1859: 940-950, 2018.
48. Fuhrmann DC and Brune B: Mitochondrial composition and function under the control of hypoxia. *Redox Biol* 12: 208-215, 2017.
49. Sawayama H, Ishimoto T, Watanabe M, Yoshida N, Sugihara H, Kurashige J, Hirashima K, Iwatsuki M, Baba Y, Oki E, *et al*: Small molecule agonists of PPAR- γ exert therapeutic effects in esophageal cancer. *Cancer Res* 74: 575-585, 2014.
50. Jin JQ, Han JS, Ha J, Baek HS and Lim DJ: Lobeglitazone, A peroxisome proliferator-activated receptor-gamma agonist, inhibits papillary thyroid cancer cell migration and invasion by suppressing p38 MAPK signaling pathway. *Endocrinol Metab (Seoul)* 36: 1095-1110, 2021.
51. Jiao HL and Zhao BL: cytotoxic effect of peroxisome proliferator fenofibrate on human HepG2 hepatoma cell line and relevant mechanisms. *Toxicol Appl Pharmacol* 185: 172-179, 2002.
52. Gao J, Liu Q, Xu Y, Gong X, Zhang R, Zhou C, Su Z, Jin J, Shi H, Shi J and Hou Y: PPAR α induces cell apoptosis by destructing Bcl2. *Oncotarget* 6: 44635-44642, 2015.
53. Chong SJ, Low IC and Pervaiz S: Mitochondrial ROS and involvement of Bcl-2 as a mitochondrial ROS regulator. *Mitochondrion* 19 Pt A: 39-48, 2014.



Copyright © 2024 Huang et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.