

Effects of 6-(methylsulfinyl)hexyl isothiocyanate on cyclooxygenase-2 expression induced by lipopolysaccharide, interferon- γ and 12-*O*-tetradecanoylphorbol-13-acetate

TAKUHIRO UTO, MAKOTO FUJII and DE-XING HOU

Department of Biochemical Science and Technology, Faculty of Agriculture,
Kagoshima University, Korimoto 1-21-24, Kagoshima City 890-0065, Japan

Received August 7, 2006; Accepted September 15, 2006

Abstract. 6-(methylsulfinyl)hexyl isothiocyanate (6-MITC) is a bioactive compound extracted from a typical Japanese spice, wasabi (*Wasabia japonica* (Miq.) Matsumura). In the present study, we found that 6-MITC suppressed the expression of cyclooxygenase-2 (COX-2) induced by lipopolysaccharide (LPS), interferon- γ (IFN- γ), but did not suppress that induced by 12-*O*-tetradecanoylphorbol-13-acetate (TPA), in murine macrophage RAW264. Molecular mechanisms were investigated by targeting the transcriptional factors including activator protein-1 (AP-1), CCAAT/enhancer-binding protein δ (C/EBP δ), CRE-binding protein (CREB) and nuclear factor κ B (NF- κ B), which bind to the core element of *COX-2* promoter. LPS induced activation of all of these factors and 6-MITC suppressed LPS-induced activation of AP-1, C/EBP δ , CREB, but not NF- κ B. IFN- γ did not induce any activation of these factors, but 6-MITC suppressed IFN- γ -induced COX-2 expression, suggesting that the upstream region of the core element is linked for this suppression. Finally, TPA stimulated the activation of CREB and AP-1, but 6-MITC did not block TPA-induced COX-2 expression. These results suggest that LPS, IFN- γ and TPA regulate COX-2 expression through different mechanisms, and 6-MITC acts as a potent inhibitor of COX-2 expression induced by LPS or IFN- γ .

Introduction

Numerous studies have demonstrated that high intake of cruciferous vegetables such as broccoli, watercress, cabbage, and cauliflower, can prevent cancer (1). The bioactive compounds involved in these vegetables are isothiocyanates (ITCs), which have been reported to have cancer-preventive effect in the carcinogenesis of lung, breast and colon (2). 6-(methylsulfinyl) hexyl isothiocyanate (6-MITC) is an ITC, present as a major allyl isothiocyanate in a typical Japanese spice, wasabi (*Wasabia japonica* (Miq.) Matsumura). Oral administration of 6-MITC showed the inhibitory effects on mouse skin tumor induced by 9,10-dimethyl-1,2-benzanthracene as an initiator and 12-*O*-tetradecanoylphorbol-13-acetate (TPA) as a promoter (3), and on mouse lung tumor induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (4). 6-MITC also has been demonstrated to specifically suppress cell proliferation of breast cancer and melanoma cell lines (5). These results suggest that 6-MITC is a potent chemopreventive compound.

Cyclooxygenase (COX-2) is an enzyme catalyzing the conversion of arachidonic acid to prostaglandins. There are two isoforms of COX, designated COX-1 and COX-2. COX-1 is constitutively expressed in most tissues to maintain normal physiological functions (6). In contrast, COX-2 is undetectable or present at very low levels under basal conditions, but is rapidly induced by inflammatory stimuli, cytokines, growth factors, tumor promoters, and carcinogens (7,8). Several lines of evidence suggest that COX-2 plays an important role in the process of carcinogenesis because COX-2 is overexpressed in a variety of transformed cells and tumors (9,10). Enhanced induction of COX-2 promotes the growth of malignant cells by increasing cell proliferation (11), promoting angiogenesis (12), and inhibiting immune surveillance (13) and apoptosis (14). Various chemopreventive strategies have focused on the identification of COX-2 inhibitors because the targeted inhibition of COX-2 is an effective approach for suppressing inflammation and carcinogenesis.

Accumulated data have indicated that there are three *cis*-acting elements including nuclear factor κ B (NF- κ B), CCAAT/enhancer-binding protein (C/EBP) and cAMP-responsive element (CRE) present in the promoter (15). The tran-

Correspondence to: Dr De-Xing Hou, Department of Biochemical Science and Technology, Faculty of Agriculture, Kagoshima University, Korimoto 1-21-24, Kagoshima 890-0065, Japan
E-mail: hou@ms.kagoshima-u.ac.jp

Abbreviations: 6-MITC, 6-(methylsulfinyl)hexyl isothiocyanate; AP-1, activator protein-1; C/EBP, CCAAT/enhancer-binding protein; COX-2, cyclooxygenase-2; CREB, CRE-binding protein; IFN- γ , interferon- γ ; LPS, lipopolysaccharide; NF- κ B, nuclear factor κ B; TPA, 12-*O*-tetradecanoylphorbol-13-acetate

Key words: 6-(methylsulfinyl)hexyl isothiocyanate, macrophages, cyclooxygenase-2, lipopolysaccharide, 12-*O*-tetradecanoylphorbol-13-acetate, interferon- γ

scriptional factors such as NF- κ B, C/EBPs, CRE binding protein (CREB) and activator protein-1 (AP-1) bind to these *cis*-acting elements and transactivate *COX-2* gene expression. Various different inflammatory stimuli have been reported to induce *COX-2* gene transcription (7,8). However, the precise mechanisms underlying *COX-2* gene expression triggered by these stimuli have not been fully elucidated.

In the present study, we investigated the effect of 6-MITC on *COX-2* expression activated by LPS, INF- γ , and TPA in murine macrophage RAW264 cells. Our study revealed that LPS, INF- γ and TPA induced *COX-2* expression through different pathways, and 6-MITC acted as a potent inhibitor for intervening LPS- and INF- γ -activated *COX-2* expression.

Materials and methods

Materials and cell culture. 6-MITC purified by reverse-phase HPLC to >99% were obtained from Hakucho Pharmaceutical Co., Ltd (Japan). 6-MITC was dissolved in dimethyl sulfoxide (DMSO, final concentration was 0.2%). Antibodies against phospho-CREB, phospho-c-Jun, CREB, c-Jun and I κ B- α were purchased from Cell Signaling Technology (Beverly, MA). Antibodies against *COX-2*, *COX-1*, C/EBP β , C/EBP δ , p65 and α -tubulin were from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Fetal bovine serum (FBS) was from Equitech-Bio (Kerrville, TX). Lipofectamine was from Life Technologies, Inc. (Grand Island, NY). LPS (*Escherichia coli* Serotype 055:B5) was from Sigma (St. Louis, MO). Murine macrophage-like RAW264 cells were obtained from RIKEN BioResource Center Cell Bank (cell No. RCB0535), and cultured at 37°C in a 5% CO₂ atmosphere in Dulbecco's modified Eagle's medium (DMEM) containing 10% FBS.

Western blot analysis. Western blot assay was performed as described previously (16). In brief, RAW264 (1x10⁶ cells) were pre-cultured in a 6-cm dish for 24 h, and then starved by being cultured in serum-free medium for another 2.5 h to eliminate the influence of FBS. The cells were treated with or without 6-MITC for 30 min before exposure to LPS, INF- γ , TPA, interleukin- β (IL- β), tumor necrosis factor- α (TNF- α), or H₂O₂ for different times. The harvested cells were lysed and the supernatants were boiled for 5 min. Protein concentration was determined using dye-binding protein assay kit (Bio-Rad Hercules, CA) according to the manufacturer's manual. Equal amounts of lysate protein (~40 μ g) were run on 10% SDS-PAGE and electrophoretically transferred to PVDF membrane (Amersham Pharmacia Biotech, Little Chalfont, UK). After Western blotting, the membrane was incubated with specific primary antibody overnight at 4°C, and further incubated for 1 h with HRP-conjugated secondary antibody. Bound antibodies were detected by ECL system with a Lumi Vision PRO machine (Taitec Co., Japan). The relative amount of proteins associated with each specific antibody was quantified using the Lumi Vision Imager software (Taitec Co.).

Plasmids. The *COX-2* promoter-luciferase deletion constructs (-1432/+59, -327/+59) were generous gifts of Dr Hiroyasu Inoue (Nara Women's University, Nara, Japan) and have been described previously (17-19).




Figure 1. Effects of inflammatory stimuli on *COX-2* expression in RAW264 cells. After RAW264 cells (1x10⁶) were starved in serum-free medium for 3 h, the cells were treated with the indicated concentrations of LPS, INF- γ , TPA, IL-1 β , TNF- α , or H₂O₂ for 12 h. *COX-2* was detected with its antibody by Western blot analysis.

Transient transfection assay. Transient transfection was performed according to the modified method as described previously (20). RAW264 cells (1x10⁵) were plated into each well of 12-well plates, and cultured for 24 h. The cells were then co-transfected with 0.5 μ g *COX-2* promoter-luciferase plasmids and 0.12 μ g CMV- β -galactosidase plasmids using Lipofectamine 2000. After 5 h of incubation, the medium was replaced and cultured for another 20.5 h. The cells were treated with or without 6-MITC for 30 min before exposure to 40 ng/ml LPS, INF- γ , or TPA for 6 h. The activities of luciferase and β -galactosidase in cell lysate were measured with a luminometer (Berthold) according to the supplier's recommendations. Luciferase activity is normalized to transfection efficiency monitored by β -galactosidase expression, and *COX-2* transcription activity was expressed as fold induction relative to the control cells without LPS, INF- γ , or TPA treatment.

Nuclear extracts. Nuclear extracts were prepared using the nuclear extract kit from Active Motif (Carlsbad, CA). Briefly, RAW264 cells (3x10⁶) were pre-cultured in a 10-cm dish for 24 h, and then starved by being cultured in serum-free medium for another 2.5 h to eliminate the influence of FBS. The cells were treated with or without 6-MITC for 30 min before exposure to 40 ng/ml LPS, INF- γ , or TPA. The cell pellet was suspended in 1 x hypotonic buffer supplemented with complete protease inhibitor cocktail, incubated for 15 min at 4°C, vortexed with detergents and centrifuged briefly. The nuclear pellet was washed twice with the cytoplasmic buffer




Figure 2. Effect of 6-MITC on COX-2 expression induced by LPS, IFN- γ or TPA. (A) After RAW264 cells (1×10^6) were starved in serum-free medium for 2.5 h, the cells were treated with 15 μ M 6-MITC for 30 min, and then exposed to 40 ng/ml LPS, IFN- γ , TPA for 12 h. (B) The cells were treated with the indicated concentrations of 6-MITC for 30 min, and then exposed to 40 ng/ml LPS or IFN- γ for 12 h. COX-2 and COX-1 were detected by Western blot analysis with their antibodies, respectively.

followed by resuspension in the lysis buffer supplemented with 1 mM DTT and protease inhibitors. The suspension was incubated on a rocking platform at 4°C for 30 min. The suspension was then vortexed briefly and centrifuged for 10 min at 14,000 x g at 4°C. The supernatant (nuclear fraction) was collected, and nuclear extracts were stored at -80°C until Western blot analysis.

Statistical analysis. Difference between the treated and the control was analyzed by Student's t-test. A probability of $P < 0.05$ was considered significant.

Results

Effects of inflammatory stimuli on COX-2 expression. To determine the effects of inflammatory stimuli on COX-2 protein expression in mouse macrophages, RAW264 cells were incubated with the indicated concentrations of LPS, INF- γ , TPA, interleukin-1 β (IL-1 β), tumor necrosis factor- α (TNF- α), or H₂O₂ for 12 h. As shown in Fig. 1, the expression of COX-2 protein was increased after incubation with LPS, INF- γ or TPA. On the other hand, IL-1 β and TNF- α did not induce COX-2 expression although the used concentrations could induce COX-2 expression in Caco-2 colon carcinoma cells (21), human airway smooth muscle (HASM) cells (22,23), L929 fibroblast cells (24), and lung adenocarcinoma A549




Figure 3. Different effects of 6-MITC on COX-2 promoter activity induced by LPS, IFN- γ or TPA. (A) Schematic diagram of COX-2 promoter-luciferase reporter constructs used. (B) RAW264 cells were transfected with 0.5 μ g of COX-2 promoter constructs (-1432/+59 or -327/+59) and 0.12 μ g CMV- β -galactosidase plasmid. After 5 h of incubation, the medium was replaced with complete medium and cultured for another 20.5 h. The cells were then treated with or without 6-MITC for 30 min before they were exposed to 40 ng/ml LPS, IFN- γ and TPA for 6 h. The luciferase activity values were normalized to transfection efficiency monitored by β -galactosidase expression, and COX-2 promoter activity was expressed as fold induction to control. Each value represents the mean \pm SD of three to four separate experiments. * $P < 0.05$ versus control.

cells (24). Thus, LPS, INF- γ and TPA are potent inducers of COX-2 expression in RAW264 cells.

Effect of 6-MITC on COX-2 expression induced by LPS, IFN- γ or TPA. To investigate the inhibitory effect of 6-MITC on COX-2 expression induced by LPS, IFN- γ , or TPA, RAW264 cells were treated with 15 μ M 6-MITC for 30 min before exposure to 40 ng/ml LPS, IFN- γ , or TPA for 12 h. As shown in Fig. 2A, 6-MITC suppressed LPS- and IFN- γ -induced




Figure 4. Effects of 6-MITC on the activation of the transcriptional factors induced by LPS, IFN- γ or TPA. (A) Cell culture and Western blot analysis were done as described in Fig. 2. RAW264 cells were treated with 15 μ M 6-MITC for 30 min, and then exposed to 40 ng/ml LPS, IFN- γ or TPA for 30 min. Phosphorylated and total c-Jun or CREB were detected by Western blot analysis with their antibodies, respectively. (B) The cells were treated with or without 15 μ M 6-MITC for 30 min, and then exposed to 40 ng/ml LPS, IFN- γ or TPA for 4 h. The nuclear proteins and whole cell lysates were extracted, and C/EBP δ and β were detected by Western blot analysis. (C) The cells were treated with 15 μ M 6-MITC for 30 min, and then exposed to 40 ng/ml LPS, IFN- γ and TPA for 30 min. The nuclear proteins and whole cell lysates were extracted, and I κ B- α , p65 and α -tubulin were detected by Western blot analysis.


COX-2 expression. However, 6-MITC did not suppress TPA-induced COX-2. The amount of COX-1, which was constitutive protein, remained unchanged. To further examine the inhibitory effects of 6-MITC on COX-2 expression induced by LPS or IFN- γ , the cells were incubated with various concentrations of 6-MITC before the treatment with LPS or IFN- γ . Fig. 2B shows that 6-MITC inhibited COX-2 expression induced by LPS or IFN- γ in a dose-dependent manner. Our previous results have shown that the concentration range of 6-MITC used in these experiments had no significant effect on the cellular viability as measured by MTT assay (data not shown) (25). Thus, the inhibitory actions of 6-MITC on COX-2 expression were not caused by its toxicity.

Different effects of 6-MITC on LPS-, IFN- γ - or TPA-induced COX-2 promoter activity. The core element (-327/-52) in COX-2 promoter contains three *cis*-acting elements including NF- κ B, C/EBP and CRE sites, which are involved in the regulation of COX-2 gene expression (15). To investigate whether the inhibition of COX-2 expression by 6-MITC is mediated through the core element, transient transfections were performed using full (-1432/+59) and core (-327/+59) COX-2 promoter-luciferase constructs (Fig. 3A). As shown in Fig. 3B, 1) LPS induced COX-2 promoter activity in both -1432/+59 and -327/+59 constructs, and 6-MITC reduced them to control level. 2) IFN- γ induced COX-2 promoter activity of -1432/+59 construct, which was suppressed by 6-MITC. IFN- γ did not induce the promoter activity of -327/+59 construct. 3) TPA also induced the COX-2 promoter activity in both -1432/+59 and -327/+59, but 6-MITC did not block the transcriptional activity. These results suggest that the core element is required for LPS- and TPA-induced COX-2 gene expression, and 6-MITC might block LPS-induced COX-2 expression by suppressing the core element. On the other hand, the upstream region of the core element may respond to the IFN- γ -induced COX-2 expression, and 6-MITC may block COX-2 expression through the upstream region.

Effects of 6-MITC on the transcriptional factors regulating COX-2 expression. Transcription factors including CREB, AP-1, C/EBP and NF- κ B can regulate COX-2 transcription by binding the core elements in COX-2 promoter (15,18,26). To elucidate the effects of 6-MITC on these transcriptional factors, we pretreated RAW264 cells with 6-MITC for 30 min before exposure to LPS, IFN- γ , or TPA, and then examined the activation of these transcription factors by Western blot analysis.

As shown in Fig. 4A, LPS and TPA strongly induced the phosphorylation of CREB and c-Jun, which is a key component of AP-1 in c-Jun/c-Fos heterodimer form. 6-MITC blocked the phosphorylation of CREB and c-Jun induced by LPS, but not TPA. In contrast, IFN- γ did not induce phosphorylation of CREB and c-Jun.

LPS also markedly induced expression and nuclear translocation of C/EBP δ , which were blocked by 6-MITC (Fig. 4B). However, IFN- γ and TPA did not induce C/EBP δ at either whole or nuclear fraction. The C/EBP β levels at either whole or nuclear were not affected by three inducers and 6-MITC. These results suggest that 6-MITC suppressed LPS-activated

 SPANDIDOS PUBLICATIONS and subsequent nuclear translocation of C/EBP δ and COX-2 expression. In contrast, C/EBP δ was not involved in IFN- γ - and TPA-induced COX-2 expression.

NF- κ B is also known as a critical factor for COX-2 expression induced by LPS or proinflammatory cytokines (17,27). NF- κ B is inactivated in the cytosol by binding to I κ B, and becomes active through translocation to the nucleus preceded by LPS-induced proteolytic degradation of I κ B (28). To determine whether 6-MITC inhibit degradation of I κ B, the level of I κ B- α protein was assessed in the treated RAW264 cells. As shown in Fig. 4C, LPS reduced I κ B- α protein, and stimulated nuclear translocation of p65, a major component of NF- κ B, into nucleus. However, 6-MITC did not prevent the degradation of I κ B- α and nuclear translocation of p65. On the other hand, IFN- γ and TPA had no influence on I κ B- α degradation and nuclear translocation of p65. These data suggest that the inhibitory effect of 6-MITC on LPS-induced COX-2 expression is not mediated through the suppression of NF- κ B activation, and IFN- γ - and TPA-induced COX-2 expression in RAW264 cells is not mediated by NF- κ B activation.

Discussion

Multiple lines of evidence suggest that COX-2 has a critical role in inflammation and carcinogenesis. Thus, the targeted inhibition of COX-2 by chemopreventive agents is a promising approach to prevent inflammation and cancer. In the present study, 6-MITC, a major compound in wasabi, strongly blocked the expression of COX-2 induced by LPS or IFN- γ , but not TPA, suggesting that LPS, IFN- γ and TPA might induce COX-2 expression through distinct pathways in mouse macrophage cells.

The inductive effects of LPS on COX-2 induction are mediated by transcriptional factors including AP-1, C/EBP and NF- κ B, which bind to the core elements of COX-2 (15). Moreover, there is redundancy in the three *cis*-acting elements, and two of these *cis*-acting elements are at least recruited to achieve maximal transcription of COX-2 gene in mouse macrophage RAW264.7 cells (29). In the present study, our data indicated that the 6-MITC blocked LPS-induced COX-2 expression by suppressing the activation of CREB, AP-1 and C/EBP δ , which lead to the binding to the core element of COX-2 gene.

IFN- γ is a potent inducer of inflammatory responses (30). The present study demonstrated that IFN- γ induced COX-2 expression, which was suppressed by 6-MITC (Fig. 2). IFN- γ did not activate the transcriptional factors including NF- κ B, C/EBP, CREB and AP-1, which bind to the core element of COX-2 promoter (Fig. 4). Blanco *et al* reported that a member of the IRF family, IRF-1, regulates IFN- γ -induced COX-2 gene expression through two ISRE sites in mouse peritoneal macrophages (31). ISRE II sites are located in the -1354/-1345 region of COX-2 promoter, and are efficiently bound with IRF-1 and IRF-2. In this study, transient transfections using COX-2-luciferase constructs indicated that IFN- γ increased COX-2 gene expression through the region between -1432 and -327 in COX-2 promoter (Fig. 3). Therefore, the inhibitory action of 6-MITC on IFN- γ -induced COX-2 expression is probably associated with IRF-1/2-ISRE II pathway, which is worthy of further study.

TPA has been shown to upregulate COX-2 expression in several cell lines. In the present study, we found that TPA activated CREB and AP-1, and induced COX-2 expression. However, 6-MITC could not block them. On the other hand, 6-MITC strongly blocked LPS-induced activation of CREB and AP-1. These results indicate that the signaling pathway leading to the activation of CREB and AP-1 by TPA is different from that induced by LPS. It has been reported that TPA acted as an activator of protein kinase C in several cell lines including RAW264.7 cells (32-34), and TPA-induced PKC activation is strongly linked with COX-2 induction. However, further study is needed to clarify this.

In summary, we report for the first time that 6-MITC suppressed COX-2 expression induced by LPS or IFN- γ , but not TPA, in murine macrophages. Molecular analysis revealed that LPS, IFN- γ and TPA might induce COX-2 expression through different signaling pathways. These findings are very helpful to further understand the property of 6-MITC as an anti-inflammatory and anti-cancer agent.

Acknowledgements

We thank Dr H. Inoue, Nara Women's University, Nara, for COX-2-luciferase constructs, and Dr Y. Fuke, Tokyo Metropolitan College, for MITCs. This work was partly supported by Kagoshima University Fund for Frontier Science Research.

References

- Murillo G and Mehta RG: Cruciferous vegetables and cancer prevention. *Nutr Cancer* 41: 17-28, 2001.
- Zhang Y: Cancer-preventive isothiocyanates: measurement of human exposure and mechanism of action. *Mutat Res* 555: 173-190, 2004.
- Fuke Y, Haga Y, Ono H, Nomura T and Ryoyama K: Anti-carcinogenic activity of 6-methylsulfinylhexyl isothiocyanates (6-MITC), and an active antiproliferative principal of wasabi (*Eutrema wasabi* Maxim.). *Cytotechnology* 25: 197-203, 1997.
- Yano T, Yajima S, Virgona N, Yano Y, Otani S, Kumagai H, Sakurai H, Kishimoto M and Ichikawa T: The effect of 6-methylthiohexyl isothiocyanate isolated from *Wasabia japonica* (wasabi) on 4-(methylnitrosamino)-1-(3-pyridyl)-1-butane-induced lung tumorigenesis in mice. *Cancer Lett* 155: 115-120, 2000.
- Nomura T, Shinoda S, Yamori T, Sawaki S, Nagata I, Ryoyama K and Fuke Y: Selective sensitivity to wasabi-derived 6-(methylsulfinyl)hexyl isothiocyanate of human breast cancer and melanoma cell lines studied *in vitro*. *Cancer Detect Prev* 29: 155-160, 2005.
- Funk CD, Funk LB, Kennedy ME, Pong AS and Fitzgerald GA: Human platelet/erythrocyte cell prostaglandin G/H synthase: cDNA cloning, expression, and gene chromosomal assignment. *FASEB J* 5: 2304-2312, 1991.
- Hempel SL, Monick MM and Hunninghake GW: Lipopolysaccharide induces prostaglandin H synthase-2 protein and mRNA in human alveolar macrophages and blood monocytes. *J Clin Invest* 93: 391-396, 1994.
- Crofford LJ, Wilder RL, Ristimaki AP, Sano H, Remmers EF, Epps HR and Hla T: Cyclooxygenase-1 and -2 expression in rheumatoid synovial tissues. Effects of interleukin-1 beta, phorbol ester, and corticosteroids. *J Clin Invest* 93: 1095-1101, 1994.
- Rolland PH, Martin PM, Jacquemier J, Rolland AM and Toga M: Prostaglandin in human breast cancer: evidence suggesting that an elevated prostaglandin production is a marker of high metastatic potential for neoplastic cells. *J Natl Cancer Inst* 64: 1061-1070, 1980.
- Lupulescu A: Prostaglandins, their inhibitors and cancer. *Prostaglandins Leukot Essent Fatty Acids* 54: 83-94, 1996.

11. Sheng H, Shao J, Morrow JD, Beauchamp RD and DuBois RN: Modulation of apoptosis and Bcl-2 expression by prostaglandin E2 in human colon cancer cells. *Cancer Res* 58: 362-366, 1998.
12. Ben-Av P, Crofford LJ, Wilder RL and Hla T: Induction of vascular endothelial growth factor expression in synovial fibroblasts by prostaglandin E and interleukin-1: a potential mechanism for inflammatory angiogenesis. *FEBS Lett* 372: 83-87, 1997.
13. Goodwin JS and Ceuppens J: Regulation of the immune response by prostaglandins. *J Clin Immunol* 3: 295-314, 1983.
14. Tsujii M and DuBois RN: Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase 2. *Cell* 83: 493-501, 1995.
15. Tanabe T and Tohnai N: Cyclooxygenase isozymes and their gene structures and expression. *Prostaglandins Other Lipid Mediat* 68-69: 95-114, 2002.
16. Hou D-X, Kai K, Li JJ, Lin S, Terahara N, Wakamatsu M, Fujii M, Young MR and Colburn N: Anthocyanidins inhibit activator protein 1 activity and cell transformation: structure-activity relationship and molecular mechanisms. *Carcinogenesis* 25: 29-36, 2004.
17. Inoue H and Tanabe T: Transcriptional role of the nuclear factor kappa B site in the induction by lipopolysaccharide and suppression by dexamethasone of cyclooxygenase-2 in U937 cells. *Biochem Biophys Res Commun* 244: 143-148, 1998.
18. Inoue H, Yokoyama C, Hara S, Tone Y and Tanabe T: Transcriptional regulation of human prostaglandin-endoperoxide synthase-2 gene by lipopolysaccharide and phorbol ester in vascular endothelial cells. *J Biol Chem* 270: 24965-24971, 1995.
19. Inoue H, Nanayama T, Hara S, Yokoyama C and Tanabe T: The cyclic AMP response element plays an essential role in the expression of the human prostaglandin-endoperoxide synthase 2 gene in differentiated U937 monocytic cells. *FEBS Lett* 350: 51-54, 1994.
20. Hou D-X, Fukuda M, Fujii M and Fuke Y: Transcriptional regulation of nicotinamide adenine dinucleotide phosphate: quinone oxidoreductase in murine hepatoma cells by 6-(methylsulfinyl)hexyl isothiocyanate, an active principle of wasabi (*Etrema wasabi* Maxim). *Cancer Lett* 20: 195-200, 2000.
21. Duque J, Díaz-Muñoz MD, Fresno M and Iñiguez MA: Up-regulation of cyclooxygenase-2 by interleukin-1beta in colon carcinoma cells. *Cell Signal* 18: 1262-1269, 2006.
22. Moore PE, Lahiri T, Laporte JD, Church T, Panettieri RA and Shore SA: Selected contribution: synergism between TNF-alpha and IL-1 beta in airway smooth muscle cells: implications for beta-adrenergic responsiveness. *J Appl Physiol* 91: 1467-1474, 2001.
23. Lin CC, Hsiao LD, Chien CS, Lee CW, Hsieh JT and Yang CM: Tumor necrosis factor-alpha-induced cyclooxygenase-2 expression in human tracheal smooth muscle cells: involvement of p42/p44 and p38 mitogen-activated protein kinases and nuclear factor-kappaB. *Cell Signal* 16: 597-607, 2004.
24. Pettus BJ, Bielawski J, Porcelli AM, Reames DL, Johnson KR, Morrow J, Chalfant CE, Obeid LM and Hannun YA: The sphingosine kinase 1/sphingosine-1-phosphate pathway mediates COX-2 induction and PGE2 production in response to TNF-alpha. *FASEB J* 17: 1411-1421, 2003.
25. Uto T, Fujii M and Hou D-X: Inhibition of lipopolysaccharide-induced cyclooxygenase-2 transcription by 6-(methylsulfinyl)hexyl isothiocyanate, a chemopreventive compound from *Wasabia japonica* (Miq.) Matsumura, in mouse macrophages. *Biochem Pharmacol* 70: 1772-1784, 2005.
26. Thomas B, Berenbaum F, Humbert L, Bian H, Bereziat G, Crofford L and Olivier JL: Critical role of C/EBPdelta and C/EBPbeta factors in the stimulation of the cyclooxygenase-2 gene transcription by interleukin-1beta in articular chondrocytes. *Eur J Biochem* 267: 6798-6809, 2000.
27. D'Acquisto F, Iuvone T, Rombola L, Sautebin L, Di Rosa M and Carnuccio R: Involvement of NF-kappaB in the regulation of cyclooxygenase-2 protein expression in LPS-stimulated J774 macrophages. *FEBS Lett* 418: 175-178, 1997.
28. Zandi E, Rothwarf DM, Delhase M, Hayakawa M and Karin M: The IkkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkkappaB phosphorylation and NF-kappaB activation. *Cell* 91: 243-252, 1997.
29. Mestre JR, Mackrell PJ, Rivadeneira DE, Stapleton PP, Tanabe T and Daly JM: Redundancy in the signaling pathways and promoter elements regulating cyclooxygenase-2 gene expression in endotoxin-treated macrophage/monocytic cells. *J Biol Chem* 276: 3977-3982, 2001.
30. Boehm U, Klamp T, Groot M and Howard JC: Cellular responses to interferon-gamma. *Annu Rev Immunol* 15: 749-795, 1997.
31. Blanco JC, Contursi C, Salkowski CA, De Witt DL, Ozato K and Vogel SN: Interferon regulatory factor (IRF)-1 and IRF-2 regulate interferon gamma-dependent cyclooxygenase 2 expression. *J Exp Med* 191: 2131-2144, 2000.
32. Chen CC, Wang JK and Chen WC: TPA induces translocation but not down-regulation of new PKC isoform eta in macrophages, MDCK cells and astrocytes. *FEBS Lett* 412: 30-34, 1997.
33. Chang MS, Chen BC, Yu MT, Sheu JR, Chen TF and Lin CH: Phorbol 12-myristate 13-acetate upregulates cyclooxygenase-2 expression in human pulmonary epithelial cells via Ras, Raf-1, ERK, and NF-kappaB, but not p38 MAPK, pathways. *Cell Signal* 17: 299-310, 2005.
34. Subbaramaiah K, Cole PA and Dannenberg AJ: Retinoids and carnosol suppress cyclooxygenase-2 transcription by CREB-binding protein/p300-dependent and -independent mechanisms. *Cancer Res* 62: 2522-2530, 2002.