

The NQO1*2/*2 polymorphism is associated with poor overall survival in patients following resection of stages II and IIIa non-small cell lung cancer

JILL M. KOLESAR^{1,2}, SUZANNE E. DAHLBERG^{3,5}, SHARON MARSH⁴, HOWARD L. McLEOD⁶,
DAVID H. JOHNSON^{7,10}, STEVEN M. KELLER⁸ and JOAN H. SCHILLER⁹

¹School of Pharmacy, ²University of Wisconsin Paul P. Carbone Comprehensive Cancer Center, University of Wisconsin, Madison, WI, USA; ³Eastern Cooperative Oncology Group, ⁴Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada; ⁵Dana-Farber Cancer Institute, Boston, MA; ⁶University of North Carolina, Chapel Hill, NC; ⁷Vanderbilt University, Nashville, TN; ⁸Albert Einstein College of Medicine, Bronx, NY; ⁹University of Texas Southwestern Medical Center, Dallas, TX, USA

Received February 1, 2011; Accepted March 15, 2011

DOI: 10.3892/or.2011.1249

Abstract. NAD(P)H:quinone oxidoreductase 1 (NQO1), is a cytosolic flavoenzyme that catalyzes the two-electron reduction of quinones into hydroquinones. A polymorphism (NQO1*2) alters enzymatic activity of NQO1 resulting in diminished NQO1 activity. Malignancies with NQO1*2 may be resistant to radiation and chemotherapy with resulting poorer survival. NQO1 allele was evaluated in subjects enrolled in ECOG 3590, a randomized comparison of radiation (RT) vs radiation and chemotherapy with cisplatin/etoposide (RCT) in patients with completely resected stages II and IIIa NSCLC. Overall survival was estimated using the Kaplan-Meier method and compared via the log-rank test. Cox models were used to assess the impact of covariates on outcomes. Among 152 patients with assessable samples, 24 (16%) had NQO1*2. Median follow-up was 139 months. The presence of NQO1*2/*2 was associated with decreased overall survival (OS) (median in the heterozygote/wild-type group 42.3 vs. 33.5 months in the variant group, $p=0.04$). In a multivariable Cox model, variant NQO1 (HR=1.58, $p=0.05$), age <60 (HR=0.67, $p=0.04$), PS 1 (HR=1.47, $p=0.05$), cardiovascular disease (HR=1.93, $p=0.003$) and alkaline phosphatase <100 mg/ml (HR=0.59, $p=0.005$) were all significant predictors of OS. NQO1*2/*2 may be an independent predictor of poor overall survival in individuals with resected stages II and IIIa NSCLC. Although the basis

for the NQO1 association with decreased survival requires additional evaluation, NQO1 may represent a biomarker for guiding individualized therapy.

Introduction

Lung cancer is the leading cause of cancer death in the United States, with 219440 new diagnoses and 159390 deaths estimated for 2009 (1). Lung cancer is the most common cause of cancer death among both men and woman in industrialized nations (2). Both chemotherapy and radiotherapy are utilized to treat locally advanced or metastatic tumors. Moderate survival improvements are reported for patients with inoperable disease who received chemotherapy as well as for patients who have received platinum based chemotherapy following surgery (3-6). Despite treatment, overall survival remains poor, not all patients respond to therapy, and patients receiving chemotherapy experience significant toxicity (4). Genetic markers are a potential method to identify patients most likely to benefit from chemotherapy or radiotherapy.

NAD(P)H:quinone oxidoreductase 1 (NQO1), is a cytosolic flavoenzyme that catalyzes the two-electron reduction of quinones into hydroquinones. NQO1 contains one characteristic polymorphism (NQO1*2), a C→T transition at bp 609 that results in a proline to serine substitution at codon 187 (7,8). Individuals homozygous for NQO1*2 have essentially no active enzyme (9), while those who are heterozygotes have intermediate activity when compared to NQO1*1 homozygotes (10). The decreased activity associated with this polymorphism is mediated by two mechanisms: i) the polymorphism is in the active site of NQO1, and ii) the NQO1*2 protein is degraded rapidly via the ubiquitin-dependent pathway (half-life of wild-type NQO1 >18 h and the half-life of variant NQO1*2 is 1.2 h) (10).

A lack of NQO1 may increase the generation of semiquinones rather than hydroquinones, from the quinines found in cigarette smoke. Polymorphisms in NQO1 have therefore

Correspondence to: Dr Jill Kolesar, University of Wisconsin Comprehensive Cancer Center K4/554, 600 Highland Avenue, Madison, WI 53792-5669, USA
E-mail: jmkolesar@pharmacy.wisc.edu

Present address: ¹⁰University of Texas Southwestern Medical Center, Dallas, TX, USA

Key words: lung cancer, NAD(P)H:quinone oxidoreductase 1, polymorphism, p53

Table I. Baseline demographic variables, stratified by NQO1 status E3590, n=152 patients.

Variable	All (n=152)	Wild-type or heterozygous (n=128)	(%)	Variant (n=24)	(%)	P-value
Treatment Arm						
RT	78	65	50.78	13	54.17	0.82
RTC	74	63	49.22	11	45.83	
Total	152	128		24		
Age (median, range)		60.5	35-77	63	35-81	0.19
Gender						
Male	58	49	38.28	9	37.50	1
Female	94	79	61.72	15	62.50	
Total	152	128		24		
p53						
Mutated	66	61	48.03	5	22.73	0.04
Wild	83	66	51.97	17	77.27	
Total	149	127		22		
Race						
White	133	110	85.94	23	95.83	0.31
Other	19	18	14.06	1	4.17	
Total	152	128		24		
PS						
0	61	49	38.28	12	50.00	0.36
1	91	79	61.72	12	50.00	
Total	152	128		24		
Weight loss (%)						
None	73	60	47.24	13	54.17	0.72
<5	43	35	27.56	8	33.33	
5-10	25	23	18.11	2	8.33	
>10	10	9	7.09	1	4.17	
Total	151	127		24		
Cardiovascular diseases						
Yes	33	29	22.83	4	16.67	0.6
No	118	98	77.17	20	83.33	
Total	151	127		24		
Respiratory diseases						
Yes	26	22	17.32	4	16.67	1
No	125	105	82.68	20	83.33	
Total	151	127		24		
Other diseases						
Yes	41	37	30.33	4	18.18	0.31
No	103	85	69.67	18	81.82	
Total	144	122		22		
Analgesics						
Yes	47	44	34.92	3	13.04	0.05
No	102	82	65.08	20	86.96	
Total	149	126		23		
Steroids						
Yes	6	6	4.92	0	0.00	0.59
No	140	116	95.08	24	100.00	
Total	146	122		24		

Table I. Continued.

Variable	All (n=152)	Wild-type or heterozygous (n=128)	(%)	Variant (n=24)	(%)	P-value
Other medicines						
Yes	80	76	62.30	4	18.18	0.0001
No	64	46	37.70	18	81.82	
Total	144	122		22		
Albumin (median, range)		40	25-53	39	31-44	0.16
Alkaline phosphatase (median, range)		98	44-284	112.5	59-416	0.24
Histology						
Squamous	53	43	33.59	10	41.67	0.8
Small cell	0	0	0.00	0	0.00	
Adenocarcinoma	81	69	53.91	12	50.00	
Large cell	12	10	7.81	2	8.33	
Other	6	6	4.69	0	0.00	
Total	152	128		24		
Nodal status						
N1	78	65	51.59	13	54.17	0.83
N2	72	61	48.41	11	45.83	
Total	150	126		24		
Lymph node dissection						
Complete	74	60	47.24	14	58.33	0.38
Sampling	77	67	52.76	10	41.67	
Total	151	127		24		
Stage						
II	71	61	47.66	10	41.67	0.66
IIIa	81	67	52.34	14	58.33	
Total	152	128		24		
TNM stage						
T1	38	32	25.00	6	25.00	0.1
T2	100	87	67.97	13	54.17	
T3	14	9	7.03	5	20.83	
Total	152	128		24		
Previous chemotherapy						
No	151	128	100.00	23	95.83	0.16
Yes, combinations	1	0	0.00	1	4.17	
Total	152	128		24		

been evaluated as modifiers of lung cancer risk and have been the subject of two meta-analysis. Chao and colleagues (11) reported that in Caucasian populations, there was an increased prevalence of the variant allele among ever smokers with lung cancer (OR, 1.08; 95% CI, 0.96-1.22), but not among never smokers (OR, 0.97; 95% CI, 0.62-1.54). However, Kiyohara *et al* (12) showed the opposite effect in a Japanese cohort where the allele was significantly associated with decreased risk of lung cancer [random effects odds ratio (OR) = 0.70]. The role of NQO1 in lung cancer pathogenesis appears small.

NQO1 has also been studied as a predictor of chemosensitivity and radiosensitivity and overexpression has typically been associated with increased chemosensitivity and radiosensitivity (13). NQO1 gene expression is induced in response to xenobiotics, antioxidants, UV light, ionizing radiation and cisplatin. Additionally, wild-type NQO1 binds to and stabilizes both mutant and wild-type p53 under conditions of oxidative stress or after exposure to γ -irradiation. Degradation of p53 is mediated by two alternative pathways, ubiquitin-independent or ubiquitin-dependent. The ubiquitin-independent pathway is

regulated by NQO1 and is mediated via the 20S proteasome (13).

Song *et al* (14) demonstrated that NQO1 genotype significantly influenced loco-regional recurrence in Korean patients with completely resected stage II or IIIa NSCLC who were treated with adjuvant radiation and sometimes adjuvant chemotherapy [wild-type (NQO1*1/*1) 6.3% (2/32), heterozygous patients 8.1% (5/62), NQO1*2/*2 27.3% (6/22)]. Multivariate analysis revealed that NQO1 genotype and pathologic nodal stage were statistically significant predictive factors for loco-regional recurrence-free survival, but not overall survival.

Individuals with NQO1*2 polymorphisms may be both chemoresistant and radiation resistant. The current study was performed to determine the effect of the NQO1 genetic polymorphisms on overall survival in stages II and IIIa NSCLC patients treated with RT or RTC.

Materials and methods

Clinical study. ECOG 3590 was a randomized, prospective trial of adjuvant therapy in patients with resected stages II or IIIa NSCLC (15). All patients had complete resection of their tumor and mediastinal lymph node sampling or dissection. Eighty-five percent were Caucasian. Following stratification by weight loss, histology, nodal biopsy technique and lymph node status, patients were randomized to receive chemotherapy (cisplatin 60 mg/m² IV on day 1 and etoposide 120 mg/m² days 1-3) with concurrent RT (50.4 Gy in 28 daily fractions) or RT alone.

Patients enrolled in ECOG 3590 were also eligible for enrollment in ECOG 4592 (16), a study designed to evaluate the prognostic value of p53 mutations in exons 5-8 which were determined by single-strand confirmation polymorphism (SSCP) analysis. Genomic DNA extracted from submitted paraffin blocks were utilized. Patients enrolled in ECOG 3590 gave permission to use samples for future studies. The current study design was reviewed and approved by the University of Wisconsin Institutional Review Board.

Genotyping. Genomic DNA extracted from the ECOG 4592 samples was stored at -80°C and variations in the NQO1 gene were analyzed by pyrosequencing. Initial polymerase chain reaction (PCR) primers were designed using Primer Express version 1.5 (ABI, Foster City, CA) and consisted of forward primer (5'-CTGATGGAATTGGTTGACTTA-3'), and reverse primer [5'(Biotin)-TCTGGTGATCACCAAGGCT-3'] to amplify a portion of the NQO1 gene. The forward pyrosequencing primer (5'-GTGGCTTCCAAGTCTTA-3') was designed using Pyrosequencing SNP Primer Design Version 1.01 software (<http://www.pyrosequencing.com>). Primers were obtained from Integrated DNA Technologies (Coralville, IA). PCR was carried out using Amplitaq Gold PCR master mix (ABI), 5 pmole of each primer, and 10 ng DNA, on an MJ Research PTC-100 thermal cycler (Waltham, MA). Reactions were denatured initially at 93°C for 20 min, then cycled at 95°C for 30 sec, 55°C for 30 sec and 72°C for 30 sec for 55 cycles, followed by a final elongation step at 72°C for 5 min. Pyrosequencing was carried out as previously described using a PSQ96 instrument and software (Biotage AB, Uppsala, Sweden) (32). Genotype was called variant if it differed from




Figure 1. Kaplan-Meier curve comparing NQO1*1/*1 and NQO1*1/*2 to NQO1*2/*2 in all patients median OS in the heterozygous/wild-type group was 42.3 vs. 33.5 months in the variant group ($p=0.04$).

the Refseq consensus sequence for the SNP position. To ensure proper amplification and the absence of contamination, each PCR and pyrosequencing run contained no template controls or cancer cell line positive controls.

Statistical analysis. Polymorphisms were classified into two groups, heterozygous/wild-type and variant. Wild-types and heterozygotes were combined as NQO1 activity is similar. OS, defined as the time from randomization to death from any cause, was estimated using the Kaplan-Meier method. Differences between the polymorphism status groups were analyzed using the log-rank test. Fisher's exact test and the Kruskal-Wallis rank sum test were used to compare baseline variables between the two polymorphism groups. Cox models were used to assess the univariate association between baseline variables and overall survival. Variables significant at the 0.10 level were fitted to a multivariable Cox proportional hazards model and backwards selection was used to select the final model.

Results

Four hundred and eighty-eight patients were enrolled in ECOG 3590, of whom 217 entered ECOG 4592. Among 152 patients with assessable samples, 24 (16%) had variant NQO1. Most demographic variables were balanced between the two NQO1 groups (Table I). However, a greater proportion of patients with mutated p53 was present in the heterozygous/wild-type NQO1 group, ($p=0.04$).

Median follow-up was 139 months. One hundred and twenty-three (81%) patients had died. Median OS in the heterozygous/wild-type group was 42.3 months (95% CI 35.9-66.3 mos) vs. 33.5 months (95% CI 16.2-80.9 mos) in the variant group ($p=0.04$, Fig. 1). Among patients receiving




Figure 2. Kaplan-Meier curve comparing NQO1^{+/+} and NQO1^{+/2} to NQO1^{2/2} in patients receiving RT only, no difference in OS was observed (p=0.49).




Figure 3. Kaplan-Meier curve comparing NQO1^{+/+} and NQO1^{+/2} to NQO1^{2/2} in patients receiving CRT, demonstrating a significant difference in median OS in favor of the heterozygous/wild-type group was observed (40.0 vs. 16.2 months, p=0.007).

RT only, no difference in OS was observed (45.2 mos (95% CI 35.9-91.8 mos) vs. 53.4 mos (95% CI 29.2-Inf); p=0.49, Fig. 2). However, among those patients who received chemoradiotherapy a significant difference in median OS in favor of the heterozygous/wild-type group was observed (40.0 months (95% CI 30.8-80.6 mos) vs. 16.2 months (13.6-Inf mos), p=0.007, Fig. 3).

Univariate Cox proportional hazards models (Table II) revealed that overall survival was associated with: NQO1 status, age, race, performance status, cardiovascular disease, alkaline phosphatase (mg/ml) and TNM stage. Treatment and NQO1 status were not statistically significant when adjusted for these two factors alone (p=0.16). A multivariable Cox model revealed variant NQO1 (HR=1.58, p=0.05), age <60 (HR 0.67, p=0.04), PS 1 (HR=1.47, p=0.05), cardiovascular disease (HR=1.93, p=0.003) and alkaline phosphatase <100 mg/ml (HR= 0.59, p=0.005) were all significant predictors of OS. A treatment by NQO1 status interaction test was not statistically significant (p=0.16).

Discussion

Our study is the first to report that NQO1^{2/2} may be an independent predictor of poor overall survival in individuals with resected stages II and IIIa NSCLC. These results are in contrast with Song and colleagues who identified a decrease in locoregional recurrence-free survival in NQO1^{2/2} patients, but not in overall survival. This may be related to the different length of median follow-up (49.2 months compared to 139 months), and chemotherapy differences (cisplatin/vinorelbine allowed vs. patients randomized to cisplatin/etoposide or no chemotherapy). Locoregional recurrence-free survival was not evaluated as part of E3590. These results suggest that NQO1^{2/2} may be considered an independent predictor of poor survival for patients with NSCLC. Individuals with NQO1^{2/2} have no NQO1 activity, and subsequently lose NQO1 stabilization of three tumor suppressors p53 (17-19), p73a (20) and p33 (21), by inhibiting their breakdown via the 20S proteasome. Lack of stabilization may lead to a more aggressive tumor biology and lack of responsiveness to chemotherapy and radiation therapy (22-25).

These data suggest that RT is the optimal therapy for individuals with NQO1^{2/2}. The lack of a difference in OS when analyzed by treatment and NQO1 status (p=0.016) may be related to inadequate power.

In this study, patients who were NQO1^{2/2} were more likely to be p53 wild-type. Since NQO1^{+/+} is known to bind to and stabilize both mutant and wild-type p53 (and these genotypes should have opposite prognostic effects), we hypothesize that p53 status becomes important only in the absence of NQO1 activity. That the role of p53 in lung cancer prognosis has been the subject of dozens of paper and two meta-analyses (25,26) without resolution of the question, lends support to this hypothesis. The association between NQO1^{2/2} and p53 wild-type suggests that these genes may be important in lung cancer pathogenesis.

Our study has several limitations, including that it was as unplanned retrospective analysis, had limited in power to detect treatment effects and used of SSCP for identifying p53 mutations. While p53 mutations were evaluated in exons 5-8, the specific mutation is not identifiable by SSCP. Therefore, the effect of the mutation on p53 protein expression and the interaction between NQO1 and the mutant p53 is unknown. However, this study does have the advantage of utilizing data from a large, randomized multi-institutional trial with mature survival data.

In summary, NQO1^{2/2} independently predicts poor survival in completely resected NSCLC stages II and III patients. In addi-

Table II. Univariate Cox model results.

Variable	No. of deaths	Median OS	HR for death	95% CI	P-value
NQO1					
Wild-type or heterozygote	99	42.3	ref		0.046
Variant	24	33.5	1.58	1.01-2.47	
Treatment Arm					
RT	62	48.5	ref		0.49
RT + chemotherapy	61	37.7	1.13	0.79-1.62	
Age (years)					
≥60	74	35.9	ref		0.011
<60	49	62.2	0.624	0.43-0.9	
Gender					
Male	77	37.1	1.2	0.83-1.74	0.32
Female	46	55.1	ref		
p53					
Mutated	57	39.9	ref		0.42
Wild-type	63	50.7	0.863	0.6-1.23	
Race					
White	110	38.2	1.64	0.92-2.92	0.091
Other	13	80.6	ref		
PS					
0	46	63.7	ref		0.091
1	77	38.3	1.37	0.95-1.98	
Weight loss (%)					
<5	94	44	0.972	0.6-1.41	0.9
≥5	28	38.3	ref		
Cardiovascular diseases					
Yes	33	30.8	1.93	1.29-2.89	0.0015
No	89	52.7	ref		
Respiratory diseases					
Yes	23	53.2	0.964	0.61-1.52	0.88
No	99	39.8	ref		
Other diseases					
Yes	32	65.4	0.782	0.52-1.18	0.24
No	83	38.2	ref		
Analgesics					
Yes	38	40.2	1.02	0.7-1.51	0.91
No	82	43.5	ref		
Steroids					
Yes	5	55.5	0.993	0.4-2.44	0.99
No	112	41	ref		
Other medicins					
Yes	67	41.8	1.15	0.81-1.65	0.45
No	49	41	ref		
Albumin					
≥40	60	39.8	ref		0.57
<40	60	40.2	1.11	0.76-1.59	

Table II. Continued.

Variable	No. of deaths	Median OS	HR for death	95% CI	P-value
Alkaline phosphatase					
≥100	65	28	ref		
<100	56	66.3	0.59	0.41-0.84	0.004
Histology					
Squamous	45	45.2	1.01	0.7-1.46	0.96
Non-squamous	78	39.9	ref		
Nodal status					
N1	60	43.5	ref		
N2	61	40	1.19	0.83-1.69	0.35
Lymph node					
Complete	62	61	ref		
Dissection					
Sampling	60	40	1.04	0.73-1.49	0.83
Stage					
II	53	41.8	ref		
IIIA	70	40.2	1.26	0.88-1.81	0.2
TNM stage					
T1/T2	109	44	ref		
T3	14	18.6	1.93	1.1-3.37	0.022

tion, NQO1*2 is associated with p53 wild-type. Prospective trials and determination of specific p53 mutations are needed to confirm and define the relationship of NQO1, p53 and NSCLC. The results of the current study demonstrate that evaluation of NQO1 polymorphisms may be an important consideration in individualizing the treatment of lung cancer.

Acknowledgements

This study was supported by the ACCP Oncology Research Award, the NIH Pharmacogenetics Research Network (GM63340), and the American Cancer Society.

References

- Jemal A, Siegel R, Ward E, Hao Y, Xu J and Thun MJ: Cancer statistics, 2009. *CA Cancer J Clin* 59: 225-249, 2009.
- Pisani P, Parkin DM and Ferlay J: Estimates of the worldwide mortality from eighteen major cancers in 1985. *Int J Cancer* 55: 891-903, 1993.
- Keller SM, Adak S, Wagner H, *et al*: A randomized trial of post-operative adjuvant therapy in patients with completely resected stage II or IIIA non-small cell lung cancer. *N Engl J Med* 343: 1217-1222, 2000.
- Rube C and Fleckenstein J: Combined-modality treatment of non-small cell lung cancer stages I-III (take home messages). *Lung Cancer* 45: S139-S141, 2004.
- Belani CP: Adjuvant and neoadjuvant therapy in non-small cell lung cancer. *Semin Oncol* 32: S9-S15, 2005.
- Bradley JD, Paulus R, Graham MV, *et al*: Phase II trial of postoperative adjuvant paclitaxel/carboplatin and thoracic radiotherapy in resected stage II and IIIA non-small cell lung cancer: promising long-term results of the radiation therapy oncology group-RTOG 9705. *J Clin Oncol* 23: 3480-3487, 2005.
- Traver RD, Horikoshi T, Danenberg KD, *et al*: NAD(P)H:quinone oxidoreductase gene expression in human colon carcinoma cells: characterization of a mutation which modulates DT-diaphorase activity and mitomycin sensitivity. *Cancer Res* 52: 797-802, 1992.
- Ross D, Traver RD, Siegel D, *et al*: A polymorphism in NAD(P)H:quinone oxidoreductase (NQO1): relationship of a homozygous mutation at position 609 of the NQO1 cDNA to NQO1 activity. *Br J Cancer* 74: 995-996, 1996.
- Traver RD, Siegel D, Beall HD, *et al*: Characterization of a polymorphism in NAD(P)H: quinone oxidoreductase (DT-diaphorase). *Br J Cancer* 75: 69-75, 1997.
- Siegel D, McGuinness SM, Winski SL, *et al*: Genotype-phenotype relationships in studies of a polymorphism in NAD(P)H:quinone oxidoreductase 1. *Pharmacogenetics* 9: 113-121, 1999.
- Chao C, Zhang ZF, Berthiller J, Boffetta P and Hashibe M: NAD(P)H:quinone oxidoreductase 1 (NQO1) Pro187Ser polymorphism and the risk of lung, bladder, and colorectal cancers: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* 5: 979-987, 2006.
- Kiyohara C, Yoshimasu K, Takayama K and Nakanishi Y: NQO1, MPO, and the risk of lung cancer: a HuGE review. *Genet Med* 7: 463-478, 2005.
- Dinkova-Kostova AT and Talalay P: NAD(P)H:quinone acceptor oxidoreductase 1 (NQO1), a multifunctional antioxidant enzyme and exceptionally versatile cytoprotector. *Arch Biochem Biophys* 1: 116-123, 2010.
- Song SY, Jeong SY, Park HJ, *et al*: Clinical significance of NQO1 C609T polymorphisms after postoperative radiation therapy in completely resected non-small cell lung cancer. *Lung Cancer* 2: 278-282, 2010.
- Schiller JH, Harrington D, Belani CP, Langer C, Sandler A, Krook J, Zhu J, Johnson DH and Eastern Cooperative Oncology Group: Comparison of four chemotherapy regimens for advanced non-small cell lung cancer. *N Engl J Med* 346: 92-98, 2002.
- Schiller JH, Adak S, Feins RH, Keller SM, Fry WA, Livingston RB, Hammond ME, Wolf B, Sabatini L, Jett J, Kohman L and Johnson DH: Lack of prognostic significance of p53 and K-ras mutations in primary resected non-small cell lung cancer on E4592: a Laboratory Ancillary Study on an Eastern Cooperative Oncology Group Prospective Randomized Trial of Postoperative Adjuvant Therapy. *J Clin Oncol* 19: 448-457, 2001.

17. Asher G, Lotem J, Sachs L, *et al*: Mdm-2 and ubiquitin-independent p53 proteasomal degradation regulated by NQO1. *Proc Natl Acad Sci USA* 99: 13125-13130, 2002.
18. Asher G, Lotem J, Kama R, *et al*: NQO1 stabilizes p53 through a distinct pathway. *Proc Natl Acad Sci USA* 99: 3099-3104, 2002.
19. Asher G, Lotem J, Cohen B, *et al*: Regulation of p53 stability and p53-dependent apoptosis by NADH quinone oxidoreductase 1. *Proc Natl Acad Sci USA* 98: 1188-1193, 2001.
20. Asher G, Tsvetkov P, Kahana C and Shaul Y: A mechanism of ubiquitin-independent proteasomal degradation of the tumor suppressors p53 and p73. *Genes Dev* 19: 316-321, 2005.
21. Garate M, Wong RP, Campos EI, Wang Y and Li G: NAD(P)H quinone oxidoreductase 1 inhibits the proteasomal degradation of the tumour suppressor p33(ING1b). *EMBO J* 6: 576-581, 2008.
22. Tsutsumi K, Yasuda M and Nishioka T: X-ray irradiation altered chemosensitivity of a p53-null non-small cell lung cancer cell line. *Cell Struct Funct* 31: 47-52, 2006.
23. Lai SL, Perng RP and Hwang J: p53 gene status modulates the chemosensitivity of non-small cell lung cancer cells. *J Biomed Sci* 7: 64-70, 2000.
24. Rusch V, Klimstra D, Venkatraman E, *et al*: Aberrant p53 expression predicts clinical resistance to cisplatin-based chemotherapy in locally advanced non-small cell lung cancer. *Cancer Res* 55: 5038-5042, 1995.
25. Tammemagi MC, McLaughlin JR and Bull SB: Meta-analyses of p53 tumor suppressor gene alterations and clinicopathological features in resected lung cancers. *Cancer Epidemiol Biomarkers Prev* 8: 625-634, 1999.
26. Mitsudomi T, Hamajima N, Ogawa M, *et al*: Prognostic significance of p53 alterations in patients with non-small cell lung cancer: a meta-analysis. *Clin Cancer Res* 6: 4055-4063, 2000.