Interleukin-4 receptor α-based hybrid peptide effectively induces antitumor activity in head and neck squamous cell carcinoma

KAHORI SETO¹, JUNICHI SHODA², TOMOHISA HORIBE³, EIJI WARABI⁴, KAZUNORI ISHIGE⁵, KENJI YAMAGATA⁶, MASAYUKI KOHNO³, TORU YANAGAWA⁶, HIROKI BUKAWA⁶ and KOJI KAWAKAMI³

¹Department of Oral and Maxillofacial Surgery, Clinical Sciences, Graduate School of Comprehensive Human Sciences, University of Tsukuba, Ibaraki 305-8575; ²Medical Science Section, Faculty of Medicine, University of Tsukuba Graduate School, Ibaraki 305-8575; ³Department of Pharmacoepidemiology, Graduate School of Medicine and Public Health, Kyoto University, Kyoto 606-8501; ⁴Division of Biomedical Science, Faculty of Medicine, University of Tsukuba, Ibaraki 305-8575; Departments of ⁵Gastroenterology, ⁶Oral and Maxillofacial Surgery, Faculty of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

Received December 28, 2012; Accepted February 21, 2013

DOI: 10.3892/or.2013.2387

Abstract. Interleukin-4 receptor α (IL-4R α) is highly expressed on the surface of various human solid tumors including head and neck squamous cell carcinoma (HNSCC). We designed a novel IL-4Ra-lytic hybrid peptide composed of a binding peptide to IL-4R α and a cell-lytic peptide. In the present study, we evaluated the antitumor activity of the IL-4Ra-lytic hybrid peptide as a novel molecular-targeted therapy in HNSCC. Immunoblot analysis revealed that IL-4R α was expressed in all tested HNSCC cell lines (HSC-2, HSC-3, HSC-4, Ca9-22 and OSC-19), but not in a human normal keratinocyte (HaCaT) cell line. Immunohistochemical expression levels of IL-4Ra in HNSCC tissues were higher compared to those in normal epithelial tissue. The IL-4Rα-lytic hybrid peptide showed cytotoxic activity in all five cancer cell lines with a concentration that killed 50% of all cells (IC₅₀) as low as 10 μ M. HaCaT cells were less sensitive to this peptide with an IC₅₀ of >30 μ M. In addition, intratumoral administration of IL-4Ra-lytic hybrid peptide significantly inhibited tumor growth in a xenograft model of human HNSCC in vivo. These results indicate that the IL-4R α -lytic hybrid peptide may serve as a potent agent to provide a novel therapy for patients with HNSCC.

Introduction

Head and neck squamous cell carcinoma (HNSCC) which includes cancers of the oral cavity, oropharynx, larynx and

E-mail: kawakami.koji.4e@kyoto-u.ac.jp

hypopharynx is the sixth most common cancer worldwide and has an incidence of approximately 600,000 cases per year (1). The current management and treatment of HNSCC involves multi-modality approaches of surgery, chemotherapy and radiotherapy (2). Despite recent advances in early detection, diagnosis and treatment, the 5-year survival for patients with HNSCC has remained at 50% for the past 30 years (3). Molecular-targeted therapy, based on molecular findings of the last 50 years, is one of the most promising gateways to the development of new strategies in oncology (4). Cetuximab, a monoclonal antibody to the epidermal growth factor receptor (EGFR), is the only molecular-targeted therapy to be routinely used in clinical practice for the treatment of recurrent and metastatic HNSCC (5). Based on a limited number of phase II and III trials that have investigated the efficacy of cetuximab in addition to cisplatin in patients who were refractory to platinum-based therapy, the combination appears to confer further benefit over anti-EGFR agents alone (6,7).

The first generation of immunotoxins developed 35 years ago, which heralded targeted therapy, employed chemical conjugations of antibodies and either intact toxins or toxins with attenuated cell-binding properties. We previously reported that recombinant fusion protein IL4 (38-37)-PE38KDEL (also termed IL4-PE), consisting of circularly permuted interleukin (IL)-4 and a mutated form of Pseudomonas exotoxin (PE), induced significant regression of established biliary tract tumors and significantly improved the survival of animals with disseminated tumors (8,9). In addition, IL4-PE was reported to be highly and specifically cytotoxic to glioma cell lines in vitro, and caused partial or complete regression of established human glioblastoma multiforme tumors in nude mice (10). IL-4 receptor α (IL-4R α)-targeted protein-based immunotoxin was tested in the clinic for the treatment of human solid tumors (11,12). However, its clinical application faced many challenges, including non-specific toxicities and immunogenicity (13).

To overcome these issues, we previously developed a 'hybrid peptide', composed of target-binding and cytotoxic

Correspondence to: Professor Koji Kawakami, Department of Pharmacoepidemiology, Graduate School of Medicine and Public Health, Kyoto University, Yoshida Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan

Key words: interleukin-4 receptor α , hybrid peptide, head and neck squamous cell carcinoma, molecular-targeted therapy

sequences containing cationic-rich D- and L-amino acids to form amphipathic partial α -helices that disrupt the cancer cell membrane selectively, and are stable when combined with a cancer-targeting moiety (14). It is known that peptide drugs are relatively easily synthesized using either recombinant or solidphase chemical synthesis techniques and the production costs are generally affordable when compared to antibody-based therapeutics (14). IL-4R α has been previously reported to have high expression on the surface of a variety of human solid tumors such as renal cell carcinoma, malignant melanoma and glioblastoma (15). Although the biological function of IL-4R α expression on solid tumors remains unclear, this receptor may be an effective candidate for a novel molecular-targeted therapy.

In the present study, we examined the expression levels of IL-4R α in both patient samples and HNSCC cell lines, and then explored the antitumor activity of the IL-4R α -lytic hybrid peptide against HNSCC.

Materials and methods

Patient samples. HNSCC specimens were obtained from 5 patients (4 males, 1 female; mean age, 56.6 years), who underwent radical surgery at the Department of Oral and Maxillofacial Surgery, Tsukuba University Hospital, Japan from 2010 to 2011. Primary tumor sites were the tongue and the gingiva. For immunoblot analysis, we obtained HNSCC tissue from the cancerous lesion (cancer) and normal tissue from the normal area (normal) in other specimens from 1 patient. For immunohistochemistry, tissue was fixed in 10% formalin and paraffin embedded. The study protocol was approved in accordance with the ethics guidelines of the Tsukuba University (H23-61). All patients provided written informed consent for use of specimens.

Peptides. The following IL-4Rα-lytic hybrid peptide, the lytic peptide and IL-4 binding peptide were purchased from Invitrogen: the IL-4Rα-lytic hybrid peptide, KQLIRFLK RLDRNGGGKLLLKLLKKLLKLLKLLKK (underlined letters are D-amino acids); the lytic peptide, KLLLKLLKKLLKLLKK; and the IL-4 binding peptide, KQLIRFLKRLDRN. All peptides were synthesized by the use of solid-phase chemistry, purified to homogeneity by reverse-phase high-pressure liquid chromatography and assessed by mass spectrometry. All peptides were dissolved in water and buffered to pH 7.4.

Cell lines and culture conditions. The HNSCC cell lines (HSC-2, HSC-3, HSC-4, Ca9-22 and OSC-19) were purchased from the Japanese Collection of Research Bioresources (Osaka, Japan). The human normal keratinocyte cell line (HaCaT) was purchased from the American Type Culture Collection (Manassas, VA, USA). These cell lines were maintained in Dulbecco's modified Eagle's medium or RPMI-1640 containing 10% heat-inactivated fetal calf serum (Nichirei Biosciences Inc., Tokyo, Japan) and 1% penicillin-streptomycin in a humidified atmosphere with 5% CO₂ at 37°C.

Immunoblot analysis. Immunoblot analysis was carried out as previously described (9). Briefly, whole-cell extracts were

obtained using buffer containing 1% (v/v) Triton X-100, 0.1% (w/v) SDS, and 0.5% (w/v) sodium deoxycholate, separated by SDS-PAGE, and transferred onto a PVDF membrane. IL-4R α antibody was used at dilution 1:200 (Santa Cruz Biotechnology, Santa Cruz, CA, USA) and actin as the internal control (Sigma-Aldrich, St. Louis, MO, USA). Proteins were visualized on Hyperfirm using an enhanced chemiluminescence/western blotting system (GE Healthcare, Piscataway, NJ, USA).

Quantitative real-time PCR. Total RNA of cells was isolated using High Pure RNA Tissue kit (Roche, Basel, Switzerland). For the reverse transcriptase reaction, 400 ng of the RNA sample was used. The reaction was carried out in a final volume of 10 μ l of reaction mixture with Takara Kit II (Takara, Shiga, Japan). Aliquots (2 μ l) of the cDNA samples were amplified in a final volume of 20 μ l of PCR mixture containing SYBR Premix Ex Taq II (Takara). Quantitative real-time PCR was carried out using PRISM 7000 (Applied Biosystems, Carlsbad, CA, USA). The following primers were used: IL-4R α forward, 5'-CTGACCTGGAGCAACCCGTATC-3' and IL-4R α reverse, 5'-GCAGACGGACAACACGATACAG-3'; GAPDH forward, 5'-GTCTTCACCACCATGGAGAAGGCT-3' and GAPDH reverse, 5'-CATGCCAGTGAGCTTCCCGTTCA-3'.

Cell viability assay. Cell viability assay was performed as previously described (14). Briefly, cells were seeded into 96-well plates at $3x10^3$ cells/well in 90 μ l of medium and incubated at 37° C for 24 h. Each peptide (IL-4R-lytic hybrid peptide, lytic peptide or IL-4 binding peptide) diluted in 10 μ l culture medium was added to the cells. After a 72-h incubation, the cell viability assay using WST-8 solution (Nacalai Tesque, Kyoto, Japan) was performed.

Immunohistochemistry. For immunostaining of IL-4R α , 2- μ m sections from patient samples were stained using the Vectastain kit according to the manufacturer's instructions with the anti-IL-4R α antibody (R&D Systems, Minneapolis, MN, USA).

Binding assay. The IL-4 binding peptide labeled with fluorescein isothiocyanate (FITC) was incubated with HSC-2 and HaCaT cells. Quantification of the binding activity of this peptide to HSC-2 and HaCaT cells treated with various concentrations for 30 min was carried out. Cells were washed twice with phosphate-buffered saline, and the peptides were detected using a flow cytometer (FACSCalibur, BD Biosciences, San Jose, CA, USA).

Antitumor activity of the IL-4R α -lytic hybrid peptide in a human tumor xenograft mouse model in vivo. Animal experiments were carried out in accordance with the guidelines of Tsukuba University. HSC-2 cells (5x10⁶) resuspended in 150 μ l of phosphate-buffered saline were inoculated subcutaneously into the flank region of 4- to 6-week-old athymic female nude mice weighing 17-20 g. When the tumors reached 20-60 mm³ in volume, animals were assigned randomly to four groups. Saline (control), the IL-4R α -lytic hybrid peptide (5 or 10 mg/ kg), or the lytic peptide alone (5 mg/kg) was injected intratumorally (50 μ l/injection) three times a week for a total of 9 times. Tumors were measured with a caliper, and the tumor

No.	Age (years)	Gender	Primary site	TNM classification	Differentiation of SCC
1	57	Male	Tongue	T1N0M0	Well
2	40	Male	Tongue	T2N0M0	Moderate
3	47	Female	Tongue	T2N0M0	Well
4	78	Male	Gingiva	T3N2bM0	Well
5	61	Male	Gingiva	T4aN2bM0	Moderate

Table I. Clinicopathological features of the HNSCC patients.

SCC, squamous cell carcinoma.



Figure 1. IL-4R α expression in HNSCC tissue specimens. (A) IL-4R α protein levels in HNSCC tissues. Tissue sections were prepared from 5 patients who underwent radical resection of oral cancer. Cancerous (cancer) and normal (normal) tissue regions were obtained as described in Materials and methods. (B-E) Immunohistochemical analysis of HNSCC samples of patient no. 2. (B) H&E staining. (C-E) Anti-IL-4R α antibody reactivity in normal area (D) and cancer area (E). Scale bars indicate 300 μ m (B and C) and 50 μ m (D and E).

volume (in mm³) was calculated using the following formula: Length x width² x 0.5. All values are expressed as means \pm SD.

Results

Expression of IL-4R α in HNSCC tissue specimens. We first analyzed the expression levels of IL-4R α in HNSCC tissue specimens. Patient characteristics including age, gender, primary site of tumor, TNM classification and differentiation are shown in Table I. Immunoblot analysis and immunohistochemistry were performed to investigate the expression levels of IL-4R α in tissue specimens from HNSCC patients. Immunoblot analysis showed that IL-4R α was expressed in all HNSCC specimens (tongue and gingival carcinomas) but not in the normal tissue specimens from the same patients (Fig. 1A). Similarly, immunohistochemical analysis using anti-IL-4R α antibody showed IL-4R α immunopositivity in the HNSCC cancerous epithelium but not in the normal epithelium in all patients. Fig. 1B-E shows the staining pattern for patient no. 2.

Expression of IL-4R α in cultured HNSCC cell lines. We next investigated the expression levels of IL-4R α in HNSCC cell lines by immunoblot and real-time PCR analyses. Immunoblot and real-time PCR analyses demonstrated that all HNSCC cell lines expressed IL-4R α but HaCaT did not (Fig. 2A). We also examined mRNA expression levels of IL-4R α by real-time PCR analysis (Fig. 2B). Relative expression levels of IL-4R α in HSC-2 and HSC-3 cells were ~4-fold higher than that of HaCaT cells. The lowest expression level of IL-4R α in HNSCC cell lines was found in OSC-19, however, this level was still 2-fold higher than that of HaCaT (Fig. 2B).



Figure 2. IL-4R α expression in cultured HNSCC cell lines. (A) Protein levels of IL-4R α in HNSCC cell lines and HaCaT normal keratinocytes. Specimens were prepared from five HNSCC cell lines: HSC-2, HSC-3, HSC-4, OSC-19 (tongue carcinoma), Ca9-22 (gingival carcinoma) and HaCaT cells. Actin was used as a loading control. (B) Expression levels of IL-4R α mRNA in HNSCC and HaCaT cell lines. Real-time PCR was performed as described in Materials and methods using IL-4R α -specific primers, and GAPDH served as an internal control.



Figure 3. Cytotoxic activity of the IL-4R α -lytic hybrid peptide in HNSCC and HaCaT cell lines *in vitro*. The five HNSCC cell lines (HSC-2, HSC-3, HSC-4, Ca9-22 and OSC-19) and HaCaT cells (3x10³ cells) were cultured with the indicated concentrations of peptides for 72 h, and cytotoxic activity was assessed using WST-8 reagent (•, IL-4R α -lytic hybrid peptide; o, lytic peptide; •, IL-4 binding peptide). Data represent means ± SD (bars) of triplicate determinations.

Cytotoxic activity of the IL-4R α -lytic hybrid peptide in HNSCC cell lines. To assess the *in vitro* cytotoxic activity of the IL-4R α -lytic hybrid peptide in HNSCC and HaCaT cells, the WST assay was performed using HNSCC cell lines treated with the IL-4R α -lytic hybrid peptide, lytic peptide or IL-4 binding peptide. HSC-2, HSC-3, HSC-4 and Ca9-22 cells were sensitive to the IL-4R α -lytic hybrid peptide; the concentration that killed 50% of all cells (IC₅₀) was <5 μ M. The OSC-19 cell line was also sensitive to the IL-4R α -lytic hybrid peptide with an IC₅₀ of <10 μ M. In contrast, optimal cell killing was not induced in HaCaT cells by either the lytic peptide or IL-4 binding peptide or IL-4R α -lytic hybrid peptide (Fig. 3). The cytotoxic activity of the hybrid peptide was strongly enhanced when compared with that of the lytic peptide. The cytotoxic activity of the IL-4R α -lytic hybrid peptide increased 4.0- to 13.2-fold when compared with that of the lytic peptide in the

	IC ₅₀ (µ1	IC ₅₀ ratio	
Cell lines	IL-4Rα-lytic hybrid peptide	Lytic peptide	Lytic/IL-4Rα-lytic hybrid peptide
HNSCC cells			
HSC-2	1.9±0.1	8.2±0.2	4.4
HSC-3	2.3±0.6	17.4±1.2	7.4
HSC-4	2.8±0.8	33.2±6.1	11.9
Ca9-22	2.2±0.7	29.6±5.8	13.2
OSC-19	8.2±5.4	33.3±13.1	4.0
Normal cells			
HaCaT	37.6±15.3	43.4±13.6	1.2

Table II. Cytotoxic activity of each peptide in HNSCC and HaCaT cell lines.

 IC_{50} values (peptide concentration inducing 50% inhibition of control cell growth) are represented as means \pm SD from triplicate determinations.



Figure 4. Binding assay of HSC-2 and HaCaT cells treated with the FITClabeled IL-4 binding peptide. Quantification of the binding activity of this peptide to HSC-2 and HaCaT cells treated with various concentrations for 30 min. IL-4 binding peptide at 0 μ M represents untreated cells used as a control. Fold-change in fluorescence intensity is the extent of binding of the FITC-labeled IL-4 binding peptide to each cell line, relative to the fluorescence intensity values. Data are presented as means ± SD (error bars) from triplicate determinations.

cancer cell lines (Table II). We examined the binding activity of the IL-4 binding peptide labeled with FITC to both HSC-2 and HaCaT cells by flow cytometry, and then found that exposure of HSC-2 to this peptide resulted in the increased binding activity of this peptide in a concentration-dependent manner (Fig. 4). These results suggest that the IL-4R α -lytic hybrid peptide selectively kills cancer cells expressing IL-4R α .

In vivo antitumor activity of the IL-4R α -lytic hybrid peptide in a human HNSCC xenograft mouse model. Following the observation that the IL-4R α -lytic hybrid peptide exhibits a marked cytotoxic effect on HNSCC cells in vitro (Fig. 3), the antitumor activity of the hybrid peptide was assessed in a xenograft model of human HNSCC. HSC-2 cells were inoculated



Figure 5. Antitumor activity of the IL-4R α -lytic hybrid peptide in a xenograft model. (A) HSC-2 cells were inoculated subcutaneously into nude Balb/c mice. Animals received an intratumoral peptide injection or saline alone [\Box , control; lytic peptide (\odot , 5 mg/kg); IL-4R α -lytic hybrid peptide (\odot , 5 mg/kg) or \blacktriangle , 10 mg/kg)] three times per week from day 3 as indicated by the arrows. Data are expressed as mean \pm SD (bars), n=6 animals per group. *P<0.01. (B-E) Images show tumor growth on day 24 after tumor inoculation: (B) saline alone; (C) lytic peptide; (D) IL-4R α -lytic hybrid peptide (5 mg/kg); (E) IL-4R α -lytic hybrid peptide (10 mg/kg).

subcutaneously into athymic nude mice, and the animals were subsequently treated with the IL-4R α -lytic hybrid peptide by intratumoral injection. As shown in Fig. 5, tumors grew aggressively in the control mice injected with saline alone, reaching a volume of >1000 mm³ by day 24. In contrast, mice treated with the IL-4Ra-lytic hybrid peptide showed significant tumor regression at both dosages: mean tumor volumes were 491 mm³ (5 mg/kg) and 283 mm³ (10 mg/kg) on day 24. Moreover, tumors in mice injected with the lytic peptide grew rapidly similar to tumor growth in the control mice with saline alone, reaching a volume of 1104 mm³ (Fig. 5). No other abnormalities, such as loss of appetite and body weight, were observed in mice injected with the IL-4Ra-lytic hybrid peptide (data not shown). Histological analysis also showed no side effects in tissues from the major organs, including the liver and kidney, which were obtained from mice treated with

intratumoral administration of the IL- $4R\alpha$ -lytic hybrid peptide (data not shown). These results demonstrated that the IL- $4R\alpha$ -lytic hybrid peptide exhibited effective antitumor activity in a mouse xenograft model of HNSCC.

Discussion

The main treatment options for patients with HNSCC currently involve surgery, radiotherapy and chemotherapy, alone or in combination. Despite significant advances in HNSCC treatment, survival rates and prognosis have improved only moderately over the years (16). Systemic chemotherapy remains the only effective treatment option, but it is associated with significant rates of toxicity in HNSCC patients, who usually have a high prevalence of co-morbidities and problematic lifestyle habits (17). Ideally, future therapies should act over the short term, to minimize damage to healthy cells and target tumor compartments that have the highest sensitivity.

The concept of a 'magic bullet' proposed by Paul Ehlrich over 100 years ago has led to the search for agents that can selectively target cancer cells (18). Immunotoxins are proteins used to treat cancer and are composed of an antibody fragment linked to a toxin (19). Several disadvantages of these conventional immunotoxins for clinical use include immunogenicity, undesirable toxicity, manufacturing difficulties, short half-lives and neutralizing antibody production (20,21). However, peptides can be produced affordably by chemical synthesis, with a cost comparable to that of producing protein drugs. Moreover, since peptides are easy to produce, a wide variety of candidate peptides combining moieties for targeting and for toxicity can be tested in preclinical settings. We previously linked two functional peptide domains to produce a novel chimerical peptide termed a 'hybrid peptide', which was designed as a bifunctional peptide that binds to receptors or proteins overexpressing in cancer cells and consequently disrupts the cancer membrane (14,22,23). In the present study, we focused on IL-4R α as recent evidence suggests that IL-4R α is preferentially expressed on the surface of a variety of solid tumors including HNSCC (24).

The high degree of antitumor activity of the IL-4R α -lytic hybrid peptide in HNSCC correlated with the expression of IL-4R α *in vitro*. All HNSCC tumor specimens showed specific immunohistochemical staining for IL-4R α , and western blot analysis revealed expression of IL-4R α . However, IL-4R α expression was not observed in normal tissue specimens from the same patients (Fig. 1). These data are consistent with previous reports that HNSCC cells express IL-4R α on their cell surface and confirm that IL-4R α is expressed *in situ* (25,26). These results also indicate that this receptor may be an attractive target for the treatment of HNSCC.

Previous results suggest that IL-4 receptor-targeted cytotoxin may provide an effective therapeutic option for HNSCC (24,26). In the present study, the *in vitro* cytotoxicity of the IL-4R α -lytic hybrid peptide was examined in five HNSCC cell lines (Fig. 3). HSC-2 cells, which showed the highest level of IL-4R α expression in western blot analysis, also showed the highest sensitivity to the IL-4R α -lytic hybrid peptide (Figs. 2 and 3). Normal HaCaT cells with low IL-4R α expression were not sensitive to the IL-4R α -lytic hybrid peptide (Fig. 3). These results suggest that the cytotoxic effect

of the IL-4R α -lytic hybrid peptide correlates well with the level of IL-4R α expression.

In the present study, although the growth rate of HSC-2 was rapid, it was found that intratumoral administration of the hybrid peptide at 10 mg/kg, dramatically inhibited the growth of HSC-2 tumors *in vivo* (Fig. 5). Histological analysis also showed no abnormal changes in the tissues of major organs obtained from the mice injected with the hybrid peptide (data not shown). For clinical use, local injection may be effective for tumors such as HNSCC, and combination with prior chemoradiotherapy should be developed. These observations indicate that abundant IL-4R α expression in HNSCC tumors would facilitate efficient targeting by the IL-4R α -lytic hybrid peptide.

In conclusion, IL-4R α was overexpressed in both tumor specimens from patients with HNSCC and in HNSCC cell lines *in vitro*. The overexpressed IL-4R α on HNSCC cells could be successfully targeted with the IL-4R α -lytic hybrid peptide *in vitro* and *in vivo*. Future investigations using cancer progenitor cells isolated from primary malignant tissues at different stages during cancer progression and metastatic disease may help to identify new biomarkers for the development of more effective diagnostic and prognostic methods and targeted therapies (27). Additional studies should be performed to reveal the antitumor activity of the IL-4R α -hybrid peptide in animal models, and perhaps a phase I clinical trial should be undertaken to study its antitumor activity.

Acknowledgements

We thank Megumi Kawamoto, Kumi Kodama, Nana Ohkubo, Aya Torisawa (Department of Pharmacoepidemiology, Kyoto University), and Airi Ueda (Department of Clinical Sciences, and Molecular Cellular Physiology, Tsukuba University) for technical assistance or advice with the tissue culturing and *in vivo* experiments. This study was supported by Grantsin-Aid for Scientific Research (B) (grant no. 24390449), Young Scientists (A) (grant no. 2368009), and Challenging Exploratory Research (grant no. 23659934) from the Japan Society for the Promotion of Science (JSPS).

References

- Kamangar F, Dores GM and Anderson WF: Patterns of cancer incidence, mortality, and prevalence across five continents: defining priorities to reduce cancer disparities in different geographic regions of the world. J Clin Oncol 24: 2137-2150, 2006.
- Seiwert TY and Cohen EE: State-of-the-art management of locally advanced head and neck cancer. Br J Cancer 92: 1341-1348, 2005.
- Forastiere AA, Goepfert H, Maor M, et al: Concurrent chemotherapy and radiotherapy for organ preservation in advanced laryngeal cancer. N Engl J Med 349: 2091-2098, 2003.
- 4. Dietz A, Boehm A, Mozet C, Wichmann G and Giannis A: Current aspects of targeted therapy in head and neck tumors. Eur Arch Otorhinolaryngol 265 (Suppl 1): S3-S12, 2008.
- Price KA and Cohen EE: Current treatment options for metastatic head and neck cancer. Curr Treat Options Oncol 13: 35-46, 2012.
- Kirby AM, A'Hern RP, D'Ambrosio C, *et al*: Gefitinib (ZD1839, Iressa) as palliative treatment in recurrent or metastatic head and neck cancer. Br J Cancer 94: 631-636, 2006.
- 7. Herbst RS, Arquette M, Shin DM, *et al*: Phase II multicenter study of the epidermal growth factor receptor antibody cetuximab and cisplatin for recurrent and refractory squamous cell carcinoma of the head and neck. J Clin Oncol 23: 5578-5587, 2005.

- 8. Shimamura T, Royal RE, Kioi M, Nakajima A, Husain SR and Puri RK: Interleukin-4 cytotoxin therapy synergizes with gemcitabine in a mouse model of pancreatic ductal adenocarcinoma. Cancer Res 67: 9903-9912, 2007.
- 9. Ishige K, Shoda J, Kawamoto T, et al: Potent in vitro and in vivo antitumor activity of interleukin-4-conjugated Pseudomonas exotoxin against human biliary tract carcinoma. Int J Cancer 123: 2915-2922, 2008.
- 10. Pastan I, Hassan R, Fitzgerald DJ and Kreitman RJ: Immunotoxin therapy of cancer. Nat Rev Cancer 6: 559-565, 2006.
- 11. Weber f, Asher A, Bucholz R, et al: Safty, tolerability, and tumor response of IL4-Pseudomonas exotoxin (NBI-3001) in patients with recurrent malignant glioma. J Neurooncol 64: 125-137, 2003.
- 12. Attia P, Powell DJ Jr, Maker AV, Kreitman RJ, Pastan I and Rosenberg SA: Selective elimination of human regulatory T lymphocytes in vitro with the recombinant immunotoxin LMB-2. J Immunother 29: 208-214, 2006.
- 13. Choudhary S, Mathew M and Verma RS: Therapeutic potential of anticancer immunotoxins. Drug Discov Today 16: 495-503, 2011.
- 14. Kohno M, Horibe T, Haramoto M, et al: A novel hybrid peptide targeting EGFR-expressing cancers. Eur J Cancer 47: 773-783, 2011.
- 15. Garland L, Gitlitz B, Ebbinghaus S, et al: Phase I trial of intravenous IL-4 Pseudomonas exotoxin protein (NBI-3001) in patients with advanced solid tumors that express the IL-4 receptor. J Immunother 28: 376-381, 2005.
- 16. Forastiere A, Koch W, Trotti A and Sidransky D: Head and neck cancer. N Engl J Med 345: 1890-1900, 2001
- 17. Goon PK, Stanley MA, Ebmeyer J, et al: HPV & head and neck cancer: a descriptive update. Head Neck Oncol 1: 36, 2009.

- 18. Bosch F and Rosich L: The contributions of Paul Ehrlich to pharmacology: a tribute on the occasion of the centenary of his Nobel Prize. Pharmacology 82: 171-179, 2008.
- 19. Pastan I, Hassan R, FitzGerald DJ and Kreitman RJ: Immunotoxin treatment of cancer. Annu Rev Med: 221-237, 2007.
- 20. Kreitman RJ: Immunotoxins for targeted cancer therapy (review). AAPS J 8: 532-551, 2006. 21. Li Z, Yu T, Zhao P and Ma J: Immunotoxins and cancer therapy.
- Cell Mol Immunol 2: 106-112, 2005.
- 22. Horibe T, Kohno M, Haramoto M, Ohara K and Kawakami K: Designed hybrid TPR peptide targeting Hsp90 as a novel anticancer agent. J Transl Med 9: 8, 2011.
- 23. Tada N, Horibe T, Haramoto M, Ohara K, Kohno M and Kawakami K: A single replacement of histidine to arginine in EGFR-lytic hybrid peptide demonstrates the improved anticancer activity. Biochem Biophys Res Commun 407: 383-388, 2011.
- 24. Strome SE, Kawakami K, Alejandro D, et al: Interleukin 4 receptor-directed cytotoxin therapy for human head and neck squamous cell carcinoma in animal models. Clin Cancer Res 8: 281-286, 2002.
- 25. Mehrotra R, Varricchio F, Husain SR and Puri RK: Head and neck cancers, but not benign lesions, express interleukin-4 receptors in situ. Oncol Rep 5: 45-48, 1998.
- 26. Kawakami K, Leland P and Puri RK: Structure, function, and targeting of interleukin 4 receptors on human head and neck cancer cells. Cancer Res 60: 2981-2987, 2000.
- 27. Mimeault M, Hauke R, Mehta PP and Batra SK: Recent advances in cancer stem/progenitor cell research: therapeutic implications for overcoming resistance to the most aggressive cancers. J Cell Mol Med 11: 981-1011, 2007.