MicroRNA-329-3p targets MAPK1 to suppress cell proliferation, migration and invasion in cervical cancer

WENFENG LI^{1,2}, JINGJING LIANG^{1,3}, ZHECHAO ZHANG^{1,3}, HONGYAN LOU^{1,3},
LIANG ZHAO^{1,4}, YUNSHENG XU^{1,5} and RONGYING OU^{1,3}

¹Laboratory for Advanced Interdisciplinary Research, Center for Personalized Medicine/Institutes of Translational Medicine, The First Affiliated Hospital of Wenzhou Medical University; Departments of ²Radiation Oncology, ³Gynaecology and Obstetrics, ⁴Division of PET/CT, Department of Radiology, and ⁵Department of Dermatovenereology, The First Affiliated Hospital of Wenzhou Medical University, Wenzhou, Zhejiang 325000, P.R. China

Received September 20, 2016; Accepted November 3, 2016

DOI: 10.3892/or.2017.5555

Abstract. Cervical cancer is the second most common gynecological cancer worldwide and remains as one of the leading causes of cancer-related death among women. Despite great progress in the treatment of cervical cancer, the 5-year overall survival rate for patients with this disease remains unsatisfactory. Over the past decade, an increasing number of studies indicate a central role for microRNAs in the initiation and progression of cervical cancer. microRNA-329-3p (miR-329-3p) has been studied in many types of human cancer; however, the expression level, biological role and the underlying mechanism of miR-329-3p in cervical cancer has not yet been investigated. In the present study, we found that the expression levels of miR-329-3p were reduced in both cervical cancer tissues and cell lines. Low miR-329-3p expression was negatively correlated with histological grade, International Federation of Gynecology and Obstetrics (FIGO) stage, and lymph node metastasis of cervical cancer patients. In addition, upregulation of miR-329-3p suppressed cell proliferation, migration and invasion of cervical cancer. Furthermore, MAPK1 was identified as a direct target gene of miR-329-3p. MAPK1 was significantly upregulated in cervical cancer tissues and was inversely correlated with miR-329-3p expression in the cervical cancer tissues. Silencing of MAPK1 by RNA interference mimicked the effects of miR-329-3p overexpression on cell proliferation, migration and invasion in cervical cancer. Moreover, rescue experiments showed that restoration of the expression of MAPK1 reversed the effects

of miR-329-3p overexpression in cervical cancer cells. Taken together, these findings suggest that miR-329-3p has a critical tumor-suppressive roles by directly targeting MAPK1 in cervical cancer, and it may be investigated as a novel therapeutic target for the treatment of patients with this disease.

Introduction

Cervical cancer, a primary cancer of the uterine cervix, is the second most common gynecological cancer worldwide and remains as one of the leading causes of cancer-related death among women (1). According to the latest estimated global cancer statistics, there are ~530,000 newly diagnosed cervical cancer cases and 275,000 deaths/year (2). More than 80% of cervical cancer cases are diagnosed in developing countries. This is mainly attributed to the unavailability of widespread screening by cervical cytology (3). Cervical cancer is histologically classified into 3 subtypes: squamous cell carcinoma, adenocarcinoma and adenosquamous carcinoma. Squamous cell carcinoma is the most common of these subtypes and accounts for ~85% of the total number of cases (4,5). Previous studies indicate that many risk factors contribute to cervical cancer carcinogenesis and progression, such as early sexual intercourse, promiscuity and infection with high-risk types of human papillomavirus (HPV) (6,7). Currently, the standard treatments for patients with cervical cancer are surgery, radiotherapy and chemotherapy (8,9). Despite great progress in the treatment of cervical cancer, the 5-year overall survival rate for patients with this disease remains unsatisfactory (10). Therefore, a full understanding of the molecular mechanisms underlying the occurrence and development of cervical cancer is important for investigating more effective therapeutic targets for the treatment of this disease.

MicroRNAs (miRNAs) are a class of single-strand, non-coding, endogenous and small RNA molecules consisting of 19-25 nucleotides (11). miRNAs regulate gene expression in a post-transcriptional pattern via its base-pairing with the 3'-untranslated regions (3'UTRs) of their target genes (12). To date, over 1,000 miRNAs have been predicted to exist in the human genome and they regulate thousands of human protein-coding genes (13). miRNAs have been identified as

---

Correspondence to: Professor Rongying Ou, Laboratory for Advanced Interdisciplinary Research, Center for Personalized Medicine/Institutes of Translational Medicine, The First Affiliated Hospital of Wenzhou Medical University, Shangcai, Nanbaixiang, Ouhai, Wenzhou, Zhejiang 325000, P.R. China
E-mail: tantang846067@163.com

Key words: microRNA-329-3p, cervical cancer, MAPK1, proliferation, migration, invasion, target

regulators of many physiological and pathological processes, including cell proliferation, differentiation, angiogenesis, morphogenesis, apoptosis, metastasis, migration and invasion (14). More than half of miRNAs are located in fragile sites and genomic regions that frequently exhibit abnormal expression in human cancer (15). Over the past decade, an increasing number of studies indicate a central role for miRNAs in tumorigenesis and tumor progression (16-18). The aberrant overexpression of miRNAs can act as oncogenes by negatively regulating tumor-suppressor genes, whereas lowly expressed miRNAs can function as tumor suppressors via directly targeting oncogenes (19). Therefore, it may be beneficial to identify novel miRNAs to serve as therapeutic targets in human cancer.

Although miR-329-3p has been reported to be frequently dysregulated in various types of tumors (20-22), there is no information available concerning miR-329-3p in cervical cancer. The aim of the present study was to elucidate the expression and effects of miR-329-3p in cervical cancer, and to investigate its underlying mechanisms.

Materials and methods

Tissue samples. The present study was approved by the Ethics Committee of The First Affiliated Hospital of Wenzhou Medical University (Wenzhou, China), and written informed consent was obtained from all patients. Cervical cancer and paired adjacent normal cervical tissues were collected from 53 cervical cancer patients who were treated with surgical operation between February 2011 and November 2014 at the Department of Gynaecology and Obstetrics, The First Affiliated Hospital of Wenzhou Medical University. None of these patients had received radiotherapy or chemotherapy prior to surgery. All fresh tissues were immediately snap-frozen in liquid nitrogen and stored at -80°C until use.

Cell lines and culture conditions. The human cervical cancer cell lines (HeLa, C33A, Caski and SiHa), an immortalized HPV-negative skin keratinocyte line (HaCaT) and the HEK293T cell line were purchased from the Shanghai Institute of Biochemistry and Cell Biology (Shanghai, China). All cell lines were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) (both from Gibco, Grand Island, NY, USA) in a humidified atmosphere containing 5% CO₂ and 100% humidity at 37°C.

RNA isolation and quantitative reverse-transcription polymerase chain reaction (RT-qPCR). Total RNA was extracted from tissues and cells using TRIzol (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. Reverse transcription was performed using M-MLV reverse transcriptase (Promega, Madison, WI, USA). Detection and quantitation of miR-329-3p and MAPK1 mRNA were performed using SYBR Premix Ex Taq™ kits (Takara, Tokyo, Japan) on Applied Biosystems® 7900HT Real-Time PCR system (Thermo Fisher Scientific, Waltham, MA, USA). U6 snRNA and GAPDH were used as reference genes for miR-329-3p and MAPK1 mRNA expression, respectively. The relative expression levels of miR-329-3p and MAPK1 mRNA were analyzed using the 2^{-ΔΔCt} method.

Transfection. The miR-329-3p and corresponding negative control mimics (miR-NC) were obtained from GenePharma (Shanghai, China). Small interfering RNA targeting MAPK1 (si-MAPK1) and its negative control (si-NC) were purchased from Ambion (Shanghai, China). The overexpression plasmid of MAPK1 (pCDNA3.1-MAPK1) and blank plasmid (pCDNA3.1) were synthesized at the Chinese Academy of Sciences (Changchun, China). For transfection, the cells were seeded in a 6-well plate until reaching 50-60% confluency. The following day, the cells were transfected with the mimics, siRNA or plasmid using Lipofectamine™ 2000 reagent (Thermo Fisher Scientific) following the manufacturer's instructions.

Cell Counting Kit-8 (CCK-8) assay. Cell proliferation was evaluated using the CCK-8 assay (Dojindo, Kumamoto, Japan). Briefly, the transfected cells were harvested, suspended and seeded in 96-well plates at a density of 3,000 cells/well. Cells were incubated in a humidified atmosphere containing 5% CO₂ and 100% humidity at 37°C for 4 consecutive days after seeding. At each time point, 10 μl CCK-8 solution was added in each well for another 4-h incubation at 37°C. Finally, the absorbance was determined at a wavelength of 450 nm using a microplate reader (Infinite® M1000 PRO; Tecan, Männedorf, Switzerland).

Migration and invasion assays. Migration assays were performed using Transwell chambers (8-μm; BD Biosciences, Franklin Lakes, NJ, USA). After transfection for 48 h, the cells were trypsinized, washed with PBS and re-suspended in FBS-free DMEM. Then, 5x10⁴ cells were seeded in the upper part of each Transwell chamber, while the lower part of each Transwell chamber was filled with 600 μl DMEM containing 20% FBS. After incubation for 48 h in a humidified atmosphere containing 5% CO₂ and 100% humidity at 37°C, the cells migrating to the bottom of the Transwell membrane were fixed with 100% methanol, stained with 0.5% crystal violet solution, dried in air and photographed under a microscope (Olympus, Tokyo, Japan). Invasion assays were carried out in a similar manner but by allowing the cells to migrate through Matrigel (BD Biosciences, San Jose, CA, USA)-coated Transwell chambers.

Bioinformatic prediction. TargetScan Human 7.0 (<http://www.targetscan.org/>) and miRanda (<http://www.microrna.org/microrna/>) were used to identify the potential target genes of miR-329-3p.

Luciferase reporter assay. To explore whether MAPK1 was a direct target gene of miR-329-3p, luciferase reporter assay was performed. For the luciferase reporter assay, luciferase reporter plasmids (pmirGLO-MAPK1-3'UTR Wt and pmirGLO-MAPK1-3'UTR Mut) were synthesized and purified by GenePharma. HEK293T cells were seeded in triplicate in 24-well plates. After incubation overnight, the cells were transfected with luciferase reporter plasmids, along with miR-329-3p mimics or miR-NC using Lipofectamine 2000. Transfected cells were collected 48 h post-transfection, and luciferase activities were detected using Dual-Luciferase Reporter Assays (Promega, Mannheim, Germany) following

Figure 1. miR-329-3p is downregulated in cervical cancer. (A) miR-329-3p expression in cervical cancer and paired adjacent normal cervical tissues was assessed using RT-qPCR. (B) Expression levels of miR-329-3p were determined in cervical cancer cell lines (HeLa, C33A, Caski and SiHa) and an immortalized HPV-negative skin keratinocyte cell line (HaCaT); *P<0.05 compared with the respective control.

manufacturer's procedures. *Renilla* luciferase activities were measured as a control.

Western blotting. For western blotting, total protein was extracted from tissues and cells using RIPA buffer (150 mM NaCl, 1% NP-40, 0.5% deoxycholate and 1% SDS) supplemented with proteinase and phosphatase inhibitors (Roche, Basel, Switzerland). The protein concentration was determined using the BCA protein assay kit (Thermo Fisher Scientific). Equal amounts of proteins were separated using 10% SDS polyacrylamide gels. The separated proteins were electrophoretically transferred to polyvinylidene difluoride (PVDF) membranes (Millipore, Billerica, MA, USA) and blocked in Tris-buffered saline (TBS) containing 0.1% Tween-20 (TBST) containing 5% skimmed milk at room temperature for 1 h. Then, the PVDF membranes were incubated with primary antibodies at 4°C overnight. Next, the membranes were washed with TBST 3 times and probed with the corresponding horseradish peroxidase (HRP)-conjugated secondary antibody (1:5,000 dilution; Santa Cruz Biotechnology, Santa Cruz, CA, USA) at room temperature for 1 h. Finally, the protein bands were visualized using ECL chemiluminescence reagents (Amersham Biosciences Corp., Piscataway, NJ, USA). Primary antibodies used in the present study included, mouse anti-human monoclonal MAPK1 antibody (1:1,000 dilution; sc-81459) and mouse anti-human monoclonal GAPDH antibody (1:1,000 dilution; sc-137179) (both from Santa Cruz Biotechnology). GAPDH was used as an internal control.

Table I. Correlation of miR-329-3p expression with the clinicopathological characteristics of the cervical cancer patients.

Clinicopathological characteristics	Cases	miR-329-3p expression		P-value
		Low	High	
Age (years)				0.269
<60	20	9	11	
≥60	33	20	13	
Tumor size (cm)				0.200
<4	25	16	9	
≥4	28	13	15	
Family history of cancer				0.504
Yes	11	7	4	
No	42	22	20	
Histological grade				0.037
Well/moderate	27	11	16	
Poor	26	18	8	
FIGO stage				0.024
I-II	22	8	14	
III-IV	31	21	10	
Lymph node metastasis				0.007
No	31	12	19	
Yes	22	17	5	
Distant metastasis				0.707
No	22	13	12	
Yes	31	16	12	

FIGO, International Federation of Gynecology and Obstetrics.

Statistical analysis. Data are expressed as mean ± SD, and the Student's t-test was used to compare differences between two groups. P-value of <0.05 was considered to indicate a statistically significant result. All analyses were carried out using SPSS version 13.0 software (SPSS, Inc., Chicago, IL, USA).

Results

miR-329-3p is downregulated in cervical cancer and negatively correlates with clinicopathological characteristics of the cervical cancer patients. To investigate whether or not miR-329-3p is abnormally expressed in cervical cancer, we analyzed its expression in cervical cancer and paired adjacent normal cervical tissues using RT-qPCR. The results showed that miR-329-3p was significantly downregulated in the cervical cancer tissues compared with that in the paired adjacent normal cervical tissues (Fig. 1A; P<0.05).

We next analyzed the correlation between miR-329-3p expression levels and clinicopathological characteristics of the cervical cancer patients. The correlations between miR-329-3p expression levels and the clinicopathological characteristics of the cervical cancer patients are shown in Table I. The

results showed that miR-329-3p was inversely correlated with histological grade ($P=0.037$), International Federation of Gynecology and Obstetrics (FIGO) stage ($P=0.024$) and lymph node metastasis ($P=0.007$). However, there were no significant association between miR-329-3p expression and age ($P=0.269$), tumor size ($P=0.200$), family history of cancer ($P=0.504$) and distant metastasis ($P=0.707$).

Further experiments were carried out using an immortalized HPV-negative skin keratinocyte line (HaCaT) and 4 cervical cancer cell lines to confirm that expression levels of miR-329-3p were reduced in cervical cancer cell lines, including HeLa, C33A, Caski and SiHa cells in comparison with HaCaT (Fig. 1B; $P<0.05$). Taken together, these results indicated that miR-329-3p was lowly expressed in the cervical cancer tissues and cell lines.

Upregulation of miR-329-3p inhibits cell proliferation, migration and invasion of cervical cancer. To further explore the roles of miR-329-3p in cervical cancer, miR-329-3p mimics were used to increase its expression in HeLa and SiHa cells (Fig. 2A; $P<0.05$). CCK-8, and migration and invasion assays were performed to test the effects of miR-329-3p overexpression on cell proliferation, migration and invasion of cervical cancer, respectively. As shown in Fig. 2B, upregulation of miR-329-3p obviously inhibited the proliferation of HeLa and SiHa cells. The results of the migration and invasion assays showed that the migration and invasion capacities of the HeLa and SiHa cells were reduced when cells were transfected with miR-329-3p mimics (Fig. 2C; $P<0.05$). These results indicated that miR-329-3p re-expression inhibited cell proliferation, migration and invasion of cervical cancer.

MAPK1 is a direct target of miR-329-3p. To explore the mechanism underlying the tumor-suppressive roles of miR-329-3p in cervical cancer, we next aimed to explore the potential targets of miR-329-3p. Bioinformatic analysis was performed with publicly available algorithms to predict the candidate targets of miR-329-3p. As shown in Fig. 3A, 3'UTR of MAPK1 contains a target sequence for miR-329-3p. Following, a luciferase reporter assay was carried out to further confirm whether MAPK1 is a direct target of miR-329-3p. HEK293T cells were co-transfected with pmirGLO-MAPK1-3'UTR Wt or pmirGLO-MAPK1-3'UTR Mut, and miR-329-3p mimics or miR-NC. Results showed that miR-329-3p overexpression significantly decreased luciferase activities in the HEK293T cells transfected with pmirGLO-MAPK1-3'UTR Wt, but no significant change in cells with pmirGLO-MAPK1-3'UTR Mut were noted (Fig. 3B; $P<0.05$). Moreover, RT-qPCR and western blotting were adopted to determine the regulatory roles of miR-329-3p on MAPK1 expression. As shown in Fig. 3C and D, restoration of the expression of miR-329-3p obviously downregulated MAPK1 expression in the HeLa and SiHa cells at the mRNA ($P<0.05$) and protein ($P<0.05$) levels. Taken together, these results demonstrated that MAPK1 is directly targeted by miR-329-3p.

MAPK1 is upregulated in cervical cancer tissues and inversely correlates with miR-329-3p expression in cervical cancer tissues. The above results indicated that MAPK1 is a direct target of miR-329-3p; therefore, we next analyzed the

Figure 2. Upregulation of miR-329-3p inhibits proliferation, migration and invasion of HeLa and SiHa cells. (A) miR-329-3p was markedly upregulated in HeLa and SiHa cells transfected with the miR-329-3p mimics. (B) miR-329-3p overexpression inhibited the proliferation of HeLa and SiHa cells. (C) Ectopic expression of miR-329-3p significantly decreased the migration and invasion abilities of the HeLa and SiHa cells; * $P<0.05$ compared with the respective control.

expression of MAPK1 in cervical cancer and paired adjacent normal cervical tissues. Results of RT-qPCR revealed that MAPK1 mRNA was significantly upregulated in cervical cancer tissues compared with that noted in the paired adjacent normal cervical tissues (Fig. 4A; $P<0.05$). Moreover, Spearman's correlation analysis showed a negative correlation between miR-329-3p and MAPK1 mRNA expression levels in the cervical cancer tissues (Fig. 4B; $r=-0.5598$; $P<0.001$). Moreover, MAPK1 protein expression in the cervical cancer and paired adjacent normal cervical tissues was determined using western blotting. As shown in Fig. 4C, MAPK1 protein was highly expressed in the cervical cancer tissues when compared with that in the paired adjacent normal cervical tissues ($P<0.05$).

Downregulation of MAPK1 mimics the effects of miR-329-3p on cell proliferation, migration and invasion of cervical

Figure 3. MAPK1 is a direct target of miR-329-3p. (A) The seed sequences of miR-329-3p in the wild-type and mutant 3'UTR of MAPK1. (B) Relative luciferase activities of luciferase reporter plasmids in the HEK293T cells co-transfected with miR-329-3p mimics or miR-NC. (C) RT-qPCR was performed to determine MAPK1 mRNA expression in the HeLa and SiHa cells transfected with miR-329-3p mimics or miR-NC. (D) Western blotting was adopted to determine MAPK1 protein expression in the HeLa and SiHa cells transfected with miR-329-3p mimics or miR-NC; *P<0.05 compared with the respective control.

Figure 4. Correlation between miR-329-3p and MAPK1 mRNA expression level in cervical cancer tissues. (A) RT-qPCR analysis of MAPK1 mRNA expression in cervical cancer tissues and paired adjacent normal cervical tissues. (B) Spearman's correlation analysis of the relationship between miR-329-3p and MAPK1 mRNA expression in cervical cancer tissues. (C) Western blot analysis of MAPK1 protein expression in cervical cancer and paired adjacent normal cervical tissues; *P<0.05 compared with respective control.

cancer. To confirm that the tumor-suppressive roles of miR-329-3p are mediated by downregulation of MAPK1, we investigated the biological roles of MAPK1 in cervical cancer. si-MAPK1 was employed to knock down MAPK1 expression in the HeLa and SiHa cells (Fig. 5A; P<0.05). Results of the CCK-8 assay showed that downregulation of MAPK1 obviously suppressed the proliferation of the HeLa and SiHa cells which was similar to the effect of miR-329-3p overexpression on cell proliferation (Fig. 5B; P<0.05). In addition, the effects of MAPK1 underexpression on migration and invasion of HeLa

and SiHa cells were similar to those induced by miR-329-3p overexpression (Fig. 5C; P<0.05). These results indicated that restoration of the expression of miR-329-3p suppressed cell proliferation, migration and invasion of cervical cancer through downregulation of MAPK1.

Restoration of the expression of MAPK1 reverses the effects of miR-329-3p overexpression on cell proliferation, migration and invasion of cervical cancer. Rescue experiments were performed to further confirm that MAPK1 is a direct and func-

Figure 5. Downregulation of MAPK1 mimics the effects of miR-329-3p overexpression on the proliferation, migration and invasion of HeLa and SiHa cells. (A) Western blot analysis of MAPK1 protein expression in the HeLa and SiHa cells transfected with si-MAPK1 or si-NC. (B) Inhibition of MAPK1 expression suppressed the proliferation of the HeLa and SiHa cells compared with the cells transfected with si-NC. (C) Migration and invasion assays of HeLa and SiHa cells after transfection with si-MAPK1 or si-NC. *P<0.05 compared with the respective control.

tional downstream target of miR-329-3p. pcDNA3.1-MAPK1 was used to increase MAPK1 expression in the HeLa and SiHa cells (Fig. 6A; P<0.05). Notably, restoration of the expression of MAPK1 significantly reversed the inhibition of HeLa and SiHa cell proliferation (Fig. 6B; P<0.05), migration and invasion (Fig. 6C; P<0.05) induced by miR-329-3p overexpression. These results indicated that miR-329-3p targets MAPK1 directly, resulting in inhibition of cell proliferation, migration and invasion of cervical cancer.

Discussion

miRNAs have drawn attention owing to their important regulatory roles in multiple biological processes related to cancer initiation, progression, diagnosis and treatment (23). Recently, an increasing number of studies have reported that miR-329-3p, located on 14q32.31, is aberrantly expressed in various types of cancers and is inversely related with clinicopathological features. For instance, in hepatocellular carcinoma, miR-329-3p was downregulated in tumor tissues and negatively correlated with

Figure 6. Restoration of the expression of MAPK1 rescues the inhibition of cell proliferation, migration and invasion induced by miR-329-3p overexpression in the HeLa and SiHa cells. (A) Western blot analysis of MAPK1 expression in the HeLa and SiHa cells after transfection with pcDNA3.1-MAPK1 or pcDNA3.1. (B) MAPK1 re-expression rescued the inhibition of cell proliferation induced by miR-329-3p in the HeLa and SiHa cells. (C) Upregulation of MAPK1 reversed the tumor-suppressive roles of miR-329-3p on migration and invasion abilities of the HeLa and SiHa cells; *P<0.05 compared with the respective control.

tumor stage and metastasis of patients with hepatocellular carcinoma (20). In osteosarcoma, miR-329-3p expression was lower in tumor tissues and was inversely associated with advanced stages (21). In glioma, miR-329-3p expression was reduced in tumor tissues and cell lines compared with non-neoplastic brain specimens and primary normal human astrocytes, respectively (22). In neuroblastoma, miR-329-3p was downregulated in metastatic tumor tissues compared with that in matched primary tumor tissues (24). Li *et al* showed that expression levels of miR-329-3p were decreased in gastric cancer tissues when compared with the adjacent controls (25). Kang *et al* revealed that miR-329-3p was lowly expressed in breast cancer tissues (26). Moreover, miR-329-3p was lowly expressed in pancreatic (27) and non-small cell lung cancer (28). These findings suggest that miR-329-3p could serve as a prognostic marker and has predictive value for poor prognosis in human cancer.

Accumulated studies have demonstrated that miR-329-3p plays a critical role in the regulation of tumor biological behaviors. Xiao *et al* reported that upregulation of miR-329-3p blocked G1/S phase transition, inhibited cell proliferation and the capacity of colony formation in glioma by directly targeting E2F1 (22). Wang *et al* found that miR-329-3p targets CD146 to suppress angiogenesis (29). Yang *et al* showed that restoration of the expression of miR-329-3p decreased cell proliferation, colony formation, migration and invasion of neuroblastoma via blockade of KDM1A (24). In gastric cancer, ectopic of miR-329-3p was found to suppress cell proliferation, migration and invasion *in vitro* through downregulation of TIAM1 (25). Liang *et al* demonstrated that miR-329-3p overexpression inhibited cellular proliferation, migration and invasion, and enhanced apoptosis of pituitary tumor by targeting PTTG1 (30). In breast cancer, restoration of expression of miR-329-3p reduced cell proliferation, migration, invasion *in vitro*, and tumor growth *in vivo* by negatively regulating p130Cas (26). Zhou *et al* indicated that enforced miR-329-3p expression suppressed cell invasion by targeting BRD4, but had no effect on cell proliferation and apoptosis in hepatocellular carcinoma (20). Jiang *et al* found that miR-329-3p re-expression suppressed cell proliferation, enhanced apoptosis, G0/G1 cell cycle arrest and decreased wound-healing and migration ability in osteosarcoma by downregulation of Rab10 (21). These findings suggest that miR-329-3p plays vital roles in human cancer and may therefore be investigated as a novel therapeutic target for antitumor treatment.

To date, several target genes of miR-329-3p have been validated, such as E2F1 (22), CD146 (29), KDM1A (24), TIAM1 (25) and PTTG1 (30). To explore the molecular mechanism underlying the suppression of cervical cancer cell growth and metastasis induced by miR-329-3p, we further predicted another miR-329-3p target. In the present study, MAPK1 was identified as a novel direct target gene of miR-329-3p. There are several lines of evidence to support this. Firstly, bioinformatic analysis predicated that MAPK1 is a theoretical target of miR-329-3p. This hypothesis was further confirmed by luciferase reporter assay. RT-qPCR and western blotting showed that MAPK1 expression at both the mRNA and protein levels was significantly downregulated in cervical cancer after transfection with miR-329-3p mimics. In addition, MAPK1 was upregulated in cervical cancer tissues and was inversely correlated with miR-329-3p expression in cervical cancer tissues. Silencing of MAPK1 by RNA interference mimicked the effects of miR-329-3p on cell proliferation, migration and invasion of cervical cancer. Moreover, rescue experiments showed that restoration of the expression of MAPK1 reversed the effects of miR-329-3p overexpression in cervical cancer cells. These results suggest that miR-329-3p exerts a tumor-suppressive role in cervical cancer, at least in part, by targeting MAPK1.

The mitogen activated protein kinase (MAPK) signaling cascade are membrane-to-nucleus signaling modules and play important roles in multiple physiological processes (31). MAPK1, a member of the MAPKs, is a well-known oncogene and is significantly upregulated in various types of human cancer, such as ovarian cancer (32), sacral chordoma (33), non-small cell lung cancer (34), myeloma (35) and gastric cancer (36). In cervical cancer, research has shown that MAPK1

is highly expressed in tumor tissues (37). Inhibition of MAPK1 by RNA interference suppressed cell proliferation, invasion, metastasis and induced apoptosis of cervical cancer (37-39). Consistent with the above observation, we found that MAPK1 was highly expressed in cervical cancer tissues. MAPK1 knockdown significantly suppressed cell proliferation, migration and invasion of cervical cancer, suggesting the oncogenic role of MAPK1 in cervical cancer. Therefore, MAPK1 could be a promising therapeutic target for the treatment of patients with cervical cancer.

In conclusion, we found that miR-329-3p is lowly expressed in cervical cancer and is inversely correlated with histological grade, FIGO stage and lymph node metastasis of cervical cancer patients. Functional studies showed that miR-329-3p inhibited cervical cancer growth and metastasis by directly targeting MAPK1. Therefore, miR-329-3p/MAPK1-based targeted therapy may be an effective therapeutic strategy for patients with cervical cancer.

Acknowledgements

The present study was supported by the National Natural Science Foundation of China (nos. 81571395, 81371748 and 81373075).

References

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D and Bray F: Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136: E359-E386, 2015.
2. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J and Jemal A: Global cancer statistics, 2012. *CA Cancer J Clin* 65: 87-108, 2015.
3. Ferlay J, Shin HR, Bray F, Forman D, Mathers C and Parkin DM: Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 127: 2893-2917, 2010.
4. Shi TY, Chen XJ, Zhu ML, Wang MY, He J, Yu KD, Shao ZM, Sun MH, Zhou XY, Cheng X, *et al*: A *pri-miR-218* variant and risk of cervical carcinoma in Chinese women. *BMC Cancer* 13: 19, 2013.
5. Zheng W, Liu Z, Zhang W and Hu X: miR-31 functions as an oncogene in cervical cancer. *Arch Gynecol Obstet* 292: 1083-1089, 2015.
6. Bosch FX and de Sanjosé S: Chapter 1: Human papillomavirus and cervical cancer - burden and assessment of causality. *J Natl Cancer Inst Monogr* 2003: 3-13, 2003.
7. Yu Y, Zhang Y and Zhang S: MicroRNA-92 regulates cervical tumorigenesis and its expression is upregulated by human papillomavirus-16 E6 in cervical cancer cells. *Oncol Lett* 6: 468-474, 2013.
8. Yee GP, de Souza P and Khachigian LM: Current and potential treatments for cervical cancer. *Curr Cancer Drug Targets* 13: 205-220, 2013.
9. Wang F, Liu M, Li X and Tang H: MiR-214 reduces cell survival and enhances cisplatin-induced cytotoxicity via down-regulation of Bcl2l2 in cervical cancer cells. *FEBS Lett* 587: 488-495, 2013.
10. Du J, Wang L, Li C, Yang H, Li Y, Hu H, Li H and Zhang Z: MicroRNA-221 targets PTEN to reduce the sensitivity of cervical cancer cells to gefitinib through the PI3K/Akt signaling pathway. *Tumour Biol* 37: 3939-3947, 2016.
11. Lagos-Quintana M, Rauhut R, Lendeckel W and Tuschl T: Identification of novel genes coding for small expressed RNAs. *Science* 294: 853-858, 2001.
12. Ambros V: The functions of animal microRNAs. *Nature* 431: 350-355, 2004.
13. Bartel DP: MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell* 116: 281-297, 2004.
14. Hwang HW and Mendell JT: MicroRNAs in cell proliferation, cell death, and tumorigenesis. *Br J Cancer* 96 (Suppl): R40-R44, 2007.

15. Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, Yendamuri S, Shimizu M, Rattan S, Bullrich F, Negrini M, *et al*: Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci USA* 101: 2999-3004, 2004.
16. McManus MT: MicroRNAs and cancer. *Semin Cancer Biol* 13: 253-258, 2003.
17. Zhu J, Zheng Z, Wang J, Sun J, Wang P, Cheng X, Fu L, Zhang L, Wang Z and Li Z: Different miRNA expression profiles between human breast cancer tumors and serum. *Front Genet* 5: 149, 2014.
18. He L and Hannon GJ: MicroRNAs: Small RNAs with a big role in gene regulation. *Nat Rev Genet* 5: 522-531, 2004.
19. Esquela-Kerscher A and Slack FJ: Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer* 6: 259-269, 2006.
20. Zhou J, Li W, Guo J, Li G, Chen F and Zhou J: Downregulation of miR-329 promotes cell invasion by regulating BRD4 and predicts poor prognosis in hepatocellular carcinoma. *Tumour Biol* 37: 3561-3569, 2016.
21. Jiang W, Liu J, Xu T and Yu X: MiR-329 suppresses osteosarcoma development by downregulating Rab10. *FEBS Lett* 590: 2973-2981, 2016.
22. Xiao B, Tan L, He B, Liu Z and Xu R: MiRNA-329 targeting E2F1 inhibits cell proliferation in glioma cells. *J Transl Med* 11: 172, 2013.
23. Tsai MM, Wang CS, Tsai CY, Huang HW, Chi HC, Lin YH, Lu PH and Lin KH: Potential diagnostic, prognostic and therapeutic targets of microRNAs in human gastric cancer. *Int J Mol Sci* 17: pii: E945, 2016.
24. Yang H, Li Q, Zhao W, Yuan D, Zhao H and Zhou Y: miR-329 suppresses the growth and motility of neuroblastoma by targeting KDM1A. *FEBS Lett* 588: 192-197, 2014.
25. Li Z, Yu X, Wang Y, Shen J, Wu WK, Liang J and Feng F: By downregulating TIAM1 expression, microRNA-329 suppresses gastric cancer invasion and growth. *Oncotarget* 6: 17559-17569, 2015.
26. Kang H, Kim C, Lee H, Rho JG, Seo JW, Nam JW, Song WK, NamSW, Kim W and Lee EK: Downregulation of microRNA-362-3p and microRNA-329 promotes tumor progression in human breast cancer. *Cell Death Differ* 23: 484-495, 2016.
27. Wang X, Lu X, Zhang T, Wen C, Shi M, Tang X, Chen H, Peng C, Li H, Fang Y, *et al*: mir-329 restricts tumor growth by targeting grb2 in pancreatic cancer. *Oncotarget* 7: 21441-21453, 2016.
28. Sun CC, Li SJ, Zhang F, Pan JY, Wang L, Yang CL, Xi YY and Li J: Hsa-miR-329 exerts tumor suppressor function through down-regulation of *MET* in non-small cell lung cancer. *Oncotarget* 7: 21510-21526, 2016.
29. Wang P, Luo Y, Duan H, Xing S, Zhang J, Lu D, Feng J, Yang D, Song L and Yan X: MicroRNA 329 suppresses angiogenesis by targeting CD146. *Mol Cell Biol* 33: 3689-3699, 2013.
30. Liang HQ, Wang RJ, Diao CF, Li JW, Su JL and Zhang S: The PTTG1-targeting miRNAs miR-329, miR-300, miR-381, and miR-655 inhibit pituitary tumor cell tumorigenesis and are involved in a p53/PTTG1 regulation feedback loop. *Oncotarget* 6: 29413-29427, 2015.
31. Seger R and Krebs EG: The MAPK signaling cascade. *FASEB J* 9: 726-735, 1995.
32. Yiwei T, Hua H, Hui G, Mao M and Xiang L: HOTAIR interacting with MAPK1 regulates ovarian cancer skov3 cell proliferation, migration, and invasion. *Med Sci Monit* 21: 1856-1863, 2015.
33. Zhang K, Chen H, Zhang B, Sun J, Lu J, Chen K and Yang H: Overexpression of Raf-1 and ERK1/2 in sacral chordoma and association with tumor recurrence. *Int J Clin Exp Pathol* 8: 608-614, 2015.
34. You B, Yang YL, Xu Z, Dai Y, Liu S, Mao JH, Tetsu O, Li H, Jablons DM and You L: Inhibition of ERK1/2 down-regulates the Hippo/YAP signaling pathway in human NSCLC cells. *Oncotarget* 6: 4357-4368, 2015.
35. Tsubaki M, Takeda T, Ogawa N, Sakamoto K, Shimaoka H, Fujita A, Itoh T, Imano M, Ishizaka T, Satou T, *et al*: Overexpression of survivin via activation of ERK1/2, Akt, and NF- κ B plays a central role in vincristine resistance in multiple myeloma cells. *Leuk Res* 39: 445-452, 2015.
36. Fei B and Wu H: MiR-378 inhibits progression of human gastric cancer MGC-803 cells by targeting MAPK1 in vitro. *Oncol Res* 20: 557-564, 2012.
37. Li XW, Tuergan M and Abulizi G: Expression of MAPK1 in cervical cancer and effect of *MAPK1* gene silencing on epithelial-mesenchymal transition, invasion and metastasis. *Asian Pac J Trop Med* 8: 937-943, 2015.
38. Huang C, Liu LY, Li ZF, Wang P, Ni L, Song LP, Xu DH and Song TS: Effects of small interfering RNAs targeting MAPK1 on gene expression profile in HeLa cells as revealed by microarray analysis. *Cell Biol Int* 32: 1081-1090, 2008.
39. Lwin WW, Park K, Wauson M, Gao Q, Finn PW, Perkins D and Khanna A: Systems biology approach to transplant tolerance: Proof of concept experiments using RNA interference (RNAi) to knock down hub genes in Jurkat and HeLa cells in vitro. *J Surg Res* 176: e41-e46, 2012.