

# Associations of *miR-146aC>G*, *miR-149C>T*, *miR-196a2C>T* and *miR-499A>G* polymorphisms with brain tumors

JAEJOON LIM<sup>1\*</sup>, JUNG OH KIM<sup>2\*</sup>, HAN SUNG PARK<sup>2</sup>, IN BO HAN<sup>1</sup>, KYUBUM KWACK<sup>2</sup>,  
NAM KEUN KIM<sup>2</sup> and KYUNGGI CHO<sup>1</sup>

<sup>1</sup>Department of Neurosurgery, CHA University, CHA Bundang Medical Center, Seongnam 134-96;

<sup>2</sup>Department of Biomedical Science, College of Life Science, CHA University, Seongnam 134-88, Republic of Korea

Received January 23, 2018; Accepted July 2, 2018

DOI: 10.3892/or.2018.6557

**Abstract.** MicroRNAs (miRNAs/miRs) are short, non-coding RNAs that are implicated in tumorigenesis, functioning as tumor suppressors and oncogenes. However, the clinical significance of miRNA expression profiles for brain tumors remains unclear. Therefore, the present study was designed to investigate the associations between miRNA genetic variants and brain tumor risk. A total 362 participants were recruited, including 179 who were healthy subjects and 183 who were patients with brain tumors confirmed as gliomas, meningiomas or schwannomas. This study investigated the single nucleotide polymorphisms *miR-146aC>G*, *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G* by polymerase chain reaction-restriction fragment length polymorphism. It was found that the dominant *miR-149* and CC genotypes were significantly more frequent in patients with glioma. The odds ratios for the C-C-C-G, C-T-C-G and G-C-T-G haplotypes (*miR-146aC>G-miR-149T>C-miR-196a2T>C-miR-499A>G*) were significantly increased in glioma, as were the odds ratios for the GCT haplotype of *miR-146aC>G, miR-149T>C* and *miR-196a2T>C*, and for the C-C-G haplotype of *miR-149T>C, miR-196a2T>C* and *miR-499A>G*. In meningioma, the odds ratios were increased in the G-T-C-G haplotype of *miR-146aC>G, miR-149T>C, miR-196a2T>C* and *miR-499A>G*. The odds ratios were also increased in the G-C-G haplotype of *miR-146aC>G, miR-196a2T>C* and *miR-499A>G*, and in the C-C-G haplotype

of *miR-149T>C, miR-196a2T>C* and *miR-499A>G*. The odds ratios for schwannoma were increased in the G-C-T-G haplotype of *miR-146aC>G, miR-149T>C, miR-196a2T>C* and *miR-499A>G*, and in the C-C-G haplotype of *miR-149T>C, miR-196a2T>C* and *miR-499A>G*. In conclusion, these results suggested that the *miR-149* polymorphism may be involved in the development of gliomas, and the C-C-G haplotype of *miR-149T>C, miR-196a2T>C* and *miR-499A>G* showed increased odds ratios for all types of brain tumors.

## Introduction

MicroRNAs (miRNAs/miRs) regulate mRNA expression through RNA interference and are known to be associated with various diseases (1-3). Abnormal miRNA expression is a well-known and crucial factor that is associated with the initiation and progression of various tumors, including brain tumors and breast cancer (1,4,5). The term brain tumor describes an inhomogeneous collection of tumors of the brain, which can be either malignant or benign and either originate in the central nervous system or represent metastases from other tumors (4,6). Among these various types of brain tumors, studies have been conducted on miRNAs in gliomas, meningiomas and schwannomas (6-10). Notably, certain previous studies have focused on the expression profiles of miRNAs in cancerous tissues and their associations with target genes involved in cancer cell formation and metastasis.

Abnormal expression of miRNAs in glioma tissues was previously reported, and miRNAs, including *miR-34a*, *146a*, *7*, *128* and *195*, were downregulated in cancer tissues compared with those in normal tissues, suggesting dysregulation of tumor suppressor genes (6). However, the expression of certain miRNAs (*miR-21*, *26a*, *10b*, *30e* and *221/222*) was increased, suggesting that miRNAs act not only as tumor suppressors, but also, dependent on the function of the targeted mRNA, as oncogenes (6,7). The miRNA expression profiling of meningiomas has shown a reduction in *miR-29c* and *miR-219* depending on the tumor grade and has indicated that high expression of *miR-190a* in meningiomas correlates with tumorigenic risk. In addition, the expression pattern of *miR-190a* is a prognostic predictor of postoperative outcome (8). In addition, the expression patterns of miRNAs, including *miR-200a* and *miR-145*, are reportedly associated with the progression of meningiomas (9).

**Correspondence to:** Professor Nam Keun Kim, Department of Biomedical Science, College of Life Science, CHA University, 335 Pangyo-ro, Bundang, Seongnam 134-88, Republic of Korea  
E-mail: nkKim@cha.ac.kr

Professor Kyunggi Cho, Department of Neurosurgery, Bundang CHA Medical Center, CHA University College of Medicine, Yatap-ro 59, Seongnam 134-96, Republic of Korea  
E-mail: sandori50@gmail.com

\*Contributed equally

**Key words:** microRNA, single nucleotide polymorphism, case-control study, brain tumor, glioma, meningioma, schwannoma

A schwannoma study also showed altered miRNA expression patterns in tumor tissues, with 8 miRNAs showing increased expression and 4 miRNAs showing decreased expression in tumor tissues compared with that in normal tissues (10).

Numerous studies have found a correlation between *miR-146a*, *149*, *196a2* and *499* and tumorigenesis and tumor suppression. These miRNAs are associated with tumor initiation, invasion, metastasis and proliferation through a variety of mechanisms, and can also function as tumor suppressor genes (4,11-13). Thus, previous studies of miRNA have focused on miRNA regulation and its influence on tumorigenesis. Nevertheless, the reasons for the widespread differential expression of miRNAs in malignant cells compared with that in normal cells are not fully elucidated.

Previous studies have reported miRNA polymorphisms in various cancer types. miRNA single nucleotide polymorphisms (SNPs) have been found to be associated with carcinogenesis, progression, development and prognosis (14,15). SNPs are known to affect miRNA expression and maturation (16). *miR-146aC>G*, *miR-149C>T*, *miR-196a2C>T* and *miR-499A>G* are well-known miRNA SNPs, and among these, three variants (*miR-146aC>G*, *miR-196a2C>T* and *miR-499A>G*) are located in the mature form of the sequence, and the remaining variant (*miR-149C>T*) is located in the precursor form of the sequence. A previous study (14) reported that increased expression of the *miR-196a2C>T* polymorphism was associated with a decreased survival rate in patients with lung cancer. In addition, other studies have reported significant associations between the *miR-196a2C>T* polymorphism and the prevalence of various cancer types and heart disease (17-22). Furthermore, previous studies have suggested that *miR-146aC>G*, *miR-149C>T* and *miR-499A>G* variants are associated with heart disease and esophageal cancer (18,19,22).

Recently, reports have found an association between miRNAs and brain tumors. However, the function of miRNA polymorphisms in brain tumors is unclear in terms of the pathogenesis, and the majority of studies have been limited to specific tumors. The results of these studies have been inconsistent (11,14-24). Thus, the present study sought to investigate the frequency of four miRNA polymorphisms (*146aC>G*, *149C>T*, *196aC>T* and *499A>G*) in brain tumors, specifically gliomas, meningiomas and schwannomas.

## Materials and methods

**Study population.** A total of 362 patients were enrolled, including 179 with brain tumors and 183 healthy controls. Patients with brain tumors were diagnosed through imaging examinations, including computed tomography and magnetic resonance imaging, and were confirmed to have gliomas, meningiomas or schwannomas through histopathological studies following surgical resection. Adults who underwent a general health care examination at CHA Bundang Hospital (Seongnam, South Korea) and who had no history of tumors, brain diseases, including Alzheimer's disease, dementia, stroke or intracerebral hemorrhage, or other underlying diseases were selected for the control group. All study participants were fully informed and received an explanation of the study, and provided written informed consent; individuals who did not agree to the study were excluded.

**Genetic analysis.** Genomic DNA was extracted using the G-DEX blood kit (iNtRON Biotechnology, Inc., Seongnam, South Korea) from anticoagulated peripheral blood, as previously described (25). The four miRNA polymorphisms of interest (*miR-146a*, *149*, *196a2* and *499*) were identified by a literature search using the key words 'miRNA' and 'brain tumor' in Pubmed (<https://www.ncbi.nlm.nih.gov/pubmed>). The genotype examination conditions outlined in our previous study protocol were used (25). Samples underwent re-genotyping examination by an additional operator for confirmation. In addition, 20% of the total samples were randomly selected and the four miRNA polymorphisms were confirmed by sequencing (ABI3730xl DNA analyzer; Applied Biosystem; Thermo Fisher Scientific, Inc., Waltham, MA, USA). PCR was performed with conditions and primer and probe sequences as detailed in our previous study (25).

**Statistical analysis.** The genotype and allele frequencies for miRNA polymorphisms in the patients with brain tumors and the controls were determined. The differences in the genotypes and allele frequencies were analyzed using the  $\chi^2$  test and Fisher's exact test, respectively. Allelic frequencies included calculated deviations based on the Hardy-Weinberg equilibrium, using  $P<0.05$  as a threshold (26). As the inferences of the present study were derived from multiple tests, the Benjamini and Hochberg strategy was adopted, which effectively reduced the potential impact of spurious significant results (27). Statistical analyses were performed that measured the efficacy of the association between brain tumor and genotype based on multivariable logistic regression and according to the statistical methods discussed by Kim and Hong (28). The multifactor dimensionality reduction method has been described in detail previously (29-33). In addition, all possible allele-allelic combinations were performed using HAPSTAT software (v.3.0; [www.bios.unc.edu/~lin/hapstat/](http://www.bios.unc.edu/~lin/hapstat/)). Survival analysis estimated the adjusted hazard ratios (HRs) and their 95% confidence intervals (CIs), with adjustment for age by multivariate Cox proportion hazards regression.

## Results

**Participant characteristics.** A total of 362 participants whose aged from 21 to 85 years were enrolled in the present study, including 183 healthy controls (age range, 24-85 years) and 179 patients (age range, 21-78 years) with brain tumors. In the brain tumor group, 79 patients had gliomas, 69 had meningiomas and 31 had schwannomas. The male:female ratio for all participants was 1:1.48 (1:1.88 for the control group and 1:1.16 for the brain tumor group). The mean age of all participants was  $48.8\pm15.9$  years (control group mean,  $45.9\pm16.6$  years; brain tumor group mean,  $51.9\pm14.7$  years) (Table I).

**Genetic analysis.** The *miR-146a* rs2910164, *miR-149* rs4846049, *miR-196a2* rs11614913 and *miR-499* rs3746444 polymorphisms were compared between the brain tumor and control groups. Participant genotype and allelic frequencies of the four miRNA polymorphisms are detailed in Table II. Analysis by tumor type found that the frequencies of the dominant *miR-149* genotype [odds ratio (OR), 1.842; 95% CI, 1.074-3.159;  $P=0.02$ ] and CC type of *miR-149* (OR, 2.771; 95% CI, 1.158-6.635;




Figure 1. Survival analysis for the *miR-196a2* polymorphism in patients with brain tumors. OS was analyzed by Cox proportional-hazards regression for the patients with brain tumors based on *miR-196a2* genotype. (A) OS curve of patients with *miR-196a2* TT vs. TC genotype. (B) OS curve of patients with *miR-196a2* TT vs. TC+CC genotype. miRNA/miR, microRNA; HR, hazard ratio; OS, overall survival.

$P=0.02$ ) were significantly increased compared with those of the TT and TC genotypes for gliomas. The frequencies of the *miR-146a*, *miR-149*, *miR-196a2* and *miR-499* genotypes were not significantly different between the control group and the meningioma or schwannoma patient groups.

**Allele combination analysis according to subgroup.** ORs for gliomas, meningiomas and schwannomas are detailed in Tables III, IV and V, respectively. The ORs for glioma were increased in the following combinations: i) CCGG, CTCG and GCTG haplotypes of *miR-146aC>G*, *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G*; ii) GCT haplotype of *miR-146aC>G*, *miR-149T>C* and *miR-196a2T>C*; iii) CCG haplotype of *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G*; iv) GC haplotype of *miR-146aC>G* and *miR-149T>C*; and v) CA haplotype of *miR-149T>C* and *miR-499A>G* (Table III). For meningiomas, the ORs were increased in the following combinations: i) GTCG haplotype of *miR-146aC>G*, *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G*;

ii) GCG haplotype of *miR-146aC>G*, *miR-196a2T>C* and *miR-499A>G*; and iii) CCG haplotype of *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G* (Table IV). The ORs for schwannomas were increased in the following combinations: i) GCTG haplotype of *miR-146aC>G*, *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G*; and ii) CCG haplotype of *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G* (Table V).

**Multivariate survival analysis according to genotypes.** Fig. 1 shows the association between the overall survival (OS) of patients with brain tumors and the four pre-miRNA SNPs. No significant association was found between OS and *miR-146a*, *miR-149* or *miR-499*; however, *miR-196a2T>C* was associated with a significantly shorter OS time (adjusted HR, 2.094; 95% CI, 1.147-3.823;  $P=0.017$ ; Fig. 1). Additionally, when OS was analyzed according to the dominant genotype (TT vs. TC+CC), a poorer OS time was associated with the *miR-196a2* C allele (HR, 1.809; 95% CI, 1.043-3.137;  $P=0.036$ ; Fig. 1).

Table I. Demographic characteristics of patients with brain tumor and control subjects.

Characteristics	Control (n=183)	Brain tumor (n=179)
Sex (male:female)	1:1.88	1:1.16
Age, years (mean $\pm$ SD)	45.9 $\pm$ 16.6	51.9 $\pm$ 14.7
Hypertension, n	-	43
Diabetes mellitus, n	-	21
FBS, mg/dl (mean $\pm$ SD)	-	168.0 $\pm$ 70.8
Dyslipidemia, n	-	21
T. chol, mg/dl (mean $\pm$ SD)	-	197.9 $\pm$ 57.3
Triglyceride, mg/dl (mean $\pm$ SD)	-	153.9 $\pm$ 145.5
HDL-C, mg/dl (mean $\pm$ SD)	-	46.4 $\pm$ 16.0
LDL-C, mg/dl (mean $\pm$ SD)	-	106.5 $\pm$ 43.9
BUN, mg/dl (mean $\pm$ SD)	-	22.3 $\pm$ 19.7
Creatinine, mg/dl (mean $\pm$ SD)	-	0.8 $\pm$ 0.5
Platelets, 10 <sup>3</sup> cell/ $\mu$ l (mean $\pm$ SD)	-	346.5 $\pm$ 854.0
Antithrombin, % (mean $\pm$ SD)	-	85.2 $\pm$ 27.5
aPTT, sec (mean $\pm$ SD)	-	34.2 $\pm$ 21.5
Prothrombin time, sec (mean $\pm$ SD)	-	12.5 $\pm$ 2.7
D-dimer, ng/ml (mean $\pm$ SD)	-	2925.8 $\pm$ 2907.5
Fibrinogen, mg/dl (mean $\pm$ SD)	-	477.8 $\pm$ 195.9
Hematocrit, % (mean $\pm$ SD)	-	28.9 $\pm$ 6.1
Hemoglobin, mg/dl (mean $\pm$ SD)	-	9.5 $\pm$ 2.6

FBS, fasting blood sugar; T. chol, total cholesterol; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; BUN, blood urea nitrogen; aPTT, activated partial thromboplastin time; SD, standard deviation.

## Discussion

Brain tumors can cause severe impairment and impose a decreased quality of life, often resulting in mortality (34,35). The overall incidence rate of all brain tumors is 10.82 per 100,000 person-years (36). In the 2013 Central Brain Tumor Registry of the United States report, the average annual age-adjusted incidence rate of glioblastoma was 3.19 per 100,000 individuals, which was the highest incidence among all types of brain and central nervous system (CNS) tumors. Meningioma is the second most common type of brain tumor (37). Sex and age standardized incidence rates range from 1.28 per 100,000 individuals to 7.80 per 100,000 individuals for cerebral meningioma. The overall incidence rate of schwannomas was 1.2 per 100,000 individuals per year in the United States in the period between 2004 and 2009 (38).

Studies on the pathogenesis of brain tumors are ongoing. For several decades, studies have identified molecular alterations characterizing gliomas and reported decreased expression of tumor suppressor genes, such as retinoblastoma transcriptional corepressor 1 and p53, or alterations of genes in pathways associated with tumor suppressors (39). In addition, genetic variants, including isocitrate dehydrogenase 1 and

phosphatase and tensin homolog (PTEN), have been reported in gliomas (40-42). Chromosomal anomalies, aberrant cellular pathways and alterations in tumor suppressor genes are associated with meningioma pathogenesis (43-46). Schwannomas are directly associated with genetic changes in neurofibromin 2 gene inactivation (47). However, no clear mechanism of the pathogenesis for all brain tumors has been identified.

miRNAs regulate target gene expression at the post-transcriptional level. miRNAs control key physiological processes, including cell growth, differentiation and apoptosis, which suggests that miRNA gene abnormality could be involved in tumorigenesis (1-3,48). The correlation between cancer and miRNAs was established in 2002 (49). Certain miRNA abnormalities have been associated with cancer types, and with carcinogenesis and progression (1,3-5). The results reported in the present study suggest that identifying miRNAs and their targets may provide potential diagnostic and prognostic tumor biomarkers and novel cancer therapeutic strategies. Studies of miRNAs associated with brain tumors are ongoing. miRNAs are associated with regulation of tumorigenic cells in CNS tumors, and certain miRNAs are oncogenes (4,50,51). miRNAs also regulate tumor invasion, metastasis and cell apoptosis, and are involved in tumor chemoresistance and radioresistance (52-54).

Recent epidemiological studies have demonstrated that miRNA variants cause altered expression and are associated with the risk of cancer (55). For example, miRNA-146a antagonizes the expression of interleukin (IL)-1 $\beta$ , tumor necrosis factor- $\alpha$  and nuclear factor- $\kappa$ B (56). *miR-146a* is known to be important for tumor proliferation and metastatic ability (23). In addition, certain studies have suggested that miR-196a dysfunction is associated with tumor abnormality (13). *miR-196a2* has a double mature strand containing a 5'-end strand (*hsa-miR-196a2-5p*) and a 3'-end strand (*hsa-miR-196a2-3p*), and the *miR-196a2* rs11614913 T>C polymorphism is located in the *hsa-miR-196a-3p* sequence. The rs11612913 SNP may influence the maturation of *hsa-miR-196a*. In other words, this polymorphism may affect or alter the expression of the target, which may be involved in regulating carcinogenesis. Several studies have identified *miR-499A>G* as a potential marker for a number of cancer types, including breast cancer, gastric cancer, squamous cell carcinoma and hepatocellular carcinoma (11,14,24). The *miR-499* rs3746444 A>G polymorphism is located in the stem loop. To date, few studies have investigated rs2292932 in *miR-149* compared with the three other genes. Li *et al* (12) suggested that *miR-149* inhibits associated fibroblasts by regulating prostaglandin E2 and IL-6 in tumor cells (12).

The present study identified the genotypic distribution of the four most common miRNAs associated with tumors (i.e., *miR-146a*, *miR-149*, *miR-196a2* and *miR-499*) in patients with glioma, meningioma or schwannoma. In this Korean population, the frequencies of the dominant *miR-149* genotype and the CC genotype were significantly increased compared with those of the TT and TC genotypes in the patients with glioma. However, the frequency of the *miR-146a* rs2910164 C>G, *196a2* rs11614913 T>C and *499* rs3746444 A>G polymorphisms were not significantly different between the control group and the tumor group. To date, there have been considerably fewer studies on *miR-149* than on other miRNAs,

Table II. Genotype frequencies of miRNA polymorphisms between patients with three different brain tumor subtypes [glioma (n=79), meningioma (n=69) and schwannoma (n=31)] and control subjects (n=183).

Genotypes	Controls, n (%)	Glioma				Meningioma				Schwannoma						
		n (%)	AOR (95% CI)	P-value <sup>a</sup>	FDR-P-value	Statistical power, %	n (%)	AOR (95% CI)	P-value <sup>a</sup>	FDR-P-value	Statistical power, %	n (%)	AOR (95% CI)	P-value <sup>a</sup>	FDR-P-value	Statistical power, %
miR-146aC>G																
CC	70 (38.3)	30 (38.0)	1.000 (reference)				28 (40.6)	1.000 (reference)				10 (32.3)	1.000 (reference)			
CG	88 (48.1)	34 (43.0)	0.902 (0.503-1.615)	0.727	0.969	5.1	32 (46.4)	0.909 (0.501-1.651)	0.754	0.898	5.7	17 (54.8)	1.352 (0.583-3.138)	0.482	0.749	7.6
GG	25 (13.7)	15 (19.0)	1.400 (0.648-3.023)	0.392	0.467	12.6	9 (13.0)	0.900 (0.374-2.168)	0.814	0.814	4.2	4 (12.9)	1.120 (0.322-3.895)	0.859	0.859	3.8
Dominant (CC vs. CG+GG)			1.012 (0.5881.742)	0.966	0.966	4.1		0.907 (0.515-1.597)	0.735	0.747	5.5		1.301 (0.579-2.924)	0.524	0.726	8.2
Recessive (CC+CG vs. GG)			1.481 (0.733-2.992)	0.273	0.364	17.5		0.948 (0.418-2.148)	0.898	0.994	3.9		0.936 (0.302-2.903)	0.909	0.909	3.0
miR-149T>C																
TT	97 (53.0)	30 (38.0)	1.000 (reference)				35 (50.7)	1.000 (reference)				18 (58.1)	1.000 (reference)			
TC	72 (39.3)	37 (46.8)	1.662 (0.940-2.938)	0.081	0.162	49.3	27 (39.1)	1.039 (0.578-1.870)	0.898	0.898	4.2	11 (35.5)	0.823 (0.366-1.850)	0.638	0.749	6.3
CC	14 (7.7)	12 (15.2)	2.771 (1.158-6.635)	0.022	0.088	67.4	7 (10.1)	1.386 (0.517-3.715)	0.517	0.814	8.7	2 (6.5)	0.770 (0.161-3.681)	0.743	0.859	1.8
Dominant (TT vs. TC+CC)			1.842 (1.074-3.159)	0.026	0.104	68.1		1.096 (0.630-1.907)	0.747	0.747	5.9		0.815 (0.377-1.760)	0.602	0.726	5.9
Recessive (TT+TC vs. CC)			2.162 (0.951-4.916)	0.066	0.264	48.8		1.363 (0.526-3.534)	0.524	0.994	8.6		0.833 (0.180-3.857)	0.815	0.909	1.9
miR-196a2T>C																
TT	46 (25.1)	22 (27.8)	1.000 (reference)				20 (29.0)	1.000 (reference)				10 (32.3)	1.000 (reference)			
TC	92 (50.3)	44 (55.7)	1.000 (0.537-1.863)	1.000	1.000	4.1	32 (46.4)	0.800 (0.413-1.550)	0.508	0.898	8.9	15 (48.4)	0.750 (0.313-1.799)	0.519	0.749	8.9
CC	45 (24.6)	13 (16.5)	0.604 (0.272-1.344)	0.216	0.432	22.6	17 (24.6)	0.869 (0.404-1.869)	0.719	0.814	4.6	6 (19.4)	0.613 (0.206-1.829)	0.380	0.859	9.6
Dominant (TT vs. TC+CC)			0.870 (0.480-1.577)	0.646	0.861	6.4		0.823 (0.443-1.526)	0.536	0.747	8.6		0.705 (0.309-1.607)	0.406	0.726	12.0
Recessive (TT+TC vs. CC)			0.604 (0.305-1.196)	0.148	0.296	27.4		1.003 (0.527-1.907)	0.994	0.994	4.1		0.736 (0.284-1.908)	0.528	0.909	6.1
miR-499A>G																
AA	112 (61.2)	58 (73.4)	1.000 (reference)				44 (63.8)	1.000 (reference)				20 (64.5)	1.000 (reference)			
AG	64 (35.0)	19 (24.1)	0.573 (0.314-1.047)	0.070	0.162	42.1	24 (34.8)	0.955 (0.532-1.713)	0.876	0.898	4.3	10 (32.3)	0.875 (0.386-1.985)	0.749	0.749	4.6

Table II. Continued.

Genotypes	Controls, n (%)	Glioma			Meningioma			Schwannoma		
		AOR (95% CI)	P-value <sup>a</sup>	FDR-P-value	Statistical power, %	n (%)	AOR (95% CI)	P-value <sup>a</sup>	FDR-P-value	Statistical power, %
GG	7 (3.8)	0.552 (0.111-2.741)	0.467	0.467	4.4	1 (1.4)	0.364 (0.044-3.042)	0.351	0.814	3.5
Dominant (AA vs. AG+GG)		0.571 (0.320-1.021)	0.059	0.118	45.2		0.896 (0.505-1.591)	0.708	0.747	5.5
Recessive (AA+AG vs. G)		0.653 (0.133-3.216)	0.600	0.600	2.9		0.370 (0.045-3.062)	0.356	0.994	3.6
								0.839	0.859	1.3
								0.726	0.726	5.4
								0.871	0.909	1.4

<sup>a</sup>Adjusted for age. AOR, adjusted odds ratio; CI, confidence interval; miR/miRNA, microRNA; FDR, false discovery rate.

particularly concerning the association of *miR-149* with brain tumors. One previous study of the association between *miR-149* and brain tumors reported that increased *miR-149* levels downregulate tumor proliferation and metastasis in glioblastoma (57). *miR-149* regulates cell cycle-related genes and controls potential proliferation and invasion activity of glioma cells through a mechanism that induces arrest at the G<sub>0</sub>/G<sub>1</sub> phase (58). However, the manner in which miRNAs affect tumorigenesis or suppression of brain tumors is not yet known, and the results differ by study and participant ethnicity. Analysis of haplotype frequencies according to brain tumor type showed that odds ratios were higher in certain haplotypes in the present study. In particular, the GCTG haplotype of *miR-146aC>G*, *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G* had increased the ORs for gliomas and schwannomas. The CCG haplotype of *miR-149T>C*, *miR-196a2T>C* and *miR-499A>G* showed increased ORs for all three types of brain tumors. However, the mechanism by which certain haplotypes simultaneously increase the risk for various tumors has not been determined.

The *miR-196a2* rs11614913C allele has been reported to be associated with the risk of various cancer types in several studies. Hu *et al* (14) showed that the *miR-196a2* rs11614913C allele was associated with a significant risk of breast cancer. In addition, Tian *et al* (17) demonstrated that the *miR-196a2* rs11614913 CC type posed a significant risk of lung cancer in Chinese individuals. Moreover, several studies suggested the association of the *miR-196a2C* allele with an elevated risk of various cancer types, including hepatoma, gastric cancer and esophageal squamous cell carcinoma (20-22). To date, there have been only two reports on the effects of an association between *miR-196a2* and brain tumor prognosis (59,60). The present study showed that the *miR-196a2T>C* polymorphism was a significant factor for mortality in Korean patients with brain tumors. However, as the survival analysis included the entire brain tumor group, care must be taken when interpreting these results.

The present study had several limitations. First, the way in which miRNA genetic variants affect brain tumor progression remains unclear. Second, in this study, the expression patterns of the four miRNAs in gliomas, meningiomas and schwannomas samples could not be identified as the biopsy samples were in poor condition and the number of tissue samples was insufficient. Therefore, the acquisition of tissue samples is currently being attempted and further studies are being planned. Third, the sample size of this study is limited in number. Lastly, the analysis was performed in a solely Korean population. Although the results present the first evidence for utilization of miRNA polymorphisms as diagnostic and prognostic markers of brain tumor risk, further research in large and diverse cohorts is necessary. Based on the results from this study, a future large-population study is required to identify the association between brain tumors and miRNAs beyond the four miRNAs tested here. In addition, the expression patterns of miRNAs in brain tumors and normal tissues require further study, as does the function of cultured cell lines derived from brain tumors, including gliomas, meningiomas and schwannomas. Additional studies will also be necessary to determine the mechanism by which miRNAs affect carcinogenesis and tumor progression. In conclusion, the

Table III. Allele combination of miRNA polymorphisms between glioma patients (2n=158) and control subjects (2n=366).

Allele combination	Controls, n (%)	Glioma, n (%)	Reference vs. Allele combination			Overall vs. Allele combination		
			OR (95% CI)	P-value <sup>a</sup>	FDR-P-value	OR (95% CI)	P-value <sup>a</sup>	FDR-P-value
miR-146aC>G/miR-149T>C/miR-196a2T>C/miR-499A>G								
C-T-T-A	51 (14.0)	28 (17.8)	1.000 (reference)			1.272 (0.773-2.091)	0.361	0.525
C-T-C-G	26 (7.0)	3 (1.7)	0.210 (0.058-0.757)	0.015	0.045	0.267 (0.080-0.896)	0.021	0.048
C-C-T-G	16 (4.4)	0 (0.0)	0.055 (0.003-0.948)	0.003	0.015	0.070 (0.004-1.176)	0.005	0.020
C-C-C-G	0 (0.0)	4 (2.6)	16.260 (0.844-313.200)	0.020	0.050	20.810 (1.113-389.100)	0.009	0.024
G-T-C-A	53 (14.6)	6 (4.0)	0.206 (0.079-0.540)	0.001	0.015	0.262 (0.110-0.623)	0.001	0.008
G-T-C-G	0 (0.0)	6 (3.8)	23.490 (1.275-432.700)	0.003	0.015	30.060 (1.682-537.200)	0.001	0.008
G-C-T-A	14 (3.8)	17 (10.9)	2.212 (0.951-5.146)	0.085	0.159	2.813 (1.353-5.847)	0.008	0.024
G-C-T-G	0 (0.1)	5 (2.9)	19.880 (1.060-372.900)	0.008	0.030	25.440 (1.397-463.100)	0.003	0.016
miR-146aC>G/miR-149T>C/miR-196a2T>C								
C-T-T	77 (21.0)	33 (20.7)	1.000 (reference)			0.993 (0.634-1.555)	1.000	1.000
G-T-C	53 (14.5)	10 (6.5)	0.440 (0.200-0.970)	0.045	0.158	0.437 (0.217-0.881)	0.018	0.072
G-C-T	14 (3.7)	17 (10.6)	2.833 (1.252-6.412)	0.018	0.126	2.813 (1.353-5.847)	0.008	0.064
miR-146aC>G/miR-149T>C/miR-499A>G								
C-T-A	116 (31.8)	57 (36.1)	1.000 (reference)			1.138 (0.788-1.645)	0.507	0.579
C-T-G	51 (13.9)	9 (5.5)	0.359 (0.165-0.781)	0.008	0.056	0.409 (0.196-0.851)	0.015	0.120
miR-146aC>G/miR-196a2T>C/miR-499A>G								
C-T-A	81 (22.2)	45 (28.2)	1.000 (reference)			1.287 (0.855-1.938)	0.240	0.384
C-T-G	42 (11.3)	6 (3.8)	0.257 (0.102-0.652)	0.003	0.021	0.331 (0.138-0.794)	0.008	0.064
G-C-A	72 (19.8)	20 (12.7)	0.500 (0.270-0.925)	0.035	0.123	0.644 (0.379-1.093)	0.106	0.212
miR-149T>C/miR-196a2T>C/miR-499A>G								
T-T-A	93 (25.3)	45 (28.3)	1.000 (reference)			1.121 (0.750-1.675)	0.604	0.690
C-C-G	0 (0.0)	8 (5.0)	34.930 (1.971-619.100)	0.0002	0.001	39.310 (2.254-685.700)	<0.0001	0.001
miR-146aC>G/miR-149T>C								
C-T	167 (45.8)	66 (41.6)	1.000 (reference)			0.916 (0.651-1.287)	0.666	0.804
G-C	39 (10.8)	33 (20.8)	2.141 (1.242-3.690)	0.009	0.027	1.960 (1.189-3.231)	0.010	0.040
miR-146aC>G/miR-499A>G								
C-A	163 (44.4)	81 (51.6)	1.000 (reference)			1.151 (0.831-1.594)	0.404	0.539
C-G	65 (17.9)	13 (7.9)	0.403 (0.210-0.773)	0.006	0.018	0.463 (0.248-0.865)	0.015	0.060
miR-149T>C/miR-499A>G								
T-A	206 (56.2)	83 (52.6)	1.000 (reference)			0.933 (0.681-1.280)	0.689	0.830
C-A	82 (22.5)	52 (32.9)	1.574 (1.023-2.422)	0.044	0.132	1.469 (0.990-2.179)	0.062	0.124
miR-196a2T>C/miR-499A>G								
T-A	134 (36.7)	78 (49.3)	1.000 (reference)			1.348 (0.964-1.887)	0.082	0.164
T-G	50 (13.6)	10 (6.4)	0.344 (0.165-0.716)	0.003	0.009	0.463 (0.229-0.937)	0.034	0.136
C-A	154 (42.0)	57 (36.2)	0.636 (0.421-0.961)	0.037	0.056	0.857 (0.600-1.225)	0.421	0.561

<sup>a</sup>P-value calculated by Fisher's exact test. OR, odds ratio; CI, confidence interval; miR/miRNA, microRNA; FDR, false discovery rate.

Table IV. Allele combination of miRNA polymorphisms between patients with meningioma (2n=138) and control subjects (2n=366).

Allele combination	Controls, n (%)	Meningioma, n (%)	Reference vs. Allele combination			Overall vs. Allele combination		
			OR (95% CI)	P-value <sup>a</sup>	FDR-P-value	OR (95% CI)	P-value <sup>a</sup>	FDR-P-value
<i>miR-146aC&gt;G/miR-149T&gt;C/miR-196a2T&gt;C/miR-499A&gt;G</i>								
C-T-T-A	51 (14.0)	26 (18.7)				1.352 (0.811-2.255)	0.277	0.462
C-T-C-G	26 (7.0)	0 (0.0)	0.037 (0.002-0.626)	0.0002	0.003	0.050 (0.003-0.826)	0.0002	0.003
G-T-C-A	53 (14.6)	6 (4.7)	0.222 (0.084-0.584)	0.002	0.014	0.300 (0.126-0.714)	0.004	0.030
G-T-C-G	0 (0.0)	4 (2.8)	17.490 (0.907-337.500)	0.017	0.079	23.820 (1.273-445.600)	0.006	0.030
<i>miR-146aC&gt;G/miR-149T&gt;C/miR-196a2T&gt;C</i>								
C-T-T	77 (21.0)	41 (29.5)	1.000 (reference)			1.412 (0.922-2.164)	0.115	0.308
C-T-C	89 (24.3)	23 (16.9)	0.485 (0.268-0.880)	0.019	0.112	0.685 (0.416-1.128)	0.154	0.308
C-C-T	46 (12.7)	10 (7.3)	0.408 (0.187-0.892)	0.032	0.112	0.577 (0.283-1.174)	0.151	0.308
<i>miR-146aC&gt;G/miR-149T&gt;C/miR-499A&gt;G</i>								
C-T-A	116 (31.8)	54 (39.3)	1.000 (reference)			1.235 (0.846-1.801)	0.281	0.664
C-T-G	51 (13.9)	10 (7.4)	0.421 (0.199-0.893)	0.029	0.203	0.520 (0.257-1.053)	0.066	0.528
<i>miR-146aC&gt;G/miR-196a2T&gt;C/miR-499A&gt;G</i>								
C-T-A	81 (22.2)	36 (25.8)	1.000 (reference)			1.179 (0.760-1.828)	0.493	0.789
G-C-G	4 (1.2)	7 (5.2)	3.938 (1.084-14.300)	0.042	0.186	4.641 (1.337-16.110)	0.014	0.112
<i>miR-149T&gt;C/miR-196a2T&gt;C/miR-499A&gt;G</i>								
T-T-A	93 (25.3)	47 (34.0)	1.000 (reference)			1.340 (0.897-2.003)	0.170	0.496
C-C-G	0 (0.0)	5 (4.0)	21.650 (1.172-400.200)	0.005	0.035	29.110 (1.598-530.300)	0.002	0.016
<i>miR-149T&gt;C/miR-196a2T&gt;C</i>								
T-T	126 (34.4)	58 (41.9)	1.000 (reference)			1.221 (0.845-1.763)	0.295	0.295
T-C	140 (38.3)	39 (28.4)	0.605 (0.378-0.970)	0.044	0.096	0.739 (0.493-1.108)	0.165	0.264

<sup>a</sup>P-value calculated by Fisher's exact test. OR, odds ratio; CI, confidence interval; miR/miRNA, microRNA; FDR, false discovery rate.



Table V. Allele combination of miRNA polymorphisms between patients with Schwannoma (2n=62) and control subjects (2n=366).

Allele combination	Controls, n (%)	Schwannoma, n (%)	Reference vs. Allele combination			Overall vs. Allele combination		
			OR (95% CI)	P-value <sup>a</sup>	FDR-P-value	OR (95% CI)	P-value <sup>a</sup>	FDR-P-value
<i>miR-146aC&gt;G/miR-149T&gt;C/miR-196a2T&gt;C/miR-499A&gt;G</i>								
C-T-T-A	51 (14.0)	18 (28.6)	1.000 (reference)			2.083 (1.142-3.801)	0.021	0.135
C-T-C-A	63 (17.3)	3 (4.6)	0.135 (0.038-0.484)	0.001	0.013	0.281 (0.086-0.923)	0.029	0.135
C-C-T-G	16 (4.4)	0 (0.0)	0.084 (0.005-1.479)	0.019	0.095	0.178 (0.011-3.002)	0.144	0.504
G-T-T-A	39 (10.6)	4 (6.3)	0.291 (0.091-0.928)	0.049	0.127	0.606 (0.209-1.754)	0.490	0.858
G-C-T-A	14 (3.8)	0 (0.0)	0.096 (0.005-1.692)	0.033	0.107	0.202 (0.012-3.435)	0.235	0.632
G-C-T-G	0 (0.1)	3 (4.3)	19.490 (0.959-395.800)	0.022	0.095	41.050 (2.093-805.000)	0.003	0.042
<i>miR-146aC&gt;G/miR-149T&gt;C/miR-196a2T&gt;C</i>								
C-T-T	77 (21.0)	20 (32.2)	1.000 (reference)			1.533 (0.875-2.687)	0.162	0.627
C-T-C	89 (24.3)	8 (12.5)	0.346 (0.144-0.830)	0.023	0.161	0.531 (0.245-1.148)	0.135	0.627
<i>miR-146aC&gt;G/miR-196a2T&gt;C/miR-499A&gt;G</i>								
C-T-A	81 (22.2)	22 (36.0)	1.000 (reference)			1.603 (0.932-2.759)	0.098	0.302
C-C-A	81 (22.2)	7 (11.0)	0.318 (0.129-0.786)	0.014	0.084	0.510 (0.225-1.156)	0.122	0.302
G-T-A	53 (14.5)	4 (6.5)	0.278 (0.091-0.852)	0.024	0.084	0.446 (0.156-1.275)	0.151	0.302
<i>miR-149T&gt;C/miR-196a2T&gt;C/miR-499A&gt;G</i>								
T-T-A	93 (25.3)	19 (31.4)	1.000 (reference)			1.206 (0.687-2.116)	0.552	0.877
T-C-G	27 (7.5)	0 (0.0)	0.087 (0.005-1.492)	0.024	0.084	0.107 (0.006-1.772)	0.037	0.148
C-C-G	0 (0.0)	3 (5.1)	33.560 (1.665-676.700)	0.006	0.042	41.050 (2.093-805.000)	0.003	0.024

OR, odds ratio; CI, confidence interval; miR/miRNA, microRNA; FDR, false discovery rate.

<sup>a</sup>P-value calculated by Fisher's exact test. OR, odds ratio; CI, confidence interval; miR/miRNA, microRNA; FDR, false discovery rate.

present study analyzed the association between the *miR-149* rs2292832 C>T polymorphism and glioma susceptibility and found allele-allelic combinations in which miRNA polymorphisms were positively associated with glioma, meningioma and schwannoma susceptibility.

### Acknowledgements

The abstract of this paper was presented at the 5th Quadrennial Meeting of the World Federation of Neuro-Oncology Societies, May 4-7, 2017, in Zurich, Switzerland, and was published as Abstract no. p08.33 in Neuro-Oncology (Suppl 3) 2017.

### Funding

This study was supported by the Industrial Technology Innovation Program through the Ministry of Trade, Industry and Energy (Korea), funded by the Ministry of Trade, Industry and Energy, Republic of Korea (grant no. 10067378), and supported by a grant of the Korea Health Technology Research and Development Project through the Korea Health Industry Development Institute, funded by the Ministry of Health and Welfare, Republic of Korea (grant no. H115C1972010015).

### Availability of data and materials

The datasets used during the present study are available from the corresponding author upon reasonable request.

### Authors' contributions

KC and NKK conceived and designed the experiments. JOK and HSP performed the experiments. JL, JOK, HSP, IBH, KK, KC and NKK analyzed the data. JL, JOK, HSP, IBH, KC and NKK were responsible for reagents, materials and analysis tools. JL and JOK wrote the paper. KC and NKK performed the article editing.

### Ethics approval and consent to participate

All study protocols were reviewed and approved by The Institutional Review Board of CHA Bundang Medical Center (Seongnam, South Korea) and followed the recommendations of the Declaration of Helsinki. All patients provided written informed consent.

### Patient consent for publication

All patients provided written informed consent for publication.

### Competing interests

The authors have no competing interests to declare.

### References

- Bandres E, Agirre X, Ramirez N, Zarate R and Garcia-Foncillas J: MicroRNAs as cancer players: Potential clinical and biological effects. *DNA Cell Biol* 26: 273-282, 2007.
- Deng S, Calin GA, Croce CM, Coukos G and Zhang L: Mechanisms of microRNA deregulation in human cancer. *Cell Cycle* 7: 2643-2646, 2008.
- Ruan K, Fang X and Ouyang G: MicroRNAs: Novel regulators in the hallmarks of human cancer. *Cancer Lett* 285: 116-126, 2009.
- Garg N, Vijayakumar T, Bakhshinyan D, Venugopal C and Singh SK: MicroRNA regulation of brain tumour initiating cells in central nervous system tumours. *Stem Cells Int* 2015: 141793, 2015.
- Iorio MV, Ferracin M, Liu CG, Veronese A, Spizzo R, Sabbioni S, Magri E, Pedriali M, Fabbri M, Campiglio M, *et al.*: MicroRNA gene expression deregulation in human breast cancer. *Cancer Res* 65: 7065-7070, 2005.
- Kreth S, Thon N and Kreth FW: Epigenetics in human gliomas. *Cancer Lett* 342: 185-192, 2014.
- Hummel R, Maurer J and Haier J: MicroRNAs in brain tumors: A new diagnostic and therapeutic perspective? *Mol Neurobiol* 44: 223-234, 2011.
- Galani V, Lampri E, Varouktsi A, Alexiou G, Mitselou A and Kyritsis AP: Genetic and epigenetic alterations in meningiomas. *Clin Neurol Neurosurg* 158: 119-125, 2017.
- Murnyák B, Bognár L, Klekner Á and Hortobágyi T: Epigenetics of meningiomas. *Biomed Res Int* 2015: 532451, 2015.
- Saydam O, Senol O, Würdinger T, Mizrak A, Ozdener GB, Stemmer-Rachamimov AO, Yi M, Stephens RM, Krichevsky AM, Saydam N, *et al.*: miRNA-7 attenuation in Schwannoma tumors stimulates growth by upregulating three oncogenic signaling pathways. *Cancer Res* 71: 852-861, 2011.
- Catucci I, Yang R, Verderio P, Pizzamiglio S, Heesen L, Hemminki K, Sutter C, Wappenschmidt B, Dick M, Arnold N, *et al.*: Evaluation of SNPs in miR-146a, miR-196a2 and miR-499 as low-penetrance alleles in German and Italian familial breast cancer cases. *Hum Mutat* 31: E1052-E1057, 2010.
- Li P, Shan JX, Chen XH, Zhang D, Su LP, Huang XY, Yu BQ, Zhi QM, Li CL, Wang YQ, *et al.*: Epigenetic silencing of microRNA-149 in cancer-associated fibroblasts mediates prostaglandin E2/interleukin-6 signaling in the tumor microenvironment. *Cell Res* 25: 588-603, 2015.
- Liu M, Du Y, Gao J, Liu J, Kong X, Gong Y, Li Z, Wu H and Chen H: Aberrant expression miR-196a is associated with abnormal apoptosis, invasion, and proliferation of pancreatic cancer cells. *Pancreas* 42: 1169-1181, 2013.
- Hu Z, Chen J, Tian T, Zhou X, Gu H, Xu L, Zeng Y, Miao R, Jin G, Ma H, *et al.*: Genetic variants of miRNA sequences and non-small cell lung cancer survival. *J Clin Invest* 118: 2600-2608, 2008.
- Yazici H, Zipprich J, Peng T, Akisik EZ, Tigli H, Isin M, Akisik EE, Terry MB, Senie RT, Li L, *et al.*: Investigation of the *miR16-1* (C>T)+7 substitution in seven different types of cancer from three ethnic groups. *J Oncol* 2009: 827532, 2009.
- Duan R, Pak C and Jin P: Single nucleotide polymorphism associated with mature miR-125a alters the processing of pri-miRNA. *Hum Mol Genet* 16: 1124-1131, 2007.
- Tian T, Shu Y, Chen J, Hu Z, Xu L, Jin G, Liang J, Liu P, Zhou X, Miao R, *et al.*: A functional genetic variant in microRNA-196a2 is associated with increased susceptibility of lung cancer in Chinese. *Cancer Epidemiol Biomarkers Prev* 18: 1183-1187, 2009.
- Xu J, Hu Z, Xu Z, Gu H, Yi L, Cao H, Chen J, Tian T, Liang J, Lin Y, *et al.*: Functional variant in microRNA-196a2 contributes to the susceptibility of congenital heart disease in a Chinese population. *Hum Mutat* 30: 1231-1236, 2009.
- Hu Z, Liang J, Wang Z, Tian T, Zhou X, Chen J, Miao R, Wang Y, Wang X and Shen H: Common genetic variants in pre-microRNAs were associated with increased risk of breast cancer in Chinese women. *Hum Mutat* 30: 79-84, 2009.
- Qi P, Dou TH, Geng L, Zhou FG, Gu X, Wang H and Gao CF: Association of a variant in MIR 196A2 with susceptibility to hepatocellular carcinoma in male Chinese patients with chronic hepatitis B virus infection. *Hum Immunol* 71: 621-626, 2010.
- Peng S, Kuang Z, Sheng C, Zhang Y, Xu H and Cheng Q: Association of microRNA-196a-2 gene polymorphism with gastric cancer risk in a Chinese population. *Dig Dis Sci* 55: 2288-2293, 2010.
- Wang K, Guo H, Hu H, Xiong G, Guan X, Li J, Xu X, Yang K and Bai Y: A functional variation in pre-microRNA-196a is associated with susceptibility of esophageal squamous cell carcinoma risk in Chinese Han. *Biomarkers* 15: 614-618, 2010.
- Jazdzewski K, Murray EL, Franssila K, Jarzab B, Schoenberg DR and de la Chapelle A: Common SNP in pre-miR-146a decreases mature miR expression and predisposes to papillary thyroid carcinoma. *Proc Natl Acad Sci USA* 105: 7269-7274, 2008.
- Liu Z, Li G, Wei S, Niu J, El-Naggar AK, Sturgis EM and Wei Q: Genetic variants in selected pre-microRNA genes and the risk of squamous cell carcinoma of the head and neck. *Cancer* 116: 4753-4760, 2010.

25. Ahn TK, Kim JO, Kumar H, Choi H, Jo MJ, Sohn S, Ropper AE, Kim NK and Han IB: Polymorphisms of miR-146a, miR-149, miR-196a2, and miR-499 are associated with osteoporotic vertebral compression fractures in Korean postmenopausal women. *J Orthop Res* 36: 244-253, 2018.
26. Han M, Yang Q, Feng K, Li R, Ren J and Wei L: Associations of MMP-2 -1306 C/T and MMP-9 -1562 C/T polymorphisms with breast cancer risk among different populations: a meta-analysis. *Genes Genom* 39: 331-340, 2017.
27. Benjamini Y and Hochberg Y: Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat* 57: 289-300, 1995.
28. Kim YR and Hong SH: Influences of -482C>T and 3238G>C polymorphisms of the apolipoprotein C3 gene on prevalence of metabolic syndrome. *Genes Genom* 38: 857-864, 2016.
29. Ritchie MD, Hahn LW, Roodi N, Bailey LR, Dupont WD, Parl FF and Moore JH: Multifactor-dimensionality reduction reveals high-order interactions among estrogen-metabolism genes in sporadic breast cancer. *Am J Hum Genet* 69: 138-147, 2001.
30. Moore JH and Williams SM: New strategies for identifying gene-gene interactions in hypertension. *Ann Med* 34: 88-95, 2002.
31. Hahn LW, Ritchie MD and Moore JH: Multifactor dimensionality reduction software for detecting gene-gene and gene-environment interactions. *Bioinformatics* 19: 376-382, 2003.
32. Ritchie MD, Hahn LW and Moore JH: Power of multifactor dimensionality reduction for detecting gene-gene interactions in the presence of genotyping error, missing data, phenocopy, and genetic heterogeneity. *Genet Epidemiol* 24: 150-157, 2003.
33. Lee SY, Chung Y, Elston RC, Kim Y and Park T: Log-linear model-based multifactor dimensionality reduction method to detect gene gene interactions. *Bioinformatics* 23: 2589-2595, 2007.
34. Jacques G and Cormac O: Central nervous system tumors. *Handb Clin Neurol* 112: 931-958, 2013.
35. Lacy J, Saadati H and Yu JB: Complications of brain tumors and their treatment. *Hematol Oncol Clin North Am* 26: 779-796, 2012.
36. de Robles P, Fiest KM, Frolkis AD, Pringsheim T, Atta C, St Germaine-Smith C, Day L, Lam D and Jette N: The worldwide incidence and prevalence of primary brain tumors: A systematic review and meta-analysis. *Neuro Oncol* 17: 776-783, 2015.
37. Choy W, Kim W, Nagasawa D, Stramotas S, Yew A, Gopen Q, Parsa AT and Yang I: The molecular genetics and tumor pathogenesis of meningiomas and the future directions of meningioma treatments. *Neurosurg Focus* 30: E6, 2011.
38. Babu R, Sharma R, Bagley JH, Hafez J, Friedman AH and Adamson C: Vestibular schwannomas in the modern era: Epidemiology, treatment trends, and disparities in management. *J Neurosurg* 119: 121-130, 2013.
39. Huse JT and Aldape KD: The evolving role of molecular markers in the diagnosis and management of diffuse glioma. *Clin Cancer Res* 20: 5601-5611, 2014.
40. Bastien JI, McNeill KA and Fine HA: Molecular characterizations of glioblastoma, targeted therapy, and clinical results to date. *Cancer* 121: 502-516, 2015.
41. Cloughesy TF, Cavenee WK and Mischel PS: Glioblastoma: From molecular pathology to targeted treatment. *Annu Rev Pathol* 9: 1-25, 2014.
42. Louis DN: Molecular pathology of malignant gliomas. *Annu Rev Pathol* 1: 97-117, 2006.
43. Liu Y, Pang JC, Dong S, Mao B, Poon WS and Ng HK: Aberrant CpG island hypermethylation profile is associated with atypical and anaplastic meningiomas. *Hum Pathol* 36: 416-425, 2005.
44. Mawrin C and Perry A: Pathological classification and molecular genetics of meningiomas. *J Neurooncol* 99: 379-391, 2010.
45. Perry A, Gutmann DH and Reifenberger G: Molecular pathogenesis of meningiomas. *J Neurooncol* 70: 183-202, 2004.
46. Yew A, Trang A, Nagasawa DT, Spasic M, Choy W, Garcia HM and Yang I: Chromosomal alterations, prognostic factors, and targeted molecular therapies for malignant meningiomas. *J Clin Neurosci* 20: 17-22, 2013.
47. Grönholm M, Teesalu T, Tyynelä J, Piltti K, Böhling T, Wartiovaara K, Vaheri A and Cárpen O: Characterization of the NF2 protein merlin and the ERM protein ezrin in human, rat, and mouse central nervous system. *Mol Cell Neurosci* 28: 683-693, 2005.
48. Giannakakis A, Coukos G, Hatzigeorgiou A, Sandaltzopoulos R and Zhang L: miRNA genetic alterations in human cancers. *Expert Opin Biol Ther* 7: 1375-1386, 2007.
49. Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, Aldler H, Rattan S, Keating M, Rai K, *et al*: Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci USA* 99: 15524-15529, 2002.
50. Guessous F, Alvarado-Velez M, Marcinkiewicz L, Zhang Y, Kim J, Heister S, Kefas B, Godlewski J, Schiff D, Purow B and Abounader R: Oncogenic effects of miR-10b in glioblastoma stem cells. *J Neurooncol* 112: 153-163, 2013.
51. Shi L, Zhang J, Pan T, Zhou J, Gong W, Liu N, Fu Z and You Y: MiR-125b is critical for the suppression of human U251 glioma stem cell proliferation. *Brain Res* 1312: 120-126, 2010.
52. Alsaidawi S, Malek E and Driscoll JJ: MicroRNAs in brain metastases: Potential role as diagnostics and therapeutics. *Int J Mol Sci* 15: 10508-10526, 2014.
53. Besse A, Sana J, Fadrus P and Slaby O: MicroRNAs involved in chemo- and radioresistance of high-grade gliomas. *Tumour Biol* 34: 1969-1978, 2013.
54. Low SY, Ho YK, Too HP, Yap CT and Ng WH: MicroRNA as potential modulators in chemoresistant high-grade gliomas. *J Clin Neurosci* 21: 395-400, 2014.
55. Lv M, Dong W, Li L, Zhang L, Su X, Wang L, Gao L and Zhang L: Association between genetic variants in pre-miRNA and colorectal cancer risk in a Chinese population. *J Cancer Res Clin Oncol* 139: 1405-1410, 2013.
56. Williams AE, Perry MM, Moschos SA, Lerner-Svensson HM and Lindsay MA: Role of miRNA-146a in the regulation of the innate immune response and cancer. *Biochem Soc Trans* 36: 1211-1215, 2008.
57. Li D, Chen P, Li XY, Zhang LY, Xiong W, Zhou M, Xiao L, Zeng F, Li XL, Wu MH, *et al*: Grade-specific expression profiles of miRNAs/mRNAs and docking study in human grade I-III astrocytomas. *OMICS* 15: 673-682, 2011.
58. Pan SJ, Zhan SK, Pei BG, Sun QF, Bian LG and Sun BM: MicroRNA-149 inhibits proliferation and invasion of glioma cells via blockade of AKT1 signaling. *Int J Immunopathol Pharmacol* 25: 871-881, 2012.
59. Yang G, Han D, Chen X, Zhang D, Wang L, Shi C, Zhang W, Li C, Chen X, Liu H, *et al*: MiR-196a exerts its oncogenic effect in glioblastoma multiforme by inhibition of IκBα both in vitro and in vivo. *Neuro Oncol* 16: 652-661, 2014.
60. Guan Y, Chen L, Bao Y, Qiu B, Pang C, Cui R and Wang Y: High miR-196a and low miR-367 cooperatively correlate with unfavorable prognosis of high-grade glioma. *Int J Clin Exp Pathol* 8: 6576-6588, 2015.