

Pilot investigation of the mutation profile of *PIK3CA/PTEN* genes (PI3K pathway) in grade 3 endometrial cancer

FRANCESCA MALENTACCHI¹, IRENE TURRINI¹, FLAVIA SORBI¹, ELISABETTA PROJETTO²,
FRANCESCA CASTIGLIONE², MASSIMILIANO FAMBRINI¹, FELICE PETRAGLIA¹,
SERENA PILLOZZI^{3*} and IVO NOCI^{1*}

¹Department of Biomedical, Experimental and Clinical Sciences, Division of Obstetrics and Gynecology;

²Department of Surgery and Translational Medicine, Division of Pathological Anatomy;

³Department of Experimental and Clinical Medicine, University of Florence, I-50134 Florence, Italy

Received April 16, 2018; Accepted October 9, 2018

DOI: 10.3892/or.2018.6939

Abstract. Endometrial cancer (EC) comprises a biological and clinical heterogeneous group of tumors. Several genetic alterations are involved in the development and progression of EC, and may be used for targeted therapy, particularly in patients with advanced-stage EC. In the present study, a combined procedure was developed based on polymerase chain reaction (PCR)-high resolution melting analysis (HRMA) and Sanger sequencing for the evaluation of somatic mutations in selected phosphoinositide 3-kinase (PI3K) catalytic subunit α (*PIK3CA*; exons 1, 9 and 21) and phosphatase and tensin homolog (*PTEN*; exons 5, 6, 7 and 8) exons. This combined procedure has the specificity and sensitivity of the two techniques, and overcomes their limitations. A pilot study was performed on 18 selected homogenous EC samples, of grade 3 endometrioid subtype (G3 EEC). First, the feasibility of the combined procedure was investigated to properly identify the presence of somatic mutations on *PIK3CA* and *PTEN*, the variations identified were analyzed using Catalogue of Somatic Mutations in Cancer,

PolyPhen-2 and Mutation Taster software, and the frequency of mutations/variations was determined in the selected samples. The evaluation of mutational load revealed that the majority of the G3 EEC samples exhibited *PIK3CA* mutations (39%) and *PTEN* mutations (67%), and the majority of the samples (83%) had mutations in at least one of the two genes, and 33% had mutations in the two genes. The results of the present pilot study suggested that the cost-effective combined PCR-HRMA and Sanger sequencing procedure may be suitable for identification of *PTEN* and *PIK3CA* mutations in G3 EEC and that their frequency was consistent in G3 EEC, indicating that the PI3K pathway serves a pivotal function that may have potential for defining targeted therapy for the treatment of G3 EEC.

Introduction

Endometrial cancer (EC) comprises a biologically and clinically heterogeneous group of tumors, generally classified as Type I (endometrioid) or Type II (non-endometrioid) on the basis of morphological characteristics. It was recently demonstrated that these two groups are characterized by different genetic alterations; consequently, the function of these variations has been widely investigated, in order to identify diagnostic and prognostic markers for improving the delineation of EC (1).

Recently, a novel classification system was published by The Cancer Genome Atlas (TCGA) on the basis of genetic and epigenetic modifications (2). In addition, a previous TCGA-based analysis has confirmed that >90% of endometrioid EC (EEC) harbor genetic alterations in the phosphoinositide 3-kinase (PI3K) signaling pathway (3), primarily concerning PI3K catalytic subunit α (*PIK3CA*), and phosphatase and tensin homolog (*PTEN*) (3-12).

Furthermore, it has been identified that grade 3 (G3) EEC exhibits clinical and pathological behavior between type I and type II, suggesting that EEC may evolve by mechanisms different from those for other EEC grades (13-16).

In EEC, the frequency of *PIK3CA* mutations is 50% (17) making it one of the most frequently mutated genes (18-21), independently of histological types (17,22). In total, >70% of *PIK3CA* mutations cluster at three 'hotspot' codons, exon 1

Correspondence to: Dr Francesca Malentacchi, Department of Biomedical, Experimental and Clinical Sciences, Division of Obstetrics and Gynecology, University of Florence, Viale G.B. Morgagni 50, I-50134 Florence, Italy
E-mail: francesca.malentacchi@gmail.com

*Joint senior authorship

Abbreviations: EC, endometrial cancer; TCGA, The Genome Cancer Atlas; PI3K, phosphoinositide 3-kinase; EEC, endometrioid EC; AKT, protein kinase B; PTEN, phosphatase and tensin homolog; PIK3CA, PI3K catalytic subunit α ; mTOR, mammalian target of rapamycin; G, grade; SNP, single nucleotide polymorphism; PCR, polymerase chain reaction; HRM, high-resolution melting; HRMA, high-resolution melting analysis; MT, Mutation Taster

Key words: endometrial endometrioid cancer, somatic mutations, phosphoinositide 3-kinase catalytic subunit α /phosphatase and tensin homolog genes, polymerase chain reaction-high-resolution melting analysis-Sanger sequencing combined procedure, mutational load

(p.R88Q), exon 9 (p.E542K, p.E545K/G/A, p.Q546R) and exon 21 (p.H1047R/L/Y, p.M1043I/V) (17,23), each of which induces its constitutive activation (3,24-26). However, the association between genetic variations and endometrial carcinogenesis remains poorly understood. Several studies have identified different correlations between *PIK3CA* variations and tumor grade, stage and type (5,6,26-29).

PTEN somatic mutations have been reported in several types of gynecological tumor; in addition, *PTEN* inactivation is present in endometriosis and endometrial hyperplasia, and may be considered an early event in cancer development (30-33). In EECs, *PTEN* mutations have been identified (16), and *PTEN* protein loss was revealed to be correlated with overall survival (OS) (34).

As suggested by TCGA, the PI3K signaling pathway may represent a potential target for targeted therapy for the treatment of EC (3,23). Indeed, several PI3K pathway inhibitors have been developed and are currently under investigation in preclinical studies, and in early clinical trials (35), with various results depending on the therapeutic agent used and treatment set-up (8,18,20,22,24,26,29,36-44). Nevertheless, it is currently unclear how *PIK3CA* and *PTEN* mutations affect the sensitivity to inhibitors in EC (45-47). Consequently, efforts should be focused on improving the characterization of the *PIK3CA* and *PTEN* mutational profiles, to allow the establishment of personalized targeted therapy.

Several approaches may be used for the characterization of mutations [e.g. next-generation sequencing (NGS), mass spectrometry (MS), Sanger sequencing, polymerase chain reaction (PCR)-high-resolution melting analysis (HRMA) and quantitative PCR (qPCR)]. Certain methods (i.e. NGS and MS) provide a comprehensive overall evaluation with high specificity and sensitivity, but require expensive instrumentation and procedures, and trained personnel; other methods (i.e. Sanger sequencing, PCR-HRMA and qPCR) are simpler and more cost-effective, but allow the analysis of only a limited number of selected exons, and the specificity and sensitivity differ (Sanger sequencing had high specificity and low sensitivity, and vice versa for PCR-HRMA, whereas qPCR is sensitive and specific, but allows the analysis of only single known hotspots) (48-51).

The present pilot study had two main aims: i) To develop an economical and efficient combined procedure on the basis of PCR-HRMA and Sanger sequencing for evaluating the presence of variants in *PIK3CA* (exons 1, 9 and 21) and *PTEN* (exons 5, 6, 7 and 8) genes with the intention to overcome the limitations of each technique; and ii) to assess the presence of the primary mutations in *PIK3CA* and *PTEN* in a selected homogeneous subgroup of G3 EECs.

Materials and methods

Tissue collection and DNA extraction. A total of 18 available and suitable tissues were selected from patients with G3 EEC and with at least 2 years of follow up (up to 1 February 2018). All patients underwent surgical treatment between May 2013 and November 2015, at the Obstetrics and Gynecology Unit, Careggi University Hospital (Florence, Italy). Patients were treated depending on International Federation of Gynecology

and Obstetrics (FIGO staging), grading and European Society of Gynecological Oncology (ESGO) risk (52). Tumor samples were collected at the time of surgery, following acquisition written informed consent from the patients. The study was approved by the Ethics Committee of Careggi University Hospital. Patients' demographic, clinical and pathological features are presented in Table I.

Tumor samples were immediately fixed in formalin and processed for paraffin embedding. The resulting formalin-fixed paraffin-embedded (FFPE) tissues were examined histologically by pathologists at Careggi University Hospital for the selection of areas containing >70% of tumor cells.

DNA was extracted from slices using a QIAamp DNA FFPE tissue kit (Qiagen GmbH, Hilden, Germany), according to the manufacturer's protocol. DNA quality (ratio R) and quantity (DNA concentration) were determined using a spectrophotometer (NanoDrop® 1000 UV spectrophotometer; Thermo Fisher Scientific, Inc., Waltham, MA, USA).

Primers and PCR conditions. Primers for selected exons of *PIK3CA* and *PTEN* were custom-made by Sigma-Aldrich; Merck KGaA (Darmstadt, Germany), and were designed to be suitable for Sanger sequencing and PCR-HRMA using primer3web software (version 4.1.0; primer3.ut.ee), checked for specificity and mismatch using Primer-BLAST (www.ncbi.nlm.nih.gov/tools/primer-blast) and using SNP Check (version 3; secure.ngml.org.uk/SNPCheck/snpcheck.htm). Primer sequences are presented in Table II.

PCR was performed using the custom primers, Taq PCR core kit (Qiagen GmbH) and the third-generation DNA-intercalating dye Syto 9 (Thermo Fisher Scientific, Inc.). Briefly, 10 ng DNA was amplified using 2 μ l master mix (10X), 0.5 μ l each primer (10 μ M), 0.4 μ l dNTPs (10 mM each), 0.4 μ l Syto9 (50 μ M), 0.1 μ l Taq DNA polymerase (5 U/ μ l) and water in a final volume of 20 μ l. Samples were subjected to incubation at 95°C for 5 min, then 40 amplification cycles of 95°C for 30 sec, 56/60°C (56°C for *PTEN* and 60°C for *PIK3CA*) for 45 sec and 72°C for 45 sec, and a final incubation at 72°C for 20 min in a GeneAmp PCR system 9700 (Thermo Fisher Scientific, Inc.).

HRMA. Amplified samples were submitted to a high-resolution melting (HRM) protocol (95°C for 5 min, 40°C for 1 min, ramping temperature from 72 to 84°C) in a Rotor-gene Q PCR cyclor (Qiagen GmbH). The melting curve of each sample was analyzed using melting and HRM software (Rotor-Gene® Q-Pure Detection series software, version 2.1.0, build 9; Qiagen GmbH).

Sanger sequencing. The amplified and melted PCR products were purified using a QIAquick PCR purification kit (Qiagen GmbH) and sequenced on an ABI 310 genetic analyzer (Thermo Fisher Scientific, Inc.). Briefly, 3 μ l purified PCR was added to 1 μ l BigDye Terminator, 3.2 μ l forward/reverse primer (1 μ M), 3.5 μ l BigDye Terminator buffer and water in a final volume of 20 μ l, and amplified for 25 cycles of 96°C for 10 sec, 59°C for 5 sec and 60°C for 2 min in the GeneAmp PCR system 9700, purified using a DyeEx 2.0 Spin column (Qiagen GmbH), according to the manufacturer's protocol, and 7 μ l was Sanger sequenced.

Table I. Demographic and clinicopathological features of patients with grade 3 endometrioid endometrial cancer.

Sample	BMI	Age at surgery, years	Menopause, status	Menopause, duration years	Myometrial, infiltration %	FIGO stage	Time disease-free, months	Total follow-up, months	LVSI	Lymph node involvement (n)	Risk
S1	Ob	57	Yes	4	>50	IB	22	58	No	No	High
S2	Nw	74	Yes	29	>50	IB	-	56	No	No	High
S3	Ob II	65	Yes	15	>50	IIIC	-	42	No	3	Advanced
S4	Ob II	78	Yes	23	>50	IB	6	38	No	No	High
S5	Ob	60	Yes	9	>50	IB	-	37	Yes	No	High
S6	Ob II	68	Yes	16	>50	IIIA	-	36	Yes	No	High
S7	Nw	54	Yes	1	>50	IB	-	36	Yes	No	High
S8	Ob I	72	Yes	20	>50	II	20	37	No	No	High
S9	Nw	80	Yes	31	>50	IB	-	36	Yes	No	High
S10	Ob III	80	Yes	30	>50	IB	-	34	Yes	No	High
S11	Ob I	79	Yes	30	>50	II	-	34	No	No	High
S12	Nw	77	Yes	24	>50	IB	25	32	Yes	No	High
S13	Nw	43	No	-	<50	IA	-	31	No	No	High-intermediate
S14	Ob II	74	Yes	15	<50	IA	-	30	Yes	No	High-intermediate
S15	Ob	69	Yes	13	<50	IA	-	28	No	No	High-intermediate

BMI, body mass index; FIGO, International Federation of Gynecology and Obstetrics; LVSI, lymphovascular space invasion; Nw, normal weight; Ob I, obesity class I; Ob II, obesity class II; Ob III, obesity class III.

Table II. Primer sequences.

Gene	Exon	Forward primer	Reverse primer	Product size, bp
<i>PIK3CA</i>	1	5'-TGTTACTCAAGAAGCAGAAAGGG-3'	5'-ACGAAGGTATTGGTTTAGACAGA-3'	231
	9	5'-AGGGAAAATGACAAAGAACA-3'	5'-ACCTGTGACTCCATAGAAA-3'	124
	21	5'-TGCTCCAAACTGACCAAACCTG-3'	5'-TGCATGCTGTTTAATTGTGTGG-3'	299
<i>PTEN</i>	5	5'-TGTGAAGATCTTGACCAATGGC-3'	5'-AAATTCTCAGATCCAGGAAGAGG-3'	231
	6	5'-ACGACCCAGTTACCATAGCA-3'	5'-TGTGAAACAACAGTGCCACT-3'	185
	7	5'-CCTCAGTTTGTGGTCTGCC-3'	5'-GCCAGAGTAAGCAAAACACCT-3'	296
	8	5'-TACCAGGACCAGAGGAAACC-3'	5'-AGCAAGTTCTTCATCAGCTGT-3'	280

To confirm identified mutations, the samples were also sequenced using reverse primers. The electropherogram of each sequenced amplicon was independently evaluated by two individuals. For validation of the combined procedure, in a subgroup of three samples, NGS was performed using Ion Torrent S5 with HotSpot cancer panel version 1 on Chip 520 (Thermo Fisher Scientific, Inc.) with a coverage (mean depth) average of 1,000X, mean reads of 400,000 and mean read length of 115 bp, according to the manufacturer's protocol.

For validation of variations identified by PCR-HRMA, but not by Sanger sequencing (sample S2 and S4), their corresponding PCR products associated with *PTEN* exon 5 were cloned using TOPO-TA cloning (Thermo Fisher Scientific, Inc.), according to the manufacturer's protocol. A total of 20 colonies was isolated and the plasmids were extracted using a QIAprep Spin Miniprep kit (Qiagen GmbH). The extracted plasmids were sequenced as aforementioned.

Mutation identification. The nucleotide changes were evaluated using the Catalogue of Somatic Mutations in Cancer (COSMIC) database (version 86; cancer.sanger.ac.uk/cosmic), ensembl (release 93; www.ensembl.org/index.html), PolyPhen-2 (version 2; genetics.bwh.harvard.edu/pph2) and Mutation Taster (MT) (version 2; current build, NCBI 37/Ensembl 69; www.mutationtaster.org) software. Single nucleotide polymorphisms (SNPs) were confirmed using UCSC Genome Browser (genome.ucsc.edu/). The clinical impact was evaluated using COSMIC by Functional Analysis Through Hidden Markov Models score (pathogenic score, range: 0, not pathogenic, to 1, pathogenic), for single amino acid substitutions by PolyPhen-2 score (range: 0, not pathogenic, to 1, pathogenic) and for synonymous single amino acid substitutions, complex variations (insertions/deletions/frameshifts) and modifications in intronic sequence by MT [evaluation: Disease-causing (DC) or SNP].

Results

Quality of DNA. The extracted DNAs had sufficient amount and quality for PCR amplification and Sanger sequencing; in particular, the concentration mean was 164.15 ± 2.00 ng/ μ l; the median was 138.00 ± 2.00 ng/ μ l; the range was between 18.00 and 299.10 ng/ μ l; the value of R was 2.15 (mean) and 2.15 (median), and the range was between 2.00 and 2.47.

Establishment of the combined PCR-HRMA and Sanger sequencing procedure. The combined procedure is based on the sequential execution of PCR-HRMA and Sanger sequencing, without other steps, and avoids double PCR if the two procedures are performed separately with a corresponding saving of DNA, time of execution and limiting the number of mistakes due to performing the analysis twice. All samples were amplified to verify the success of amplification and to perform the HRMA. An appropriate curve was obtained for all the samples and for each analyzed exon. The HRMA suggested the presence of nucleotide alteration(s) without providing direct information on the specific nucleotide change(s), which induced changes in the shape of the HRM and melting curves. In particular, the presence of nucleotide change(s) induced or a shift in the peak or the appearance of additional peak(s) in the melting curve and a change in the shape of the HRM curve. In order to identify associations of the curve shape with a specific nucleotide sequence, the amplicons were sequenced using the Sanger procedure.

Figs. 1 and 2 present representative results of the HRM and melting curves, and the corresponding specific base modification identified by Sanger sequencing, for *PIK3CA* exon 1 and for *PTEN* exon 5, respectively.

The comparison of HRMA and sequencing data confirmed that samples with the same HRM curve shared the same nucleotide sequence. Differences in HRM curves for the same amplicon suggested the presence of different nucleotide sequences. Figs. 3 and 4 present the principal variations detected by HRM curves along with the corresponding nucleotide change(s) for each exon of *PIK3CA* and *PTEN* analyzed. Notably, for *PTEN* exon 5, it was observed that three samples shared the same HRM and melting curves (Fig. 5), but the corresponding Sanger sequences revealed the presence of a specific variant only for one sample (sample S9; p.Y155S), suggesting that the two samples (samples S2 and S4) may have the same variation as sample S9, but at lower levels. To verify this hypothesis, the PCR products of samples S2 and S4 were cloned into a plasmid and sequenced, which confirmed the presence of the suggested variant at low level (two and three positive colonies, respectively, in samples S2 and S4).

This combined approach allowed the avoidance of a double PCR (if the two methodologies were applied separately), retaining DNA for use in further analysis. Furthermore, the use of HRMA alone suggested the presence of homoduplex




Figure 1. Investigation of the principal genetic variants in phosphoinositide 3-kinase catalytic subunit α exon 1. Each curve represents an exemplificative sample. The same color is used to represent the sample and the corresponding identified variation(s); 'wt' corresponds to a sample(s) that presented no variation in the specific amplicon in comparison with the GenBank reference sequence. (A) High-resolution melting curves. (B) Melting curves (C) Sanger hotspot sequence. If the mutation was present in more than one sample, only one sequence is presented. wt, wild-type.




Figure 2. Investigation of the principal genetic variants in phosphatase and tensin homolog exon 5. Each curve represents an exemplificative sample. The same color is used to represent the sample and the corresponding identified variation(s); 'wt' corresponds to a sample(s) that presented no variation in the specific amplicon in comparison with the GenBank reference sequence. (A) High-resolution melting curves. (B) Melting curves. (C) Sanger hotspot sequence. If the mutation was present in more than one sample, only one sequence is presented. wt, wild-type.

or heteroduplex variations in the analyzed exons [*PIK3CA* exon 1 (61%), exon 9 (6%) and exon 21 (33%), and *PTEN* exon 5 (44%), exon 6 (78%), exon 8 (50%) and none in exon 7),

but provided information on the type of variation. Sanger sequencing provided this specific information on the type of alteration, allowing the characterization of the nucleotide




Figure 3. High-resolution melting curves associated with to the principal identified phosphoinositide 3-kinase catalytic subunit α gene variations. Each curve represents an exemplificative sample. The color key indicates the sample and the corresponding principal variation identified. The term 'wt' corresponds to a sample(s) that presented no variation in the specific amplicon in comparison with the GenBank reference sequence. (A) Exon 1. (B) Exon 9. (C) Exon 21. For each sample, the corresponding identified variants are presented in Table III. wt, wild-type.

variation, but with low sensitivity. The merged information provided the sensitivity of HRMA and the specificity of Sanger sequencing (for example, for *PTEN* exon 5, the mutation detection frequency increased from 33 to 44%).

To further evaluate this approach, three samples were validated using this combined procedure and by NGS technology in which all the *PTEN* and *PIK3CA* exons were analyzed, as well as 48 other cancer-associated genes. The two technologies yielded the same results: For sample S16 only, a single-base mutation in *PTEN* exon 5 was detected.

Evaluation of variants on *PIK3CA* and *PTEN*.

***PIK3CA*.** A total of seven different variations in 11 samples (61%) were identified, all of which were heterozygous, except for IVS1+40A>G (intronic) which in three samples was homozygous and 7/11 were pathogenic (Table III). In particular, in exon 1, two mutations were identified, namely p.L89R (6%) and p.R88G (18%), as well as the SNP IVS1+40A>G (rs3729674; 44%); in exon 9, the mutation p.Q546R (cosm12459; 6%), was identified; and in exon 21, the mutation p.H1047P (cosm249874; 22%) and the variant p.E1037Q (12%) were detected. The highest number of mutations (three mutations and one variant) was observed in exon 21 compared with exon 1 (two mutations) and exon 9 (one mutation). All five samples at FIGO stage III (IIIA and IIIC) did not exhibit any variation in *PIK3CA*.

***PTEN*.** A total of 12 different variations were identified and 16/18 (89%) samples exhibited at least one variation, among

which 12 (67%) presented a pathological variation (Table IV). In particular, exon 6 exhibited the highest number of variations. The most represented alteration was p.Q171Q (78%), followed by p.R189K (cosm1745951; 44%). Exon 5 exhibited pathogenic mutations in six samples (33%): p.Y155S (18%), p.D162H (cosm5274; 12%), p.Y155H (cosm5038; 6%), p.G143fs*4 (cosm30623; 6%), R130* (cosm5152; 6%) and G129Stop (cosm18663; 6%). Exon 8 had an intronic variant (IVS8+32T>G; 44%) and in one sample it was associated with p.I300fs*2. Exon 7 did not present any mutation according to Kafshooz *et al* (53). No particular trend was observed for the *PTEN* mutation, except for the non-menopausal patient that had *PTEN* R130* and the presence of the *PTEN* T167fs* in the obese class III patient.

Variant/mutational load. Regarding the overall variant evaluation, the majority of the samples (89%) exhibited at least one variation (Table V). Regarding SNPs, 50% of the samples exhibited one SNP and 18% presented with two SNPs.

Regarding the mutations, 67% of the samples had confirmed mutations at least in one of the two genes, whereas 18% had mutations in the two genes, and these values increased to 72 and 33%, respectively, if the mutations predicted with PolyPhen-2 were considered, and to 82 and 39%, respectively, if the MT prediction was considered.

It was observed that 28% of the samples exhibited a single confirmed mutation, 16% exhibited two mutations, 22% exhibited three mutations and 6% exhibited four mutations. Considering the overall variant load (excluding SNPs), 18% of




Figure 4. High-resolution melting curves associated with the principal identified phosphatase and tensin homolog gene variants. Each curve represents an exemplificative sample. 'wt' corresponds to a sample(s) that presented no variation in the specific amplicon in comparison with the GenBank reference sequence. (A) Exon 5. (B) Exon 6. (C) Exon 6. (D) Exon 7. (E) Exon 8. For each sample, the corresponding identified variants are presented in Table IV. wt, wild-type.




Figure 5. Investigation of phosphatase and tensin homolog exon 5 variant c.464A>C by HRM analysis in samples S9 (red), S2 (light blue) and S4 (blue). S11 represents the reference 'wt' (sample with no variations in this amplicon). (A) HRM curves. (B) Melting curves. HRM, high-resolution melting; wt, wild-type.

Table III. *PIK3CA* variants.

Sample	FIGO stage	PIK3CA variants			Pathogenic effect		Reported by COSMIC	
		Location	Nucleotide change	Predicted protein change	Yes/no	Predicted by	Other types of cancer	In EC
S1	IB	Exon 1	IVS1+40A>G		No (SNP)	UCSC/dbSNP/ClinVar	-	-
S2	IB	Exon 1	c.262C>G	p.R88G	Yes	PolyPhen-2 (score 0.994)	-	-
S3	IIIC	-	-	-	-	-	-	-
S4	IB	Exon 1	IVS1+40A>G		No (SNP)	UCSC/dbSNP/ClinVar	-	-
S5	IB	Exon 1	IVS1+40A>G		No (SNP)	UCSC/dbSNP/ClinVar	-	-
		Exon 21	c.3073A>G	p.T1025A	Yes	COSMIC (FATHMM score 1.00)	Yes	Yes (first) ^a
		Exon 21	c.3140A>C	p.H1047P	No (SNP)	Cosmic (FATHMM score 0.96)	Yes	No
S6	IIIA	-	-	-	-	-	-	-
S7	IB	Exon 1	IVS1+40A>G		No (SNP)	UCSC/dbSNP/ClinVar	-	-
		Exon 21	c.3140A>C	p.H1047P	Yes	Cosmic (FATHMM score 0.96)	Yes	No
S8	II	Exon 1	IVS1+40A>G homo		No (SNP)	UCSC/dbSNP/ClinVar	-	-
		Exon 21	c.3140A>C	p.H1047P	Yes	Cosmic (FATHMM score 0.96)	Yes	No
S9	IB	Exon 1	IVS1+40A>G homo		No (SNP)	UCSC/dbSNP/ClinVar	-	-
S10	IB	Exon 1	c.266T>G	p.L89R	Yes	PolyPhen-2 (score 1.000)	-	-
		Exon 9	c.1637A>G	p.Q546R	Yes	Cosmic (FATHMM score 0.97)	Yes	Yes (third) ^a
		Exon 21	c.3140A>C	p.H1047P	Yes	Cosmic (FATHMM score 0.96)	Yes	No
S11	II	Exon 1	c.262C>G	p.R88G	Yes	PolyPhen-2 (score 0.994)	-	-
		Exon 1	IVS1+40A>G		No (SNP)	UCSC/dbSNP/ClinVar	-	-
		Exon 21	c.3109G>C	p.E1037Q	Yes	PolyPhen-2 (score 0.011)	-	-
						Mutation Taster (disease-causing)	-	-
S12	IB	Exon 1	IVS1+40A>G homo		No (SNP)	UCSC/dbSNP/ClinVar	-	-
S13	IA	Exon 1	c.262C>G	p.R88G	Yes	PolyPhen-2 (score 0.994)	-	-
		Exon 21	c.3109G>C	p.E1037Q	No	PolyPhen-2 (score 0.011)	-	-
						Mutation Taster (disease-causing)	-	-
S14	IA	-	-	-	-	-	-	-
S15	IA	-	-	-	-	-	-	-
S16	IIIA	-	-	-	-	-	-	-
S17	IIIC	-	-	-	-	-	-	-
S18	IIIC	-	-	-	-	-	-	-

All the variants were heterozygous, excepted those indicated 'homo' (homozygosity). ^aThis value represents the position of EC in the cosmic ranking for the mutation investigated. FIGO, International Federation of Gynecology and Obstetrics; PIK3CA, phosphoinositide 3-kinase catalytic subunit α ; COSMIC, Catalogue of Somatic Mutations in Cancer; EC, endometrial cancer; SNP, single nucleotide polymorphism; FATHMM, Functional Analysis Through Hidden Markov Models.

Table IV. *PTEN* variants.

Sample	FIGO stage	PTEN variants			Pathogenic effect		Reported by COSMIC	
		Location	Nucleotide change	Predicted protein change	Yes/no	Predicted by	Other types of cancer	In EC
S1	IB	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
S2	IB	Exon 8	IVS8+32T>G		No (SNP)	UCSC/dbSNP/Clin Var	-	-
		Exon 5	c.464A>C	p.Y155S	Yes	PolyPhen-2 (score 1.000)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
S3	IIIC	Exon 8	IVS8+32T>G homo		No (SNP)	UCSC/dbSNP/Clin Var	-	-
		Exon 5	c.484G>C	p.D162H	Yes	COSMIC (FATHMM score 0.99)	Yes	No
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 5	c.464A>C	p.Y155S	Yes	PolyPhen-2 (score 1.000)	-	-
S4	IB	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
S5	IB	Exon 8	IVS8+32T>G homo		No (SNP)	UCSC/dbSNP/Clin Var	-	-
		Exon 5	c.484G>C	p.D162H	Yes	COSMIC (FATHMM score 0.99)	Yes	No
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
S6	IIIA	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 8	IVS8+32T>G		No (SNP)	UCSC/dbSNP/Clin Var	-	-
		Exon 5	c.425delG	p.G143Afs*4	No	COSMIC (FATHMM score no.)	(ovary)	No
		Exon 5	c.463T>C	p.Y155H	Yes	COSMIC (FATHMM score 0.97)	Yes	Yes (first) ^a
S7	IB	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
S8	II	Exon 5	c.464A>C	p.Y155S	Yes	PolyPhen-2 (score 1.000)	-	-
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
		Exon 8	c.899_900delCT	p.I300Nfs*2	Yes	PolyPhen-2 (score 0.059), associated with an amino acid alteration	-	-
S9	IB	Exon 8	IVS8+32T>G		No (SNP)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.497_498insT	p.T167Nfs*13	Yes	UCSC/dbSNP/Clin Var	-	-
		Exon 6	c.497_498insT	p.T167Nfs*13	Yes	PolyPhen-2 (score 0.999), associated with an amino acid alteration	-	-
		Exon 6	c.497_498insT	p.T167Nfs*13	Yes	Mutation Taster (disease-causing)	-	-
S10	IB	Exon 6	c.497_498insT	p.T167Nfs*13	Yes	UCSC/dbSNP/Clin Var	-	-
		Exon 6	c.497_498insT	p.T167Nfs*13	Yes	PolyPhen-2 (score 0.999), associated with an amino acid alteration	-	-
		Exon 6	c.497_498insT	p.T167Nfs*13	Yes	Mutation Taster (disease-causing)	-	-
		Exon 6	c.497_498insT	p.T167Nfs*13	Yes	Mutation Taster (disease-causing)	-	-

Table IV. Continued.

Sample	FIGO stage	PTEN variants			Pathogenic effect		Reported by COSMIC	
		Location	Nucleotide change	Predicted protein change	Yes/no	Predicted by	Other types of cancer	In EC
S11	II	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 8	IVS8+32T>G homo		No (SNP)	UCSC/dbSNP/Clin Var		
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
S12	IB	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
S13	IA	Exon 5	c.388C>T	p.R130*		COSMIC (FATHMM score 0.95)	Yes	Yes (first) ^a
S14	IA	Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)		
		Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
		Exon 8	IVS8+32T>G		No (SNP)	UCSC/dbSNP/Clin Var	-	-
		Exon 6	c.513G>A	p.Q171Q	(Yes)	Mutation Taster (disease-causing)	-	-
S15	IA	Exon 6	c.549G>A	p.K183K	(Yes)	Mutation Taster (disease-causing)	-	-
S16	IIIA	Exon 6	c.566G>A	p.R189K	Yes	COSMIC (FATHMM score 0.88)	Yes	No
		Exon 8	IVS8+32T>G			UCSC/dbSNP/Clin Var	-	-
		-	-	-	-	-	-	-
		Exon 5	c.385G>T	p.G129Stop	Yes	COSMIC (FATHMM score 0.99)	Yes	Yes (third) ^a
S17	IIIC	-	-	-	-	-	-	-
S18	IIIC	-	-	-	-	-	-	-

All the variants were heterozygous, excepted those indicated 'homo' (homozygosity). ^aThis value represents the position of EC in the cosmic ranking for the mutation investigated. FIGO, International Federation of Gynecology and Obstetrics; PTEN, phosphatase and tensin homolog; COSMIC, Catalogue of Somatic Mutations in Cancer; EC, endometrial cancer; SNP, single nucleotide polymorphism; FATHMM, Functional Analysis Through Hidden Markov Models.

Table V. Overall evaluation of *PIK3CA* and *PTEN* variants in the samples analyzed.

Sample	<i>PIK3CA</i> variants	<i>PTEN</i> variants
S1	<i>IVS1+40A>G</i>	p.Q171Q, <i>IVS8+32T>G</i>
S2	p.R88G	p.R189K , p.Y155S , p.Q171Q, <i>IVS8+32T>G</i>
S3		p.D162H , p.Q171Q
S4	<i>IVS1+40A>G</i>	p.R189K , p.Y155S , p.Q171Q
S5	p.T1025A , p.H1047P , <i>IVS1+40A>G</i>	p.R189K , p.Q171Q, <i>IVS8+32T>G</i>
S6		p.D162H , p.Q171Q, <i>IVS8+32T>G</i>
S7	p.H1047P , <i>IVS1+40A>G</i>	p.Y155H , p.R189K , <i>p.G143Afs*4</i> , p.Q171Q
S8	p.H1047P , <i>IVS1+40A>G</i>	p.R189K , p.Q171Q
S9	<i>IVS1+40A>G</i>	p.R189K , p.Q171Q, p.Y155S , <i>p.I300Nfs*2</i> , <i>IVS8+32T>G</i>
S10	p.Q546R , p.H1047P , p.L89R	p.T167Nfs*13 , p.Q171Q, <i>IVS8+32T>G</i>
S11	p.R88G , p.E1037Q, <i>IVS1+40A>G</i>	p.Q171Q
S12	<i>IVS1+40A>G</i>	p.Q171Q
S13	p.R88G , p.E1037Q	p.R130* , p.R189K , p.Q171Q
S14		<i>IVS8+32T>G</i>
S15		p.R189K , p.Q171Q, p.K183K, <i>IVS8+32T>G</i>
S16		
S17		p.G129Stop
S18		

Italic represents single nucleotide polymorphisms. Bold represents variations predicted as 'pathogenic' using PolyPhen-2. Underlining represents variation predicted as 'pathogenic' using Mutation Taster. Bold and underlining represents variation confirmed as 'pathogenic' by Catalogue of Somatic Mutations in Cancer. *PIK3CA*, phosphoinositide 3-kinase catalytic subunit α ; *PTEN*, phosphatase and tensin homolog.

the samples exhibited none, one or four mutations, 10% exhibited two or five mutations, and 26% exhibited three mutations.

No specific mutation/variant load distribution was observed according to clinicopathological characteristics.

Discussion

Currently, there is increasing interest in investigating the genome alterations in tumors, with such research made possible by the development of specific and sensitive technologies (e.g. NGS or MassArray). These promising technologies are useful for understanding the mechanisms underlying carcinogenesis and cancer evolution, but are costly and require computational support for data analysis. For clinical application, the identification of genetic variations must be performed rapidly and cost-effectively, as usually performed by Sanger sequencing/pyrosequencing or PCR-HRMA procedures. Typically, qPCR is performed for confirmation analysis.

It is established that PCR-HRMA has a lower cost and is an accurate tool for mutation scanning, allowing the identification of variations through comparison of HRM curves; consequently, it is used as screening method or for the detection of known hotspot mutations. Its ability in discriminating known mutations is due to the use of reference material, or if this material is not available or novel variations are present in the samples analyzed, the technique is not able to identify the nucleotide change, but only to detect the presence (54-56). The HRMA curve shape depends on the number and type of nucleotide changes, with different nucleotide changes

yielding different curves. The procedure alone is highly sensitive (overall sensitivity 0.1-5%) (50,51), but lacks specificity, since it does not allow for the direct evaluation of the specific nucleotide changes. Consequently, for the identification of nucleotide variations, the procedure requires the use of reference materials with the same alteration to be investigated along with the target samples, or further investigations by other techniques (such as sequencing or qPCR).

Sanger sequencing is the reference procedure for the identification of nucleotide changes for clinical purposes; even if this technology is unable to reveal variations at low levels (overall sensitivity ~20%) (48,49), sometimes pyrosequencing is used due its increased sensitivity. However, this methodology is relatively expensive compared with Sanger sequencing, even if more sensitive (57).

Currently, other high-throughput methodologies are beginning to be used in the clinic for the identification of genetic variation in clinical samples, such as NGS and MS that allow the evaluation of multiple genes/exons and several samples in the same run. Usually these procedures improve specificity and sensitivity, but are expensive either for their required instrumentation and methodological procedures, provide abundant information that is not useful for clinical purposes, and require trained personnel for data analysis. In particular, MS allows the evaluation of known variations on specific exons/genes; instead NGS allows the evaluation of genes or genome and the identifications of new genetic variants. qPCR, which has high sensitivity and specificity, allows the identification of well-known and characterized variations, but is expensive and

Table VI. Principal characteristics of the main procedures used for identification of variants in DNA sequence.

	Procedure						
	High-throughput		Single amplicon				
	MassArray	NGS	qPCR	PCR-HRMA	Sanger sequencing	Pyrosequencing	PCR-HRMA+Sanger sequencing
Cost per sample	Medium	High	Medium	Low	Medium	Medium	Medium
Sensitivity	High	High	High	High	Low	Medium	High
Specificity	High	High	High	Low	High	High	High
Discovery of novel variant	No	Yes	No	Yes	Yes	Yes/no ^a	Yes
Identification of sequence change	Yes	Yes	Yes	No	Yes	Yes	Yes
Multiple hotspot within amplicon length	No in the same well/yes in a run	Yes	No/yes ^b	Yes	Yes	Yes	Yes
Multi-exon analysis in the same preparation	Yes	Yes	No	No	No	No	No
Multi-sample analysis in the same preparation	Yes	Yes	No	No	No	No	No
Trained personnel	Yes	Yes	No	No	No	Medium	No
Cost of instrumentation	High	High	No/medium ^c	No	No	Medium	No
Data analysis	Yes	Yes	No	No	No	No	No

^aIt depends on whether the run is set for detection of targeted variants (easy and less time-consuming) or for identification of a novel mutation (more time-consuming and expensive compared with targeted approach). ^bIt depends on whether a multiplex or singleplex assay is used. ^cSeveral instruments are present on the market with a broad range of prices. NGS, next-generation sequencing; PCR, polymerase chain reaction; qPCR, quantitative PCR; HRMA, high-resolution melting analysis.

time-consuming if several genes or variations require analysis, and is not suitable for the detection of unknown variations. It is typically used for confirmation when Sanger sensitivity fails.

The present study establishes a combined procedure that overcomes the limitations of each technique alone (low sensitivity of Sanger sequencing and low specificity of HRMA) and combines the strengths (high specificity of Sanger sequencing and high sensitivity of HRMA) of these techniques. It also allows a double check for the presence of variants in the main exons of *PIK3CA* and *PTEN*, in order to improve the reliability of the results. The target of this approach is the identification of variants in single samples and in amplicons of between 100 and 200 bp in length; consequently, it cannot be compared with high-throughput procedures. Furthermore, this combined approach, in comparison with the application of either method alone, may decrease the amount of DNA used in the analysis (a single PCR is performed for the two methodologies) and the number of samples analyzed by Sanger sequencing. In particular, if a reference wild-type (wt) sample is used in the PCR-HRMA step, only the samples presenting HRM curves different from the wt should be analyzed by Sanger sequencing. Furthermore, PCR-HRMA can also only suggest the presence of novel variants, but they cannot be identified;

by contrast, the combined use of PCR-HRMA and Sanger analysis can immediately confirm and identify the novel variations. This combined procedure may even be useful for defining patient eligibility for targeted therapy; it is known that targeted therapy is dependent on the identification of specific mutations by Sanger sequencing, and its efficacy is determined by the number of mutated cells and by the development of any new mutations that can originate from tumor heterogeneity or *de novo*. Owing to the high sensitivity of HRMA, suspected cases with a low proportion of mutations may still be identified and monitored (Table VI).

The present study focused on the analysis of specific exons of *PIK3CA* and *PTEN*, since these are established as the main genes involved in alterations in the PI3K signaling pathway, and they have been widely investigated for the characterization of EC as specific targets for personalized therapies on the basis of PI3K pathway inhibitors (17,18,20,29,35,40). Indeed, early clinical data on several tumor types suggested that *PIK3CA* and *PTEN* mutations may affect the success of PI3K/protein kinase B (AKT)/mammalian target of rapamycin (mTOR)-targeted therapies (58-61). Patients with gynecological tumors and *PIK3CA* mutations have demonstrated a 30% response rate (RR) in early-phase clinical trials

with PI3K/AKT/mTOR inhibitors, compared with a 10% RR in patients lacking *PIK3CA* mutations (42). It is conceivable that loss of PTEN function can be similarly predictive for therapy efficacy, whereas simultaneous mutations in the mitogen-activated protein kinase signaling pathway may lead to resistance to treatment (58,60). Identifying actionable molecular aberrations has been a critical step for several major therapeutic advances in cancer medicine.

A combined technique was established in the present study for the examination of G3 EEC in order to investigate the involvement of major mutations of the PI3K signaling pathway in this EC subtype, and to ultimately implement the clinical and pathological knowledge of this grade of EEC regardless of differing characteristics (e.g. FIGO stage and ESGO risk). The present pilot study demonstrated that the approach was successful in analyzing G3 EEC, specifically for the genetic profile of selected *PIK3CA* and *PTEN* exons.

It was observed, in the cohort of the present study, that several samples were wt for specific exons (from 22% of *PTEN* exon 6 to 100% of *PTEN* exon 7) and that the overall number of variations was broad (in a single sample there were up to three variations for *PTEN* exon 6 and up to seven variations overall).

The focus was on the *PIK3CA* exons 1, 9, and 21, and on the *PTEN* exons 5, 6, 7 and 8 since these are typically the more frequently mutated exons in EECs; indeed, in a previous study, Rudd *et al* (17), when investigating all *PIK3CA* and *PTEN* exons, identified that no other exons were involved in G3 EEC. Furthermore, to verify this result and to partially validate the combined methodological approach, three samples were analyzed by NGS obtained the same results. Nevertheless, not examining all exons of *PTEN* and *PIK3CA* is a limitation of the present study, and perhaps something to perform in the future.

Using the approach developed in the present study, the overall evaluation revealed that 89% of samples exhibited at least one variation in either of the two genes, and that 83% had mutations. Regarding *PIK3CA*, 39% of samples exhibited mutations, among which the most frequent were H1047P (27%), p.T1025A (7%) and p.Q546R (7%) (7,17,21,22). With regard to *PTEN*, 67% of the samples had mutations, including p.R189K (53%), p.D132H (13%), p.Y155H (7%) and p.R130* (7%).

As aforementioned, the overall mutational load was determined as at least one mutation in 83% of the samples and coexistence of mutations in the two genes in 33% of the samples. An increased mutation frequency was identified for each single gene compared with the coexistence of mutations, in contrast with results reported previously (27), particularly those reported by Oda *et al* (6), which may be due to the specific evaluation of the G3 subgroup of EEC in the present study.

It was observed that the FIGO stage III tumors did not exhibit any mutation in *PIK3CA*, and that the patient with class III obesity had *PIK3CA* p.Q546 and *PTEN* p.T167fs* mutations, suggesting an association supported by a previous study of the association of obesity and EC through the involvement of the PI3K signaling pathway (62).

The preliminary data of the present study indicated potential for the use of a combined technological approach for the

identification of variants in *PIK3CA* and *PTEN*. Although they are preliminary results owing to the limited number of samples analyzed, the data confirm the involvement of PI3K pathway alterations in G3 EEC; however, further investigation is required.

In conclusion, the present study has demonstrated the suitability and reliability of a combined approach (PCR-HRMA and Sanger sequencing) for the evaluation of variants in selected exons of *PIK3CA* and *PTEN* in G3 EEC, suggesting that this approach may be useful for improving the classification and personalized treatment of patients with EC.

Acknowledgements

Not applicable.

Funding

The present study was supported by grants from the Tuscan Tumor Institute, Florence (Italy) and the University of Florence (Italy).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

FM performed Sanger sequencing, evaluation of Sanger electropherograms, analyses and interpretation of HRMA and Sanger data, identification of nucleotide changes using databases (COSMIC, PolyPhen-2 and mutation taster), and was a major contributor in writing the manuscript. IT collected the clinical and pathophysiological data for each patient, evaluated the criteria for tissue collection and selection, and set up a database containing patient information. FS was involved in patient recruitment, collection of the clinical and pathophysiological data for each patient, and evaluation of the criteria for tissue collection and selection. EP performed the FFPE sample selection, the histological examination of selected FFPE slices, and the selection of appropriate slices for molecular analysis. FC performed the FFPE sample collection, FFPE slices and the histological examination of selected FFPE slices. MF performed the patient recruitment, the surgery and treatment on enrolled women, and collection of informed consent. FP co-ordinated the clinical team for patient recruitment and supervised the data analysis. SP performed DNA extraction and PCR, evaluation of Sanger electropherograms and contributed in editing of the manuscript. IN performed the patient recruitment, the surgery and treatment on enrolled women, the collection of informed consents, follow-up of patients and contributed in editing of the manuscript. All authors read, revised and approved the final manuscript.

Ethics approval and consent to participate

Research is performed on humans following international and national regulations in accordance with The Declaration of Helsinki, or any other relevant set of ethical principles. This

research involved human subjects or tissues, and the authors state that informed consent for participation in the study or use of their tissue was obtained from all participants.

Patient consent for publication

The manuscript did not contain identifying information, including names, initials, date of birth, hospital numbers or images related to patients.

Competing interests

The authors declare that they have no competing interests.

References

- Murali R, Soslow RA and Weigelt B: Classification of endometrial carcinoma: More than two types. *Lancet Oncol* 15: e268-e278, 2014.
- Cancer Genome Atlas Research Network; Kandoth C, Schultz N, Cherniack AD, Akbani R, Liu Y, Shen H, Robertson AG, Pashtan I, Shen R, *et al*: Integrated genomic characterization of endometrial carcinoma. *Nature* 497: 67-73, 2013.
- Marchiò C, De Filippo MR, Ng CK, Piscuoglio S, Soslow RA, Reis-Filho JS and Weigelt B: PIKING the type and pattern of PI3K pathway mutations in endometrioid endometrial carcinomas. *Gynecol Oncol* 137: 321-328, 2015.
- McConechy MK, Ding J, Cheang MCU, Wiegand K, Senz J, Tone A, Yang W, Prentice L, Tse K, Zeng T, *et al*: Use of mutation profiles to refine the classification of endometrial carcinomas. *J Pathol* 228: 20-30, 2012.
- Salvesen HB, Carter SL, Mannelqvist M, Dutt A, Getz G, Stefansson IM, Raeder MB, Sosh ML, Engelsens IB, Trovik J, *et al*: Integrated genomic profiling of endometrial carcinoma associates aggressive tumors with indicators of PI3 kinase activation. *Proc Natl Acad Sci USA* 106: 4834-4839, 2009.
- Oda K, Stokoe D, Taketani Y and McCormick F: High frequency of coexistent mutations of *PIK3CA* and *PTEN* genes in endometrial carcinoma. *Cancer Res* 65: 10669-10673, 2005.
- Catus L, D'Angelo E, Pons C, Espinosa I and Prat J: Expression profiling of 22 genes involved in the PI3K-AKT pathway identifies two subgroups of high-grade endometrial carcinomas with different molecular alterations. *Mod Pathol* 23: 694-702, 2010.
- Hayes MP, Wang H, Espinal-Witter R, Douglas W, Solomon GJ, Baker SJ and Ellenson HL: *PIK3CA* and *PTEN* mutations in uterine endometrioid carcinoma and complex atypical hyperplasia. *Clin Cancer Res* 12: 5932-5935, 2006.
- van der Zee M, Jia Y, Wang Y, Heijmans-Antonissen C, Ewing PC, Franken P, DeMayo FJ, Lydon JP, Burger CW, Fodde R, *et al*: Alterations in Wnt- β -catenin and Pten signalling play distinct roles in endometrial cancer initiation and progression. *J Pathol* 230: 48-58, 2013.
- Sal V, Demirkiran F, Erenel H, Tokgozoglu N, Kahramanoglu I, Bese T, Turan H, Sofiyeva N, Calay Z, Arvas M and Guralp O: Expression of PTEN and β -catenin and their relationship with clinicopathological and prognostic factors in endometrioid type endometrial cancer. *Int J Gynecol Cancer* 26: 512-520, 2016.
- McConechy MK, Ding J, Senz J, Yang W, Melnyk N, Tone AA, Prentice LM, Wiegand KC, McAlpine JN, Shah SP, *et al*: Ovarian and endometrial endometrioid carcinomas have distinct CTNNB1 and PTEN mutation profiles. *Mod Pathol* 27: 128-134, 2014.
- Machin P, Catus L, Pons C, Muñoz J, Matias-Guiu X and Prat J: CTNNB1 mutations and beta-catenin expression in endometrial carcinomas. *Hum Pathol* 33: 206-212, 2002.
- Tejerizo-García A, Jiménez-López JS, Muñoz-González JL, Bartolomé-Sotillos S, Marqueta-Marqués L, López-González G and Gómez JF: Overall survival and disease-free survival in endometrial cancer: Prognostic factors in 276 patients. *Onco Targets Ther* 6: 1305-1313, 2013.
- Mang C, Birkenmaier A, Cathomas G and Humburg J: Endometrioid endometrial adenocarcinoma: An increase of G3 cancers? *Arch Gynecol Obstet* 295: 1435-1440, 2017.
- Voss MA, Ganesan R, Ludeman L, McCarthy K, Gornall R, Schaller G, Wei W and Sundar S: Should grade 3 endometrioid endometrial carcinoma be considered a type 2 cancer-a clinical and pathological evaluation. *Gynecol Oncol* 124: 15-20, 2012.
- Yasuda M: Immunohistochemical characterization of endometrial carcinomas: Endometrioid, serous and clear cell adenocarcinomas in association with genetic analysis. *J Obstet Gynecol Res* 40: 2137-2176, 2014.
- Rudd ML, Price JC, Fogoros S, Godwin AK, Sgroi DC, Merino MJ and Bell DW: A unique spectrum of somatic *PIK3CA* (p110A α) mutations within primary endometrial carcinomas. *Clin Cancer Res* 17: 1331-1340, 2011.
- Mackay HJ, Eisenhauer EA, Kamel-Reid S, Tsao M, Clarke B, Karakasis K, Werner HM, Trovik J, Akslen LA, Salvesen HB, *et al*: Molecular determinants of outcome with mammalian target of rapamycin inhibition in endometrial cancer. *Cancer* 15: 603-610, 2014.
- Mori N, Kyo S, Sakaguchi J, Mizumoto Y, Ohno S, Maida Y, Hashimoto M, Takakura M and Inoue M: Concomitant activation of AKT with extracellular-regulated kinase 1/2 occurs independently of *PTEN* or *PIK3CA* mutations in endometrial cancer and may be associated with favorable prognosis. *Cancer Sci* 98: 1881-1888, 2007.
- Shoji K, Oda K, Kashiya T, Ikeda Y, Nakagawa S, Sone K, Miyamoto Y, Hiraike H, Tanikawa M, Miyasaka A, *et al*: Genotype-dependent efficacy of a dual PI3K/mTOR inhibitor, NVP-BEZ235, and an mTOR inhibitor, RAD001, in endometrial carcinomas. *PLoS One* 7: e37431, 2012.
- Millis SZ, Ikeda S, Reddy S, Gatalica Z and Kurzrock R: Landscape of phosphatidylinositol-3-kinase pathway alterations across 19784 diverse solid tumors. *JAMA Oncol* 2: 1565-1573, 2016.
- Catus L, Gallardo A, Cuatrecasas M and Prat J: Concomitant PI3K-AKT and p53 alterations in endometrial carcinomas are associated with poor prognosis. *Mod Pathol* 22: 522-529, 2009.
- Konopka B, Janiec-Jankowska A, Kwiatkowska E, Najmola U, Bidziński M, Olszewski W and Goluda C: *PIK3CA* mutations and amplification in endometrioid endometrial carcinomas: Relation to other genetic defects and clinicopathologic status of the tumors. *Hum Pathol* 42: 1710-1719, 2011.
- German S, Aslam HM, Saleem S, Raees A, Anum T, Alvi AA and Haseeb A: Carcinogenesis of *PIK3CA*. *Hered Cancer Clin Pract* 11: 5, 2013.
- English DP, Bellone S, Cocco E, Bortolomai I, Pecorelli S, Lopez S, Silasi DA, Schwartz PE, Rutherford T and Santin AD: Oncogenic *PIK3CA* gene mutations and HER2/neu gene amplifications determine the sensitivity of uterine serous carcinoma cell lines to GDC-0980, a selective inhibitor of Class I PI3 kinase and mTOR kinase (TORC1/2). *Am J Obstet Gynecol* 209: 465.e1-e9, 2013.
- Dong Y, Yang X, Wong O, Zhang X, Liang Y, Zhang Y, Wong W, Nong L, Liao Q and Li T: *PIK3CA* mutations in endometrial carcinomas in Chinese women: Phosphatidylinositol 3'-kinase pathway alterations might be associated with favorable prognosis. *Hum Pathol* 43: 1197-1205, 2012.
- Herrero-Gonzalez S and Di Cristofano A: New routes to old places: *PIK3R1* and *PIK3R2* join *PIK3CA* and *PTEN* as endometrial cancer genes. *Cancer Discov* 1: 106-107, 2011.
- Lin DI: Improved survival associated with somatic *PIK3CA* mutations in copy-number low endometrioid endometrial adenocarcinoma. *Oncol Lett* 10: 2743-2748, 2015.
- Wik E, Raeder MB, Krakstad C, Trovik J, Birkeland E, Hoivik EA, Mjos S, Werner HM, Mannelqvist M, Stefansson IM, *et al*: Lack of estrogen receptor- α is associated with epithelial-mesenchymal transition and PI3K alterations in endometrial carcinoma. *Clin Cancer Res* 19: 1094-1105, 2013.
- Smith IN and Briggs JM: Structural mutation analysis of PTEN and its genotype-phenotype correlations in endometriosis and cancer. *Proteins* 84: 1625-1643, 2016.
- Akiyama-Abe A, Minaguchi T, Nakamura Y, Michikami H, Shikama A, Nakao S, Sakurai M, Ochi H, Onuki M, Matsumoto K, *et al*: Loss of PTEN expression is an independent predictor of favourable survival in endometrial carcinomas. *Br J Cancer* 109: 1703-1710, 2013.
- Djordjevic B, Hennessy BT, Li J, Barkoh BA, Luthra R, Mills GB and Broaddus RR: Clinical assessment of PTEN loss in endometrial carcinoma: Immunohistochemistry outperforms gene sequencing. *Mod Pathol* 25: 699-708, 2012.

33. Catusas L, Gallardo A, Cuatrecasas M and Prat J: *PIK3CA* mutations in the kinase domain (exon 20) of uterine endometrial adenocarcinomas are associated with adverse prognostic parameters. *Mod Pathol* 21: 131-139, 2008.
34. Li W, Wang Y, Fang X, Zhou M, Li Y, Dong Y and Wang R: Differential expression and clinical significance of DNA methyltransferase 3B (DNMT3B), phosphatase and tensin homolog (PTEN) and human MutL homologs 1 (hMLH1) in endometrial carcinomas. *Med Sci Monit* 23: 938-947, 2017.
35. Bradford LS, Rauh-Hain A, Clark RM, Groeneweg JW, Zhang L, Borger D, Zuckerberg LR, Growdon WB, Foster R and Rueda BR: Assessing the efficacy of targeting the phosphatidylinositol 3-kinase/AKT/mTOR signaling pathway in endometrial cancer. *Gynecol Oncol* 133: 346-352, 2014.
36. Morice P, Leary A, Creutzberg C, Abu-Rustum N and Darai E: Endometrial cancer. *Lancet* 387: 1094-1108, 2016.
37. Matias-Guiu X and Prat J: Molecular pathology of endometrial carcinoma. *Histopathology* 62: 111-123, 2013.
38. McIntyre JB, Nelson GS, Ghatage P, Morris D, Duggan MA, Lee CH, Doll CM and Köbel M: *PIK3CA* missense mutation is associated with unfavorable outcome in grade 3 endometrioid carcinoma but not in serous endometrial carcinoma. *Gynecol Oncol* 132: 188-193, 2014.
39. Miyake T, Yoshino K, Enomoto T, Takata T, Ugaki H, Kim A, Fujiwara K, Miyatake T, Fujita M and Kimura T: *PIK3CA* gene mutations and amplifications in uterine cancers, identified by methods that avoid confounding by *PIK3CA* pseudogene sequences. *Cancer Lett* 261: 120-126, 2008.
40. Garcia-Dios DA, Lambrechts D, Coenegrachts L, Vandenput I, Capoen A, Webb PM, Ferguson K, ANECS, Akslen LA, Claes B, *et al*: High-throughput interrogation of *PIK3CA*, *PTEN*, *KRAS*, *FBXW7* and *TP53* mutations in primary endometrial carcinoma. *Gynecol Oncol* 128: 327-334, 2013.
41. Tsikouras P, Bouchlariotou S, Vrachnis N, Dafopoulos A, Galazios G, Csorba R and von Tempelhoff GF: Endometrial cancer: Molecular and therapeutic aspects. *Eu J Obstet Gynecol Reprod Biol* 169: 1-9, 2013.
42. Janku F, Wheler JJ, Westin SN, Moulder SL, Naing A, Tsimberidou AM, Fu S, Falchook GS, Hong DS, Garrido-Laguna I, *et al*: PI3K/AKT/mTOR inhibitors in patients with breast and gynecologic malignancies harboring *PIK3CA* mutations. *J Clin Oncol* 30: 777-782, 2012.
43. Bian X, Gao J, Luo F, Rui C, Zheng T, Wang D, Wang Y, Roberts TM, Liu P, Zhao JJ, *et al*: *PTEN* deficiency sensitizes endometrioid endometrial cancer to compound PARP-PI3K inhibition but not PARP inhibition as monotherapy. *Oncogene* 37: 341-351, 2018.
44. Philip CA, Laskov I, Beauchamp MC, Marques M, Amin O, Bitharas J, Kessous R, Kogan L, Baloch T, Gotlieb WH, *et al*: Inhibition of PI3K-AKT-mTOR pathway sensitizes endometrial cancer cell lines to PARP inhibitors. *BMC Cancer* 17: 638, 2017.
45. Hasegawa K, Kagabu M, Mizuno M, Oda K, Aoki D, Mabuchi S, Kamiura S, Yamaguchi S, Aoki Y, Saito T, *et al*: Phase II basket trial of perifosine monotherapy for recurrent gynecologic cancer with or without *PIK3CA* mutations. *Invest New Drugs* 35: 800-812, 2017.
46. Mirantes C, Dosil MA, Eritja N, Felip I, Gatiús S, Santacana M, Matias-Guiu X and Dolcet X: Effects of the multikinase inhibitors Sorafenib and Regorafenib in *PTEN* deficient neoplasias. *Eur J Cancer* 63: 74-87, 2016.
47. McCormick A, Earp E, Leeson C, Dixon M, O'Donnell R, Kaufmann A and Edmondson RJ: Phosphatase and Tensin Homolog is a potential target for ovarian cancer sensitization to cytotoxic agents. *Int J Gynecol Cancer* 26: 632-639, 2016.
48. Tsiatis AC, Norris-Kirby A, Rich RG, Hafez MJ, Gocke CG, Eshleman JR and Murphy KM: Comparison of Sanger sequencing, pyrosequencing, and melting curve analysis for the detection of *KRAS* mutations: Diagnostic and clinical implications. *J Mol Diagn* 12: 425-432, 2010.
49. Ihle MA, Fassunke J, König K, Grünewald I, Schlaak M, Kreuzberg N, Tietze L, Schildhaus HU, Büttner R and Merkelbach-Bruse S: Comparison of high resolution melting analysis, pyrosequencing, next generation sequencing and immunohistochemistry to conventional Sanger sequencing for the detection of p.V600E and non-p V600E *BRAF* mutations. *BMC Cancer* 14: 13, 2014.
50. Li BS, Wang XY, Ma FL, Jiang B, Song XX and Xu AG: Is high resolution melting analysis (HRMA) accurate for detection of human disease-associated mutations? A meta analysis. *PLoS One* 6: e28078, 2011.
51. Vossen RH, Aten E, Roos A and den Dunnen JT: High-resolution melting analysis (HRMA): More than just sequence variant screening. *Hum Mutat* 30: 860-866, 2009.
52. Colombo N, Creutzberg C, Amant F, Bosse T, González-Martín A, Ledermann J, Marth C, Nout R, Querleu D, Mirza MR, *et al*: ESMO-ESGO-ESTRO consensus conference on endometrial cancer diagnosis, treatment and follow-up. *Int J Gynecol Cancer* 26: 2-30, 2016.
53. Kafshooz L, Kafshdooz T, Tabrizi AD, Mohaddes Ardabili SM, Ardabili M, Arkbazadeh A, Gharepouran J, Ghojzadeh M and Farajnia S: Role of exon 7 *PTEN* gene in endometrial carcinoma. *Asian Pac J cancer Prev* 16: 4521-4524, 2015.
54. Simi L, Pratesi N, Vignoli M, Sestini R, Cianchi F, Valanzano R, Nobili S, Mini E, Pazzagli M and Orlando C: High-resolution melting analysis for rapid detection of *KRAS*, *BRAF*, and *PIK3CA* gene mutations in colorectal cancer. *Am J Clin Pathol* 130: 247-253, 2008.
55. Ousati Ashtiani Z, Mehraei AR, Pourmand MR and Pourmand GR: High resolution melting analysis for rapid detection of *PIK3CA* gene mutations in bladder cancer: A mutated target for cancer therapy. *Urol J* 15: 26-31, 2018.
56. Martínez-Carretero C, Pascual FI, Rus A and Bernardo I: Detection of *EGFR* mutations in patients with non-small cell lung cancer by high resolution melting. Comparison with other methods. *Clin Chem Lab Med* 55: 1970-1978, 2017.
57. Tsiatis AC, Norris-Kirby A, Rich G, Hafez MJ, Gocke CG, Eshleman JR and Murphy KM: Comparison of Sanger sequencing, pyrosequencing, and melting curve analysis for the detection of *KRAS* mutations: Diagnostic and clinical implications. *J Mol Diagn* 12: 425-432, 2010.
58. Engelman A: Targeting PI3K signaling in cancer: Opportunities, challenges and limitations. *Nat Rev Cancer* 8: 550-562, 2009.
59. Di Nicolantonio F, Arena S, Tabernero J, Grosso S, Molinari F, Macarulla T, Russo M, Cancelliere C, Zecchin D, Mazzucchelli L, *et al*: Deregulation of the PI3K and *KRAS* signaling pathways in human cancer cells determines their response to everolimus. *J Clin Invest* 120: 2558-2866, 2010.
60. Ni X, Zhang W and Huang RS: Pharmacogenomics discovery and implementation in genome-wide association studies era. *Wiley Interdiscip Rev Syst Biol Med* 5: 1-9, 2013.
61. Tsimberidou AM, Iskander NG, Hong DS, Wheler JJ, Falchook GS, Fu S, Piha-Paul S, Naing A, Janku F, Luthra R, *et al*: Personalized medicine in a phase I clinical trials program: The MD Anderson Cancer Center initiative. *Clin Cancer Res* 15: 6373-6383, 2012.
62. Westin SN, Ju Z, Broaddus RR, Krakstad C, Li J, Pal N, Lu KH, Coleman RL, Hennesy BT, Klempner SJ, *et al*: *PTEN* loss is a context-dependent outcome determinant in obese and non-obese endometrioid endometrial cancer patients. *Mol Oncol* 9: 1694-1703, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.