

# Interaction between C2ORF68 and HuR in human colorectal cancer

ZHAOXU LV, KUNYAN HE, LIHONG SHI, KAI SHI, TINGTING JIANG and YAO CHEN

Department of Human Anatomy, West China School of Basic Medical Sciences and Forensic Medicine, Sichuan University, Chengdu, Sichuan 610041, P.R. China

Received April 26, 2018; Accepted October 31, 2018

DOI: 10.3892/or.2019.6973

**Abstract.** The detailed molecular mechanisms underlying the carcinogenesis of colorectal carcinoma (CRC) remain unknown. Therefore, the present study was designed to investigate the effect of the relationship between C2ORF68 and HuR in regards to the carcinogenesis of CRC. Immunohistochemistry, immunofluorescence, flow cytometry, Transwell migration and CCK-8 assays, co-immunoprecipitation, qRT-PCR and western blot analysis were performed. The results revealed that expression of C2ORF68 was significantly upregulated in the cytoplasm and nucleus in rectal cancer, and upregulation of the expression of C2ORF68 was associated with lymph node metastasis and pathological grade. C2ORF68 and HuR were found to be mainly localized in the nucleus in both SW480 and LoVo cells. In LoVo<sup>+c2orf68, -HuR</sup> and LoVo<sup>+c2orf68</sup> cells, the cell apoptosis rate was significantly decreased, cell proliferation rate was significantly increased, and the cell migration rate was only significantly increased in the LoVo<sup>+c2orf68</sup> cells. In SW480<sup>-c2orf68, -HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup>, the cell apoptosis rate was significantly increased. At the same time, cell proliferation and the cell migration rate were significantly decreased. The mRNA and protein expression levels of *C2orf68*, *HuR*, *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* were upregulated, while the expression of *Bax* was downregulated in LoVo<sup>+c2orf68</sup> and LoVo<sup>+c2orf68, -HuR</sup> cells. Expression levels of *C2orf68*, *HuR*, *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* were downregulated while *Bax* was upregulated in the SW480<sup>-c2orf68, -HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells. In conclusion, it is suggested that *c2orf68* is a potential carcinogenesis factor in rectal cancer. Furthermore, *c2orf68* may have a synergistic effect with *HuR* in the onset and development of CRC.

## Introduction

Colorectal cancer (CRC) is the third most commonly diagnosed malignant tumor and is the fourth leading cause of cancer-related death worldwide (1). It is largely diagnosed in individuals >50 years of age (2). CRC patients in the early stage [tumor-node-metastasis (TNM) stage I and II] present with a prolonged 5-year survival following surgical excision; the 5-year survival is up to 95% and 60-80% for stage I and II, respectively. However, existing therapies for CRC usually exhibit a limited effect on patient prognosis, and the 5-year survival of patients in stages III and IV can be as low as 35 and 10% (3). Research has shown that CRC incidence and mortality can be decreased significantly through screening programs, while CRC screening is only offered to very few individuals worldwide based on the CRC incidence rate, national economic level and medical security system (4). A previous study demonstrated that germline mutations enable next generation hereditary susceptibility to CRC accounting for 6-7%. In addition, mutations in DNA repair genes and signal transduction genes also contribute to the occurrence of CRC. In addition to inherited genetic mutations, environmental factors, such as the heavy consumption of alcohol, smoking habit, increased body fat and diets high in fat, salt and red and processed meat, also play important roles (5). However, more and more studies have shown that CRC displays accumulated defects in the activation of oncogenes and the inactivation of tumor-suppressor genes (TSGs) (6). Except for classical CRC risk factors, such as: *KRAS*, *TP53*, *APC* and markers for microsatellite instability (MSI) (7,8), an impressive body of literature indicates that multiple factors such as hsa-miR-19a (9), *PROK2* (10), *B7-H3* (11), *DSCC1* (12) and microRNAs (13) are involved in the occurrence of CRC. A recent study revealed that the dysbiosis of microbial communities in the human body are also associated with gastrointestinal tract cancer, such as gastric and colorectal cancer. Moreover, a hypothesis called 'Alpha bug' considers that some bacteria could alter the primary bacterial community and the remodeled bacterial community could promote CRC by strengthening the mucosal immune response (14). Although numerous studies have shown that the pathogenesis of CRC is multifactorial, the detailed mechanisms remain unclear at present.

C2ORF68, which belongs to the UPF0561 family, contains 166 amino acids, and the monoisotopic molecular weight is

---

**Correspondence to:** Professor Yao Chen, Department of Human Anatomy, West China School of Basic Medical Sciences and Forensic Medicine, Sichuan University, 17 People's South Road, Chengdu, Sichuan 610041, P.R. China  
E-mail: chenya062@scu.edu.cn

**Key words:** C2ORF68, HuR, colorectal cancer, interaction

18.6033 kDa (15). The predicted potential NLS and NES sequences of C2ORF68 and the domain of the EF hand indicate that *c2orf68* could function as a cell cycle regulator in the nucleus, which may explain its possible role in CRC pathogenesis. Our early research suggests that C2ORF68 may regulate colon cancer cell apoptosis, proliferation, migration and cell cycle distribution through the PI3K/Akt/mTOR pathway (16). We also suggested that C2ORF68 may promote the carcinogenesis of CRC and play a vital role in the pathogenesis of CRC through activation of the Wnt/ $\beta$ -catenin signaling pathway (17), such as the upregulation of  $\beta$ -catenin, *survivin*, *cyclin D1* and *c-Myc*, and the downregulation of *GSK-3 $\beta$*  in CRC cells. Although it was demonstrated that C2ORF68 may play a role in the occurrence of CRC and may be a potential oncogene in CRC, its specific molecular mechanism remains dim.

Our previous study indicated that C2ORF68 may play an important role in the occurrence and development of CRC through the PI3K/AKT/mTOR and Wnt/ $\beta$ -catenin signaling pathways. However, its exact mechanisms are still mysterious. To elucidate the role and status of C2ORF68 in the carcinogenesis of colorectal adenocarcinoma, we carried out bioinformatic analyses. It was found that 12 proteins interact with *c2orf68*, including KLHL15, GMNN, SMG6, GNE, DDHD2, PIR, ELAVL1 (HuR), NAGK, XPO1, HSPA9, JOSD2 and GUK1. HuR is an RNA-binding protein whose expression level is widely upregulated in many types of human cancer, including CRC. A previous study showed that the HuR expression level is closely related to AKT phosphorylation and increased cytoplasmic abundance of HuR in human cancer may be associated with oncogenic activation of AKT signaling (18). To determine the interaction between C2ORF68 and HuR, and their roles in the occurrence of CRC, immunohistochemistry (IHC), immunofluorescence (IF), flow cytometry, Transwell migration and CCK-8 assays, co-immunoprecipitation (co-IP), qRT-PCR and western blot analysis were performed. The results revealed that C2ORF68 has a synergistic effect with HuR, and their role in the onset and the development of CRC may be through the upregulated expression of *Bcl-2*, *c-Myc*, *cyclin A* and *cyclin D1* and the downregulation of *Bax*, consequently promoting cell proliferation and inhibiting cell apoptosis.

## Materials and methods

**Bioinformatic analysis.** BioGRID is an interaction database with data based on comprehensive curation efforts. The interactive proteins for C2ORF68 were predicted through the BioGrid repository (<https://thebiogrid.org/>).

**Tissue microarray.** A tissue microarray including 90 rectal cancer tissues and adjacent normal tissues were purchased from Shanghai Xinchao (Shanghai Xinchao Biological Co. Ltd., Shanghai, China). The Medical Research Ethics Committee of Sichuan University approved the sample acquisition (Chengdu, China), and a written informed consent was also obtained from all patients.

**Immunohistochemistry (IHC) of the rectal cancer tissue microarray.** IHC was performed as previously described (19). Clinicopathological parameters of the CRC patients are shown

in Table I. The primary antibody (Ab) used for IHC was mouse monoclonal C2ORF68 Ab (1:200; BIO014915; Beacombio, Birmingham, UK). The expression level of C2ORF68 protein in rectal cancer tissues were scored by three independent examiners. The level of C2ORF68 staining pattern was scored according to four subgroups: i) Negative (-); ii) weak (+); iii) moderate (++); and iv) strong (+++).

**Cell lines and cell culture.** Human colon cancer cell lines, SW480 and LoVo [American Type Culture Collection (ATCC) Manassas, VA, USA], were cultured in Dulbecco's modified Eagle's medium (DMEM; HyClone; GE Healthcare Life Sciences, Logan, UT, USA) which contained 10% fetal bovine serum (FBS), penicillin (100 IU/ml) and streptomycin (100 IU/ml). The cells were grown at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>. The cells were harvested when they were in the exponential growth phase and then the experiments stated above were performed.

**Immunofluorescence (IF).** For immunostaining, the SW480 and LoVo cells were fixed with 4% paraformaldehyde for 20 min, permeabilized with 0.5% Triton X-100 for 15 min and blocked with 3% bovine serum albumin (BSA) for 30 min at room temperature. The treated cells were then incubated with mouse monoclonal anti-C2ORF68 (1:100) and rabbit polyclonal anti-HuR (dilution 1:100; cat. no. ab200342; Abcam, Cambridge, UK) overnight at 4°C, and finally incubated with FITC-labeled and TRITC-labeled secondary Ab for 30 min under the conditions of protection from light at room temperature. Each step was followed by two 5-min washes in PBS. The nuclei were counterstained using DAPI, and observed using an Olympus BX53 fluorescence microscope (Olympus, Hamburg, Germany).

**siRNA selection and transient transfection.** siRNAs for *c2orf68* (siRNA sequences, 5'-CUAUGAAGAGUCCGGUGAAdTdT-3' and 3'-dTdTTCUUCUCCAUAACUUAACG AU-5') and *HuR* (siRNA sequences, 5'-GGUUGCGUUUAUCCGGUUAUdTdT-3' and 3'-dTdTCCAACGCAAAUAGGCCAAA-5') were designed and synthesized by RiboBio (Guangzhou, China). Using the blank and negative control groups, the transfection was performed with 100 nM of siRNA and Invitrogen™ Lipofectamine 2000™ (Thermo Fisher Scientific, Inc., Waltham, MA, USA) to induce the knockdown of *c2orf68*/HuR expression. After transfection for 6 h, the cells were respectively incubated in fresh DMEM for 48 h to detect the mRNA expression level and for 72 h to detect the protein expression level.

**Strain, plasmid, plasmid extraction and overexpression.** The *c2orf68* gene was synthesized and inserted into the *XhoI/EcoRI* site of the pcDNA3.1 eukaryotic expression vector (Pharma Co., Shanghai, China), which was verified by restriction digestion followed by sequencing (Beijing Genomic Institute, Beijing, China) in our previous study (17). The microbial strain and plasmids used in the present study were *Escherichia coli* (*E. coli*), DH5 $\alpha$  and the pcDNA3.1-*c2orf68* eukaryotic expression vector. The plasmid extraction process was carried out by TIANprep Plasmid Mini kit introduction (cat. no. DP103-02; Tiangen Biotech Co., Ltd., Beijing, China).

Table I. Association of the expression of C2ORF68 with the clinicopathological parameters of the CRC samples.

Parameter	Expression intensity (n)				$\chi^2$	P-value
	+++	++	+	-		
Sex						
Female	16	25	11	2	1.568	0.698
Male	14	16	4	2		
Age, years						
≤60	13	11	8	2	4.283	0.233
>60	17	30	7	2		
TNM stage						
I	4	13	2	1	13.882	0.115
II	13	19	3	1		
III	12	9	10	2		
IV	1	0	0	0		
Differentiation degree						
Well	1	7	2	1	13.047	0.043
Moderate	19	32	10	3		
Low	10	2	3	0		
Lymph node metastasis						
Yes	13	9	10	2	10.343	0.013
No	17	32	5	2		
Survival status						
Surviving	15	25	10	3	1.310	0.727
Deceased	15	16	5	1		

CRC, colorectal cancer; TNM, tumor-node-metastasis.

The LoVo cells at a density of  $3 \times 10^5$  cells/well in a 6-well plate transfected with 5  $\mu$ l interference fragment or negative control (NC) vector using Lipofectamine 2000 which was then replaced with fresh growth medium after 6 h. Following culture for 48 h, the transfected LoVo cells were treated with 600 lg/ml of G418 (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany). After 14 days, the monoclonal cells were cultured in the presence of 300 lg/ml of G418.

**Co-immunoprecipitation (co-IP).** For co-IP, pcDNA3.1-c2orf68 eukaryotic expression vector was transfected for 60 h in SW620 cells. Then the cells with overexpression of *c2orf68* were lysed with mild lysis buffer containing several protease inhibitors for 30 min, and the cell lysates were separated by centrifugation at  $15,000 \times g$  for 20 min. Next, a modicum of cell lysates was reserved and utilized for western blot analysis. Next, the mouse anti-C2ORF68 Ab and rabbit anti-HuR Ab were added into the cell lysates for one night at 4°C. On the following day, 100  $\mu$ l protein A+G were added into the compound and incubated for 4 h. Subsequently, the sediments were gathered and the beads were washed twice using mild lysis buffer. After that, the same volume of 2X SDS-PAGE loading buffer was used to elute the protein which was absorbed on sepharose beads. Finally, the protein was used for SDS-PAGE.

**Cell proliferation assay.** The cell proliferation assay was performed as previously described (19). Plates were read at an absorbance wavelength of 450 nm with the help of a microplate reader (Model 680; Bio-Rad Laboratories, Hercules, CA, USA). Each transfection group had six replicates, and the experiment was repeated three times.

**Flow cytometry.** Flow cytometry was performed as previously described (19). Then, the results were analyzed by flow cytometry (FACS Aria II Cell Sorter; BD Biosciences, Franklin Lakes, NJ, USA). Each transfection group had three replicates, and the experiment was repeated three times.

**Cell migration assay.** Cell migration assay was performed as previously described (19). Migrated cells were quantified by counting the stained cells under a microscope (Model 680; Bio-Rad Laboratories) at x200 magnification. For each well, five random fields were selected to determine the total number of migrated cells. The assay was performed in triplicate and repeated three times.

**qRT-PCR analysis.** qRT-PCR analysis was performed as previously described (19). The sequences of forward and reverse primers are shown in Table II. The amplification was performed on a Bio-Rad C1000 Touch Thermal Cycler

Table II. Primer sequences for reverse transcription-quantitative polymerase chain reaction.

Gene	Primer sequence (5'→3')	Location	Product length, bp
<i>c2orf68</i>	F: GAAGAGTCCGGTCAAAGCAG R: TACGCAACTTGAGGGCTTCT	315-336 483-462	169
<i>HuR</i>	F: ACCCAGGATGAGTTACGA R: GCCCAAACCGAGAGAACAT	245-264 369-349	125
<i>Bax</i>	F: AAGCTGAGCGAGTGTCTCAAG R: CAAAGTAGAAAAGGGCGACAAC	172-192 349-328	178
<i>Bcl2</i>	F: GTTTGATTTCTCCTGGCTGTCTC R: GAACCTTTTGCATATTTGTTTGG	1-23 649-627	133
<i>c-Myc</i>	F: TCAAGAGGCGAACACACAAC R: GGCCTTTTCATTGTTTCCA	1,631-1,550 1,740-1,721	110
<i>Cyclin D1</i>	F: GTGGCTCTAAGATGAAGGAGA R: GGAAGTGTTCAATGAAATCGTG	534-555 702-681	169
<i>Cyclin A</i>	F: TGTCTCATGGACCTTCACCA R: CTCTGGTGGGTTGAGGAGAG	1,541-1,560 1,657-1,638	117
<i>GAPDH</i>	F: GGAAGGTGAAGGTCGGAGT R: TGAGGTCAATGAAGGGGTC	179-197 295-277	117

F, forward; R, reverse.

(Bio-Rad Laboratories). The GAPDH gene was used as an endogenous control, and the  $\Delta\Delta Cq$  (20) method was used to quantify the data, and the experiment was repeated three times.

**Western blot analysis.** Western blotting was performed as previously described (19). The primary antibodies used for western blotting were as follows: C2ORF68 (cat. no. ab81363; Abcam, Cambridge, UK), Bcl-2 (cat. no. sc-492; Santa Cruz Biotechnology, Inc., Dallas, TX, USA), Bax (cat. no. sc-623; Santa Cruz Biotechnology, Inc.), c-Myc (cat. no. ab32072; Abcam), (cat. no. ab200342; Abcam), cyclin D (cat. no. ab134175; Abcam), cyclin A (cat. no. ab181591; Abcam) and  $\beta$ -actin (cat. no. sc-8432; Santa Cruz Biotechnology, Inc.). All the primary antibodies used in this step at 1:1,000 dilution. Signals detection were performed through a Gel Imaging system (ProteinSimple, Santa Clara, CA, USA). Subsequent densitometry analysis were conducted using ImageJ (version 1.52g; National Institutes of Health, Bethesda, MD, USA).

**Statistical analysis.** Statistical analysis was performed by SPSS software (version 19.0; IBM Corp., Armonk, NY, USA). All experiments were performed three times, and all data are expressed as mean  $\pm$  SD. Student's t-test and one-way ANOVA were used for statistical analysis. Multiple comparison between the groups was performed using the S-N-K method. A P-value <0.05 was considered to indicate a statistically significant result.

## Results

From the BioGrid database, it is found that there are 12 proteins which may interact with C2ORF68, including KLHL15, GMNN, SMG6, GNE, DDHD2, PIR, ELAVL1(HuR), NAGK, XPO1, HSPA9, JOSD2 and GUK1 (Fig. 1).




Figure 1. Interactive proteins for C2ORF68.

**Expression of C2ORF68 in the rectal cancer tissue microarray.** The representative cytoplasmic staining of C2ORF68 in the rectal cancer tissue microarray is shown in Fig. 2A-C (magnification, x400). Our previous study (15) demonstrated that C2ORF68 presents two different staining patterns, including nuclear staining and cytoplasmic staining. In line with the observation in IF, C2ORF68 is a predominantly nuclear protein, but cytoplasmic C2ORF68 localization may play a vital role in the occurrence of CRC. In the rectal cancer tissue microarray (Fig. 2A), C2ORF68 protein expression was detected in 95.56% (86/90) of the cancer samples. Among these, 4.44% (4/90), 16.67% (15/90), 45.56% (41/90) and




Figure 2. Representative immunohistochemical expression patterns of c2orf68 in (A) rectal cancer tissues and (B) normal rectal tissues. (C) The immunostaining result of C2ORF68 in 90 rectal cancer and adjacent normal tissues.

33.33% (30/90) of these cases exhibited negative (-), weak (+), moderate (++) and strong (+++) C2ORF68 protein staining, respectively. In contrast, 5.56% (5/90), 25.56% (23/90), 58.89% (53/90) and 10% (9/90) of normal rectal specimens exhibited negative (-), weak (+), moderate (++) and strong (+++) C2ORF68 protein staining, respectively (Fig. 2C). We selected five nonoverlapping views randomly from each image and calculated the mean optical density. Then, the difference in mean optical density between rectal cancer and adjacent normal tissue was analyzed by statistical analysis. It was showed that compared with the adjacent normal rectal tissues, the expression of C2ORF68 was significantly increased in the rectal cancer tissues ( $P < 0.05$ ). In addition, we also researched the relationship between the expression of C2ORF68 and clinical parameters. Significant associations were noted between




Figure 3. (A) Double immunostaining for the indicated localization of c2orf68 (red) and HuR (green) in SW480 cells and LoVo cells. (B) Co-IP for direct interaction between C2ORF68 and HuR. co-IP, co-immunoprecipitation.

C2ORF68 expression and pathological grade ( $P < 0.05$ ) and lymph node metastasis ( $P < 0.05$ ). However, the association between C2ORF68 expression and age, sex and TNM stage was not statistically significant.

**Cellular localization of C2ORF68 and HuR in SW480 and LoVo cells.** The results revealed that C2ORF68 and HuR were localized mainly in the nucleus in both SW480 and LoVo cells (Fig. 3A) (magnification,  $\times 100$ ). BioGrid predicted that there is an interaction between C2ORF68 and HuR. It was shown that C2ORF68 interacts with HuR by Co-IP (Fig. 3B).

**Cell apoptosis, cell proliferation and cell migration in LoVo<sup>+c2orf68</sup>, HuR and LoVo<sup>+c2orf68</sup> cells.** Following transfection for 24 h, flow cytometry was performed to analyze c2orf68 and HuR-induced cell cycle arrest and apoptosis in colon cancer cells. The cell apoptosis rate of LoVo<sup>+c2orf68</sup> and LoVo<sup>+c2orf68</sup> cells was significantly decreased compared to that of the LoVo<sup>-NC</sup> and LoVo cells (Fig. 4B,  $P < 0.05$ ). There was no statistical significance noted between LoVo<sup>+c2orf68</sup>, HuR and LoVo<sup>+c2orf68</sup> cells, LoVo<sup>-NC</sup> and LoVo cells. Compared with the control group, LoVo<sup>+c2orf68</sup> and LoVo<sup>+c2orf68</sup>, HuR cell proliferation was significantly increased ( $P < 0.05$ ); and the cell proliferation of LoVo<sup>+c2orf68</sup> cells was statistically significantly different when compared with that of LoVo<sup>+c2orf68</sup>, HuR cells (Fig. 4C,  $P < 0.05$ ). Following transfection for 24 h, the number of migrated LoVo<sup>+c2orf68</sup>, HuR, LoVo<sup>+c2orf68</sup>, LoVo<sup>-NC</sup> and LoVo cells was  $223 \pm 24$ ,  $379 \pm 33$ ,  $239 \pm 31$  and  $244 \pm 26$ , respectively. The number of migrated LoVo<sup>+c2orf68</sup> cells was




Figure 4. (A) Cell migration was observably increased in LoVo<sup>+</sup>c2orf68 cells. The number of migrated LoVo<sup>+</sup>c2orf68 cells was significantly higher than that of the LoVo<sup>+</sup>c2orf68,-HuR cells, while there was no observable difference between LoVo<sup>+</sup>c2orf68,-HuR and LoVo cells. (B) In the LoVo<sup>+</sup>c2orf68 and LoVo<sup>+</sup>c2orf68,-HuR cells, cell apoptosis was significantly decreased ( $P<0.05$ ). There was no statistical significance noted between LoVo<sup>+</sup>c2orf68,-HuR and LoVo<sup>+</sup>c2orf68 cells. (C) In LoVo<sup>+</sup>c2orf68 and LoVo<sup>+</sup>c2orf68,-HuR cells, cell proliferation was increased. The cell proliferation of LoVo<sup>+</sup>c2orf68 cells was statistically significant when compared with that of LoVo<sup>+</sup>c2orf68,-HuR cells ( $P<0.05$ ). \* $P<0.05$ .




Figure 5. (A) Compared with the blank group, the number of migrated cells was observably inhibited in the SW480<sup>-</sup>c2orf68,-HuR, SW480<sup>-</sup>c2orf68 and SW480<sup>-</sup>HuR cells. However, the number of SW480<sup>-</sup>c2orf68,-HuR cells that migrated was observably less than that of SW480<sup>-</sup>c2orf68 and SW480<sup>-</sup>HuR cells ( $P<0.05$ ). (B) The apoptosis rates of SW480<sup>-</sup>c2orf68,-HuR, SW480<sup>-</sup>c2orf68 and SW480<sup>-</sup>HuR cells were significantly increased ( $P<0.05$ ). (C) Compared with the blank group, cell proliferation was significantly decreased in the SW480<sup>-</sup>c2orf68,-HuR, SW480<sup>-</sup>c2orf68 and SW480<sup>-</sup>HuR cells ( $P<0.05$ ). Furthermore, the proliferation of SW480<sup>-</sup>c2orf68,-HuR cells was significantly lesser than that of SW480<sup>-</sup>c2orf68 and SW480<sup>-</sup>HuR cells ( $P<0.05$ ). \* $P<0.05$ .




Figure 6. (A and B) The protein expression levels of C2ORF68, Bcl-2, c-Myc, cyclin D and cyclin A were overexpressed in LoVo<sup>+c2orf68, -HuR</sup> and LoVo<sup>+c2orf68</sup> cells; while HuR was overexpressed in LoVo<sup>+c2orf68</sup> cells when compared to the LoVo cells. Compared with the LoVo<sup>+c2orf68, -HuR</sup> cells, the protein expression of C2ORF68, HuR, Bcl-2, c-Myc, cyclin D and cyclin A in LoVo<sup>+c2orf68</sup> cells was significantly increased. The protein expression level of Bax was decreased in LoVo<sup>+c2orf68, -HuR</sup> and LoVo<sup>+c2orf68</sup> cells when compared to the LoVo cells. Compared with LoVo<sup>+c2orf68, -HuR</sup> cells, the protein expression of Bax in LoVo<sup>+c2orf68</sup> cells was significantly decreased. (C) In LoVo<sup>+c2orf68, -HuR</sup> and LoVo<sup>+c2orf68</sup> cells, the mRNA levels of c2orf68, HuR, Bcl-2, c-Myc were significantly overexpressed in both LoVo<sup>+c2orf68</sup> and LoVo<sup>+c2orf68, -HuR</sup> cells. Cyclin D and cyclin A were significantly overexpressed in LoVo<sup>+c2orf68</sup> cells, while the upregulation of cyclin D and cyclin A mRNA expression levels in LoVo<sup>+c2orf68, -HuR</sup> cells exhibited no significant difference. \*P<0.05, \*\*P<0.01.

significantly higher than that of LoVo<sup>+c2orf68, -HuR</sup>, LoVo<sup>-NC</sup> and LoVo cells (Fig. 4A, P<0.05).

**Cell apoptosis, cell proliferation and cell migration of SW480<sup>-c2orf68, -HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells.** The apoptosis rate of SW480<sup>-c2orf68, -HuR</sup> cells was significantly higher than that of SW480<sup>-c2orf68</sup>, SW480<sup>-HuR</sup>, SW480<sup>-NC</sup> and SW480 cells (Fig. 5B, P<0.05). In addition, the apoptosis rate of SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells was significantly higher than that of SW480<sup>-NC</sup> and SW480 cells, respectively (Fig. 5B, P<0.05). Compared with the control group, cell proliferation was significantly decreased in the SW480<sup>-c2orf68, -HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells (Fig. 5C, P<0.05); cell proliferation in the SW480<sup>-c2orf68, -HuR</sup> cells was significantly less than that of the SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells (Fig. 5C, P<0.05). The number of migrated cells of SW480<sup>-c2orf68, -HuR</sup>, SW480<sup>-c2orf68</sup>, SW480<sup>-HuR</sup>, SW480<sup>-NC</sup> and SW480 were 122±16, 234±51, 233±48, 381±42 and 401±24, respectively. Compared with the control group, the number of migrated cells were significantly inhibited in the SW480<sup>-c2orf68, -HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells (Fig. 5A, P<0.05). However, the number of migrated SW480<sup>-c2orf68, -HuR</sup> cells was significantly lower than that of the SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells (Fig. 5A, P<0.05).

**mRNA and protein expression of C2orf68, HuR, Bcl-2, Bax, c-Myc, cyclin D and cyclin A in LoVo<sup>+c2orf68, -HuR</sup> and LoVo<sup>+c2orf68</sup> cells.** As shown in Fig. 6C, following transfection for 48 h, c2orf68 gene expression was significantly overexpressed in the LoVo<sup>+c2orf68</sup> (P<0.01) and LoVo<sup>+c2orf68, -HuR</sup> cells (P<0.05) when compared with the control group. Similarly, HuR, Bcl-2 and c-Myc

were significantly overexpressed in the LoVo<sup>+c2orf68</sup> (P<0.01, P<0.01 and P<0.05, respectively) and LoVo<sup>+c2orf68, -HuR</sup> cells (all P<0.05). Cyclin D and Cyclin A were significantly overexpressed in the LoVo<sup>+c2orf68</sup> cells (P<0.05), while the upregulation of Cyclin D and Cyclin A mRNA expression levels in LoVo<sup>+c2orf68, -HuR</sup> cells exhibited no significance. The mRNA expression of c2orf68, HuR, Bcl-2, c-Myc and Cyclin A in LoVo<sup>+c2orf68</sup> cells was significantly increased, while cyclin D in LoVo<sup>+c2orf68, -HuR</sup> cells exhibited no statistical difference with LoVo<sup>+c2orf68</sup> cells. In contrast, the mRNA expression level of Bax was decreased in the LoVo<sup>+c2orf68</sup> (P<0.05) and LoVo<sup>+c2orf68, -HuR</sup> (P<0.01) cells, when compared to the control. Compared with the LoVo<sup>+c2orf68, -HuR</sup> cells, the mRNA expression of Bax in the LOVO<sup>+c2orf68</sup> cells was significantly decreased (P<0.05).

As shown in Fig. 6A and B, following transfection for 48-72 h, compared to the blank group, the protein expression level of C2ORF68 was significantly increased in the LoVo<sup>+c2orf68, -HuR</sup> (P<0.05) and LoVo<sup>+c2orf68</sup> (P<0.01) cells. In addition, HuR, BCL-2, C-MYC and Cyclin A protein expression levels were overexpressed in the LoVo<sup>+c2orf68, -HuR</sup> (NS, P<0.05, P<0.01 and P<0.01, respectively) and LoVo<sup>+c2orf68</sup> cells (P<0.05, P<0.01, P<0.01 and P<0.01, respectively) when compared to the blank group; compared with the LoVo<sup>+c2orf68, -HuR</sup> cells, the protein expression of C2ORF68, HuR, BCL-2, C-MYC and Cyclin A in LoVo<sup>+c2orf68</sup> cells was increased (P<0.05, P<0.05, P<0.01, P<0.01 and P<0.01, respectively). In contrast, the protein expression level of BAX was decreased in the LoVo<sup>+c2orf68, -HuR</sup> (P<0.05) and LoVo<sup>+c2orf68</sup> cells (P<0.01). Compared with LoVo<sup>+c2orf68, -HuR</sup> cells, the protein expression of BAX (P<0.01) in LoVo<sup>+c2orf68</sup> cells was significantly decreased.




Figure 7. (A and B) The protein expression levels of C2ORF68, HuR, Bcl-2, c-Myc, cyclin D and cyclin A were significantly decreased. The inhibition rate of C2ORF68, HuR, Bcl-2, c-Myc, cyclin D and cyclin A was significantly higher in the SW480-*c2orf68*, HuR cells than in SW480-*c2orf68* and SW480-*HuR* cells. However, the inhibition rate of CYCLIND was not statistically significant in SW480-*c2orf68*, HuR cells, compared with SW480-*c2orf68* and SW480-*HuR* cells. Meanwhile, the protein expression level of BAX significantly increased in SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells. (C) In the SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells, the mRNA expression levels of *c2orf68*, HuR, Bcl-2, c-Myc, cyclin D and cyclin A were significantly decreased. Furthermore, the inhibition rate of *c2orf68*, HuR, Bcl 2 and cyclin A in SW480-*c2orf68*, HuR cells was significantly higher than that in the SW480-*c2orf68* and SW480-*HuR* cells. The inhibition rate of cyclin D in SW480-*c2orf68*, HuR cells compared with SW480-*HuR* cells, was statistically significant. However, compared with SW480-*c2orf68* cells, the inhibition of cyclin D was not statistically significant. Furthermore, the mRNA expression level of Bax was significantly increased in the SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells ( $P<0.01$ ,  $P<0.01$ ,  $P<0.01$ ), and the expression of BAX was also increased in the SW480-*c2orf68*, HuR cells when compared with the SW480-*c2orf68* ( $P<0.05$ ) and SW480-*HuR* ( $P<0.05$ ) cells. \* $P<0.05$ , \*\* $P<0.01$ , \*\*\* $P<0.001$ .

*mRNA and protein expression of c2orf68, HuR, Bcl-2, Bax, c-Myc, cyclin D and cyclin A in SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells.* As shown in Fig. 7C, in SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells, the mRNA expression level of *c2orf68*, HuR, Bcl-2, c-Myc, cyclin D and cyclin A significantly decreased ( $P<0.05$ ). Furthermore, the inhibition rate of *c2orf68*, HuR, Bcl-2, c-Myc and cyclin A in SW480-*c2orf68*, HuR cells was significantly higher than that in SW480-*c2orf68* and SW480-*HuR* cells ( $P<0.05$ ). The inhibition rate of cyclin D in SW480-*c2orf68*, HuR cells compared with SW480-*HuR* cells, was statistically significant ( $P<0.05$ ). However, compared with SW480-*c2orf68* cells, the inhibition of cyclin D was not statistically significant. Meanwhile, the increase in Bax mRNA expression was significantly higher in SW480-*c2orf68*, HuR cells than in SW480-*c2orf68* and SW480-*HuR* cells ( $P<0.05$ ). Furthermore, the mRNA expression level of Bax was increased in SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells.

As shown in Fig. 7A and B, following transfection for 48-72 h, compared to the blank group, the protein expression level of C2ORF68 was significantly decreased in the SW480-*c2orf68*, HuR ( $P<0.01$ ), SW480-*c2orf68* ( $P<0.01$ ) and SW480-*HuR* cells ( $P<0.01$ ). Similarly, the protein expression levels of HuR, BCL-2, C-MYC, Cyclin D and Cyclin A were also decreased in the SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells. Furthermore, the inhibition rate of C2ORF68 ( $P<0.01$ ,  $P<0.01$ ), HuR ( $P<0.05$ ,  $P<0.05$ ), C-MYC ( $P<0.05$ ,  $P<0.01$ ) and Cyclin A ( $P<0.05$ ,  $P<0.05$ ) was

significantly higher in the SW480-*c2orf68*, HuR cells than that in the SW480-*c2orf68* and SW480-*HuR* cells. However, the inhibition rate of Cyclin D in SW480-*c2orf68*, HuR cells, compared with SW480-*c2orf68* and SW480-*HuR* cells, was not statistically significant. Meanwhile, the protein expression level of BAX was significantly increased in the SW480-*c2orf68*, HuR, SW480-*c2orf68* and SW480-*HuR* cells when compared with the blank group. Compared with the SW480-*c2orf68* and SW480-*HuR* cells, the protein expression level of BAX was significantly higher in the SW480-*c2orf68*, HuR cells ( $P<0.01$ ,  $P<0.01$ ).

## Discussion

In the present study, it was shown that the expression level of C2ORF68 was significantly upregulated in rectal cancer tissues compared with its adjacent normal tissues by IHC. This indicates that *c2orf68* may be involved in the occurrence of rectal cancer. In addition, upregulated expression of C2ORF68 was significantly correlated with a variety of important clinicopathological parameters, including pathological grade and lymph node metastasis. It was suggested that C2ORF68 may play a role in the development and metastasis of CRC. As a result, a statistical significance was found between the expression of C2ORF68 and pathological grade. This indicates that the expression level of C2ORF68 may be associated with the malignant potential of cancer. That is, the higher the expression level of C2ORF68, the higher is the degree of malignancy of

rectal carcinoma. These present results suggest that the *c2orf68* gene is associated with the occurrence and development of rectal cancer and may be a potential carcinogenic factor in rectal cancer. However, the detailed mechanism for *c2orf68* upregulation in rectal cancer remains to be clarified.

Through bioinformatic analyses, we discovered that there are 12 proteins which interact with C2ORF68, including KLHL15, GMNN, SMG6, GNE, DDHD2, PIR, ELAVL1 (HuR), NAGK, XPO1, HSPA9, JOSD2 and GUK1 (BioGrid database). At the same time, by our experiments, including IF and Co-IP, we verified that C2ORF68 co-localized with HuR in SW480 and LoVo cell lines, and C2ORF68 could interact with HuR. HuR, a member of the Hu/ELAV family, is predominantly located in the nucleus and translocates to the cytoplasm when cells are stimulated by endogenous factors or external stimuli (21,22). HuR (ELAV1) is an RNA binding protein, which has been shown to regulate the expression of multiple genes by different post-transcriptional mechanisms, such as mRNA decay and protein translation (23). HuR modulates posttranscriptional processing of target premRNAs or mRNA stabilization and translation through interaction with AU-rich elements (ARE) within 3'-untranslated regions (UTRs) of the target mRNAs to transformation (24). Furthermore, it has been shown that HuR stabilizes mRNAs that encode *p53* and *WEE1* (25), activates *ATF2* (26), *Jun D* (27) and *XIAP* (28) and enhances the translation of mRNAs that encode *c-Myc* (29), *ICH-1* (30) and *IL-1 $\beta$*  (31). Many of these transcripts are reported to participate in certain key cellular processes including cell proliferation, cell apoptosis, angiogenesis, immune response and metastasis. HuR is also increased in malignant cells when compared with corresponding normal cells, and it has been found to be associated with adverse clinicopathological factors in several different cancer types, such as gastric, gallbladder breast, urothelial and non-small cell lung cancer (32).

Our study revealed that cell apoptosis increased, cell proliferation and cell migration decreased when *c2orf68* was inhibited in SW480 cells. These results are consistent with our previous study (16). In addition, we also demonstrated that cell apoptosis increased while cell proliferation and cell migration decreased in SW480<sup>-HuR</sup> and SW480<sup>-c2orf68,-HuR</sup> cells. This indicates that both *c2orf68* and HuR can regulate cell apoptosis and proliferation in CRC cells. The cell apoptosis rate, cell proliferation and cell migration in SW480<sup>-c2orf68,-HuR</sup> cells which has more significant results than that in SW480<sup>-HuR</sup> cells revealed that *c2orf68* and HuR may have a synergistic effect in regulating cell apoptosis, cell proliferation and cell migration. This study differed from our previous study (16), which focused on the PI3K/Akt/mTOR signaling pathway and its downstream molecules, such as *Akt*, *PI3K*, *Bcl-2*, *c-Myc*, *cyclin D1* and *bax* when *c2orf68* was inhibited. A recent study showed that the HuR expression level is closely related to AKT phosphorylation and PI3K/AKT/NF- $\kappa$ B signaling can notably elevate HuR gene transcription (18). This study focused on the relationship between C2ORF68 and HuR and the downstream molecules of HuR, such as *Bcl-2*, *Bax*, *c-Myc*, *cyclin D* and *cyclin A* in SW480<sup>-c2orf68,-HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells. According to our results, *Bcl-2*, *c-Myc*, *cyclin D* and *Cyclin A* decreased, and *Bax* increased in the SW480<sup>-c2orf68,-HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells. In addition, *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* was significantly decreased and Bax was

significantly increased in SW480<sup>-c2orf68,-HuR</sup> cells. This shows that *c2orf68* and HuR may co-regulate *Bcl-2*, *c-Myc*, *cyclin D*, *cyclin A* and *Bax*, resulting in the cell apoptosis and cell proliferation CRC cells.

In the present study, the mRNA and protein expression of HuR was downregulated when *c2orf68* was inhibited in SW480 cells, and the mRNA and protein expression of HuR was upregulated when *c2orf68* was overexpressed in LoVo cells. Furthermore, when HuR was inhibited, the mRNA and protein expression levels of *c2orf68* were also decreased. That is, HuR and *c2orf68* had a synergistic effect.

The present study revealed that cell apoptosis decreased while cell proliferation and cell migration were increased in LoVo<sup>+c2orf68</sup> cells. These results are consistent with our previous study (17), and it was confirmed that *c2orf68* can regulate cell apoptosis and proliferation. In addition, it was also revealed that cell apoptosis was decreased when cell proliferation and cell migration were increased in LoVo<sup>+c2orf68,-HuR</sup> cells. However, cell apoptosis was significantly lower and cell proliferation and cell migration were significantly higher in LoVo<sup>+c2orf68</sup> cells, compared with LoVo<sup>+c2orf68,-HuR</sup> cells. All these results suggest again that *c2orf68* and HuR may have a synergistic effect in promoting cell proliferation and migration, and in inhibiting cell apoptosis in the role of CRC. Unlike our previous study, which focused on the Wnt signaling pathway and its molecules such as  $\beta$ -catenin, survivin, *cyclin D1*, *c-Myc* and *GSK-3 $\beta$*  (17), the present study focused on the relationship between C2ORF68 and HuR and the downstream molecules of HuR such as *Bcl-2*, *Bax*, *c-Myc*, *cyclin D* and *cyclin A*. In LoVo<sup>+c2orf68</sup> and LoVo<sup>+c2orf68,-HuR</sup> cells, *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* increased, while *Bax* decreased. In addition, *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* were significantly upregulated and *Bax* was significantly decreased in LoVo<sup>+c2orf68</sup> cells, compared with LoVo<sup>+c2orf68,-HuR</sup> cells. Previously in this manuscript, we described that *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* were decreased, while *Bax* was increased in the SW480<sup>-c2orf68,-HuR</sup>, SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells. In addition, *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* were significantly decreased and Bax was significantly increased in the SW480<sup>-c2orf68,-HuR</sup> cells, compared with the SW480<sup>-c2orf68</sup> and SW480<sup>-HuR</sup> cells. It is known that all these genes are involved in key cellular processes such as cell proliferation and cell cycle. The expression of cyclin D1 is increased in many tumors and it promotes cell proliferation by regulating cell cycle progression through the G1/S restriction point (33). The transcription factor c-Myc, which is a leucine zipper protein regulating the expression of 10-15% of human genes, plays an important role in cell proliferation, differentiation, growth and survival. Its overexpression is associated with cancer occurrence and development (34). BCL-2, an integral outer mitochondrial membrane protein, serving as an anti-apoptosis protein, belongs to the Bcl-2 family, and was firstly discovered in B cell malignancies and regulates the intrinsic mitochondrial apoptosis pathway (35). Activated Bax then oligomerizes at the mitochondria to induce outer mitochondrial membrane (OMM) permeabilization and releases into the cytosol apoptotic factors which promote caspase activation and subsequent apoptosis execution (36). Cyclins are fundamental regulators of the cell cycle, playing an important role in tumorigenesis, Cyclin A is required for cells to progress through the S phase (37). As a result, it is believed that *c2orf68*

and *HuR* may have a synergistic effect and may co-regulate cell proliferation and apoptosis by regulating the downstream molecules of *HuR*, such as upregulating *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* gene expression and downregulating *Bax* gene expression, resulting in the development of cancer.

In conclusion, the *c2orf68* gene may be a potential oncogene and may play a potential carcinogenic effect in the pathogenesis of colorectal cancer. Furthermore, it may be associated with lymph node metastasis of colorectal cancer. The carcinogenesis of *c2orf68* may be related to the promotion of cell proliferation and inhibition of cell apoptosis. *C2orf68* may have a synergistic effect with *HuR*, and the possible mechanism of *c2orf68* and *HuR* involves the co-promotion of cell proliferation and migration, and the co-inhibition of cell apoptosis. However, the exact mechanism remains mysterious. *C2orf68* and *HuR* may co-regulate cell proliferation and apoptosis by upregulating *Bcl-2*, *c-Myc*, *cyclin D* and *cyclin A* gene expression and downregulating *Bax* gene expression, resulting in the development of colorectal cancer. Our previous study indicated that *C2ORF68* can regulate cancer cell proliferation and apoptosis through PI3K/AKT/mTOR signaling (16). At the same time, a recent study (18) showed that the *HuR* expression level is closely related to AKT phosphorylation and cytoplasmic abundance of *HuR* in human cancer may be associated with oncogenic activation of AKT signaling. According to the above conclusions, we may hypothesize that *C2ORF68*, *HuR* and AKT signaling could make up a mutually reinforcing loop, regulating cancer cell proliferation and apoptosis. However, the specific mechanism warrants further research.

## Acknowledgements

I need to express my sincere gratitude to all authors for the assistance with this article, especially to Professor Yao Chen, whose constant encouragement directed me through all stages of the writing of this manuscript. Without her help, I could not have gotten to this point. Furthermore, I also extend my heartfelt gratitude to the other authors, and thanks for their efforts.

## Funding

The present study was supported by Chengdu Department of Science and Technology (2018-YF05-00-038-SN), Sichuan Province, China.

## Availability of data and materials

The datasets used during the present study are available from the corresponding author upon reasonable request.

## Authors' contributions

LS, KS, TJ and YC conceived and designed the study. ZL and KH performed the experiments. ZL and KH wrote the paper. LS, KS, TJ and YC reviewed and edited the manuscript. All authors read and approved the manuscript and agree to be accountable for all aspects of the research in ensuring that the accuracy or integrity of any part of the work are appropriately investigated and resolved.

## Ethics approval and consent to participate

The Medical Research Ethics Committee of Sichuan University approved the sample acquisition (Chengdu, China), and a written informed consent was also obtained from all patients.

## Patient consent for publication

Not applicable.

## Competing interests

The authors state that they have no competing interests.

## References

1. Siegel RL, Miller KD and Jemal A: Cancer statistics, 2017. *CA Cancer J Clin* 67: 7-30, 2017.
2. Lin J, Chuang CC and Zuo L: Potential roles of microRNAs and ROS in colorectal cancer: Diagnostic biomarkers and therapeutic targets. *Oncotarget* 8: 17328-17346, 2017.
3. Fang Y, Liang X, Jiang W, Li J, Xu J and Cai X: Cyclin B1 suppresses colorectal cancer invasion and metastasis by regulating e-cadherin. *PLoS One* 10: e0126875, 2015.
4. Schreuders EH, Ruco A, Rabeneck L, Schoen RE, Sung JJ, Young GP and Kuipers EJ: Colorectal cancer screening: A global overview of existing programmes. *Gut* 64: 1637-1649, 2015.
5. Jeyakumar A, Dissabandara L and Gopalan V: A critical overview on the biological and molecular features of red and processed meat in colorectal carcinogenesis. *J Gastroenterol* 52: 407-418, 2017.
6. Sakamoto N, Feng Y, Stolfi C, Kurosu Y, Green M, Lin J, Green ME, Sentani K, Yasui W, McMahon M, *et al*: BRAF<sup>V600E</sup> cooperates with CDX2 inactivation to promote serrated colorectal tumorigenesis. *Elife* 6: e20331, 2017.
7. Raskov H, Pommergaard HC, Burchard J and Rosenberg J: Colorectal carcinogenesis-update and perspectives. *World J Gastroenterol* 20: 18151-18164, 2014.
8. Lech G, Słotwiński R, Słodkowski M and Krasnodebski IW: Colorectal cancer tumour markers and biomarkers: Recent therapeutic advances. *World J Gastroenterol* 22: 1745-1755, 2016.
9. Huang L, Wang X, Wen C, Yang X, Song M, Chen J, Wang C, Zhang B, Wang L, Iwamoto A, *et al*: Hsa-miR-19a is associated with lymph metastasis and mediates the TNF- $\alpha$  induced epithelial-to-mesenchymal transition in colorectal cancer. *Sci Rep* 5: 13350, 2015.
10. Kurebayashi H, Goi T, Shimada M, Tagai N, Naruse T, Nakazawa T, Kimura Y, Hirono Y and Yamaguchi A: Prokineticin 2 (PROK2) is an important factor for angiogenesis in colorectal cancer. *Oncotarget* 6: 26242-26251, 2015.
11. Ingebrigtsen VA, Boye K, Nesland JM, Nesbakken A, Flatmark K and Fodstad Ø: B7-H3 expression in colorectal cancer: Associations with clinicopathological parameters and patient outcome. *BMC Cancer* 14: 602, 2014.
12. Yamaguchi K, Yamaguchi R, Takahashi N, Ikenoue T, Fujii T, Shinozaki M, Tsurita G, Hata K, Niida A, Imoto S, *et al*: Overexpression of cohesion establishment factor DSCC1 through E2F in colorectal cancer. *PLoS One* 9: e85750, 2014.
13. Thomas J, Ohtsuka M, Pichler M and Ling H: MicroRNAs: Clinical relevance in colorectal cancer. *Int J Mol Sci* 16: 28063-28076, 2015.
14. Lam SY, Yu J, Wong SH, Peppelenbosch MP and Fuhler GM: The gastrointestinal microbiota and its role in oncogenesis. *Best Pract Res Clin Gastroenterol* 31: 607-618, 2017.
15. Wang K and Chen Y: Analysis of a novel protein in human colorectal adenocarcinoma. *Mol Med Rep* 8: 529-534, 2013.
16. Wen X, Zhu J, Dong L and Chen Y: The role of *c2orf68* and PI3K/Akt/mTOR pathway in human colorectal cancer. *Med Oncol* 31: 92, 2014.
17. Chen Y, Jiang T, Shi L and He K: *hcrn81* promotes cell proliferation through Wnt signaling pathway in colorectal cancer. *Med Oncol* 33: 3, 2016.

18. Kang MJ, Ryu BK, Lee MG, Han J, Lee JH, Ha TK, Byun DS, Chae KS, Lee BH, Chun HS, *et al*: NF-kappaB activates transcription of the RNA-binding factor HuR, via PI3K-AKT signaling, to promote gastric tumorigenesis. *Gastroenterology* 135: 2030-2042.e1-e3, 2008.
19. He K, Shi L, Jiang T, Li Q, Chen Y and Meng C: Association between SET expression and glioblastoma cell apoptosis and proliferation. *Oncol Lett* 12: 2435-2444, 2016.
20. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* 25: 402-408, 2001.
21. Woodhoo A, Iruarizaga-Lejarreta M, Beraza N, García-Rodríguez JL, Embade N, Fernández-Ramos D, Martínez-López N, Gutiérrez-De Juan V, Arteta B, Caballeria J, *et al*: Human antigen R contributes to hepatic stellate cell activation and liver fibrosis. *Hepatology* 56: 1870-1882, 2012.
22. Zhang LF, Lou JT, Lu MH, Gao C, Zhao S, Li B, Liang S, Li Y, Li D and Liu MF: Suppression of miR-199a maturation by HuR is crucial for hypoxia-induced glycolytic switch in hepatocellular carcinoma. *EMBO J* 34: 2671-2685, 2015.
23. Jakstaite A, Maziukienė A, Silkunienė G, Kmieliute K, Gulbinas A and Dambrauskas Z: HuR mediated post-transcriptional regulation as a new potential adjuvant therapeutic target in chemotherapy for pancreatic cancer. *World J Gastroenterol* 21: 13004-13019, 2015.
24. Zhu H, Berkova Z, Mathur R, Sehgal L, Khashab T, Tao RH, Ao X, Feng L, Sabichi AL, Blechacz B, *et al*: HuR suppresses fas expression and correlates with patient outcome in liver cancer. *Mol Cancer Res* 13: 809-819, 2015.
25. Lal S, Burkhart RA, Beeharry N, Bhattacharjee V, Londin ER, Cozzitorto JA, Romeo C, Jimbo M, Norris ZA, Yeo CJ, *et al*: HuR posttranscriptionally regulates WEE1: Implications for the DNA damage response in pancreatic cancer cells. *Cancer Res* 74: 1128-1140, 2014.
26. Xiao L, Rao JN, Zou T, Liu L, Marasa BS, Chen J, Turner DJ, Zhou H, Gorospe M and Wang JY: Polyamines regulate the stability of activating transcription factor-2 mRNA through RNA-binding Protein HuR in intestinal epithelial cells. *Mol Biol Cell* 18: 4579-4590, 2007.
27. Zou T, Rao JN, Liu L, Xiao L, Yu TX, Jiang P, Gorospe M and Wang JY: Polyamines regulate the stability of JunD mRNA by modulating the competitive binding of its 3' untranslated region to HuR and AUF1. *Mol Cell Biol* 30: 5021-5033, 2010.
28. Zhang X, Zou T, Rao JN, Liu L, Xiao L, Wang PY, Cui YH, Gorospe M and Wang JY: Stabilization of XIAP mRNA through the RNA binding protein HuR regulated by cellular polyamines. *Nucleic Acids Res* 42: 4143, 2014.
29. Liu L, Rao JN, Zou T, Xiao L, Wang PY, Turner DJ, Gorospe M and Wang JY: Polyamines regulate c-Myc translation through Chk2-dependent HuR phosphorylation. *Mol Biol Cell* 20: 4885-4898, 2009.
30. Winkler C, Doller A, Imre G, Badawi A, Schmid T, Schulz S, Steinmeyer N, Pfeilschifter J, Rajalingam K and Eberhardt W: Attenuation of the ELAV1-like protein HuR sensitizes adenocarcinoma cells to the intrinsic apoptotic pathway by increasing the translation of caspase-2L. *Cell Death Dis* 5: e1321, 2014.
31. Aguado A, Rodríguez C, Martínez-Revelles S, Avendaño MS, Zhenyukh O, Orriols M, Martínez-González J, Alonso MJ, Briones AM, Dixon DA and Salaices M: HuR mediates the synergistic effects of angiotensin II and IL-1 $\beta$  on vascular COX-2 expression and cell migration. *Br J Pharmacol* 172: 3028-3042, 2015.
32. Elebro J, Ben Dror L, Heby M, Nodin B, Jirstrom K and Eberhard J: Prognostic effect of hENT1, dCK and HuR expression by morphological type in periampullary adenocarcinoma, including pancreatic cancer. *Acta Oncol* 55: 286-296, 2015.
33. Li X, Zhang Q, Fan K, Li B, Li H, Qi H, Guo J, Cao Y and Sun H: Overexpression of *TRPV3* correlates with tumor progression in non-small cell lung cancer. *Int J Mol Sci* 17: 437, 2016.
34. Sharma T, Bansal R, Haokip DT, Goel I and Muthuswami R: SMARCA11 negatively regulates C-Myc transcription by altering the conformation of the promoter region. *Sci Rep* 9: 17910, 2015.
35. Choi JE, Kang SH, Lee SJ and Bae YK: Prognostic significance of Bcl-2 expression in non-basal triple-negative breast cancer patients treated with anthracycline-based chemotherapy. *Tumor Biol* 35: 12255-12263, 2014.
36. Ouyang YB and Giffard RG: microRNAs affect BCL-2 family proteins in the setting of cerebral ischemia. *Neurochem Int* 77: 2-8, 2014.
37. Kokontis JM, Lin HP, Jiang SS, Lin CY, Fukuchi J, Hiipakka RA, Chung CJ, Chan TM, Liao S, Chang CH, *et al*: Androgen suppresses the proliferation of androgen receptor-positive castration-resistant prostate cancer cells via inhibition of Cdk2, CyclinA, and Skp2. *PLoS One* 9: e109170, 2014.