

Diagnostic and prognostic value of mRNA expression of phospholipase C β family genes in hepatitis B virus-associated hepatocellular carcinoma

XIANGKUN WANG, KETUAN HUANG, XIANMIN ZENG, ZHENGQIAN LIU, XIWEN LIAO, CHENGKUN YANG, TINGDONG YU, CHUANGYE HAN, GUANGZHI ZHU, WEI QIN and TAO PENG

Department of Hepatobiliary Surgery, The First Affiliated Hospital of Guangxi Medical University, Nanning, Guangxi Zhuang Autonomous Region 530021, P.R. China

Received August 15, 2018; Accepted March 5, 2019

DOI: 10.3892/or.2019.7066

Abstract. Four phospholipase C β (PLCB) isoforms, *PLCB1*, *PLCB2*, *PLCB3* and *PLCB4*, have been previously investigated regarding their roles in the metabolism of inositol lipids and cancer. The present study aimed to explore the association between *PLCB1-4* and hepatocellular carcinoma (HCC). Data from 212 patients with hepatitis B virus-associated HCC were used to analyze the diagnostic and prognostic significance of *PLCB* genes in. A nomogram predicted the survival probability. Gene set enrichment analysis explored gene ontology terms and the metabolic pathways associated with *PLCB* genes. Validation of the prognostic values of *PLCB* genes was performed using the Gene Expression Profiling Interactive Analysis website. *PLCB1* and *PLCB2* were revealed to have diagnostic value for HCC (0.869 and 0.836 area under the curve, respectively; both $P \leq 0.05$). The combination analysis of these genes had an advantage over each alone (0.905 *PLCB1* and *PLCB2*, and 0.877 *PLCB1* and *PLCB3* area under the curve; $P \leq 0.05$). *PLCB1* was associated with overall survival (OS) and recurrence-free survival (RFS; adjusted $P=0.002$ and $P=0.001$, respectively). A nomogram predicted survival

probability of patients with HCC at 1, 3- and 5-years. Gene set enrichment analysis indicated that *PLCB1* and *PLCB2* are involved in the cell cycle, cell division and the PPAR signaling pathway, among other functions. Validation using GEPIA revealed that *PLCB1* and *PLCB2* were associated with OS and *PLCB1* and *PLCB4* were associated with RFS. *PLCB1* and *PLCB2* exhibited diagnostic value for HCC and their combination had an advantage over each individually. *PLCB1* has OS and RFS prognostic value for patients with HCC.

Introduction

Hepatocellular carcinoma (HCC) is one of the main causes of tumor-associated mortality, being the fifth most common malignancy worldwide (1). In 2018, the number of new cases and liver cancer-associated mortalities was 841,080 and 781,631, respectively, worldwide (2). Many risk factors, including dietary aflatoxin exposure (3), hepatitis B and C virus (HBV) infection (4) and cirrhosis, contribute to the initiation and progression of HCC. To date, many diagnostic and treatment procedures, including ultrasound, computed tomography, liver resection, liver transplantation, radiofrequency, thermal and non-thermal ablation, trans-arterial chemoembolization (5), immunotherapies and therapeutic cancer vaccines (4), have been used for patients with HCC. However, the prognosis of HCC is remains poor and the 5-year relative survival rate is ~12% due to tumor metastasis and recurrence (6,7). Due to the characteristics of systemic disease, the evolution and progression of HCC involves deregulation of genes, cells and tissues (1). Therefore, it is crucial to identify novel biomarkers that may be involved in the course of tumor metastasis and recurrence, for early diagnosis and recurrence prediction for HCC.

Phospholipase C (PLC) is encoded by four genes, *PLCA*, *PLCB*, *PLCC* and *PLCD*, and is involved in the pathogenesis of several bacterial infections, including *Clostridium perfringens*, *Listeria monocytogenes*, and *Pseudomonas aeruginosa* (8,9). The activity of *PLCA* and *PLCB* in *L. monocytogenes* appears to overlap in the course of intracellular infection (10). In *Listeria*, three genes, *PLCA*, *PLCB* and *PLCC*, are clustered together on the same chromosome, whereas the *PLCD* gene

Correspondence to: Dr Professor Tao Peng, Department of Hepatobiliary Surgery, The First Affiliated Hospital of Guangxi Medical University, 22 Shuangyong Road, Nanning, Guangxi Zhuang Autonomous Region 530021, P.R. China
E-mail: pengtaogmu@163.com

Abbreviations: HCC, hepatocellular carcinoma; *PLCB*, phospholipase C β ; HBV, hepatitis B virus; GEO, gene expression omnibus; GEPIA, gene expression profiling interactive analysis; GSEA, gene set enrichment analysis; BP, biological process; CC, cellular component; MF, molecular function; GO, gene ontology; OS, overall survival; RFS, recurrence-free survival; MST, median survival time; CI, confidence interval; HR, hazard ratio; ROC, receiver operating characteristic; PPI, protein-protein interaction

Key words: hepatocellular carcinoma, diagnosis, prognosis, mRNA expression, *PLCB1*, *PLCB2*

is located in another region (11,12). Under the transcriptional control of PrfA regulator, *PLCA*, *PLCB* and *HLY* (encoding listeriolysin O precursor) have a role encoding the *Listeria* Pathogenicity Island 1, leading to the escape from endocytic and secondary vacuoles (13-15). *PLCB* isoforms in mice include *PLCB1*, *PLCB2*, *PLCB3* and *PLCB4*, which are stimulated by G protein activation ($G\alpha_{q/11}$ and/or $G\beta\gamma$) (16,17). The roles of *PLCB* isoforms in immune defense and escape, and their functions in tumors are currently being investigated. *PLCB1* has been reported to be associated with HCC prognosis in tumor proliferation (1) and an aberrant expression pattern has been reported in patients with schizophrenia (18). The *PLCB2* and *PLCB4* genes were found to be differentially expressed in human breast cancer MCF-7 cells, and to be associated with multidrug resistance using RNA-seq technology (19). *PLCB3* has been reported to be regulated by multiple protein kinases and to control hormonal signaling (20).

HBV infection is regarded as a main risk factor for the development of HCC (4). HBV is classified into ten genotypes, from A to J, and >40 associated sub-genotypes (21). The 10 genotypes are based on an intergroup divergence of $\geq 8\%$ in the complete nucleotide sequence; whereas the sub-genotypes are based on a divergence of 4-7.5% (22,23). Notably, genotypes A and B are associated with earlier hepatitis B e antigen seroconversion, less active liver disease, and a slower rate of progression to cirrhosis and HCC compared with genotypes C and D (24-27).

Some *PLCB* isoforms have been explored with regard their associations with tumor development; therefore, the present study aimed to explore the association between four *PLCB* genes and HCC.

Materials and methods

Patient data collection. The GSE14520 dataset was used for analysis (ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE14520; accessed June 10th, 2018) (28,29). This dataset contains two platforms: GPL571 (GeneChip® Human Genome U133A 2.0 Array; Thermo Fisher Scientific, Inc., Waltham, MA, USA) and GPL3921 (GeneChip® HT Human Genome U133 Array Plate Set; Thermo Fisher Scientific, Inc.). To avoid a batch effect, patients from GPL3921 were used. Patients with HBV infection were used, including a total of 212 patients. In addition, patient survival, including overall survival (OS) and recurrence-free survival (RFS), validated findings in the GSE14520 dataset using the Gene Expression Profiling Interactive Analysis (GEPIA; gepia.cancer-pku.cn/index.html; accessed June 10th, 2018) website with data from The Cancer Genome Atlas (TCGA) database (30).

Gene, protein and tissue expression, and the body map. Gene expression, the body map and transcripts per million of the *PLCB* genes were collected from the GEPIA website (gepia.cancer-pku.cn/index.html; accessed June 12th, 2018). Tissue and protein expression of the *PLCB* genes were collected from the GTEx portal (gtexportal.org/home/; accessed June 12th, 2018) (31) and The Human Protein Atlas (proteinatlas.org/; accessed June 12th, 2018) (32) websites, respectively.

Gene set enrichment analysis (GSEA). GSEA (software, broadinstitute.org/gsea/index.jsp) was performed to explore

potential mechanisms that *PLCB* genes are involved in, including biological processes and metabolic pathways. Datasets of c2.cp.kegg.v6.1.symbols.gmt, c5.bp.b6.1.symbols.gmt, c5.cc.v6.1.symbols.gmt, c5.mf.v6.1.symbols.gmt and c5.all.v6.1.symbols.gmt were used to analyze statistically significant Gene Ontology (GO) terms, including biological process (BP), cellular component (CC), and molecular function (MF), and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways (33,34).

Association and interaction analysis. The Pearson correlation matrix among *PLCB* genes was constructed using R version 3.5.0 (r-project.org/). Pearson correlation and associations between *PLCB* gene expression and tumor stage were validated using the GEPIA website. The co-expression interactive network of gene-gene interactions was constructed using the geneMANIA plugin of Cytoscape software version 3.6.0 (35,36). The protein-protein interaction (PPI) network was constructed using the STRING (string-db.org/cgi/input.pl, accessed June 20th, 2018) website (37). GO enrichment analysis was visualized using the BiNGO plugin of Cytoscape software version 3.6.0 (38).

Diagnostic and prognostic analysis and stratified, joint-effect analysis. Diagnostic receiver operating characteristic (ROC) curves were constructed using the expression of *PLCB* genes in tumor and non-tumor tissues. Gene expressions were categorized into two groups of low and high expression at a cut-off value of median expression levels. OS and RFS were calculated using the Kaplan-Meier and Cox proportional hazards regression models. Statistically significant clinical factors were adjusted for multivariate Cox models. Then, prognosis-associated genes were further stratified for analysis by clinical factors. In addition, prognosis-associated genes were combined for a joint-effect analysis with α -fetoprotein (AFP) based low and high expression.

Expression model and nomogram construction. To further explore prognosis-associated genes for HCC survival, expression models for OS and RFS prediction were constructed. Gene expression, patient survival status, expression heatmaps and prognostic ROC curves were constructed in the model (39-42). Nomograms for OS and RFS were also constructed using clinical factors and genes to predict patient survival probability at 1, 3 and 5 years.

Genome-wide analysis of prognosis-associated genes. Prognosis-associated genes were further explored in genome-wide analysis. A cut-off value of 0.4 was determined for further analysis. The cut-off 0.4 can filter a lot of genes with weak relationships with *PLCB1* and leads to a better presentation of GO and pathway results compared with other cut-off values. Gene-gene interactions, and BP, CC and MF were constructed using Cytoscape software.

Statistical analysis. Unpaired t test was used to analyze expressions of *PLCBs* in tumor and non-tumor tissues. Box plots and survival plots were generated using GraphPad software version 7.0 (GraphPad Software, Inc., La Jolla, CA, USA). Survival analyses were performed using SPSS software




Figure 1. Relative mRNA expressions and transcriptional levels of PLCB1-4 in tumor and non-tumor tissues. Relative mRNA expressions of (A) PLCB1, (B) PLCB2, (C) PLCB3 and (D) PLCB4 in tumor and non-tumor tissues. Transcriptional levels of (E) PLCB1, (F) PLCB2, (G) PLCB3 and (H) PLCB4 in tumor and non-tumor tissues. PLCB, phospholipase C β .

version 16.0 (SPSS, Inc., Chicago, IL, USA). Median survival time and log-rank P-value were calculated by the Kaplan-Meier method, and the 95% confidence interval (CI) and hazard ratio (HR) were calculated by univariate and multivariate Cox proportional hazards regression models, respectively. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Demographic and clinical characteristics. Data from 212 patients (GSE14520) with HBV-associated HCC were used in the study. AFP, BCLC stage, tumor size and cirrhosis were associated with OS ($P = 0.049$, $P < 0.0001$, $P = 0.002$ and $P = 0.041$, respectively). Gender, cirrhosis and BCLC stage were associated with RFS ($P = 0.002$, $P = 0.036$ and $P < 0.0001$, respectively). Other factors were not associated with prognosis ($P > 0.05$; Table SI).

Gene, protein, tissue expressions and transcription analysis. *PLCB1* and *PLCB3* were highly expressed in tumor tissues compared with normal tissue, whereas *PLCB2* had the opposite result (all $P \leq 0.05$; Fig. 1A-C). However, there was no difference in *PLCB4* expression between the tumor and normal tissue (Fig. 1D). Transcriptional analysis indicated that *PLCB1*, *PLCB3* and *PLCB4* consistently exhibited higher




Figure 2. Relative mRNA expressions of PLCB1-4 in tumor and normal tissues and low, high expression groups. (A) Relative mRNA expressions of PLCB1-4 in tumor and normal tissues; (B) Relative mRNA expressions of PLCB1-4 in low and high expression groups. PLCB, phospholipase C β .

transcripts per millions in tumor tissues compared with normal tissues (Fig. 1E-H). Tissue and protein expression of the *PLCB* genes were collected from the GTEx portal.




Figure 3. Diagnostic ROC curves of PLCB1-4. A-D: Diagnostic ROC curves of (A) PLCB1, (B) PLCB2, (C) PLCB3 and (D) PLCB4; Diagnostic ROC curves of combination of (E) PLCB1 and PLCB2, (F) PLCB1 and PLCB3, (G) PLCB2 and PLCB3, and (H) PLCB1, PLCB2 and PLCB3. ROC, receiver operating characteristics; PLCB, phospholipase C β ; AUC, area under the curve; CI, confidence interval.




Figure 4. Overall survival ROC curves of PLCB1-4 at 1, 3 and 5 years. ROC curves of (A) PLCB1, (B) PLCB2, (C) PLCB3 and (D) PLCB4 at 1 year; ROC curves of (E) PLCB1, (F) PLCB2, (G) PLCB3 and (H) PLCB4 at 3 years; ROC curves of (I) PLCB1, (J) PLCB2, (K) PLCB3 and (L) PLCB4 at 5 years. PLCB, phospholipase C β ; ROC, receiver operating characteristics.

Table I. Prognostic analysis of PLCB genes for overall survival.

| Variable | Patients (n=212) | No. of events | MST (months) | HR (95% CI) | Crude P-value | HR (95% CI) | Adjusted P-value ^a |
|-----------------|------------------|---------------|--------------|---------------------|------------------|----------------------------|-------------------------------|
| <i>PLCB1</i> | | | | | | | |
| Low expression | 106 | 29 | NA | Ref. | | Ref. | |
| High expression | 106 | 53 | 53 | 2.246 (1.426-3.536) | <0.001 | 2.100 (1.310-3.367) | 0.002 |
| <i>PLCB2</i> | | | | | | | |
| Low expression | 106 | 41 | NA | Ref. | | Ref. | |
| High expression | 106 | 41 | NA | 0.902 (0.585-1.391) | 0.641 | 1.041 (0.660-1.641) | 0.863 |
| <i>PLCB3</i> | | | | | | | |
| Low expression | 106 | 36 | NA | Ref. | | Ref. | |
| High expression | 106 | 46 | NA | 1.394 (0.900-2.159) | 0.137 | 1.035 (0.659-1.625) | 0.882 |
| <i>PLCB4</i> | | | | | | | |
| Low expression | 106 | 44 | NA | Ref. | | Ref. | |
| High expression | 106 | 38 | NA | 0.877 (0.568-1.354) | 0.555 | 0.870 (0.555-1.363) | 0.534 |

^aP-values were adjusted for tumor size, cirrhosis, Barcelona Clinic Liver Cancer stage and α -fetoprotein; bold indicates significant P-values. Ref., reference value (1); NA, not available; MST, median survival time; HR, hazard ratio; 95% CI, 95% confidence interval; PLCB, phospholipase B.




Figure 5. Recurrence-free survival ROC curves of PLCB1-4 at 1, 3 and 5 years. ROC curves of (A) PLCB1, (B) PLCB2, (C) PLCB3 and (D) PLCB4 at 1 year; ROC curves of (E) PLCB1, (F) PLCB2, (G) PLCB3 and (H) PLCB4 at 3 years; ROC curves of (I) PLCB1, (J) PLCB2, (K) PLCB3 and (L) PLCB4 at 5 years. PLCB, phospholipase C β ; ROC, receiver operating characteristics.

Gene expression levels in 212 patients with HBV-HCC (GSE14520) indicated that there were significant differences in

PLCB1 and *PLCB2* expression between tumor and non-tumor tissues, whereas there was not difference in *PLCB3* and

Table II. Prognostic analysis of PLCB genes for recurrence-free survival.

| Variable | Patients (n=112) | No. of events | MST (months) | HR (95% CI) | Crude P-value | HR (95% CI) | Adjusted P-value ^a |
|-----------------|------------------|---------------|--------------|----------------------------|---------------|----------------------------|-------------------------------|
| <i>PLCB1</i> | | | | | | | |
| Low expression | 106 | 46 | NA | Ref. | | Ref. | |
| High expression | 106 | 70 | 26.9 | 1.914 (1.318-2.781) | 0.001 | 1.861 (1.273-2.271) | 0.001 |
| <i>PLCB2</i> | | | | | | | |
| Low expression | 106 | 59 | 36.0 | Ref. | | Ref. | |
| High expression | 106 | 57 | 51.1 | 0.863 (0.599-1.243) | 0.429 | 0.956 (0.654-1.398) | 0.817 |
| <i>PLCB3</i> | | | | | | | |
| Low expression | 106 | 52 | 54.8 | Ref. | | Ref. | |
| High expression | 106 | 64 | 29.9 | 1.466 (1.015-2.118) | 0.042 | 1.244 (0.853-1.814) | 0.257 |
| <i>PLCB4</i> | | | | | | | |
| Low expression | 106 | 59 | 46.3 | Ref. | | Ref. | |
| High expression | 106 | 57 | 43.2 | 1.015 (0.705-1.461) | 0.936 | 0.962 (0.664-1.395) | 0.840 |

^aP-values were adjusted for gender, cirrhosis and Barcelona Clinic Liver Cancer stage. Ref., reference value (1); MST, median survival time; HR, hazard ratio; 95% CI, 95% confidence interval; PLCB, phospholipase B. Bold indicates significant P-values.




Figure 6. Overall survival and recurrence-free survival analysis plots of PLCB1-4. Overall survival analysis plot of (A) PLCB1, (B) PLCB2, (C) PLCB3 and (D) PLCB4; recurrence-free survival analysis plot (E) PLCB1, (F) PLCB2, (G) PLCB3 and (H) PLCB4; joint-effects analysis of (I) α -fetoprotein and (J) PLCB1 for overall survival and recurrence-free survival. Group I, AFP low expression and PLCB1 low expression; Group 2, AFP low expression and PLCB1 high expression, and AFP high expression and PLCB1 low expression; Group 3, AFP high expression and PLCB1 high expression; Group I, AFP low expression and PLCB1 low expression; Group II, AFP low expression and PLCB1 high expression, and AFP high expression and PLCB1 low expression; Group III, AFP high expression and PLCB1 high expression. PLCB, phospholipase C β .

PLCB4 between the samples (Fig. 2A). In addition, when tumor samples were divided into high and low expression groups using the median as the cutoff there were significant

differences in *PLCB1*, *PLCB2* and *PLCB4*; whereas *PLCB3* did not exhibit significance (Fig. 2B). The bodymap distribution of PLCB genes in different organs is shown in Fig. S1. Protein

Table III. Stratified analysis of *PLCB1* for overall survival and recurrence-free survival.

| Variable | Overall survival | | | | Recurrence-free survival | | | |
|-----------------|------------------|------|----------------------------|------------------|--------------------------|------|----------------------------|------------------|
| | Low | High | Adjusted HR (95% CI) | Adjusted P-value | Low | High | Adjusted HR (95% CI) | Adjusted P-value |
| Sex | | | | | | | | |
| Male | 86 | 89 | 1.967 (1.174-3.24) | 0.010 | 86 | 89 | 1.877 (1.249-2.820) | 0.002 |
| Female | 20 | 17 | 2.619 (0.711-9.652) | 0.148 | 20 | 17 | 0.754 (0.168-3.382) | 0.713 |
| Age (years) | | | | | | | | |
| ≤60 | 91 | 92 | 2.252 (1.370-3.702) | 0.001 | 91 | 92 | 1.736 (1.129-2.670) | 0.012 |
| >60 | 15 | 14 | 0.850 (0.140-5.148) | 0.860 | 15 | 14 | 2.043 (0.701-5.953) | 0.191 |
| HBV | | | | | | | | |
| AVR-CC | 20 | 36 | 1.987 (0.695-5.687) | 0.200 | 20 | 36 | 1.486 (0.663-3.332) | 0.336 |
| CC | 86 | 70 | 1.957 (1.114-3.438) | 0.020 | 86 | 70 | 1.864 (1.166-2.979) | 0.009 |
| Tumor size (cm) | | | | | | | | |
| ≤5 | 75 | 62 | 2.100 (1.149-3.838) | 0.016 | 75 | 62 | 1.627 (1.012-2.616) | 0.045 |
| >5 | 30 | 44 | 1.790 (0.821-3.901) | 0.143 | 30 | 44 | 2.204 (1.049-4.629) | 0.037 |
| Cirrhosis | | | | | | | | |
| Yes | 93 | 102 | 1.922 (1.196-3.091) | 0.007 | 93 | 102 | 1.678 (1.124-2.503) | 0.011 |
| No | 13 | 4 | 476.586 (5.21E-12-4.36E16) | 0.707 | 13 | 4 | 3.758 (0.379-37.311) | 0.258 |
| Multinodular | | | | | | | | |
| Yes | 21 | 24 | 1.399 (0.544-3.598) | 0.487 | 21 | 24 | 1.186 (0.474-2.965) | 0.716 |
| No | 85 | 82 | 2.662 (1.522-4.656) | 0.001 | 85 | 82 | 2.163 (1.395-3.355) | 0.001 |
| AFP (ng/ml) | | | | | | | | |
| ≤300 | 68 | 47 | 2.098 (1.097-4.015) | 0.025 | 68 | 47 | 2.180 (1.307-3.635) | 0.003 |
| >300 | 35 | 59 | 1.886 (0.934-3.806) | 0.077 | 35 | 59 | 1.294 (0.710-2.359) | 0.399 |
| BCLC stage | | | | | | | | |
| 0 | 8 | 12 | 0.535 (0.033-8.559) | 0.658 | 8 | 12 | 0.597 (0.097-3.665) | 0.577 |
| A | 79 | 64 | 2.214 (1.210-4.051) | 0.010 | 79 | 64 | 1.928 (1.200-3.097) | 0.007 |
| B | 10 | 12 | 0.746 (0.225-2.478) | 0.633 | 10 | 12 | 0.903 (0.310-2.627) | 0.851 |
| C | 9 | 18 | 2.746 (0.836-9.021) | 0.096 | 9 | 18 | 2.370 (0.774-7.252) | 0.131 |

Ref., reference value (1); PLCB, phospholipase B; HR, hazard ratio; 95% CI, 95% confidence interval; HBV, hepatitis B virus; AVR-CC, acute viral replication-chronic carrier; CC, chronic carrier; AFP, AFP, α -fetoprotein; BCLC, Barcelona Clinic Liver Cancer. Bold indicates significant P-values.

expression levels demonstrated that PLCB2 is the most highly expressed of the PLCB family (Fig. S2). The different tissue expression levels of PLCB family members demonstrated that all were expressed at low levels in the liver (Fig. S3).

Diagnostic and prognostic analysis. In the diagnostic analysis of PLCB genes, PLCB1 and PLCB2 exhibited diagnostic value for HCC, while PLCB3 showed potential diagnostic value [$P<0.0001$, $P<0.0001$ and $P=0.018$, respectively; area under the curve (AUC), 0.869, 0.836 and 0.567, respectively; Fig. 3A-C]. However, PLCB4 did not have any diagnostic value ($P=0.811$; Fig. 3D). In the combined diagnostic analysis for PLCB1, PLCB2 and PLCB3, the combinations of PLCB1 + PLCB2, PLCB1 + PLCB3, and PLCB1 + PLCB2 + PLCB3 exhibited diagnostic value for HCC with an advantage over PLCB1, PLCB2 or PLCB3 alone (AUC, 0.905, 0.877 and 0.920, respectively; all $P<0.05$; Fig. 3E, F and H). The combination of PLCB2 and PLCB3 exhibited potential diagnostic value for HCC

(AUC, 0.604; $P=0.0003$; Fig. 3G). In the prognostic analysis (Figs. 4 and 5), only PLCB1 expression was associated with patient OS at 1-, 3- and 5-years (all AUC >0.6 ; Fig. 4A, E and I). In addition, PLCB1 expression was associated with patient RFS at 3- and 5-years (both AUC >0.6 ; Fig. 5E and I).

In the univariate analysis (Tables I and II; Fig. 6), *PLCB1* expression was associated with OS (crude $P=0.002$; Fig. 6A); *PLCB1* and *PLCB3* expression was associated with RFS (crude $P=0.001$ and $P=0.042$, respectively; Fig. 6E and G). In the multivariate analysis, *PLCB1* expression was associated with OS and RFS (adjusted $P=0.002$ and 0.001 , respectively; Tables I and II). Other genes were not associated with prognosis (adjusted $P>0.05$; Tables I and II).

Stratified and joint-effect survival analysis. Stratification analysis was performed for *PLCB1* on OS and RFS. Male gender, age <60 years, chronic carrying of HBV, cirrhosis, single nodular, AFP levels <300 ng/ml, and A stage in the

Table IV. Joint-effect analysis of *PLCB1* and AFP for overall survival and recurrence-free survival.

| A, Overall survival | | | | | | |
|-----------------------------|----------------|-------------------------|--------------|--------------|-----------------------------|------------------|
| Group | AFP expression | <i>PLCB1</i> expression | Events/total | MST (months) | Adjusted HR (95% CI) | Adjusted P-value |
| 1 | Low | Low | 18/68 | NA | Ref. | 0.008 |
| 2 | Low | High | 32/82 | NA | 2.162 (1.143-4.089) | 0.018 |
| 3 | High | High | 32/59 | 36.4 | 4.382 (1.703-11.276) | 0.002 |
| B, Recurrence-free survival | | | | | | |
| Group | AFP expression | <i>PLCB1</i> expression | Events/total | MST (months) | Adjusted HR (95% CI) | Adjusted P-value |
| I | Low | Low | 29/68 | NA | Ref. | 0.075 |
| II | Low | High | 50/82 | 40.1 | 1.613 (1.019-2.555) | 0.041 |
| III | High | High | 37/59 | 23.0 | 1.670 (1.012-2.755) | 0.045 |

Group 1, AFP low expression and *PLCB1* low expression; Group 2, AFP low expression and *PLCB1* high expression, and AFP high expression and *PLCB1* low expression; Group 3, AFP high expression and *PLCB1* high expression; Group I, AFP low expression and *PLCB1* low expression; Group II, AFP low expression and *PLCB1* high expression, and AFP high expression and *PLCB1* low expression; Group III, AFP high expression and *PLCB1* high expression. Ref., reference value (1); *PLCB*, phospholipase B; AFP, α -fetoprotein; MST, median survival time; HR, hazard ratio; 95% CI, 95% confidence interval. Bold indicates significant P-values.

BCLC staging system were associated with OS and RFS (all adjusted $P \leq 0.05$; Table III). Tumor size < 5 cm was associated with OS and any group of tumor size was associated with RFS (all adjusted $P \leq 0.05$; Table III).

In the joint-effect analysis (OS/RFS: group 1/I, AFP low + *PLCB1* low; group 2/II, AFP low + *PLCB1* high, and AFP high + *PLCB1* low; groups 3/III, AFP high + *PLCB1* high), when combining *PLCB1* and AFP, prognostic significance was observed among the three groups for OS (adjusted $P = 0.008$; Table IV); group 3 exhibited the worst prognosis [adjusted $P = 0.002$, adjusted HR (95% CI) = 4.382 (1.703-11.276); Table IV]. Prognostic significance was not observed among the three groups in RFS (adjusted $P = 0.075$; Table IV). However, group III exhibited the worst prognosis [adjusted $P = 0.045$, adjusted HR (95% CI) = 1.670 (1.012-2.755); Table IV].

GSEA. Both diagnostic- and prognostic-associated genes were explored to investigate the mechanisms that *PLCB*s are involved in. Enriched GO terms and KEGG pathways annotated with *PLCB1* included 'G protein coupled receptor activity', 'sodium channel activity', 'extracellular ligand gated ion channel activity' and 'taste transduction pathway', among others (Fig. 7). Enriched GO terms and KEGG pathways annotated with *PLCB2* included 'mRNA processing', 'cell division', 'cell cycle checkpoints', 'DNA repair', 'PPAR signaling pathway', 'metabolism of xenobiotics by cytochrome P450' and 'adipocytokine signaling pathway' among others (Fig. 8).

Expression model and nomogram construction. An expression model was constructed for OS and RFS prognosis prediction (Fig. 9). *PLCB1* expression, OS and RFS survival status, and

PLCB1 expression heatmaps are shown in Fig. 9A, and prognostic ROC curves demonstrated that *PLCB1* expression has prognostic value for OS and RFS (Fig. 9B and C).

Furthermore, nomograms were constructed for clinical factors and *PLCB1*. High expression always led to low points. The same points indicated a higher probability of survival at 1 year, yet a lower probability of survival at 5 years for both OS and RFS. Survival probability at 3 years was seated in the middle (Fig. 10).

Interaction and co-expression networks and enrichment analysis. Associations between gene expression and TNM stage (I, II, III) were visualized; *PLCB1* gene expression of 212 HBV-HCC was significantly different in early (I, II) stages compared with advanced (III) stage in ($P \leq 0.01$; Fig. 11A). Associations between gene expressions and TNM stage (I, II and III) in GEPIA indicated that *PLCB1* expression was different in different tumor stages (Fig. 11B). Gene-gene co-expression interactions and PPI networks demonstrated interactions between *PLCB* members (Fig. 11C and D). The Pearson correlation matrix showed an association between *PLCB* members (Fig. 11E).

Furthermore, enriched GO terms are presented in Fig. S4A-C. KEGG pathways that *PLCB* members are involved in are presented in Fig. S5. All members were involved in diacylglycerol and IP3 metabolism and finally induced sustained angiogenesis, thus evading apoptosis and proliferation effects.

Analysis of *PLCB1* and associated genes genome-wide. Pearson correlation analysis was performed for *PLCB1* genome-wide. A total of 53 genes were identified at $r \geq 0.4$. Gene-gene interaction analysis was constructed and presented




Figure 7. Gene set enrichment analysis results of *phospholipase C beta 1* gene. Results of gene ontologies: (A) G-protein coupled receptor activity; (B) sodium channel activity; (C) neurotransmitter receptor activity; (D) extracellular ligand gated ion channel activity. GO, gene ontology; NES, normalized enrichment score; FDR, false discovery rate; KEGG, Kyoto Encyclopedia of Genes and Genomes.

in Fig. 12. Networks of BP, CC and MF terms were constructed (Fig. 13). Enriched GO terms and KEGG pathways annotated by *PLCB1* and correlated genes are presented in Table V.

Validation of prognostic values of *PLCB* genes. *PLCB* genes were further validated in GEPIA for OS and RFS (Fig. 14). *PLCB1* and *PLCB2* were associated with OS ($P=0.0075$ and $P=0.041$, respectively; Fig. 14A and B). In addition, *PLCB1* and *PLCB4* were associated with RFS ($P<0.0001$ and $P<0.018$, respectively; Fig. 14E and H). Other genes were not associated with OS or RFS (all $P>0.05$; Fig. 14). Pearson correlation in GEPIA (Fig. 15) indicated that *PLCB1* was positively correlated with *PLC3* and *PLCB4*, while *PLCB3* was positively correlated with *PLCB4*, which is consistent with Fig. 11E.

Discussion

In the current study, it was identified that *PLCB1* and *PLCB2* genes are differently expressed in tumor and normal tissues.

PLCB1 and *PLCB2* have diagnostic value for HCC, while *PLCB3* has potential diagnostic value for HCC. Combinations of these genes have an advantage over *PLCB1*, *PLCB2* or *PLCB3* alone with regard to HCC diagnosis. In addition, *PLCB1* has prognostic value of OS and RFS for HCC. Combining *PLCB1* and AFP had an advantage over *PLCB1* alone for OS and RFS. GSEA indicated that *PLCB1* and *PLCB2* were involved in 'G protein coupled receptor activity', 'sodium channel activity', 'cell division', 'cell cycle checkpoint', 'DNA repair', 'PPAR signaling pathway', 'metabolism of xenobiotics by cytochrome P450' and 'adipocytokine signaling pathway', among others. Nomograms and gene expression models were constructed for HCC prognosis prediction. The validation of the prognostic values of *PLCB* genes revealed that *PLCB1* and *PLCB2* were associated with OS, and *PLCB1* and *PLCB4* were associated with RFS.

PLC proteins are key enzymes that metabolize inositol lipids and have a pivotal role in multiple transmembrane signaling transduction pathways that modulate a series of cellular




Figure 7. Continued. Gene set enrichment analysis results of *phospholipase C β1* gene. Results of gene ontologies: (E) sodium ion transmembrane transporter; (F) passive transmembrane transporter; (G) gated channel activity; (H) cation channel activity; (I) monovalent inorganic cation transmembrane transporter activity; (J) excitatory extracellular ligand gated ion channel activity; (K) ligand gated channel activity. (L) Taste transduction KEGG pathway. GO, gene ontology; NES, normalized enrichment score; FDR, false discovery rate; KEGG, Kyoto Encyclopedia of Genes and Genomes.




Figure 8. Gene set enrichment analysis results of *phospholipase C beta 2* gene. Results of gene ontologies: (A) mRNA processing; (B) cell division; (C) negative regulation of mitotic cell cycle; (D) cell cycle checkpoint. GO, gene ontology; NES, normalized enrichment score; FDR, false discovery rate; KEGG, Kyoto Encyclopedia of Genes and Genomes; PPAR peroxisome proliferator-activated receptor.

processes, including cell proliferation and mobility (16). In mammalian cells, there are four PLCB isoforms: PLCB1, PLCB2, PLCB3 and PLCB4. PLCB2 and PLCB3 are activated by G $\beta\gamma$ dimers, which are released upon the activation of G α protein coupled receptor families (43). PLCB2 can also be activated by Rho family members of monomeric G proteins, with the strongest activation by Rac1; these participate in the cytoskeletal rearrangements that accompany cell mobility (44).

The PLCB1 enzyme is encoded by the *PLCB1* gene, which is located at chromosome of 20p12 (1). It was originally identified as a G protein coupled receptor-associated PLCB isoform that is able produce inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate (45). The deregulation of signaling transduction pathways always

leads to advantages for tumor patients (1). *PLCB1* is activated by G α and induces a variety of events, which may increase the total intracellular calcium levels (46); one possible result of this process is aberrant proliferation in the cell (1). PLCB1 has been reported to have a role in promoting cell cycle progression by targeting cyclin-cyclin kinase complexes (47).

In addition, *PLCB1* has been documented to have a pivotal role in myoblast differentiation, regulating the delayed differentiation of skeletal muscle in myotonic dystrophy myoblasts (48). *PLCB1* may also reduce cell damage under oxidative conditions and prevent α -synuclein aggregation (49). The amplification of *PLCB1* increased K562 cell viability and enables cells to evade apoptosis (50,51); the overexpression of *PLCB1* keeps Swiss 3T3 cells in the S phase of the cell cycle (52). Li *et al* (1) reported that




Figure 8. Continued. Gene set enrichment analysis results of *phospholipase C β2* gene. Results of gene ontologies: (E) DNA repair; (F) negative regulation of cell cycle phase transition (G) mitotic nuclear division; (H) iron ion binding. Results of KEGG pathways: (I) metabolism of xenobiotics by cytochrome P450; (J) PPAR signaling pathway; (K) adipocytokine signaling pathway; (L) steroid hormone biosynthesis. GO, gene ontology; NES, normalized enrichment score; FDR, false discovery rate; KEGG, Kyoto Encyclopedia of Genes and Genomes; PPAR peroxisome proliferator-activated receptor.




Figure 9. Expression model constructed using *PLCB1* gene. (A) Expression model including expression, overall survival status, recurrence-free survival status and heatmap. (B) Time dependent ROC curves of overall survival at 1, 3- and 5- years. (C) Time dependent ROC curves of recurrence-free survival at 1, 3- and 5- years. *PLCB1*, phospholipase C β 1; ROC, receiver operating characteristics; AUC, area under the curve.




Figure 10. Nomograms constructed using overall survival and recurrence-free survival-related clinical factors and genes. (A) Nomogram of OS-associated genes and clinical factors. (B) Nomogram of RFS-associated genes and clinical factors. BCLC, Barcelona Clinic Liver Cancer; AFP, α -fetoprotein; *PLCB1*, phospholipase C β 1; OS, overall survival; RFS, recurrence-free survival.

upregulated *PLCB1* expression is associated with tumor cell proliferation and infers a poor prognosis for HCC. The present study revealed that high expression has is undesirable for HCC prognosis (OS and RFS), which is consistent with the results of Li *et al* (1). Furthermore, *PLCB1* had diagnostic value for HCC.

PLCB2 mediates mitogenic, proliferative and migratory events by interacting with heterotrimeric and monomeric G proteins, and can interact with γ -synuclein to regulate G protein activation (43). Bertagnolo *et al* (53) reported that *PLCB2* induces cell cycle transition from G0/G1 to the S/G2/M phases, which is critical for tumor progression, without affecting cell cycle-associated enzymes. They also indicate that *PLCB2*, by modifying the phospholipase pool, may be responsible for the inositol lipid-associated modifications of the cytoskeleton architecture that occur in the course of division, motility and invasion of tumor cells (53). The current findings with regard to the role of *PLCB2* in the cell cycle and cell division are consistent with the results of Bertagnolo *et al* (53).

PLCB2 has been reported to promote mitosis and the migration of human breast cancer-derived cells (54), is highly expressed in breast cancer and associated with poor prognosis (55); however, little is known about HCC *PLCB2* expression, and the role in HCC diagnosis and prognosis. In the current study, *PLCB2* expression as not associated with HCC prognosis, but may be a diagnostic signature for HCC.




Figure 11. Scatter plots, matrix and interaction networks analysis. (A) Scatter plot and (B) violin plot of *PLCB1* expressions. (C) Co-expression network of *PLCB1-4* genes. (D) Protein-protein interaction network of *PLCB1-4*. (E) Pearson correlation matrix of *PLCB1-4*. *PLCB*, phospholipase C β .

PLCB3 is located on chromosome 11q13 in the vicinity of the multiple endocrine neoplasia type 1 gene; its loss leads to the development of neuroendocrine tumors (56). The transfection of *PLCB3* to a human endocrine pancreatic tumor

cell line can induce the activation of the human mismatch repair protein 3 gene (56). *PLCB3* interacts with Na(+)/H(+) exchange regulatory cofactor NHERF-1, providing a structural basis for CXCR2 signaling in pancreatic cancer (57).




Figure 12. Co-expression network of *PLCB1* gene with correlation-associated genes in genome-wide analysis. *PLCB1*, phospholipase C β 1.

Hoeppner *et al* (58) identified a novel role for *PLCB3*, functioning as a negative regulator of vascular endothelial growth factor-mediated vascular permeability by regulating intracellular Ca^{2+} release. *PLCB3* may have a tumor suppressor role via SHP-1-mediated dephosphorylation of Stat5 (59). Ju *et al* (60) reported that *PLCB* and G_{α} may have important roles in scar remodeling, cardiac hypertrophy and fibrosis following myocardial infarction rat hearts. In the present study, *PLCB3* expression exhibited potential diagnostic value for HCC and without association with HCC prognosis. *PLCB3* may have a weak role in HCC if at all, which requires further investigation.

Compared with other *PLCB* genes, *PLCB4* is less well characterized, and associations between *PLCB4* and cancer are unclear. The expression of *PLCB4* and *PLCB3* was previously explored in Purkinje cell subsets of the mouse cerebellum (61). *PLCB4* and *PLCB3* are differentially expressed in microarray databases of non-small cell lung cancer, but neither are associated with the prognosis and development of lung cancer (62). Orchel *et al* (63) reported that *PLCB4* is differentially expressed in 50 endometrium samples from women with endometrial cancer, but is not associated with the treatment of endometrial cancer. Furthermore, the present study did not find any association

between *PLCB4* and HCC. Therefore, further studies are required to explore the relationship between *PLCB4* expression and malignancy.

The findings of the present study indicate that *PLCB1* expression was associated with OS, whereas *PLCB1* and *PLCB3* expression was associated with RFS in univariate analysis. In multivariate analysis, *PLCB1* expression was associated with OS and RFS. Multivariate cox analysis contains several significant clinicopathological characteristics, which produces new adjusted results and conclusions. In addition, *PLCB1* expression was associated with OS, whereas *PLCB1* and *PLCB3* expression was associated with RFS in univariate analysis. However, *PLCB1* expression was associated with OS and RFS in multivariate analysis. Different results may be due to varied clinicopathological characteristics in the GSE14520 and TCGA dataset. Of course, HBV is a pivotal factor associated with HCC.

There are some limitations to the present study that should be recognized. Firstly, larger sample cohorts are required to validate these findings. Additionally, the results are based on a HBV-associated HCC population; therefore, further explorations are needed in a study including HBV-infected and non-infected patients. Finally, functional trials are required to further explore the roles of *PLCB* genes in HCC initiation,

Table V. Enrichment results of gene ontologies and KEGG pathways of phospholipase B1 with genome-wide associated genes.

| Category | Term | Count | P-value | False discovery rate | Genes |
|--------------------|---|-------|----------|----------------------|---|
| Biological process | Oxidation-reduction process | 11 | 1.15E-06 | 0.001521 | FMO4, CBR1, MSRA, PLOD2, BLVBR, F8, SMOX, GRHPR, NDUFA10, HPD, HSD17B8 |
| Molecular function | Long-chain fatty acid-CoA ligase activity | 3 | 0.000399 | 0.455909 | ACSL4, ACSL3, SLC27A5 |
| | Extracellular exosome | 16 | 0.000748 | 0.826682 | NACA, FCER2, CAPZA1, FBP1, AXL, SPINK1, GRHPR, CBR1, MSRA, RPL7, PLOD2, BLVBR, SNRPB, ACSL4, PLCB1, HPD |
| Biological process | Long-chain fatty acid metabolic process | 3 | 0.001186 | 1.559498 | ACSL4, ACSL3, SLC27A5 |
| | Cytosol | 17 | 0.001409 | 1.552995 | CAPZA1, FBP1, ARHGAP28, TRIB3, SAE1, GRHPR, CBR1, MSRA, RPL7, NCAPG, BLVBR, SNRPB, SMOX, PLCB1, SNRPF, NUP43, HPD |
| Cellular component | Endoplasmic reticulum membrane | 7 | 0.012018 | 12.55815 | FMO4, HMOX2, PLOD2, ACSL4, ACSL3, SLC27A5, HPD |
| Cellular component | Actin cytoskeleton | 4 | 0.012886 | 13.40731 | MSRA, SORBS2, NCAPG, CAPZA1 |
| Cellular component | U7 snRNP | 2 | 0.015645 | 16.05626 | SNRPB, SNRPF |
| Biological process | Heme catabolic process | 2 | 0.01697 | 20.28229 | HMOX2, BLVBR |
| Molecular function | Decanoate-CoA ligase activity | 2 | 0.018337 | 19.08795 | ACSL4, ACSL3 |
| Molecular function | Very long-chain fatty acid-CoA ligase activity | 2 | 0.02287 | 23.26214 | ACSL4, SLC27A5 |
| Cellular component | U4 snRNP | 2 | 0.024478 | 24.04788 | SNRPB, SNRPF |
| Cellular component | Methylosome | 2 | 0.026674 | 25.92421 | SNRPB, SNRPF |
| Biological process | Cellular protein modification process | 3 | 0.027107 | 30.50786 | MSRA, PLOD2, SAE1 |
| Cellular component | Small nucleolar ribonucleoprotein complex | 2 | 0.028865 | 27.75428 | SNRPB, SNRPF |
| Biological process | Histone mRNA metabolic process | 2 | 0.028919 | 32.20282 | SNRPB, SNRPF |
| Biological process | Positive regulation of nitric-oxide synthase biosynthetic process | 2 | 0.031291 | 34.36424 | CCL20, FCER2 |
| Molecular function | RNA binding | 5 | 0.036659 | 34.78158 | RPL7, SNRPB, CPSF6, PAPOLG, SNRPF |
| | Intracellular ribonucleoprotein complex | 3 | 0.037507 | 34.57769 | RPL7, SNRPB, CPSF6 |
| Cellular component | Small nuclear ribonucleoprotein complex | 2 | 0.037582 | 34.63438 | SNRPB, SNRPF |
| Cellular component | SMN-Sm protein complex | 2 | 0.037582 | 34.63438 | SNRPB, SNRPF |
| Cellular component | U1 snRNP | 2 | 0.041912 | 37.82519 | SNRPB, SNRPF |
| Biological process | Nuclear import | 2 | 0.043071 | 44.18227 | SNRPB, SNRPF |
| Cellular component | U12-type spliceosomal complex | 2 | 0.056916 | 47.81769 | SNRPB, SNRPF |
| Biological process | Metabolic process | 3 | 0.063294 | 57.93576 | GRHPR, ACSL4, ACSL3 |
| Biological process | Drug metabolic process | 2 | 0.063922 | 58.30767 | FMO4, CBR1 |
| Biological process | Spliceosomal snRNP assembly | 2 | 0.066211 | 59.63802 | SNRPB, SNRPF |
| Biological process | mRNA polyadenylation | 2 | 0.066211 | 59.63802 | CPSF6, PAPOLG |
| Biological process | Regulation of glucose transport | 2 | 0.077575 | 65.68049 | TRIB3, NUP43 |

Table V. Continued.

| Category | Term | Count | P-value | False discovery rate | Genes |
|--|--|-------|----------|----------------------|--|
| Biological process | Regulation of G-protein coupled receptor protein signaling pathway | 2 | 0.091035 | 71.75149 | RAMP3, PLCB1 |
| Biological process | Long-chain fatty-acyl-CoA biosynthetic process | 2 | 0.097693 | 74.37219 | ACSL4, ACSL3 |
| KEGG pathway | PPAR signaling pathway | 3 | 0.031778 | 29.78018 | ACSL4, ACSL3, SLC27A5 |
| KEGG pathway | Fatty acid biosynthesis | 2 | 0.053251 | 45.0671 | ACSL4, ACSL3 |
| KEGG pathway | Metabolic pathways | 10 | 0.058505 | 48.31368 | CBR1, FBPI, GRHPR, ACSL4, PLCB1, NDUFA10, ACSL3, SLC27A5, HPD, HSD17B8 |
| KEGG, Kyoto Encyclopedia of Genes and Genomes. | | | | | |




Figure 13. Visualized gene ontologies of *phospholipase C beta 1* and correlation-associated genes in genome-wide analysis. (A) Biological process, (B) cellular component and (C) molecular function.

development, metastasis, proliferation and angiogenesis. BCLC stage is an important factor associated with HCC and treatments concerning BCLC stage should mentioned in the material section.

The present study demonstrated that the *PLCB1* and *PLCB2* genes are differentially expressed between tumor and normal tissues and have diagnostic values for HCC. *PLCB3* has a potential diagnostic value for HCC. The combinations of these genes have an advantage over *PLCB1*, *PLCB2* or *PLCB3* used alone for HCC diagnosis. In addition, *PLCB1* has OS and RFS prognostic value for HCC. Combining *PLCB1* and AFP was advantageous over *PLCB1* alone for predicting OS and RFS. Nomogram and gene expression models were used to construct and predict HCC prognosis. GO terms and metabolic pathways associated with *PLCB1* and *PLCB2* are include 'G protein coupled receptor activity', 'cell division', 'cell cycle checkpoint', 'DNA repair', 'PPAR signaling pathway' and 'metabolism of xenobiotics by cytochrome P450'. Validation of the prognostic value of the PLCB genes revealed that *PLCB1*, *PLCB2* and *PLCB4* are associated with HCC prognosis.




Figure 14. Overall survival and disease recurrence-free survival analysis plots of PLCB1-4. Overall survival analysis plots of (A) PLCB1, (B) PLCB2, (C) PLCB3 and (D) PLCB4. Disease recurrence-free survival analysis plot (E) PLCB1, (F) PLCB2, (G) PLCB3 and (H) PLCB4. PLCB, phospholipase C β ; TPM, transcripts per million; HR, hazard ratio.




Figure 15. Pearson correlation plots of PLCB1-4 genes. (A) PLCB1 vs. PLCB2; (B) PLCB3 vs. PLCB1; (C) PLCB4 vs. PLCB1; (D) PLCB3 vs. PLCB2; (E) PLCB4 vs. PLCB2; (F) PLCB3 vs. PLCB4. PLCB, phospholipase C β ; TPM, transcripts per million.

Acknowledgements

The authors would like to acknowledge researchers for their contribution to open access data available via GTEx portal, GEPIA, Kaplan-Meier Plotter, STRING and The Human Protein Atlas websites. In addition, the authors would like to acknowledge invaluable help from peer reviewers.

Funding

This work was supported in part by the National Nature Science Foundation of China (grant no. 81560535, 81072321, 30760243, 30460143, 30560133 and 81802874), Natural Science Foundation of Guangxi Province of China (grant no. 2017JJB140189y), Key Laboratory of High-Incidence-Tumor Prevention and Treatment (Guangxi Medical University), Ministry of Education (GKE2018-01), 2009 Program for New Century Excellent Talents in University (NCET), Guangxi Nature Sciences Foundation (grant no. GuiKeGong 1104003A-7), and Guangxi Health Ministry Medicine Grant (Key-Scientific Research-Grant; grant no. Z201018). The present study is also partly supported by Scientific Research Fund of the Health and Family Planning Commission of Guangxi Zhuang Autonomous Region (grant no. Z2016318), The Basic Ability Improvement Project for Middle-aged and Young Teachers in Colleges and Universities in Guangxi (grant no. 2018KY0110). As well as, the present study is also partly supported by Research Institute of Innovative Think-tank in Guangxi Medical University (The gene-environment interaction in hepatocarcinogenesis in Guangxi HCCs and its

translational applications in the HCC prevention). We also acknowledge the supported by the National Key Clinical Specialty Programs (General Surgery & Oncology) and the Key Laboratory of Early Prevention & Treatment for Regional High-Incidence-Tumor (Guangxi Medical University), Ministry of Education, China.

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

XW and TP designed this study and the manuscript. KH, XZ, ZL, XL, CY, TY, CH, GZ, WQ and TP conducted the study and analyzed the data. XW wrote the manuscript and TP guided the writing. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Li J, Zhao X, Wang D, He W, Zhang S, Cao W, Huang Y, Wang L, Zhou S and Luo K: Up-regulated expression of phospholipase C, $\beta 1$ is associated with tumor cell proliferation and poor prognosis in hepatocellular carcinoma. *Onco Targets Ther* 9: 1697-1706, 2016.
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68: 394-424, 2018.
- Poon D, Anderson BO, Chen LT, Tanaka K, Lau WY, Van Cutsem E, Singh H, Chow WC, Ooi LL, Chow P, *et al*: Management of hepatocellular carcinoma in Asia: Consensus statement from the Asian Oncology Summit 2009. *Lancet Oncol* 10: 1119-1127, 2009.
- Tagliamonte M, Petrizzo A, Napolitano M, Luciano A, Arra C, Maiolino P, Izzo F, Tornesello ML, Aurisicchio L, Ciliberto G, *et al*: Novel metronomic chemotherapy and cancer vaccine combinatorial strategy for hepatocellular carcinoma in a mouse model. *Cancer Immunol Immunother* 64: 1305-1314, 2015.
- Lencioni R and Crocetti L: Local-regional treatment of hepatocellular carcinoma. *Radiology* 262: 43-58, 2012.
- Kanwal F, El-Serag HB and Ross D: Surveillance for hepatocellular carcinoma: Can we focus on the mission? *Clin Gastroenterol Hepatol* 13: 805-807, 2015.
- Gao Q, Wang XY, Zhou J and Fan J: Heterogeneity of intermediate-stage HCC necessitates personalized management including surgery. *Nat Rev Clin Oncol* 12: 10, 2015.
- Songer JG: Bacterial phospholipases and their role in virulence. *Trends Microbiol* 5: 156-161, 1997.
- Talarico S, Durmaz R and Yang Z: Insertion- and deletion-associated genetic diversity of *Mycobacterium tuberculosis* phospholipase C-encoding genes among 106 clinical isolates from Turkey. *J Clin Microbiol* 43: 533-538, 2005.
- Smith GA, Marquis H, Jones S, Johnston NC, Portnoy DA and Goldfine H: The two distinct phospholipases C of *Listeria monocytogenes* have overlapping roles in escape from a vacuole and cell-to-cell spread. *Infect Immun* 63: 4231-4237, 1995.
- Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV, Eiglmeier K, Gas S, Barry CE III, *et al*: Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* 393: 537-544, 1998.
- Fleischmann RD, Alland D, Eisen JA, Carpenter L, White O, Peterson J, DeBoy R, Dodson R, Gwinn M, Haft D, *et al*: Whole-genome comparison of *Mycobacterium tuberculosis* clinical and laboratory strains. *J Bacteriol* 184: 5479-5490, 2002.
- Kocks C, Gouin E, Tabouret M, Berche P, Ohayon H and Cossart P: *L. monocytogenes*-induced actin assembly requires the actA gene product, a surface protein. *Cell* 68: 521-531, 1992.
- Vazquezboland JA, Kocks C, Dramsi S, Ohayon H, Geoffroy C, Mengaud J and Cossart P: Nucleotide sequence of the lecithinase operon of *Listeria monocytogenes* and possible role of lecithinase in cell-to-cell spread. *Infect Immun* 60: 219-230, 1992.
- Cossart P: Illuminating the landscape of host-pathogen interactions with the bacterium *Listeria monocytogenes*. *Proc Natl Acad Sci USA* 108: 19484-19491, 2011.
- Rhee SG: Regulation of phosphoinositide-specific phospholipase C. *Annu Rev Biochem* 70: 281-312, 2001.
- Nakahara M, Shimozaawa M, Nakamura Y, Irino Y, Morita M, Kudo Y and Fukami K: A novel phospholipase C, PLC(η)2, is a neuron-specific isozyme. *J Biol Chem* 280: 29128-29134, 2005.
- Lo Vasco VR, Cardinale G and Polonia P: Deletion of PLCB1 gene in schizophrenia-affected patients. *J Cell Mol Med* 16: 844-851, 2012.
- Yang M, Li H, Li Y, Ruan Y and Quan C: Identification of genes and pathways associated with MDR in MCF-7/MDR breast cancer cells by RNA-seq analysis. *Mol Med Rep* 17: 6211-6226, 2018.
- Zhong M, Murtazina DA, Phillips J, Ku CY and Sanborn BM: Multiple signals regulate phospholipase CBeta3 in human myometrial cells. *Biol Reprod* 78: 1007-1017, 2008.
- Wu IC, Liu WC and Chang TT: Applications of next-generation sequencing analysis for the detection of hepatocellular carcinoma-associated hepatitis B virus mutations. *J Biomed Sci* 25: 51, 2018.
- Kramvis A, Arakawa K, Yu MC, Nogueira R, Stram DO and Kew MC: Relationship of serological subtype, basic core promoter and precore mutations to genotypes/subgenotypes of hepatitis B virus. *J Med Virol* 80: 27-46, 2008.
- Chu CJ, Keeffe EB, Han SH, Perrillo RP, Min AD, Soldevila-Pico C, Carey W, Brown RS Jr, Luketic VA, Terrault N and Lok AS: Hepatitis B virus genotypes in the United States: Results of a nationwide study. *Gastroenterology* 125: 444-451, 2003.
- Lin CL and Kao JH: The clinical implications of hepatitis B virus genotype: Recent advances. *J Gastroenterol Hepatol* 26 (Suppl 1): S123-S130, 2011.
- Chan HL, Hui AY, Wong ML, Tse AM, Hung LC, Wong VW and Sung JJ: Genotype C hepatitis B virus infection is associated with an increased risk of hepatocellular carcinoma. *Gut* 53: 1494-1498, 2004.
- Chu CJ, Hussain M and Lok AS: Hepatitis B virus genotype B is associated with earlier HBeAg seroconversion compared with hepatitis B virus genotype C. *Gastroenterology* 122: 1756-1762, 2002.
- Sumi H, Yokosuka O, Seki N, Arai M, Imazeki F, Kurihara T, Kanda T, Fukai K, Kato M and Saisho H: Influence of hepatitis B virus genotypes on the progression of chronic type B liver disease. *Hepatology* 37: 19-26, 2003.
- Roessler S, Jia HL, Budhu A, Forgues M, Ye QH, Lee JS, Thorgeirsson SS, Sun Z, Tang ZY, Qin LX and Wang XW: A unique metastasis gene signature enables prediction of tumor relapse in early-stage hepatocellular carcinoma patients. *Cancer Res* 70: 10202-10212, 2010.
- Roessler S, Long EL, Budhu A, Chen Y, Zhao X, Ji J, Walker R, Jia HL, Ye QH, Qin LX, *et al*: Integrative genomic identification of genes on 8p associated with hepatocellular carcinoma progression and patient survival. *Gastroenterology* 142: 957-966.e912, 2012.
- Tang Z, Li C, Kang B, Gao G, Li C and Zhang Z: GEPIA: A web server for cancer and normal gene expression profiling and interactive analyses. *Nucleic Acids Res* 45: W98-W102, 2017.
- Carithers LJ, Ardlie K, Barcus M, Branton PA, Britton A, Buia SA, Compton CC, DeLuca DS, Peter-Demchok J, Gelfand ET, *et al*: A novel approach to high-quality postmortem tissue procurement: The GTEx project. *Biopreserv Biobank* 13: 311-319, 2015.
- Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, *et al*: Proteomics. Tissue-based map of the human proteome. *Science* 347: 1260419, 2015.
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES and Mesirov JP: Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* 102: 15545-15550, 2005.
- Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstråle M, Laurila E, *et al*: PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 34: 267-273, 2003.
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B and Ideker T: Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13: 2498-2504, 2003.
- Montejo J, Zuberi K, Rodriguez H, Kazi F, Wright G, Donaldson SL, Morris Q and Bader GD: GeneMANIA Cytoscape plugin: Fast gene function predictions on the desktop. *Bioinformatics* 26: 2927-2928, 2010.
- Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, Santos A, Doncheva NT, Roth A, Bork P, *et al*: The STRING database in 2017: Quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res* 45: D362-D368, 2017.
- Maere S, Heymans K and Kuiper M: BiNGO: A Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics* 21: 3448-3449, 2005.
- Lossos IS, Czerwinski DK, Alizadeh AA, Wechsler MA, Tibshirani R, Botstein D and Levy R: Prediction of survival in diffuse large-B-cell lymphoma based on the expression of six genes. *N Engl J Med* 350: 1828-1837, 2004.
- Alizadeh AA, Gentles AJ, Alencar AJ, Liu CL, Kohrt HE, Houot R, Goldstein MJ, Zhao S, Natkunam Y, Advani RH, *et al*: Prediction of survival in diffuse large B-cell lymphoma based on the expression of 2 genes reflecting tumor and microenvironment. *Blood* 118: 1350-1358, 2011.
- Liao X, Han C, Wang X, Huang K, Yu T, Yang C, Huang R, Liu Z, Han Q and Peng T: Prognostic value of minichromosome maintenance mRNA expression in early-stage pancreatic ductal adenocarcinoma patients after pancreaticoduodenectomy. *Cancer Manag Res* 10: 3255-3271, 2018.

42. Liao X, Liu X, Yang C, Wang X, Yu T, Han C, Huang K, Zhu G, Su H, Qin W, *et al*: Distinct diagnostic and prognostic values of minichromosome maintenance gene expression in patients with hepatocellular carcinoma. *J Cancer* 9: 2357-2373, 2018.
43. Golebiewska U, Guo Y, Khalikaprassad N, Zurawsky C, Yerramilli VS and Scarlata S: γ -Synuclein interacts with phospholipase C β 2 to modulate G protein activation. *PLoS One* 7: e41067, 2012.
44. Harden TK and Sondek J: Regulation of phospholipase C isozymes by ras superfamily GTPases. *Annu Rev Pharmacol Toxicol* 46: 355-379, 2006.
45. Martelli AM, Fiume R, Faenza I, Tabellini G, Evangelista C, Bortul R, Follo MY, Falà F and Cocco L: Nuclear phosphoinositide specific phospholipase C (PI-PLC)-beta 1: A central intermediary in nuclear lipid-dependent signal transduction. *Histol Histopathol* 20: 1251-1260, 2005.
46. Ngho A, Mctague A, Wentzensen IM, Meyer E, Applegate C, Kossoff EH, Batista DA, Wang T and Kurian MA: Severe infantile epileptic encephalopathy due to mutations in PLCB1: Expansion of the genotypic and phenotypic disease spectrum. *Dev Med Child Neurol* 56: 1124-1128, 2014.
47. Faenza I, Matteucci A, Manzoli L, Billi AM, Aluigi M, Peruzzi D, Vitale M, Castorina S, Suh PG and Cocco L: A role for nuclear phospholipase Cbeta 1 in cell cycle control. *J Biol Chem* 275: 30520-30524, 2000.
48. Cocco L, Finelli C, Mongiorgi S, Clissa C, Russo D, Bosi C, Quaranta M, Malagola M, Parisi S, Stanzani M, *et al*: An increased expression of PI-PLC β 1 is associated with myeloid differentiation and a longer response to azacitidine in myelodysplastic syndromes. *J Leukoc Biol* 98: 769-780, 2015.
49. Guo Y and Scarlata S: A loss in cellular protein partners promotes α -synuclein aggregation in cells resulting from oxidative stress. *Biochemistry* 52: 3913-3920, 2013.
50. Poli A, Faenza I, Chiarini F, Matteucci A, McCubrey JA and Cocco L: K562 cell proliferation is modulated by PLC β 1 through a PKC α -mediated pathway. *Cell Cycle* 12: 1713-1721, 2013.
51. Bavelloni A, Poli A, Fiume R, Blalock W, Matteucci A, Ramazzotti G, McCubrey JA, Cocco L and Faenza I: PLC-beta 1 regulates the expression of miR-210 during mithramycin-mediated erythroid differentiation in K562 cells. *Oncotarget* 5: 4222-4231, 2014.
52. Ho KK and Mann DJ: Nuclear signalling through phospholipase C and phosphatidylinositol 4,5-bisphosphate. *Signal Transduction* 6: 92-100, 2006.
53. Bertagnolo V, Benedusi M, Brugnoli F, Lanuti P, Marchisio M, Querzoli P and Capitani S: Phospholipase C- β 2 promotes mitosis and migration of human breast cancer-derived cells. *Carcinogenesis* 28: 1638-1645, 2007.
54. Bertagnolo V, Benedusi M, Brugnoli F, Lanuti P, Marchisio M, Querzoli P and Capitani S: Phospholipase C-beta 2 promotes mitosis and migration of human breast cancer-derived cells. *Carcinogenesis* 28: 1638-1645, 2007.
55. Bertagnolo V, Benedusi M, Querzoli P, Pedriali M, Magri E, Brugnoli F and Capitani S: PLC-beta2 is highly expressed in breast cancer and is associated with a poor outcome: A study on tissue microarrays. *Int J Oncol* 28: 863-872, 2006.
56. Ståhlberg P, Lopez-Egido JR, Wang S, Gobl A, Oberg K and Skogseid B: Differentially expressed cDNAs in PLCbeta3-induced tumor suppression in a human endocrine pancreatic tumor cell line: Activation of the human mismatch repair protein 3 gene. *Biochem Biophys Res Commun* 281: 227-231, 2001.
57. Jiang Y, Wang S, Holcomb J, Trescott L, Guan X, Hou Y, Brunzelle J, Sirinpong N, Li C and Yang Z: Crystallographic analysis of NHERF1-PLC β 3 interaction provides structural basis for CXCR2 signaling in pancreatic cancer. *Biochem Biophys Res Commun* 446: 638-643, 2014.
58. Hoepfner LH, Phoenix KN, Clark KJ, Bhattacharya R, Gong X, Sciuto TE, Vohra P, Suresh S, Bhattacharya S, Dvorak AM, *et al*: Revealing the role of phospholipase C β 3 in the regulation of VEGF-induced vascular permeability. *Blood* 120: 2167-2173, 2012.
59. Xiao W, Hong H, Kawakami Y, Kato Y, Wu D, Yasudo H, Kimura A, Kubagawa H, Bertoli LF, Davis RS, *et al*: Tumor suppression by phospholipase C-beta3 via SHP-1-mediated dephosphorylation of Stat5. *Cancer Cell* 16: 161-171, 2009.
60. Ju H, Zhao S, Tappia PS, Panagia V and Dixon IM: Expression of Gq alpha and PLC-beta in scar and border tissue in heart failure due to myocardial infarction. *Circulation* 97: 892-899, 1998.
61. Sarna JR, Marzban H, Watanabe M and Hawkes R: Complementary stripes of phospholipase Cbeta3 and Cbeta4 expression by Purkinje cell subsets in the mouse cerebellum. *J Comp Neurol* 496: 303-313, 2006.
62. Tan X and Chen M: MYLK and MYL9 expression in non-small cell lung cancer identified by bioinformatics analysis of public expression data. *Tumour Biol* 35: 12189-12200, 2014.
63. Orchel J, Witek L, Kimsa M, Strzalka-Mrozik B, Kimsa M, Olejek A and Mazurek U: Expression patterns of kinin-dependent genes in endometrial cancer. *Int J Gynecol Cancer* 22: 937-944, 2012.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.