

Ginsenoside CK induces apoptosis and suppresses proliferation and invasion of human osteosarcoma cells through the PI3K/mTOR/p70S6K1 pathway

KANG CHEN¹, JIANBAO JIAO¹, JINWEI XUE¹, TAOPING CHEN¹, YU HOU¹, YUANTAO JIANG¹,
LIGANG QIAN¹, YUNFEI WANG¹, ZHENG MA¹, ZHIXING LIANG¹, BO SUN¹ and QIAO REN²

Departments of ¹Orthopedics and ²Rheumatism and Immunology,
Affiliated Hospital of Hebei University, Baoding, Hebei 071000, P.R. China

Received June 21, 2019; Accepted December 4, 2019

DOI: 10.3892/or.2020.7460

Abstract. Osteosarcoma is one of the most malignant bone tumors, and its major threats are aggressive invasion and early tumor metastasis, which result in a poor prognosis and high mortality. Accumulating evidence indicates that ginsenoside compound K (CK) has a significant antitumor effect, particularly on the inhibition of proliferation and invasion of numerous human tumors. In the present study, it was revealed that CK inhibited the viability and proliferation of osteosarcoma cells. Moreover, it was demonstrated that CK induced apoptosis and inhibited the migration and invasion of osteosarcoma cells via apoptotic staining, Annexin V/PI staining, and Transwell invasion assays. Furthermore, at the molecular level, the present results confirmed that apoptosis and invasion-related proteins were regulated by CK, which was possibly related to the blockade of the PI3K/mTOR/p70S6K1 signaling pathway. In summary, the present findings indicated that CK inhibited viability and proliferation, induced apoptosis, and inhibited the migration and invasion of osteosarcoma cells through the PI3K/mTOR/p70S6K1 signaling pathway.

Introduction

Osteosarcoma mainly occurs in teenagers and healthy young patients, and is one of the most common fatal primary malignant bone tumors (1,2), accounting for 5% of all pediatric tumors (3). The major threats of osteosarcoma are aggressive invasion and early multi-organ metastasis, especially pulmonary metastasis (4). Although survival has been greatly increased by surgical treatment combined with chemotherapy,

the 5 year survival rate of patients with metastasis is still less than 35% (5). Therefore, the main strategy for the treatment of osteosarcoma is to block its invasiveness and metastasis to improve its prognosis. However, there are currently no effective drugs for osteosarcoma and the molecular mechanism remains obscure. Therefore, exploring targeted, highly effective treatments for osteosarcoma is urgently required.

The PI3K/mTOR/p70S6K1 pathway is an essential signaling pathway in cells, and has an extremely important biological function in cell growth, proliferation, apoptosis, angiogenesis and autophagy, as well as in other processes. Disorders of the pathway can cause a range of diseases, including cancer, neuropathy, and autoimmune diseases (6). The ribosomal 40S small subunit S6 protein kinase 1 (p70S6K1) is a direct substrate downstream of p-mTOR. P70S6K1 is phosphorylated by p-mTOR to promote the production of ribosomes, initiation factors, and elongation factors in cells, and to promote proliferation and invasion of tumor cells (7,8).

Ginsenoside compound K (20-O-D-glucopyranosyl-20(S)-protopanaxadiol; CK; as shown in Fig. 1), is a major member of the original ginseng diol, saponins, and its study has been a popular research topic in the fields of anti-inflammatory and antitumor effects. Studies have revealed that ginsenoside CK is a compound with a variety of targets and pharmacological activities, which include anti-inflammatory (9), antitumor (10), anti-diabetic (11) and liver protection (12). However, to date, the antitumor mechanism of ginsenoside CK on osteosarcoma cells remains unclear. Therefore, in the present study, it was investigated whether CK exerted cellular antiproliferative effects and induced apoptosis in osteosarcoma cells. It was further explored whether the PI3K/mTOR/p70S6K1 signaling pathway played a decisive role in this process.

Materials and methods

Antibodies and reagents. Dulbecco's Modified Eagle's Medium (DMEM), high glucose medium, and fetal bovine serum (FBS) were obtained from Gibco; Thermo Fisher scientific, Inc. Human osteosarcoma cell lines, MG63 and U2-OS, were purchased from the Cell Bank of Shanghai Institutes for Biological Sciences. Ginsenoside CK (purity: >98%) was

Correspondence to: Professor Qiao Ren, Department of Rheumatism and Immunology, Affiliated Hospital of Hebei University, 212 Yuhua East Road, Baoding, Hebei 071000, P.R. China
E-mail: chkcmu@163.com

Key words: osteosarcoma, ginsenoside CK, proliferation, apoptosis, invasion

purchased from Beijing Solarbio Science & Technology Co., Ltd. MTT, dimethylsulfoxide (DMSO), and trypsin were purchased from Sigma-Aldrich; Merck KGaA. The acridine orange/ethidium bromide (AO/ET) staining kit was purchased from Beijing Solarbio Science & Technology Co., Ltd. The BrdU cell proliferation kit was purchased from EMD Millipore. The TUNEL apoptosis kit was purchased from R&D Systems. Matrigel was purchased from Collaborative Biomedical Products, Inc. Transwell invasion chambers were purchased from Corning Costar. Rabbit antibodies to Bax (cat. no. ab32503), Bcl-2 (cat. no. ab182858), and caspase-3 (cat. no. ab2302) were purchased from Abcam. Rabbit antibodies to MMP-2 (cat. no. 10373-2-AP) and MMP-9 (cat. no. 10375-2-AP) were purchased from ProteinTech Group, Inc. Rabbit antibodies to PTEN (cat. no. 9188), serine/threonine kinase (Akt) (cat. no. 4691), phosphor-Akt (p-Akt) (cat. no. 4060), mTOR (cat. no. 2983), phosphor-mTOR (p-mTOR) (cat. no. 5536), and p70S6K1 (cat. no. 2708) were purchased from Cell Signaling Technology, Inc. RAD001 was purchased from Selleck Chemicals.

Cell culture. Human osteosarcoma cell lines, MG63 and U2-OS, were both cultured in DMEM high glucose medium containing 10% FBS in a humidified incubator with 5% CO₂ at 37°C. When the cells were ~80% confluent, the cells were harvested with 0.25% trypsin plus 0.02% EDTA and subcultured. Both MG63 and U2-OS cells were divided into two groups: The negative control groups (NC groups) and the ginsenoside CK treatment groups (CK groups).

Cell proliferation analysis-MTT and BrdU. The viability of cells was evaluated by the MTT method. The two groups of osteosarcoma cells (3x10³) were incubated in a 96-well plate for 24 h, then the cells were treated with various concentrations of drugs (CK 0-30 µM or PBS), each concentration in parallel 6-wells. After both groups were incubated for 1-5 days, 20 µl MTT solution was added to each well. After 4 h, the medium was discarded and 150 µl DMSO was added to dissolve the formazan crystals. The absorbance was measured using a wavelength of 490 nm. The proliferation of cells was analyzed with a BrdU cell proliferation kit following the protocol of the manufacturer, similar to the aforementioned method, and the absorbance was measured at a wavelength of 450 nm.

AO/EB apoptosis assay. Apoptosis-related morphological changes were detected using an AO/EB staining method. After treatment, both groups of cells were seeded on cover slides in a 24-well plate according to the manual, then 500 µl of freshly-prepared dual stain containing 10 mg/ml AO and 10 mg/ml EB was added to each well to stain for 10 min in the dark at room temperature. The morphological changes of the cell nuclei were observed using a fluorescence microscope, and the apoptotic rate was calculated.

TUNEL apoptosis assay. A TUNEL apoptosis kit was used to further detect the morphological changes of the apoptotic nuclei. After induction of apoptosis, the cells were fixed in 4% paraformaldehyde for 30 min, then PBS containing 0.3% Triton X-100 was added and incubated for 5 min at room temperature. After washing twice with PBS, 50 µl of TUNEL

assay solution and 20 µl DAPI were added to each well of a 24-well plate and incubated at 37°C for 60 min in the dark. Then, the apoptotic nuclei were immediately observed using a fluorescence microscope, and the apoptotic rate was calculated.

Annexin V/PI apoptosis detection. Cell apoptosis was detected using an Annexin V/PI apoptosis detection kit according to the manufacturer's protocols. Briefly, after induction of apoptosis, the two cell types were harvested by trypsinization and washed twice in ice-cold PBS. The cells were centrifuged at 500 x g for 5 min and resuspended in 500 µl binding buffer, then incubated with 5 µl Annexin V-FITC and 10 µl propidium iodide (PI) in the dark at room temperature for 15 min. The samples were immediately analyzed using a FACSCalibur (BD Biosciences) flow cytometer and the data were analyzed using CellQuest software, version 5.1 (BD Biosciences).

Cell cycle analysis. After treatment with CK (0-30 µM) for 24 h, both groups of osteosarcoma cells were fixed in 70% ethanol for 24 h after washing twice with cold PBS at 4°C before analysis. After centrifugation (1,000 g), the cells were stained with 50 µg/ml PI and 100 µg/ml RNase A for 30 min in the dark at room temperature. The samples were analyzed by a BD FACSCalibur flow cytometer and the CellQuest software was used to analyze the data.

Cell migration and invasion detection. The migration and invasive abilities of both groups of osteosarcoma cells were evaluated by Transwell migration and invasion systems. For the invasion assay, 20 µl Matrigel was applied to cover the bottom of the chamber after being washed with serum-free medium to create a Matrigel membrane by incubating at 37°C for 30 min. Cells (5x10⁴) with 200 µl of serum-free DMEM medium were seeded in the upper chamber of the Transwell system, and 500 µl of DMEM high glucose medium containing 10% FBS was added to the lower chamber as a chemoattractant. The Transwell system was incubated in a cell culture incubator for 24 h, and then the upper chamber was removed and the cells from the upper surface of the membrane were removed. Then, the cells invading to the lower surface of the membrane were stained with 0.1% crystal violet for 5 min at room temperature and observed using an inverted phase-contrast microscope. The procedure for the migration assay was the same as above, but a Matrigel membrane was not required.

Western blot analysis. Cells were lysed in a radioimmuno-precipitation assay (RIPA) lysis buffer (Solarbio Science & Technology) and the BCA method was used for protein determination. The protein samples from each group were resolved by SDS-PAGE and transferred to PVDF membranes (proteins were loaded by 50 µg/lane and the percentage of the gel was 10%). After transfer, the PVDF membranes were blocked with 5% (w/v) nonfat milk in TBST at room temperature for 2 h, and then incubated with the primary antibodies (1:2,000) in TBST buffer containing 1% (w/v) BSA at 4°C for 16 h. Following three washes in TBST, the membranes were incubated with the appropriate HRP-linked secondary antibody (goat anti-rabbit 1:1,000; cat. no. SA00001-2; ProteinTech Group) in TBST for 2 h at room temperature. Proteins were detected by an ECL system (GE Healthcare).




Figure 1. The chemical structure of CK. CK, ginsenoside compound K.

Statistical analysis. All statistical analyses were performed with one-way analysis of variance (ANOVA) and Newman-Keuls post hoc test using SPSS statistical software for Windows, version 13.0 (SPSS, Inc.), and GraphPad Prism 6 software (GraphPad Software, Inc.) was used to draw statistical charts. Data are presented as the mean \pm SEM from a minimum of three independent experiments performed in triplicate. Differences with $P < 0.05$ were considered to indicate a statistically significant difference.

Results

CK inhibits the viability and proliferation and induces G2/M arrest of osteosarcoma cells. The MTT assay revealed that there was no significant difference in cell viability between the CK and normal control (NC) groups of MG-63 and U2-OS cells on days 0-2 after drug treatment ($P > 0.05$). However, after 3 days, CK exhibited significant inhibition of viability and proliferation against MG-63 and U2-OS cells ($P < 0.05$) in a time-dependent manner (Fig. 2A). Cells were treated with different concentrations of CK (0, 5, 10, 15, 20, 25 and 30 μ M). The IC_{50} value for 72 h was ~ 20 μ M, which was used as the effective drug concentration in subsequent experiments (Fig. 2A). The results of the BrdU assay further confirmed this conclusion; cells in the CK group exhibited significant inhibition of viability, and at the same drug concentration (20 μ M), the inhibition of MG-63 viability was more pronounced than that of U2-OS cells (Fig. 2B). Furthermore, it was also revealed that compared to the control group, cells in the CK group exhibited G2/M-phase cell cycle arrest, which was possibly related to cell proliferation and apoptosis (Fig. 2C and D).

Apoptotic changes of cell nuclei dual-stained with AO/EB. The abovementioned results confirmed that CK inhibited the viability and proliferation of osteosarcoma cells. To further analyze the apoptosis induction effect of CK in MG-63 and U2-OS cells, AO/EB nuclear staining was performed to observe the morphological changes of apoptotic nuclei. As revealed in Fig. 3A and B, when cells were treated with CK (20 μ M), MG-63 and U2-OS cells exhibited different degrees of apoptosis. Based on analyses of the histograms in Fig. 3, the apoptosis rates in the CK groups (MG-63, $24.16 \pm 2.25\%$; U2-OS, $17.66 \pm 1.37\%$) were significantly higher than those in

the NC groups (MG-63, $3.34 \pm 1.16\%$; U2-OS, $1.65 \pm 0.84\%$) ($P < 0.05$).

CK induces apoptosis of osteosarcoma cells detected by TUNEL assays. A TUNEL apoptosis kit was used to further detect the morphological changes of apoptotic nuclei. As revealed in Fig. 4, in the MG-63 group, the apoptosis rate of the NC group was only $1.52 \pm 0.63\%$, and the value increased to $25.33 \pm 2.68\%$ after treatment with CK (20 μ M). Similar results were also obtained in the U2-OS group [$0.45 \pm 0.22\%$ vs. $24.83 \pm 1.92\%$ (NC group)]. These results indicated that CK induced apoptosis in both MG-63 and U2-OS cells.

CK induces apoptosis in osteosarcoma cells. The Annexin V/propidium iodide (PI) staining assay was employed to further confirm the apoptotic effect of CK. As revealed in Fig. 5A and B, after treatment with CK, the percentage of apoptosis in MG-63 and U2-OS cells in the experimental group was significantly higher than that in the control group ($P < 0.05$). In the MG-63 group, both early apoptosis ($5.42 \pm 1.90\%$) and late apoptosis ($16.71 \pm 1.65\%$) were higher than that in the control groups (3.32 ± 0.85 and $5.83 \pm 1.08\%$, respectively), and the total apoptosis was more pronounced (22.17 ± 2.45 vs. $9.15 \pm 1.85\%$). U2-OS cells revealed a similar result (9.74 ± 2.20 vs. $3.59 \pm 2.17\%$ for total apoptosis). To validate the aforementioned conclusions at the cellular pathway level, the expression levels of apoptosis-related proteins were detected by western blot analyses. As revealed in Fig. 5, compared with the control group, the expression levels of the proapoptotic proteins cleaved caspase-3 and BAX were significantly upregulated in the CK group in MG-63 and U2-OS cells, while the expression of the inhibitory protein, Bcl-2, was downregulated ($P < 0.05$). From the aforementioned results, it was concluded that CK significantly induced apoptosis in osteosarcoma cells.

CK inhibits the migration and invasion of osteosarcoma cells. In the previous results, it was demonstrated that CK inhibited proliferation and induced apoptosis of osteosarcoma cells. To further confirm the effect of CK on the migration and invasion of osteosarcoma cells, a Transwell assay was used. As revealed in Fig. 6A, the number of cells in the CK group passing through the Transwell chamber membrane were significantly lower in the MG-63 and U2-OS CK-treated groups than in the control group [invasion: MG63, 75.72 ± 19.13 vs. 318.16 ± 26.37 (control), $P < 0.05$; U2-OS, 45.83 ± 14.22 vs. 362.37 ± 21.71 (control), $P < 0.05$; migration: MG63, 123.15 ± 12.39 vs. 398.63 ± 31.40 (control), $P < 0.05$; U2-OS, 132.94 ± 29.91 vs. 453.33 ± 34.17 (control), $P < 0.05$], indicating that CK inhibited both the migration and invasion of MG-63 and U2-OS cells. To further characterize the relationship between CK and the mTOR signaling pathway, RAD001, a specific inhibitor of mTOR, was added. As revealed in Fig. 6B, the number of cells in the CK or RAD001 group passing through the Transwell chamber membrane was significantly lower than in the NC group, and this trend was more pronounced when cells were treated with both CK and RAD001. Therefore, it was hypothesized that CK played a similar role as an mTOR inhibitor, and that this effect was related to the mTOR signaling pathway. Furthermore, the expression of the invasion-related proteins,




Figure 2. CK inhibits viability and proliferation of MG63 and U2-OS cells. (A) Cells were treated with different concentrations of CK (0, 5, 10, 15, 20, 25 and 30 μ M). CK exhibited significant inhibitory effects on MG63 and U2-OS cells, as revealed by MTT assay. (B) BrdU cell proliferation kits were used to evaluate the anti-proliferation effect of CK on MG63 and U2-OS cells. (C) Propidium iodide staining cell cycle analyses were performed to detect the distribution of cells at each state of the cell cycle following CK treatment. (D) Cell cycle distribution of CK-treated cells revealed G2/M-phase arrest. * P <0.05 vs. the NC group. Experiments were repeated three times and the results are expressed as the mean \pm SEM. CK, ginsenoside compound K; NC, normal control.

MMP-2 and MMP-9 were detected, in each group by western blot analyses. As indicated in Fig. 6C, the results revealed that after treatment with CK, MMP-2 and MMP-9 expression levels were significantly downregulated in MG-63 and U2-OS cells (P <0.05), which was consistent with the aforementioned conclusions, and confirmed that CK inhibited the migration and invasion of osteosarcoma cells.

CK is involved in the PI3K/mTOR/p70S6K1 signaling pathway in osteosarcoma cells. To further investigate the antitumor

mechanism of CK on MG-63 and U2-OS cells, the expression levels of related proteins in the PI3K/mTOR/p70S6K1 signaling pathway were investigated. The results of western blot analyses revealed that after treatment with CK, PTEN the major blocking protein of the PI3K/mTOR/p70S6K1 pathway was significantly upregulated in MG-63 and U2-OS cells (P <0.05; Fig. 7A). However, the core action proteins of the PI3K/mTOR/p70S6K1 pathway, including p-AKT and p-mTOR, were significantly downregulated in the CK group (P <0.05; Fig. 7A and B), indicating that the PI3K/mTOR/p70S6K1 pathway was blocked




Figure 3. Apoptosis-related morphological changes of cell nuclei induced by CK in MG63 and U2-OS cells. (A and B) Cells were treated with or without CK (20 μ M) and cell nuclei were stained with AO/EB and observed under a fluorescence microscope. Cells with condensed or fragmented nuclei were considered apoptotic, and early apoptotic nuclei presented as green or light yellow; late apoptotic nuclei presented as orange or red (N, normal nuclei; A, apoptotic nuclei). Experiments were repeated three times. * $P < 0.05$ vs. the NC group. CK, ginsenoside compound K; AO/EB, acridine orange/ethidium bromide; NC, normal control.




Figure 4. Apoptotic changes of cell nuclei induced by CK in MG63 and U2-OS cells detected by TUNEL assays. (A and B) Cells were treated with or without CK (20 μ M) and cell nuclei were stained with DAPI and TUNEL assay solution and observed under a fluorescence microscope. Normal nuclei exhibited regular elliptical shapes and were stained blue by DAPI; apoptotic nuclei presented as condensed or fragmented and were stained green by TUNEL assay solution. Compared with the control group, the CK group exhibited significant apoptosis. Data are expressed as the means \pm SEM. All experiments were repeated three times. * $P < 0.05$ vs. the NC group. CK, ginsenoside compound K; NC, normal control.

after treatment with CK. The downregulation of p70S6K1 expression in the CK group ($P < 0.05$; Fig. 7B) further confirmed

this conclusion. Furthermore, the expression levels of p-mTOR and p70S6K1 were downregulated when treated with RAD001




Figure 5. Determination of apoptosis in MG63 and U2-OS cells via Annexin V/PI staining and western blot analyses. (A) Annexin V/PI apoptosis detection kits were used to detect the induction of apoptosis. Early apoptosis is localized in the bottom right area and late apoptosis in the top right area. (B) Whether early or late apoptosis, the apoptotic ratio in the CK groups was significantly higher than that in the control group. * $P < 0.05$ vs. the NC group. (C) Detection of the expression of apoptosis-related proteins (cleaved caspase-3, Bcl-2, and BAX) by western blot analyses. Cleaved caspase-3 and BAX expression levels were upregulated, while Bcl-2 expression was downregulated, in both MG63 and U2-OS cells after treatment with CK. In addition, the ratio of BAX/Bcl-2 was upregulated in both groups. * $P < 0.05$ vs. the NC. All data represent the mean \pm SEM of three independent experiments. CK, ginsenoside compound K; NC, normal control.

in U2-OS cells, which was consistent with the effect of CK. When cells were treated with both CK and RAD001, they played a synergistic role and the downregulation of the expression levels of p-mTOR and p70S6K1 were more pronounced (Fig. 7C). Overall, these conclusions confirmed that CK blocked the PI3K/mTOR/p70S6K1 signaling pathway of MG-63 and U2-OS osteosarcoma cells.

Discussion

Ginsenoside CK is one of the main metabolites of ginsenoside, which is a glycoside compound composed of sugars and aglycones, and belongs to a triterpenoid compound family. According to the structure of sapogenin, it can be divided into three categories: One is an oleanane-type pentacyclic triterpenoid sapogenin, the sapogenin is oleanolic acid, and the other two are dammarane-type tetracyclic triterpenoid sapogenins, which are the majority of ginsenosides and are the primary active ingredients of ginseng. According to different aglycons,

the dammarane type tetracyclic triterpenoid sapogenin can be divided into the original ginseng diol sapogenin and the original ginseng triol sapogenin. Ginsenoside CK, a major member of the originally reported ginseng diol sapogenin, has been a topic of research in the fields of anti-inflammatory and antitumor effects. Recent studies have revealed that ginsenoside CK has anti-inflammatory, antioxidative, and neuroprotective effects (13-16), and of particular interest are its antitumor effects.

Studies have confirmed that CK has an inhibitory effect on various tumor cells, including liver, lung, breast and colon cancer as well as leukemia (12,17-19). The antitumor effect of CK is mainly reflected in its ability to significantly attenuate the proliferation, invasion, and migration of tumor cells. Lee *et al* (20) revealed that CK significantly inhibited the proliferation and invasion of malignant glioma cells by blocking the PI3K/AKT/mTOR signaling pathway. Kang *et al* (21) reported that CK inhibited colon cancer cell proliferation and induced apoptosis by inhibiting histone deacetylase activity.




Figure 6. Migration and invasion of MG63 and U2-OS cells after treatment with CK. (A) Cell invasion and migration abilities were detected by Transwell migration assays and observed under an inverted phase-contrast microscope. Histogram analyses revealed that CK effectively inhibited both the migration and invasion of MG63 and U2-OS cells. * $P < 0.05$ vs. the NC group. (B) RAD001, a specific inhibitor of mTOR, was added. The number of U2-OS cells in the CK or RAD001 group passing through the Transwell chamber membrane was significantly lower than in the NC group, and this trend was more pronounced when cells were treated with both CK and RAD001. * $P < 0.05$ vs. the NC group. (C) Western blots revealed that CK downregulated the expression of invasion-related proteins (MMP-2 and MMP-9) in both MG63 and U2-OS cells. * $P < 0.05$ vs. the NC group. CK, ginsenoside compound K; NC, normal control.

Osteosarcoma is one of the most malignant bone tumors, and its lethality is mainly reflected in the malignant, diffuse proliferative capacity, and early tumor metastasis. Therefore, it was speculated whether CK also had an inhibitory effect on the proliferation and invasion of osteosarcoma cells. To demonstrate the effect of CK on the *in vitro* viability and proliferation of osteosarcoma cells in this study, MG-63 and U2-OS cells were treated with CK. Both MTT and BrdU assay results confirmed that CK significantly reduced the

viability and proliferation of MG-63 and U2-OS cells *in vitro*. Furthermore, it was revealed that the effect of CK on cell proliferation may be related to cell cycle distribution. Using PI staining cell cycle analyses in both MG-63 and U2-OS cells, it was determined that CK altered the distribution of cell cycle stages. After treatment with CK, the S-phase ratio of MG-63 cells decreased from 51.06 ± 3.31 to $36.96 \pm 3.07\%$, compared to the control group, while the G2/M-phase ratio increased from 16.39 ± 2.59 to $35.23 \pm 2.33\%$. Similarly, the proportion




Figure 7. Western blot analyses of proteins in the PI3K/mTOR/p70S6K1 signaling pathway of MG63 and U2-OS cells. (A) PTEN expression was upregulated, while p-Akt expression and the ratio of p-Akt/Akt were downregulated in both MG63 and U2-OS cells after treatment with CK ($P < 0.05$ vs. the NC group). (B) p-mTOR, the ratio of p-mTOR/mTOR and p-70S6k1 expression levels were downregulated in both MG63 and U2-OS cells after treatment with CK ($P < 0.05$ vs. the NC group). (C) The expression levels of p-mTOR, the ratio of p-mTOR/mTOR and p70S6K1 were downregulated when treated with RAD001 in U2-OS cells, and the downregulation was more pronounced when treated with both CK and RAD001. * $P < 0.05$ vs. the NC group. All data represent the mean \pm SEM of three independent experiments. CK, ginsenoside compound K; NC, normal control.

of the S phase decreased from 38.00 ± 3.57 to $28.73 \pm 4.08\%$, while the proportion of the G2/M phase increased from 36.25 ± 2.77 to $45.96 \pm 3.41\%$ in U2-OS cells. Therefore, it was demonstrated that CK induced G2/M-phase cell cycle arrest, and inhibited the viability and proliferation of MG-63 and U2-OS cells.

In order to analyze the apoptosis-inducing effect of CK, apoptotic cell nuclear morphology assays, Annexin V/PI staining assays, and western blot analyses were performed. It was determined that the cell nuclei of the CK-treated group exhibited typical apoptotic morphological changes. Similar results were detected by TUNEL assays, whether using MG-63 or U2-OS cells. The apoptotic nuclei were stained green, while the normal nuclei without apoptosis were stained blue by DAPI staining. The aforementioned results demonstrated that CK had an apoptosis-inducing effect on both MG-63 and U2-OS cells using nuclear apoptotic morphological assays.

Therefore, using nuclear apoptotic morphological assays, it was confirmed that CK had a significant apoptosis-inducing effect on osteosarcoma cells, and the conclusions were further

confirmed by Annexin V/PI staining and western blot analyses. In both MG-63 and U2-OS cells, early apoptosis (5.42 ± 1.90 and $5.66 \pm 0.88\%$) and late apoptosis (16.71 ± 1.65 and $4.17 \pm 0.60\%$) were significantly higher in the CK-treated group than in the control group. The expression levels of cleaved caspase-3 and BAX, key proteins in the apoptotic pathway, were significantly upregulated in the CK group, and the downregulation of the anti-apoptotic protein, Bcl-2, further confirmed the activation of apoptosis. These results confirmed that CK-induced apoptosis in MG-63 and U2-OS cells at the level of cellular signaling pathways.

The aforementioned results revealed that CK inhibited proliferation and induced apoptosis of MG-63 and U2-OS cells, and the results are consistent with previous studies which revealed that CK has antitumor effects on various tumor cells, including liver, lung, breast and colon cancer as well as leukemia (17-21); however, another characteristic feature of osteosarcoma cells is their aggressive invasiveness and early metastasis. Since osteosarcoma is prone to early metastasis (most common in lung), most patients lose the opportunity for surgical treatment, and there is no effective

radiotherapy and chemotherapy for osteosarcoma in the clinic. Thus, patients usually have a poor prognosis and a high mortality rate (22,23).

To investigate the effects of CK on the migration and invasion of MG-63 and U2-OS cells, a Transwell assay was used and the expression levels of the invasion-related proteins, MMP-2 and MMP-9, were detected. The migration and invasion of MG-63 and U2-OS cells significantly decreased after treatment with CK, and downregulation of the expression of MMP-2 and MMP-9 also demonstrated that CK inhibited the migration and invasion of MG-63 and U2-OS cells. CK played a similar role to RAD001, a specific inhibitor of mTOR, and together they played a synergistic role in the effects of the migration and invasion. Chawla *et al* (24) reported that the mTOR inhibitor, Ridaforolimus, inhibited the phosphorylation of the mTOR effector protein, S6K, to block the PI3K/AKT pathway. Such inhibition also effectively inhibited the tumor characteristics of osteosarcoma, and achieved significant clinical effects. Moriceau *et al* (25) reported that the mTOR inhibitor, RAD001 (Everolimus), inhibited osteosarcoma cell proliferation in a dose- and time-dependent manner. Manara *et al* (26) reported that NVP-BEZ235, another mTOR inhibitor, significantly inhibited the proliferation and invasion of osteosarcoma cells, and was a possible novel potential targeted drug for the treatment of osteosarcoma. A number of previous studies have demonstrated that blocking the PI3K/mTOR/p70S6K1 signaling pathway by mTOR inhibitors inhibited osteosarcoma cell activity. Therefore, it was speculated that osteosarcoma cells may play a pathogenic role through the PI3K/mTOR/p70S6K1 pathway.

PI3K/mTOR/p70S6K1 studies have been a popular research topic in recent years. As an essential signaling pathway in cells, it plays an important biological function in cell growth, proliferation, apoptosis, angiogenesis, and autophagy. Disorders of the pathway can cause a range of diseases, including cancer, neuropathy, and autoimmune diseases (27). The phosphatidylinositol 3-kinase (PI3K) protein family is involved in the regulation of various cellular functions such as cell proliferation, differentiation, apoptosis, and glucose transport. Increases in PI3K activity are often associated with a variety of cancers (28). Cytokines such as fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), human growth factor (HGF), vascular protein I (Ang1), and insulin activate PI3Ks, and the SH2 and SH3 domains of the p85 subunit of PI3Ks bind to the adaptor protein at a phosphorylation site. PI3K initiates phosphorylation of various PI intermediates after recruitment of activated receptors. Following this, PI3K converts PIP2 into PIP3, a process that is particularly relevant to tumors (29). The result of PI3K activation is the generation of a second messenger, PIP3, on the plasma membrane. PIP3 binds to the PH domain-containing signaling proteins, AKT and phosphoinositide dependent kinase-1 (PDK1), which promotes PDK1 phosphorylation of AKT Ser308 to activate AKT (30,31). Phosphorylated AKT activates the mTOR complex (mTORC1), which activates the translation of proteins and enhances cell growth. AKT exerts anti-apoptotic effects by phosphorylating target proteins through various downstream pathways. AKT activates IκB kinase (IKKα), which leads to the degradation of the NF-κB inhibitor, IκB, following which, NF-κB is released from the

cytoplasm for nuclear translocation, and its target gene is activated to promote cell survival. AKT phosphorylates the Bcl-2 family member, BAD, which binds to 14-3-3 and prevents it from binding to Bcl-XL to initiate apoptosis (32,33).

PTEN is a PIP3-phosphatase that, in contrast to PI3K, converts PIP3 to PI-4,5-P2 by dephosphorylation. PTEN reduces AKT activation and blocks all downstream signaling events regulated by AKT (34). Previous studies have confirmed that PTEN expression in osteosarcoma cells is significantly decreased compared to normal tissues (7,35), indicating that the pathogenicity of osteosarcoma is related to PTEN expression. In the present study, the expression of PTEN in MG-63 and U2-OS cells was significantly upregulated in CK-treated cells ($P<0.05$). In contrast, the expression levels of p-Akt and p-mTOR were downregulated compared with the control group ($P<0.05$). However, the expression of Akt and mTOR was not significantly altered in the two groups ($P>0.05$), indicating blockage of the PI3K/AKT pathway, as described previously. The phosphorylation of Akt was inhibited and the downstream mTOR complex could not be activated normally; thus, inhibiting cell viability and proliferation. Concurrently, the apoptotic proteins, caspase-3 and BAX were activated, which blocked apoptosis inhibition and triggered apoptosis.

p70S6K1 is a direct substrate downstream of p-mTOR. Phosphorylation of p70s6k1 by p-mTOR promotes the formation of ribosomes, initiation factors, and elongation factors in cells, and promotes the invasion and metastasis of tumor cells (8). The expression of p70s6k1 in MG-63 and U2-OS cells was significantly downregulated after CK treatment ($P<0.05$), indicating that the PI3K/mTOR/p70S6K1 signaling pathway was blocked, and the invasion and metastasis of tumor cells was inhibited. When RAD001 was added, the PI3K/mTOR/p70S6K1 signaling pathway was blocked, the expression levels of p-mTOR and p70S6K1 were downregulated, and this result was consistent with the effect of CK. Furthermore, CK and RAD001 played a synergistic role in regulating the expression of proteins in the PI3K/mTOR/p70S6K1 signaling pathway.

In summary, the present study revealed that CK inhibited the viability and proliferation of osteosarcoma cells, which was related to G2/M-phase cell cycle arrest. CK induced apoptosis and inhibited the migration and invasion of osteosarcoma cells, and blocked the PI3K/mTOR/p70S6K1 signaling pathway of MG-63 and U2-OS osteosarcoma cells, which exhibited anti-tumor properties in osteosarcoma cells. However, this study also has some limitations. Cell viability assays on normal osteocyte cells as a healthy control group should be performed. Research on PARP activation status and the EGFR/RAS/MAPK pathways needs to be performed, to demonstrate the pro-oxidant or antioxidant effects of Ginsenoside CK in MG63 and U2-OS cells to further confirm our conclusion. In addition, apoptosis-related proteins, such as cleaved caspase-9 and cleavage of PARP-1 should also be investigated to research the specific mechanism of mitochondrial apoptosis induced by CK. The MAPK pathway is an important signaling pathway regulating osteosarcoma cells, and its relationship with CK and the PI3K/mTOR/p70S6K1 signaling pathway still requires further study.

Acknowledgements

Not applicable.

Funding

Not applicable.

Availability of data and materials

The analyzed data sets generated during the study are available from the corresponding author on reasonable request.

Authors' contributions

KC and QR conceived and designed all the experiments. YW, ZM, ZL, and BS participated in the cell experiments. JJ, JX, TC and YH analyzed the results. YJ and LQ participated in the writing of the manuscript and revising it critically for important intellectual content. All the authors reviewed and approved the final manuscript and agree to be accountable for all aspects of the research in ensuring that the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Nagao-Kitamoto H, Setoguchi T, Kitamoto S, Nakamura S, Tsuru A, Nagata M, Nagano S, Ishidou Y, Yokouchi M, Kitajima S, *et al*: Ribosomal protein S3 regulates GLI2-mediated osteosarcoma invasion. *Cancer Lett* 356: 855-861, 2015.
- Sevelde F, Mayr L, Kubista B, Lötsch D, van Schoonhoven S, Windhager R, Pirker C, Micksche M and Berger W: EGFR is not a major driver for osteosarcoma cell growth in vitro but contributes to starvation and chemotherapy resistance. *J Exp Clin Cancer Res* 34: 134, 2015.
- Lewis VO: What's new in musculoskeletal oncology. *J Bone Joint Surg Am* 91: 1546-1556, 2009.
- Judson I, Verweij J, Gelderblom H, Hartmann JT, Schöffski P, Blay JY, Kerst JM, Sufliarsky J, Whelan J and Hohenberger P: Doxorubicin alone versus intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic soft-tissue sarcoma: A randomised controlled phase 3 trial. *Lancet Oncol* 15: 415-423, 2014.
- Miller BJ, Gao Y and Duchman KR: Socioeconomic measures influence survival in osteosarcoma: An analysis of the national cancer data base. *Cancer Epidemiol* 49: 112-117, 2017.
- Zhang Y, Kwok-Shing Ng P, Kucherlapati M, Chen F, Liu Y, Tsang YH, de Velasco G, Jeong KJ, Akbani R, Hadjipanayis A, *et al*: A pan-cancer proteogenomic atlas of PI3K/AKT/mTOR pathway alterations. *Cancer Cell* 31: 820-832, 2017.
- Dowling RJ, Topisirovic I, Fonseca BD and Sonenberg N: Dissecting the role of mTOR: Lessons from mTOR inhibitors. *Biochim Biophys Acta* 1804: 433-439, 2010.
- Fan QW and Weiss WA: Inhibition of PI3K-Akt-mTOR signaling in glioblastoma by mTORC1/2 inhibitors. *Methods Mol Biol* 821: 349-359, 2012.
- Hossen MJ, Hong YD, Baek KS, Yoo S, Hong YH, Kim JH, Lee JO, Kim D, Park J and Cho JY: In vitro antioxidative and anti-inflammatory effects of the compound k-rich fraction BIOGF1K, prepared from panax ginseng. *J Ginseng Res* 41: 43-51, 2017.
- Chen L, Meng Y, Sun Q, Zhang Z, Guo X, Sheng X, Tai G, Cheng H and Zhou Y: Ginsenoside compound K sensitizes human colon cancer cells to TRAIL-induced apoptosis via autophagy-dependent and -independent DR5 upregulation. *Cell Death Dis* 7: e2334, 2016.
- Li W, Zhang M, Gu J, Meng ZJ, Zhao LC, Zheng YN, Chen L and Yang GL: Hypoglycemic effect of protopanaxadiol-type ginsenosides and compound k on type 2 diabetes mice induced by high-fat diet combining with streptozotocin via suppression of hepatic gluconeogenesis. *Fitoterapia* 83: 192-198, 2012.
- Wei S, Li W, Yu Y, Yao F, A L, Lan X, Guan F, Zhang M and Chen L: Ginsenoside compound K suppresses the hepatic gluconeogenesis via activating adenosine-5' monophosphate kinase: A study in vitro and in vivo. *Life Sci* 139: 8-15, 2015.
- Ahuja A, Kim JH, Kim JH, Yi YS and Cho JY: Functional role of ginseng-derived compounds in cancer. *J Ginseng Res* 42: 248-254, 2018.
- Muthukumar T, Aravinthan A, Sharmila J, Kim NS and Kim JH: Collagen/chitosan porous bone tissue engineering composite scaffold incorporated with ginseng compound K. *Carbohydr Polym* 152: 566-574, 2016.
- Yang L, Zhang Z, Hou J, Jin X, Ke Z, Liu D, Du M, Jia X and Lv H: Targeted delivery of ginsenoside compound K using TPGS/PEG-PCL mixed micelles for effective treatment of lung cancer. *Int J Nanomedicine* 12: 7653-7667, 2017.
- Yang XD, Yang YY, Ouyang DS and Yang GP: A review of biotransformation and pharmacology of ginsenoside compound K. *Fitoterapia* 100: 208-220, 2015.
- Li Y, Zhou T, Ma C, Song W, Zhang J and Yu Z: Ginsenoside metabolite compound K enhances the efficacy of cisplatin in lung cancer cells. *J Thorac Dis* 7: 400-406, 2015.
- Yang L, Xin J, Zhang Z, Yan H, Wang J, Sun E, Hou J, Jia X and Lv H: TPGS-modified liposomes for the delivery of ginsenoside compound K against non-small cell lung cancer: Formulation design and its evaluation in vitro and in vivo. *J Pharm Pharmacol* 68: 1109-1118, 2016.
- Zhang K and Li Y: Effects of ginsenoside compound k combined with cisplatin on the proliferation, apoptosis and epithelial mesenchymal transition in MCF-7 cells of human breast cancer. *Pharm Biol* 54: 561-568, 2016.
- Lee S, Kwon MC, Jang JP, Sohng JK and Jung HJ: The ginsenoside metabolite compound K inhibits growth, migration and stemness of glioblastoma cells. *Int J Oncol* 51: 414-424, 2017.
- Kang KA, Piao MJ, Kim KC, Zheng J, Yao CW, Cha JW, Kim HS, Kim DH, Bae SC and Hyun JW: Compound k, a metabolite of ginseng saponin, inhibits colorectal cancer cell growth and induces apoptosis through inhibition of histone deacetylase activity. *Int J Oncol* 43: 1907-1914, 2013.
- Ren T, Zheng B, Huang Y, Wang S, Bao X, Liu K and Guo W: Osteosarcoma cell intrinsic PD-L2 signals promote invasion and metastasis via the RhoA-ROCK-LIMK2 and autophagy pathways. *Cell Death Dis* 10: 261, 2019.
- Yu X, Hu L, Li S, Shen J, Wang D, Xu R and Yang H: Long non-coding RNA taurine upregulated gene 1 promotes osteosarcoma cell metastasis by mediating HIF-1 α via miR-143-5p. *Cell Death Dis* 10: 280, 2019.
- Chawla SP, Staddon AP, Baker LH, Schuetze SM, Tolcher AW, D'Amato GZ, Blay JY, Mita MM, Sankhala KK, Berk L, *et al*: Phase II study of the mammalian target of rapamycin inhibitor ridaforolimus in patients with advanced bone and soft tissue sarcomas. *J Clin Oncol* 30: 78-84, 2012.
- Moriceau G, Ory B, Mitrofan L, Riganti C, Blanchard F, Brion R, Charrier C, Battaglia S, Pilet P, Denis MG, *et al*: Zoledronic acid potentiates mTOR inhibition and abolishes the resistance of osteosarcoma cells to RAD001 (Everolimus): Pivotal role of the prenylation process. *Cancer Res* 70: 10329-10339, 2010.
- Manara MC, Nicoletti G, Zambelli D, Ventura S, Guerzoni C, Landuzzi L, Lollini PL, Maira SM, García-Echeverría C, Mercuri M, *et al*: NVP-BEZ235 as a new therapeutic option for sarcomas. *Clin Cancer Res* 16: 530-540, 2010.

27. Engelman JA, Luo J and Cantley LC: The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet* 7: 606-619, 2006.
28. Fruman DA and Rommel C: PI3K and cancer: Lessons, challenges and opportunities. *Nat Rev Drug Discov* 13: 140-156, 2014.
29. Gewinher C, Wang ZC, Richardson A, Teruya-Feldstein J, Etemadmoghadam D, Bowtell D, Barretina J, Lin WM, Rameh L, Salmena L, *et al*: Evidence that inositol polyphosphate 4-phosphatase type II is a tumor suppressor that inhibits PI3K signaling. *Cancer Cell* 16: 115-125, 2009.
30. Hay N: Interplay between FOXO, TOR, and AKT. *Biochim Biophys Acta* 1813: 1965-1970, 2011.
31. Laplante M and Sabatini DM: MTOR signaling in growth control and disease. *Cell* 149: 274-293, 2012.
32. Mabuchi S, Hisamatsu T and Kimura T: Targeting mTOR signaling pathway in ovarian cancer. *Curr Med Chem* 18: 2960-2968, 2011.
33. Wu SH, Bi JF, Cloughesy T, Cavenee WK and Mischel PS: Emerging function of mTORC2 as a core regulator in glioblastoma: Metabolic reprogramming and drug resistance. *Cancer Biol Med* 11: 255-263, 2014.
34. Yuan H and Gao Y: MicroRNA-1908 is upregulated in human osteosarcoma and regulates cell proliferation and migration by repressing PTEN expression. *Oncol Rep* 34: 2706-2714, 2015.
35. Kawano M, Tanaka K, Itonaga I, Ikeda S, Iwasaki T and Tsumura H: MicroRNA-93 promotes cell proliferation via targeting of PTEN in Osteosarcoma cells. *J Exp Clin Cancer Res* 34: 76, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.