

# Long non-coding RNAs interact with RNA-binding proteins to regulate genomic instability in cancer cells (Review)

KAI YANG, XIAOXIANG LIANG and KUNMING WEN

Department of General Surgery, Affiliated Hospital of Zunyi Medical University, Zunyi, Guizhou 563000, P.R. China

Received June 20, 2022; Accepted July 27, 2022

DOI: 10.3892/or.2022.8390

**Abstract.** Genomic instability, a feature of most cancers, contributes to malignant cell transformation and cancer progression due to the accumulation of genetic alterations. Genomic instability is reflected at numerous levels, from single nucleotide to the chromosome levels. However, the exact molecular mechanisms and regulators of genomic instability in cancer remain unclear. Growing evidence indicates that the binding of long non-coding RNAs (lncRNAs) to protein chaperones confers a variety of regulatory functions, including managing of genomic instability. The aim of the present review was to examine the roles of mitosis, telomeres, DNA repair, and epigenetics in genomic instability, and the mechanisms by which lncRNAs regulate them by binding proteins in cancer cells. This review contributes to our understanding of the role of lncRNAs and genomic instability in cancer and can potentially provide entry points and molecular targets for cancer therapies.

## Contents

1. Introduction
2. LncRNAs affect chromosome instability via RBPs
3. LncRNAs are involved in DNA repair through RBPs
4. LncRNAs regulate other epigenetic modalities through RBPs
5. Summary and prospects

## 1. Introduction

Genomic instability refers to genetic alterations that occur at a higher-than-normal frequency and are caused by dysfunctional genome maintenance programs. The term includes changes at

numerous levels, from single nucleotides to chromosomes (1), mainly in the form of microsatellite instability and chromosomal instability (CIN) (2). Genomic instability, one of the most prevalent features of human cancers, can cause cells to exhibit the cancer phenotype through mutations in oncogenes and cancer suppressor genes (3). Persistent genomic instability allows cancer cells to survive under selective pressure and adapt to their microenvironment by evolving to resist different therapies (4), and affects patient prognoses (5). Although genomic instability can promote cancer development and drug resistance, it can cause cancer cell death when genomic instability continues to increase to a limiting level; thus, genomic instability has therapeutic potential for treating cancer (5-7), and elucidating the specific regulatory mechanisms involved is of great significance. Several molecular mechanisms work together to maintain genomic stability under normal physiological conditions. For example, the precise segregation of chromosomes during mitosis and the protection of chromosome ends by telomeres ensures chromosomal stability (8,9), whereas DNA repair, the most important process in the DNA damage response, prevents genomic instability by efficiently repairing DNA damage (3,10). Dysregulation of these processes may lead to genomic instability and the development of cancer. In addition, epigenetic aberrations have been suggested as mechanisms underlying genomic instability (11).

Recent studies have shown that long non-coding RNAs (lncRNAs) are aberrantly expressed in various cancers and are involved in regulating genomic instability in cancer cells (12-14). LncRNAs are transcripts greater than 200 nt in length that do not encode proteins (15) and were considered to have no biological function (16). However, later studies have shown that some lncRNAs can encode polypeptides (17) and interact with proteins, DNA, and RNA to form functional complexes and perform a variety of functions (18). Proteins are the main partners of lncRNAs (16), and the proteins that bind to RNAs are called RNA-binding proteins (RBPs), which bind various RNAs, including lncRNAs, through their RNA-binding domains (19). This interaction between lncRNAs and RBPs plays a critical role in the genomic instability of cancer cells, and by targeting the lncRNA-RBP axis, cancer progression can be inhibited, showing some potential in cancer therapy (20-25).

The present review summarizes the involvement of lncRNAs in regulating genomic instability in cancer by binding proteins that affect mitosis, telomere function, DNA repair,

---

*Correspondence to:* Professor Kunming Wen, Department of General Surgery, Affiliated Hospital of Zunyi Medical University, 149 Dalian Road, Zunyi, Guizhou 563000, P.R. China  
E-mail: 381224619@qq.com

**Key words:** genomic instability, cancer, long non-coding RNA, RNA-binding protein, chromosomal instability

and epigenetics. The study aimed to elucidate the regulatory networks involved in genomic instability in cancer, which may contribute to the development of novel cancer therapies. A systematic literature search using PubMed was performed. The following key words were used for the literature search: 'lncRNA', 'RBP', 'genomic instability', 'telomeres', 'mitosis', 'DNA repair', and 'epigenetic'. The articles in which lncRNAs regulate genomic instability through RBPs in cancer cells were selected.

## 2. lncRNAs affect chromosome instability via RBPs

As the most common form of genomic instability in cancer, CIN is present in 60-80% of human tumors (26,27). It is closely associated with the occurrence and development of human cancers. On the one hand, CIN can promote tumor metastasis and recurrence, accelerate the development of multi-drug resistance in tumors, and be associated with poorer prognoses (28). On the other hand, exceedingly high levels of CIN lead to sensitivity or even death of cancer cells after exposure to cytotoxic drugs and radiotherapy (29). Both abnormal chromosome segregation during mitosis and defects in telomere function contribute to CIN; therefore, the role of lncRNA-protein binding in these processes is reviewed (Fig. 1).

**Mitosis.** During mitosis, precise chromosome segregation depends heavily on the precise binding of microtubules to each sister chromatids (30). Ndc80 is directly attached to microtubules and plays a central role in stable kinetochore-microtubule junctions (31). In the case of incorrect kinetochore-microtubule binding, Aurora B, phosphorylates Ndc80, causing the kinetochore-microtubule binding to become unstable or completely lose the ability to bind to microtubules (32). Concurrently, the spindle assembly checkpoint detects the binding of the kinetochore and microtubules, and transmits an unstable binding signal to the cell cycle. By generating the mitotic checkpoint complex (MCC), it inhibits the anaphase-promoting complex/cyclosome (APC/P). Through this mechanism, mitotic cells are prevented from entering anaphase and cell division until all kinetochore microtubules are stably bound (33), thus preventing chromosome missegregation and CIN.

lncRNAs can directly or indirectly affect proteins involved in this sophisticated process through RBPs (Table I). The level of lncRNA CDKN2B-AS1 is markedly upregulated in renal clear cell carcinoma and is significantly correlated with prognosis. Xie *et al* found that CDKN2B-AS1 can bind directly to IGF2BP3 protein to stabilize it while serving as a scaffold to bind to CBP and SMYD3 epigenetic modification complexes to recruit them to the NUF2 promoter. This mechanism stimulates NUF2 transcription and enhances its cancer-promoting function (34). NUF2 is a component of human Ndc80 that is required for stable microtubule-positive end-binding sites in kinetochores; the precise stoichiometry of the Ndc80 complex may play an important role in microtubule binding (31,35). Stojic *et al* screened for lncRNA linc00899 in HeLa cells by quantifying the effect of lncRNA deletion on cell division. In this study it was determined that linc00899 maintained genomic stability by inhibiting the expression of

microtubule-binding protein TPPP (a protein that stabilizes microtubule networks and its overexpression inhibits microtubule dynamics) through binding to chromatin-modifying complexes (36). Moreover, an elevated level of lncRNA CCAT2 in microsatellite stable colon cancer was revealed to prolong the half-life of BOP1 by directly binding to BOP1; overexpressed BOP1 increased the active form of Aurora B, and the direct binding of CCAT2 to Aurora B also increased active Aurora B. This was demonstrated to lead to incorrect segregation of chromosomes and the occurrence of CIN, thus promoting the progression of colon cancer. Colony formation ability and migration ability of colon cancer cells were effectively inhibited by knockout of BOP1 (20). Cdc20 and Bub3 are components of MCC, and lncRNA CRYBG3 acts as a protein decoy directly binding to Bub3, preventing Bub3 interaction with CDC20, and thus activating APC/P and promoting abnormal mitosis. This then leads to aneuploidy and the development of non-small cell lung cancer. Inhibition of CRYBG3 was revealed to reduce the ability of cancer to migrate *in vitro* and *in vivo* (21). Similarly, lncRNA NORAD induced after DNA damage in HCT116 and other human cell lines can maintain normal mitosis and chromosomal stability by binding to PUMILIO, thereby interfering with PUMILIO binding and inhibiting its target mRNAs (mainly including mRNAs such as chromosome cohesion complexes and centromere complexes) (37). In conclusion, the abovementioned evidence suggests that lncRNAs are involved in the mitotic process by binding proteins that regulate mitosis; however, the levels of induced CIN and the specific biological functions of lncRNAs in cancer require further validation.

**Telomeres.** Human telomeres are DNA-protein complexes present at the ends of chromosomes that consist of the non-coding DNA repeat sequence TTAGGG and shelterin complexes (TRF1, TRF2, POT1, TIN2, TPP1, and RAP1) to which they are bound. Its integrity is critical to the stability of chromosomes (38). Telomeres are subsequently shortened as cells divide due to end replication problems, and short telomeres and defects in the shelterin complex fail to protect chromosome ends leading to CIN. Cancer cells maintain telomere length via the function of telomerase and use of the alternative lengthening of telomeres (ALT) (9,39). In addition, lncRNAs also play an important role in the maintenance of telomeres via RBPs (Table II).

**Telomeric repeat-containing RNA (TERRA).** TERRA is a long non-coding RNA transcribed from subtelomeric- and telomeric-derived sequences containing UUAGGG repeats (40). Like telomeric DNA, TERRA can also form a G-quadruplex structure and bind to the GAR structural domain of TRF2, which is essential for TERRA localization to telomeres. In the absence of the TERRA G-quadruplex structure, telomeres bind more tightly to TRF2, which can promote the formation of the telomeric T-loop. The quino-line derivative CK-14 binds to the TERRA G-quadruplex to form a complex. This complex binds to TRF2 and acts as an allosteric regulator of TRF2, thereby preventing TRF2 from binding to telomeric DNA and ultimately initiating the DNA damage response (DDR). In addition, TERRA can bind to both the origin recognition complex (ORC) and TRF2,

Table I. LncRNAs interact with RBPs to affect mitosis.

LncRNAs	RBP	Mechanism	Effect	(Refs.)
CDKN2B-AS1	SMYD3 and CBP	Promotes the expression of NUF2	Interferes with normal mitosis	(34)
linc00899	Chromatin-modifying complexes	Inhibits TPPP expression	Interferes with normal mitosis	(36)
CCAT2	BOP1 and Aurora B	Increases the active form of Aurora B	Interferes with normal mitosis	(20)
CRYBG3	Bub3	Inhibits the binding of Bub3 and CDC20	Interferes with normal mitosis	(21)
NORAD	PUMILIO	Inhibits the binding of PUMILIO to its target mRNA	Protects normal mitosis	(37)

LncRNAs, long non-coding RNAs; RBPs, RNA-binding proteins; CDKN2B-AS1, cyclin-dependent kinase inhibitor 2B antisense; SMYD3, SET and MYND domain-containing protein 3; CBP, CREB binding protein; CCAT2, colon cancer associated transcript 2 gene; BOP1, block of proliferation 1; NORAD, noncoding RNA activated by DNA damage.




Figure 1. LncRNAs regulate mitosis and telomeres (such as telomere length, telomere capping, R-loop formation, etc.) by binding RBPs and ultimately participate in the regulation of chromosomal instability. LncRNAs, long non-coding RNAs; RBPs, RNA-binding proteins.

enabling formation of a stable ternary complex, which is involved in telomeric DNA replication and facilitates telomeric heterochromatin formation and maintenance (41-43). Similarly, translocated in liposarcoma (TLS) protein can bind to both telomeric DNA and TERRA G-quadruplex structures, while TERRA binds to histone-modifying enzymes, HP1 $\alpha$  and  $\beta$ , and H3K9me3, which play important roles in telomeric heterochromatin formation (41,44). By contrast, telomeric heterochromatin is a negative regulator of telomerase and ALT elongation of telomeres (45,46). Telomeric G-quadruplexes can also regulate telomere length

by preventing the binding of telomerase to telomeric DNA substrates. Previous experiments have demonstrated that hnRNP A1 can facilitate telomerase function by disrupting this high-level structure via binding to telomeric DNA (47). Interestingly, Redon *et al* determined that telomerase can only function when TERRA and hnRNP A1 levels are balanced and bound to form an inert complex. Moreover, excessive hnRNP A1 interferes with telomerase activity by binding to telomeric DNA substrates (48). In cancer cells lacking telomerase, cells maintain telomere length primarily through ALT, and the R-loop formed by telomeric DNA and

Table II. LncRNAs regulate telomere function through RBPs.

LncRNAs	RBPs	Mechanism	Effect	(Refs.)
TERRA	TRF2	Promotes the localization of TERRA in telomeres, and prevents TRF2 from binding to telomeres	Regulates telomere stability	(41,42)
	TRF2, ORC	Promotes the formation and maintenance of telomeric heterochromatin	Promotes telomere elongation	(41)
	TLS, Histone-modifying enzyme, HP1 $\alpha$ and $\beta$	Promotes telomeric heterochromatin formation	Promotes telomere elongation	(41,44)
	HnRNP A1	Regulates telomerase activity in a dose-dependent manner	Regulates telomere length	(48)
	BRCA1	Reduces telomeric R-loop formation	Enhances telomere stability	(49)
	RAD51	Promotes telomeric R-loop formation	Reduces telomere stability	(50)
	TERT	Inhibits telomerase activity	Prevents telomere lengthening	(53)
HTR	TERT	Constitutes the main active part of telomerase	Promotes telomere elongation	(52)
	Dyskerin, NOP10, NHP2, TCAB1 and GAR1	Components of telomerase holoenzymes, maturation and localization of helper group telomerase	Promotes telomere elongation	(52)
	PinX1	Inhibits telomerase activity	Prevents telomere lengthening	(54)
	Ku70/80	Promotes telomere capping	Enhances telomere stability	(55,56)
CUDR	Cyclin D1	Promotes telomerase activity	Promotes telomere elongation	(57)
	P53 (N340Q/L344R)	Promotes telomerase activity	Promotes telomere elongation	(58)
HULC/ MALAT1	TRF2	Promotes telomere capping	Enhances telomere stability	(59)
HULC	P53	Inhibits telomere capping	Reduces telomere stability	(60)

LncRNAs, long non-coding RNAs; RBPs, RNA-binding proteins; TERRA, telomeric repeat-containing RNA; TRF2, telomeric-repeat binding factor 2; ORC, origin recognition complex; TLS, translocated in liposarcoma; HP1, heterochromatin protein 1; hnRNP A1, heterogeneous nuclear ribonucleoprotein A1; BRCA1, breast-cancer susceptibility gene 1; TERT, telomerase reverse transcriptase; hTR, human telomerase RNA; CUDR, cancer upregulated drug resistant; HULC, highly upregulated in liver cancer; MALAT1, metastasis-associated lung adenocarcinoma transcript 1.

TERRA facilitates this process. BRCA1 binds directly to TERRA in an R-loop-dependent manner and reduces R-loop formation; interference with its binding leads to increased R-loops and telomere abnormalities. By contrast, RAD51 can promote R-loop formation by binding TERRA (49,50). Thus, TERRA plays an important role in various aspects of telomere protein capping, heterochromatin formation, secondary structure formation, and telomere lengthening. Owing to the complex role of TERRA in telomeres, targeting its secondary structure or binding proteins may be useful for cancer therapy.

*hTR*. Telomerase is a ribonucleoprotein complex consisting of an RNA component (TERC) and the catalytic subunit of telomerase reverse transcriptase (TERT) as the major active component. The mature human TERC (hTR) is 451 nucleotides long and folds into a highly conserved structural domain. It binds to TERT via CR4/CR5 and template/pseudoknot domains. The H/ACA structural domain of hTR binds to a protein complex composed of dyskerin, NOP10, NHP2, and

GAR1, and facilitates hTR processing and maturation. TCAB1 binds to the CR7 structural domain of hTR to localize telomerase (51,52). TERRA can act as a natural ligand that binds directly to TERT and hTR, thereby inhibiting telomerase activity. The binding of TERRA to TERC does not depend on the presence of hTR. By contrast, PinX1, a telomerase inhibitor, acts by binding directly to hTR and TERT, but the binding of PinX1 to hTR in the intracellular environment is dependent on the presence of TERT (53,54). In addition to providing a template for telomeric DNA replication, hTR plays an important role in telomere shelterin protein capping. The Ku70/80 heterodimer and DNA-dependent protein kinase catalytic subunit (DNA-PKcs) constitute the DNA-dependent protein kinase holoenzyme. The CR7 motif of hTR interacts with Ku70/80 to enhance the phosphorylation activity of hnRNPA1. The phosphorylation of hnRNPA1 increases its affinity for single-stranded telomeric DNA, thereby replacing the replication protein A (RPA) at the telomere ends. Subsequently, hnRNPA1 interacts with protein phosphatase 2A to undergo dephosphorylation, thereby stripping it from

telomeres and allowing POT1 to bind to telomeres. The binding of POT1 ensures telomere capping and inhibits the DDR (55,56).

**Other lncRNAs.** In addition to TERRA and hTR, which play important roles in telomere regulation, other lncRNAs also play a role in telomere physiology. PTEN is one of the most lost tumor suppressors in human cancers. In hepatocellular carcinoma stem cells, decreased PTEN levels lead to increased binding of the lncRNA CUDR to the cell cycle protein cyclin D1; the CUDR-cyclin D1 complex then loads into the lncRNA H19 promoter region and reduces DNA methylation in the H19 promoter region, thereby enhancing H19 expression. H19 overexpression increases TERT binding to TERC while reducing TERT binding to TERRA. This process results in increased cellular telomerase activity and extended telomere length and promoting the malignant proliferation of hepatocellular carcinoma stem cells (57). P53, another tumor suppressor, is frequently mutated in cancer cells to promote cancer progression. In hepatocellular carcinoma cells, the double mutant p53 (N340Q/L344R) binds to CUDR and promotes telomerase activity and lengthening of telomeres through a cascade reaction that enhances TERT expression and reduces TERRA expression (58). In addition to regulating telomerase activity, lncRNAs aberrantly expressed in cancer are involved in telomeric protein capping. Overexpression of lncRNAs HULC and MALAT1 results in increased RNAPII and P300 loading onto the TRF2 promoter region, enhancing TRF2 transcription at the transcriptional level. The increased TRF2 binds to HULC and MALAT1 to form a complex that is loaded onto telomeres, replacing CST/AAF and recruiting telomere-associated proteins, such as POT1, pPOT1, ExoI, and SNM1B, to maintain telomere length and stability. By contrast, lncRNA MEG3 promotes the binding of HULC to p53, thereby inhibiting the binding of telomere-associated proteins to HULC and decreasing telomere stability (59,60).

### 3. lncRNAs are involved in DNA repair through RBPs

The genome of an organism is subjected to endogenous and exogenous damage, causing each cell to produce up to  $10^5$  times the amount of DNA damage per cell per day. Under normal physiological conditions, cells have six main DNA repair pathways by which DNA damage can be precisely repaired to maintain genomic stability (61-64). DNA double-strand breaks (DSBs) are the most cytotoxic type of DNA damage and require complex repair mechanisms. They are repaired by homologous recombination (HR) and non-homologous end joining (NHEJ). Both not repairing DSBs and selecting the wrong way to repair DSBs lead to genomic instability (65-67). Therefore, lncRNAs that regulate DSB repair by binding key proteins during NHEJ and HR were mainly examined (Fig. 2; Table III).

**NHEJ.** Classical NHEJ (cNHEJ) is the primary repair mechanism for DSBs. It does not require a homologous template, requires minor or no processing of DSB ends, and is then directly ligated by enzymatic action, making it an efficient but error-prone repair modality (68,69). In this repair process, Ku70-Ku80 first binds to the DSB and acts as a recruitment platform for other cNHEJ proteins, such as the XRCC4.




Figure 2. lncRNAs are involved in the regulation and selection of different DSB repair pathways by binding RBPs. lncRNAs, long non-coding RNAs; DSB, DNA double-strand break; RBPs, RNA-binding proteins; RPA, replication protein A; PARP1, poly ADP ribose polymerase1; HR, homologous recombination; alt-NHEJ, alternative nonhomologous end joining; cNHEJ, classical nonhomologous end joining; MRN, MRE11-Rad50-Nbs1; DNA-Pkcs, DNA-dependent protein kinase catalytic subunit.

Ku70/80 binds to DNA-PKcs activating their kinase function, which leads to the phosphorylation of Ku and other cNHEJ factors, such as Artemis. The activated Artemis allows the processing of DNA ends. Finally, end linkage is catalyzed by a complex consisting of LIG4 and XRCC4 (70,71).

Wang *et al* reported a novel lncRNA, LRIK, induced by DSB in HeLa cells, which enhances the binding of the Ku heterodimer to DSB through direct binding to the Ku70 subunit. This process promotes assembly of downstream NHEJ factors and the formation of  $\gamma$ -H2AX, ultimately promoting the efficiency of cNHEJ (72). Similarly, lncRNA LINP1, activated by the epidermal growth factor in triple-negative breast cancer, can be recruited to the DSB by binding directly to Ku80. LINP1 also binds to DNA-PKcs through a different region and acts as a molecular scaffold to enhance the interaction between Ku heterodimers and DNA-PKcs. This in turn enhances cNHEJ-mediated DNA repair activity and reduces cancer sensitivity to radiotherapy. Downregulation of LINP1 expression sensitizes cancer cells to radiotherapy due to defective repair activity (22). Thapar *et al* performed further studies and found that Ku binds to the LINP1 stem-loop and G-quadruplex structures (73); the Ku-LINP1 interaction replaces the NHEJ cofactor PAXX protein more efficiently, increasing the stability and net concentration of NHEJ factors at the DSB. Moreover, it bridges the Ku heterodimer at both ends of the DSB to better promote DSB end-joining (71,73). Conversely, lncRNA linc00312, which is expressed at low levels in nasopharyngeal carcinoma, can act as a protein decoy to bind to DNA-PKcs, thereby blocking the recruitment of Ku to DNA-PKcs and inhibiting cNHEJ and resistance to radiotherapy (74).

Table III. LncRNAs involved in DSB repair via RBPs.

LncRNAs	RBPs	Mechanism	Effect	(Refs.)
LRIK	Ku70	Enhances the binding of Ku heterodimer to DSB	Promotes cNHEJ	(72)
LINP1	Ku80 and DNA-PKcs	Enhances the interaction between Ku heterodimers and DNA-PKcs	Promotes cNHEJ	(22,73)
Linc00312	DNA-PKcs	Inhibits the recruitment of Ku to DNA-PKcs	Inhibits cNHEJ	(74)
MALAT1	PARP1	Promotes co-localization between LIG3 and $\gamma$ H2A.X	Activates alt-NHEJ	(77)
PRLH1	RNF169	Promotes RNF169 to replace 53BP1	Promotes HR and inhibits cNHEJ	(78)
SNHG17	NONO	Promotes the formation of the NHEJ repair complex	Promotes cNHEJ and inhibits HR	(79)
HITTERS	MRE11 and Rad50	Promotes the interaction between MRE11 and Rad50	Promotes HR	(82)
HITT	ATM	Prevents ATM recruitment by the MRN complex	Inhibits HR	(23)
GUARDIN	BRCA1 and BARD1	Enhances the interaction between BRCA1 and BARD1	Promotes HR	(84)
BGL3	PARP1 and BARD1	Promotes BRCA1-BARD1 retention at the DSB	Promotes HR	(85)
DDSR1	BRCA1 and hnRNPUL1	Prevents the formation of the BRCA1-RPA80 complex	Promotes HR	(86,87)

LncRNAs, long non-coding RNAs; DSB, DNA double-strand break; RBPs, RNA-binding proteins; LRIK, lncRNA interacting with Ku; cNHEJ, classical nonhomologous end joining; LINP1, lncRNA in non-homologous end joining (NHEJ) pathway 1; DNA-PKcs, DNA-dependent protein kinase catalytic subunit; MALAT1, metastasis-associated lung adenocarcinoma transcript 1; PARP1, poly ADP ribose polymerase1; alt-NHEJ, alternative nonhomologous end joining; SNHG17, small nucleolar RNA host gene 17; HR, homologous recombination; PRLH1, p53-regulated lncRNA for homologous recombination repair 1; HITTERS, HERPUD1 intronic transcript of ER stress; MRE11, meiotic recombination 11 homolog 1; RAD50, ATP-binding cassette-ATPase; HITT, HIF-1 $\alpha$  inhibitor at translation level; ATM, Ataxia-telangiectasia mutated; BARD1, BRCA1-associated RING domain; DDSR1, DNA damage-sensitive RNA1; hnRNPUL1, heterogeneous nuclear ribonucleoprotein U-like 1.

In the case of cNHEJ damage, end linkage without the involvement of cNHEJ core factors is referred to as alt-NHEJ, as an alternate DNA repair pathway to cNHEJ that is more prone to chromosomal alterations (75,76). The alt-NHEJ pathway requires rapid recruitment of the MRN complex by poly ADP ribose polymerase1 (PARP1), which triggers end resection and is dependent on polymerase theta and LIG3 for microhomologous sequence annealing and ligation. In multiple myeloma, the lncRNA MALAT1 can bind directly to PARP1 and indirectly to LIG3. MALAT1 knockdown did not affect LIG3/PARP1 co-localization but disrupted co-localization between LIG3 and  $\gamma$ H2A.X, suggesting that MALAT1 is important for PARP1/LIG3 complex recognition of the  $\gamma$ H2A.X on DSB and activating alt-NHEJ repair, which promotes MM mutagenesis and drug resistance (77).

HR and NHEJ are two competing pathways in the early stages of DSB repair. The selection and balance between the two repair modalities are crucial for genomic stability (78), and lncRNAs are involved in the selection between them. Infection of normal gastric epithelial cells with *Helicobacter pylori* was demonstrated to induce high expression of lncRNA SNHG17, and lncRNA SNHG17 in the nucleus interacted directly with NONO, thus enhancing the interaction between NONO and Ku,

which promoted the formation of the NHEJ repair complex. SNHG17 in the cytoplasm was shown to bind to miR-3909 as a competing endogenous RNA (ceRNA), thereby inhibiting HR, shifting the balance of DSB repair to NHEJ, and ultimately promoting gastric cancer development (79). The E3 ubiquitin ligase RNF169 can replace 53BP1, which inhibits end resection at DSB to promote NHEJ to enhance HR. In hepatocellular carcinoma, lncRNA PRLH1 can bind to RNF169 to form a stable complex that enhances the stability of RNF19 and the affinity of this protein for DSB to replace 53BP1 more efficiently, shifting the balance of repair to HR (78).

**HR.** Because HR is performed using sister chromatids as templates, the process occurs in the late S and G2 phases and facilitates precise repairs. The starting step of HR is the sensing of the damaged site by the MRE11-Rad50-Nbs1 (MRN) complex and producing a free 3' end single-strand overhang (80). Rad51 recombinase is the final effector of the HR cascade reaction, and its binding to single-stranded DNA depends on BRCA2 as well as the interaction of the BRCA1-BARD1 complex and PALB2. Once Rad51 binds to single-stranded DNA at the DSB, it begins the subsequent homology search and strand invasion to initiate DNA repair (81).



The lncRNA HITTERS, which is highly expressed in oral squamous cell carcinoma cells, induced by endoplasmic reticulum stress, can directly bind MRE11 and Rad50, thus promoting their interaction, while also increasing MRE11 and Nbs1 protein levels and promoting the formation of the MRN complex. Ultimately, the function of HITTERS facilitates DNA repair via multiple pathways including HR (82). Like DNA-PKcs in NHEJ, the capillary dilation ataxia mutated gene (ATM) protein kinase is the apical kinase in HR; MRN serves as a protein platform to promote autophosphorylation of ATM to stimulate its activity (83). The lncRNA HITT, which is expressed at low levels in several cancers due to hypoxic contingency, binds directly to the binding site of ATM-binding Nbs1 and thereby prevents ATM recruitment by the MRN and antagonizes HR-mediated DNA repair. Through this mechanism, overexpression of HITT can enhance the sensitivity of cancer cells to genotoxic therapies (23).

BRCA1 and BARD1 form a heterodimer, which plays a role in HR. The lncRNA GUARDIN acts as a molecular scaffold that directly binds to BRCA1 and BARD1, enhancing their interaction to promote HR (84). Similarly, lncRNA BGL3 is recruited to the DSB early by binding to PARP1, whereas direct binding to BARD1 promotes BRCA1-BARD1 retention at the DSB and facilitates the interaction between BARD1 and Rad51 (85). The lncRNA DDSR1 is known to interact with BRCA1 and hnRNPUL1; Sharma *et al* suggested that this interaction prevents the formation of the BRCA1-RPA80 complex and the binding of this complex at the DSB (86). In turn, binding of the BRCA1-RPA80 complex to the DSB restricts DNA end excision and thus limits HR (87).

Thus, lncRNAs play an important role in the different repair pathways of DSBs by binding proteins, whereas the role of lncRNAs in other repair modalities is poorly understood and warrants further investigation.

#### 4. lncRNAs regulate other epigenetic modalities through RBPs

Epigenetic inheritance refers to the production of heritable changes in gene expression without changes to the DNA nucleotide sequence, and such epigenetic aberrations are considered a form of genomic instability. Like mutations, epigenetic inheritance plays a key role in cancer development by altering the expression of oncogenes and cancer suppressor genes (11). In addition, epigenetics may serve as an advantageous biological marker for cancer diagnosis, prognosis, and treatment. Epigenetic regulatory mechanisms mainly include DNA methylation, chromatin remodeling, and non-coding RNA (88,89). These regulatory mechanisms crosstalk: for example, lncRNAs can act as miRNA sponges to inhibit miRNAs (90). Therefore, the regulation of chromatin remodeling and DNA methylation by lncRNAs via binding to multiple enzymes (Fig. 3) was investigated.

**Chromatin remodeling.** The nucleosome is the basic unit of chromatin, and consists mainly of the core histones H2A, H2B, H3, and H4 that form an octamer wrapped around 147 base pairs of DNAs in humans (91); nucleosomes further




Figure 3. lncRNAs regulate gene expression by binding RBPs to regulate histone modifications and DNA methylation. lncRNAs, long non-coding RNAs; RBPs, RNA-binding proteins.

assemble into higher-order chromatin (92). This highly folded state of chromatin prevents the binding of DNA-binding proteins to promoters, thereby inhibiting transcription (93). Histone-modifying enzymes and ATP-dependent chromatin remodeling complexes mediate chromatin remodeling, which controls gene expression by altering the accessibility of local chromatin DNA (93-95).

**Histone modification.** The amino-terminal ends of core histones can extend outside the nucleosome and be covalently modified by various histone-modifying enzymes via methylation, acetylation, ubiquitination, and phosphorylation. These modifications can affect the affinity of histones for DNA duplexes, alter the loose or condensed state of chromatin, and mediate DNA accessibility and protein-chromatin interactions, ultimately affecting gene expression (91,96,97). Methylation and acetylation are among the most intensively studied processes.

lncRNAs can play a role in cancer development by directing histone-modifying enzymes to regulate the methylation and acetylation status of histones and cis-regulating the expression of nearby genes (24,98,99). For example, lncRNA EZR-AS1 recruits H3K4 methyltransferase SMYD3 to catalyze Tri-methylation of lysine 4 on histone H3 (H3K4me3) at the EZR promoter. This process promotes EZR transcription, thereby enhancing the metastasis and invasion of the esophageal squamous cell carcinoma. Conversely, interfering with the expression of EZR-AS1 has an inhibitory effect on cancer cells (24). Similarly, lncRNAs, such as HOTAIR and

AS1DHRS4, can have a *trans*-regulatory role in gene expression through histone modifications (100,101). LncRNAs can also serve as protein scaffolds for histone-modifying enzymes. For example, lncRNA AGAP2-AS1 acts as a protein scaffold and binds to the polycomb repressive complex 2 (PRC2) core catalytic subunit EZH2 and lysine-specific demethylase LSD1 to promote histone modifications in pancreatic adenocarcinoma (102), glioma (103), non-small cell lung cancer (104), and gastric cancer (105). Through this mechanism the expression of target genes is suppressed, and cancer progression is promoted. The lncRNA CDKN2B-AS1 can bind to both histone acetyltransferases CBP and SMYD3 to promote acetylation of lysine 27 on histone 3 (H3K27ac) and H3K4me3 at the NUF2 promoter, which further enhances NUF2 expression (34). Conversely, lncRNAs can act as a protein decoy to regulate histone modification. For example, LINC00261 binds to the acetylase P300/CBP complex, inhibiting its binding to the c-Myc gene promoter to reduce H3K27ac and inhibit pancreatic cancer progression by suppressing c-Myc expression (106). Similarly, lncRNA DLEU2 can also act as a protein decoy for EZH2 (107).

**ATP-dependent chromatin remodeling.** ATP-dependent chromatin remodeling complexes are divided into four major classes: SWI/SNF, ISWI, CHD, and INO80 (108). The complexes can alter the accessibility of transcription factors to DNA by disrupting the interaction between DNA and histones using the energy generated by ATP hydrolysis, altering the position of nucleosomes along the DNA or replacing histones (94,95). LncRNAs are also involved in this process. Subunits of the INO80 complex, INO80 and RUVBL2, bind to the lncRNA HAND2-AS1. This process activates BMPR1A expression and ultimately stimulates the self-renewal of hepatocellular carcinoma stem cells by recruiting INO80 to the BMPR1A promoter and forming the formation complex (109). Similarly, Tang *et al* reported that many lncRNAs in cancer interact with SWI/SNF and thus participate in the regulation of chromatin remodeling (110).

**DNA methylation.** In humans, DNA methylation occurs mainly at the cytosine 5' carbon atom of CpG dinucleotides. DNA methylation is catalyzed by the active DNA methyltransferases, DNMT1, DNMT3a, and DNMT3b that add methyl groups to cytosine to form 5-methylcytosine (5mC) (111,112). CpG dinucleotides usually exist in CpG islands in the promoter region of the human genome (113). Methylation of CpG islands can directly or indirectly inhibit the binding of transcription factors to promoters (114). LncRNAs can directly bind to these active DNA methylases and regulate gene expression, thereby interfering with cancer development (115-118). LncRNA LINC01270 is highly expressed in esophageal cancer and can act as a protein scaffold to simultaneously bind to DNMT1, DNMT3a, and DNMT3b. This mechanism mediates the hypermethylation of GSTP1 promoter, inhibiting its expression and promoting esophageal cancer progression and drug resistance (119). Interestingly, in colon cancer cells, lncRNA lnc-LALC can also recruit DNMT1, DNMT3a, and DNMT3b to the LZTS1 promoter simultaneously, but this requires direct binding of lnc-LALC to EZH2 (120). The lncRNA PARTICLE, which is highly expressed in response to

low irradiation, promotes both DNA and histone methylation by binding to DNMT1 and the PRC2 core subunit SUZ12 and suppresses the expression of tumor suppressor MAT2A in *cis* as well as the tumor suppressor WWOX in *trans* (121,122).

DNA methylation is a stable modification process; however, ten-eleven translocation (TET) family proteins catalyze the conversion of 5mC to 5-hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC), and 5-carboxycytosine (5caC). 5hmC is diluted during DNA replication, while 5fC and 5caC are removed by thymine DNA glycosylase (TDG), resulting in DNA demethylation (112,123). LncRNA TARID can bind to growth arrest and DNA damage-inducible 45A (GADD45A), a protein that mediates DNA demethylation. Through this interaction, TARID directs GADD45A *cis* to the tumor suppressor TCF21 promoter and indirectly recruits TETs as well as TDG, which can bind to GADD45A and co-mediate DNA demethylation (123). Similarly, lncRNA ZNF667-AS1 can bind to TET1 and histone H3K27 demethylase to promote both DNA and histone demethylation at the ZNF667 and E-cadherin promoters, thereby inhibiting the development of esophageal squamous epithelial carcinoma (25).

## 5. Summary and prospects

Genomic instability is a feature of most cancers that undoubtedly contributes to cancer progression and heterogeneity through the accumulation of oncogenic and cancer suppressor genetic mutations, regardless of whether it acts as a 'passenger' or a 'driver' in cancer. Although the exact mechanism is unknown, cancer cells have a tolerance limit to genomic instability, which indicates the potential of genomic destabilization as a therapeutic approach to cancer. Investigating the mechanisms of tolerance to genomic instability in cancer cells may provide new insights into cancer treatment. The aberrantly expressed lncRNAs in cancer cells and their binding proteins form networks that regulate genomic instability. Exploitation of lncRNA-RBP networks may provide new biological markers for cancer diagnosis and prognosis as well as new molecular targets and entry points for driving genomic instability to the limit of cellular tolerance or suppressing it in cancer therapies. However, the causal relationship between dysregulated lncRNA expression and genomic instability in many cancers has not yet been verified, and the exact molecular mechanism between the two remains unclear, which warrants further investigations.

## Acknowledgements

Not applicable.

## Funding

The present review was supported by the National Natural Science Foundation of China (grant no. 82160575) and the Outstanding Young Technological and Innovative Talent Cultivation Project of Zunyi Municipal Science and Technology Bureau, 2021 (no. 10).

## Availability of data and materials

Not applicable.



## Authors' contributions

KY and KW conceived the study. KY drafted the manuscript. KY, KW and XL made substantial contributions to the interpretation, drafting the manuscript and revising it critically for important intellectual content. Data authentication is not applicable. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

Not applicable.

## Patient consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

- Lee JK, Choi YL, Kwon M and Park PJ: Mechanisms and consequences of cancer genome instability: Lessons from genome sequencing studies. *Annu Rev Pathol* 11: 283-312, 2016.
- Aguilera A and Gómez-González B: Genome instability: A mechanistic view of its causes and consequences. *Nat Rev Genet* 9: 204-217, 2008.
- Salmaninejad A, Ilkhani K, Marzban H, Navashenaq JG, Rahimirad S, Radnia F, Yousefi M, Bahmanpour Z, Azhdari S and Sahebkar A: Genomic instability in cancer: Molecular mechanisms and therapeutic potentials. *Curr Pharm Des* 27: 3161-3169, 2021.
- Mehrotra S and Mittra I: Origin of genome instability and determinants of mutational landscape in cancer cells. *Genes (Basel)* 11: 1101, 2020.
- Andor N, Maley CC and Ji HP: Genomic instability in cancer: Teetering on the limit of tolerance. *Cancer Res* 77: 2179-2185, 2017.
- Abbas T, Keaton MA and Dutta A: Genomic instability in cancer. *Cold Spring Harb Perspect Biol* 5: a012914, 2013.
- O'Connor MJ: Targeting the DNA damage response in cancer. *Mol Cell* 60: 547-560, 2015.
- Sansregret L, Vanhaesebroeck B and Swanton C: Determinants and clinical implications of chromosomal instability in cancer. *Nat Rev Clin Oncol* 15: 139-150, 2018.
- Maciejowski J and de Lange T: Telomeres in cancer: Tumour suppression and genome instability. *Nat Rev Mol Cell Biol* 18: 175-186, 2017.
- Lord CJ and Ashworth A: The DNA damage response and cancer therapy. *Nature* 481: 287-294, 2012.
- Choi JD and Lee JS: Interplay between epigenetics and genetics in cancer. *Genomics Inform* 11: 164-173, 2013.
- Zhang W, Guan X and Tang J: The long non-coding RNA landscape in triple-negative breast cancer. *Cell Prolif* 54: e12966, 2021.
- Huang L, Xie Y, Jiang S, Han W, Zeng F and Li D: The lncRNA signatures of genome instability to predict survival in patients with renal cancer. *J Healthc Eng* 2021: 1090698, 2021.
- Yin T, Zhao D and Yao S: Identification of a genome instability-associated lncRNA signature for prognosis prediction in colon cancer. *Front Genet* 12: 679150, 2021.
- Kopp F and Mendell JT: Functional classification and experimental dissection of long noncoding RNAs. *Cell* 172: 393-407, 2018.
- Mercer TR and Mattick JS: Structure and function of long noncoding RNAs in epigenetic regulation. *Nat Struct Mol Biol* 20: 300-307, 2013.
- Matsumoto A, Pasut A, Matsumoto M, Yamashita R, Fung J, Monteleone E, Saghatelian A, Nakayama KI, Clohessy JG and Pandolfi PP: mTORC1 and muscle regeneration are regulated by the LINC00961-encoded SPAR polypeptide. *Nature* 541: 228-232, 2017.
- Guttman M and Rinn JL: Modular regulatory principles of large non-coding RNAs. *Nature* 482: 339-346, 2012.
- Hentze MW, Castello A, Schwarzl T and Preiss T: A brave new world of RNA-binding proteins. *Nat Rev Mol Cell Biol* 19: 327-341, 2018.
- Chen B, Dragomir MP, Fabris L, Bayraktar R, Knutsen E, Liu X, Tang C, Li Y, Shimura T, Ivkovic TC, *et al*: The long noncoding RNA CCAT2 induces chromosomal instability through BOP1-AURKB signaling. *Gastroenterology* 159: 2146-2162.e33, 2020.
- Guo Z, Dai Y, Hu W, Zhang Y, Cao Z, Pei W, Liu N, Nie J, Wu A, Mao W, *et al*: The long noncoding RNA CRYBG3 induces aneuploidy by interfering with spindle assembly checkpoint via direct binding with Bub3. *Oncogene* 40: 1821-1835, 2021.
- Zhang Y, He Q, Hu Z, Feng Y, Fan L, Tang Z, Yuan J, Shan W, Li C, Hu X, *et al*: Long noncoding RNA LINP1 regulates repair of DNA double-strand breaks in triple-negative breast cancer. *Nat Struct Mol Biol* 23: 522-530, 2016.
- Zhao K, Wang X, Xue X, Li L and Hu Y: A long noncoding RNA sensitizes genotoxic treatment by attenuating ATM activation and homologous recombination repair in cancers. *PLoS Biol* 18: e3000666, 2020.
- Zhang XD, Huang GW, Xie YH, He JZ, Guo JC, Xu XE, Liao LD, Xie YM, Song YM, Li EM and Xu LY: The interaction of lncRNA EZR-AS1 with SMYD3 maintains overexpression of EZR in ESCC cells. *Nucleic Acids Res* 46: 1793-1809, 2018.
- Dong Z, Li S, Wu X, Niu Y, Liang X, Yang L, Guo Y, Shen S, Liang J and Guo W: Aberrant hypermethylation-mediated downregulation of antisense lncRNA ZNF667-AS1 and its sense gene ZNF667 correlate with progression and prognosis of esophageal squamous cell carcinoma. *Cell Death Dis* 10: 930, 2019.
- Carter SL, Cibulskis K, Helman E, McKenna A, Shen H, Zack T, Laird PW, Onofrio RC, Winckler W, Weir BA, *et al*: Absolute quantification of somatic DNA alterations in human cancer. *Nat Biotechnol* 30: 413-421, 2012.
- Negrini S, Gorgoulis VG and Halazonetis TD: Genomic instability-an evolving hallmark of cancer. *Nat Rev Mol Cell Biol* 11: 220-228, 2010.
- Jo M, Kusano Y and Hirota T: Unraveling pathologies underlying chromosomal instability in cancers. *Cancer Sci* 112: 2975-2983, 2021.
- Piemonte KM, Anstine LJ and Keri RA: Centrosome aberrations as drivers of chromosomal instability in breast cancer. *Endocrinology* 162: bqab208, 2021.
- Hara M and Fukagawa T: Dynamics of kinetochore structure and its regulations during mitotic progression. *Cell Mol Life Sci* 77: 2981-2995, 2020.
- Monda JK and Cheeseman IM: The kinetochore-microtubule interface at a glance. *J Cell Sci* 131: jcs214577, 2018.
- Welburn JP, Vleugel M, Liu D, Yates JR III, Lampson MA, Fukagawa T and Cheeseman IM: Aurora B phosphorylates spatially distinct targets to differentially regulate the kinetochore-microtubule interface. *Mol Cell* 38: 383-392, 2010.
- Sacristan C and Kops GJ: Joined at the hip: Kinetochore, microtubules, and spindle assembly checkpoint signaling. *Trends Cell Biol* 25: 21-28, 2015.
- Xie X, Lin J, Fan X, Zhong Y, Chen Y, Liu K, Ren Y, Chen X, Lai D, Li X, *et al*: lncRNA CDKN2B-AS1 stabilized by IGF2BP3 drives the malignancy of renal clear cell carcinoma through epigenetically activating NUF2 transcription. *Cell Death Dis* 12: 201, 2021.
- DeLuca JG, Dong Y, Hergert P, Strauss J, Hickey JM, Salmon ED and McEwen BF: Hec1 and nuf2 are core components of the kinetochore outer plate essential for organizing microtubule attachment sites. *Mol Biol Cell* 16: 519-531, 2005.
- Stojic L, Lun ATL, Mascialchi P, Ernst C, Redmond AM, Mangei J, Barr AR, Bousgouni V, Bakal C, Marioni JC, *et al*: A high-content RNAi screen reveals multiple roles for long noncoding RNAs in cell division. *Nat Commun* 11: 1851, 2020.
- Lee S, Kopp F, Chang TC, Sataluri A, Chen B, Sivakumar S, Yu H, Xie Y and Mendell JT: Noncoding RNA NORAD regulates genomic stability by sequestering PUMILIO proteins. *Cell* 164: 69-80, 2016.
- Schmidt JC and Cech TR: Human telomerase: Biogenesis, trafficking, recruitment, and activation. *Genes Dev* 29: 1095-1105, 2015.
- Frias C, Pampalona J, Genesca A and Tusell L: Telomere dysfunction and genome instability. *Front Biosci (Landmark Ed)* 17: 2181-2196, 2012.

40. Schoeftner S and Blasco MA: Developmentally regulated transcription of mammalian telomeres by DNA-dependent RNA polymerase II. *Nat Cell Biol* 10: 228-236, 2008.
41. Deng Z, Norseen J, Wiedmer A, Riethman H and Lieberman PM: TERRA RNA binding to TRF2 facilitates heterochromatin formation and ORC recruitment at telomeres. *Mol Cell* 35: 403-413, 2009.
42. Mei Y, Deng Z, Vladimirova O, Gulve N, Johnson FB, Drosopoulos WC, Schildkraut CL and Lieberman PM: TERRA G-quadruplex RNA interaction with TRF2 GAR domain is required for telomere integrity. *Sci Rep* 11: 3509, 2021.
43. Zhang Y, Zeng D, Cao J, Wang M, Shu B, Kuang G, Ou TM, Tan JH, Gu LQ, Huang ZS and Li D: Interaction of Quindoline derivative with telomeric repeat-containing RNA induces telomeric DNA-damage response in cancer cells through inhibition of telomeric repeat factor 2. *Biochim Biophys Acta Gen Subj* 1861: 3246-3256, 2017.
44. Takahama K, Takada A, Tada S, Shimizu M, Sayama K, Kurokawa R and Oyoshi T: Regulation of telomere length by G-quadruplex telomere DNA- and TERRA-binding protein TLS/FUS. *Chem Biol* 20: 341-350, 2013.
45. Blasco MA: Telomeres and human disease: Ageing, cancer and beyond. *Nat Rev Genet* 6: 611-622, 2005.
46. Benetti R, Garcia-Cao M and Blasco MA: Telomere length regulates the epigenetic status of mammalian telomeres and subtelomeres. *Nat Genet* 39: 243-250, 2007.
47. Zhang QS, Manche L, Xu RM and Krainer AR: hnRNP A1 associates with telomere ends and stimulates telomerase activity. *RNA* 12: 1116-1128, 2006.
48. Redon S, Zemp I and Lingner J: A three-state model for the regulation of telomerase by TERRA and hnRNPA1. *Nucleic Acids Res* 41: 9117-9128, 2013.
49. Vohhodina J, Goehring LJ, Liu B, Kong Q, Botchkarev VV Jr, Huynh M, Liu Z, Abderazzaq FO, Clark AP, Ficarro SB, *et al*: BRCA1 binds TERRA RNA and suppresses R-Loop-based telomeric DNA damage. *Nat Commun* 12: 3542, 2021.
50. Feretaki M, Pospisilova M, Valador Fernandes R, Lunardi T, Krejci L and Lingner J: RAD51-dependent recruitment of TERRA lncRNA to telomeres through R-loops. *Nature* 587: 303-308, 2020.
51. Gala K and Khattar E: Long non-coding RNAs at work on telomeres: Functions and implications in cancer therapy. *Cancer Lett* 502: 120-132, 2021.
52. Podlevsky JD and Chen JJ: Evolutionary perspectives of telomerase RNA structure and function. *RNA Biol* 13: 720-732, 2016.
53. Redon S, Reichenbach P and Lingner J: The non-coding RNA TERRA is a natural ligand and direct inhibitor of human telomerase. *Nucleic Acids Res* 38: 5797-5806, 2010.
54. Banik SS and Counter CM: Characterization of interactions between PinX1 and human telomerase subunits hTERT and hTR. *J Biol Chem* 279: 51745-51748, 2004.
55. Raghunandan M and Decottignies A: The multifaceted hTR telomerase RNA from a structural perspective: Distinct domains of hTR differentially interact with protein partners to orchestrate its telomerase-independent functions. *Bioessays* 43: e2100099, 2021.
56. Sui JD, Tang Z, Chen BPC, Huang P, Yang MQ, Wang NH, Yang HN, Tu HL, Jiang QM, Zhang J, *et al*: Protein phosphatase 2A-dependent mitotic hnRNPA1 dephosphorylation and TERRA formation facilitate telomere capping. *Mol Cancer Res* 20: 583-595, 2022.
57. Pu H, Zheng Q, Li H, Wu M, An J, Gui X, Li T and Lu D: CUDR promotes liver cancer stem cell growth through upregulating TERT and C-Myc. *Oncotarget* 6: 40775-40798, 2015.
58. Wu M, An J, Zheng Q, Xin X, Lin Z, Li X, Li H and Lu D: Double mutant P53 (N340Q/L344R) promotes hepatocarcinogenesis through upregulation of Pim1 mediated by PKM2 and LncRNA CUDR. *Oncotarget* 7: 66525-66539, 2016.
59. Wu M, Lin Z, Li X, Xin X, An J, Zheng Q, Yang Y and Lu D: HULC cooperates with MALAT1 to aggravate liver cancer stem cells growth through telomere repeat-binding factor 2. *Sci Rep* 6: 36045, 2016.
60. Jiang X, Wang L, Xie S, Chen Y, Song S, Lu Y and Lu D: Long noncoding RNA MEG3 blocks telomerase activity in human liver cancer stem cells epigenetically. *Stem Cell Res Ther* 11: 518, 2020.
61. Hoeijmakers JH: DNA damage, aging, and cancer. *N Engl J Med* 361: 1475-1485, 2009.
62. Zhao H, Fuemmeler BF and Shen J: DNA repair in cancer development and aging. *Aging (Albany NY)* 13: 23435-23436, 2021.
63. Tian H, Gao Z, Li H, Zhang B, Wang G, Zhang Q, Pei D and Zheng J: DNA damage response-a double-edged sword in cancer prevention and cancer therapy. *Cancer Lett* 358: 8-16, 2015.
64. Bever KM and Le DT: DNA repair defects and implications for immunotherapy. *J Clin Invest* 128: 4236-4242, 2018.
65. Zhao Y and Chen S: Targeting DNA double-strand break (DSB) repair to counteract tumor radio-resistance. *Curr Drug Targets* 20: 891-902, 2019.
66. Ceccaldi R, Rondinelli B and D'Andrea AD: Repair pathway choices and consequences at the double-strand break. *Trends Cell Biol* 26: 52-64, 2016.
67. Scully R, Panday A, Elango R and Willis NA: DNA double-strand break repair-pathway choice in somatic mammalian cells. *Nat Rev Mol Cell Biol* 20: 698-714, 2019.
68. Burma S, Chen BP and Chen DJ: Role of non-homologous end joining (NHEJ) in maintaining genomic integrity. *DNA Repair (Amst)* 5: 1042-1048, 2006.
69. Shibata A, Conrad S, Birraux J, Geuting V, Barton O, Ismail A, Kakarougkas A, Meek K, Taucher-Scholz G, Löbrich M and Jeggo PA: Factors determining DNA double-strand break repair pathway choice in G2 phase. *EMBO J* 30: 1079-1092, 2011.
70. Stinson BM and Loparo JJ: Repair of DNA double-strand breaks by the nonhomologous end joining pathway. *Annu Rev Biochem* 90: 137-164, 2021.
71. Ghosh D and Raghavan SC: Nonhomologous end joining: New accessory factors fine tune the machinery. *Trends Genet* 37: 582-599, 2021.
72. Wang D, Zhou Z, Wu E, Ouyang C, Wei G, Wang Y, He D, Cui Y, Zhang D, Chen X, *et al*: LRIK interacts with the Ku70-Ku80 heterodimer enhancing the efficiency of NHEJ repair. *Cell Death Differ* 27: 3337-3353, 2020.
73. Thapar R, Wang JL, Hammel M, Ye R, Liang K, Sun C, Hnizda A, Liang S, Maw SS, Lee L, *et al*: Mechanism of efficient double-strand break repair by a long non-coding RNA. *Nucleic Acids Res* 48: 10953-10972, 2020.
74. Guo Z, Wang YH, Xu H, Yuan CS, Zhou HH, Huang WH, Wang H and Zhang W: LncRNA linc00312 suppresses radiotherapy resistance by targeting DNA-PKcs and impairing DNA damage repair in nasopharyngeal carcinoma. *Cell Death Dis* 12: 69, 2021.
75. Decottignies A: Alternative end-joining mechanisms: A historical perspective. *Front Genet* 4: 48, 2013.
76. Chiruvella KK, Liang Z and Wilson TE: Repair of double-strand breaks by end joining. *Cold Spring Harb Perspect Biol* 5: a012757, 2013.
77. Hu Y, Lin J, Fang H, Fang J, Li C, Chen W, Liu S, Ondrejka S, Gong Z, Reu F, *et al*: Targeting the MALAT1/PARP1/LIG3 complex induces DNA damage and apoptosis in multiple myeloma. *Leukemia* 32: 2250-2262, 2018.
78. Deng B, Xu W, Wang Z, Liu C, Lin P, Li B, Huang Q, Yang J, Zhou H and Qu L: An LTR retrotransposon-derived lncRNA interacts with RNF169 to promote homologous recombination. *EMBO Rep* 20: e47650, 2019.
79. Han T, Jing X, Bao J, Zhao L, Zhang A, Miao R, Guo H, Zhou B, Zhang S, Sun J and Shi J: H. pylori infection alters repair of DNA double-strand breaks via SNHG17. *J Clin Invest* 130: 3901-3918, 2020.
80. Ranjha L, Howard SM and Cejka P: Main steps in DNA double-strand break repair: An introduction to homologous recombination and related processes. *Chromosoma* 127: 187-214, 2018.
81. Yamamoto H and Hirasawa A: Homologous recombination deficiencies and hereditary tumors. *Int J Mol Sci* 23: 348, 2021.
82. Wu C, Chen W, Yu F, Yuan Y, Chen Y, Hurst DR, Li Y, Li L and Liu Z: Long noncoding RNA HITTERS protects oral squamous cell carcinoma cells from endoplasmic reticulum stress-induced apoptosis via promoting MRE11-RAD50-NBS1 complex formation. *Adv Sci (Weinh)* 7: 2002747, 2020.
83. Paull TT: Mechanisms of ATM activation. *Annu Rev Biochem* 84: 711-738, 2015.
84. Hu WL, Jin L, Xu A, Wang YF, Thorne RF, Zhang XD and Wu M: GUARDIN is a p53-responsive long non-coding RNA that is essential for genomic stability. *Nat Cell Biol* 20: 492-502, 2018.
85. Hu Z, Mi S, Zhao T, Peng Y, Chen L, Zhu W, Yao Y, Song Q, Li X, Li X, *et al*: BGL3 lncRNA mediates retention of the BRCA1/BARD1 complex at DNA damage sites. *EMBO J* 39: e104133, 2020.

86. Sharma V, Khurana S, Kubben N, Abdelmohsen K, Oberdoerffer P, Gorospe M and Misteli T: A BRCA1-interacting lncRNA regulates homologous recombination. *EMBO Rep* 16: 1520-1534, 2015.
87. Coleman KA and Greenberg RA: The BRCA1-RAP80 complex regulates DNA repair mechanism utilization by restricting end resection. *J Biol Chem* 286: 13669-13680, 2011.
88. Dawson MA and Kouzarides T: Cancer epigenetics: From mechanism to therapy. *Cell* 150: 12-27, 2012.
89. Costa-Pinheiro P, Montezuma D, Henrique R and Jerónimo C: Diagnostic and prognostic epigenetic biomarkers in cancer. *Epigenomics* 7: 1003-1015, 2015.
90. Wang H, Huo X, Yang XR, He J, Cheng L, Wang N, Deng X, Jin H, Wang N, Wang C, *et al*: STAT3-mediated upregulation of lncRNA HOXD-AS1 as a ceRNA facilitates liver cancer metastasis by regulating SOX4. *Mol Cancer* 16: 136, 2017.
91. Kim JJ, Lee SY and Miller KM: Preserving genome integrity and function: The DNA damage response and histone modifications. *Crit Rev Biochem Mol Biol* 54: 208-241, 2019.
92. Iacobuzio-Donahue CA: Epigenetic changes in cancer. *Annu Rev Pathol* 4: 229-249, 2009.
93. Luo RX and Dean DC: Chromatin remodeling and transcriptional regulation. *J Natl Cancer Inst* 91: 1288-1294, 1999.
94. Wang Z, Liu S and Tao Y: Regulation of chromatin remodeling through RNA polymerase II stalling in the immune system. *Mol Immunol* 108: 75-80, 2019.
95. Koreman E, Sun X and Lu QR: Chromatin remodeling and epigenetic regulation of oligodendrocyte myelination and myelin repair. *Mol Cell Neurosci* 87: 18-26, 2018.
96. Strahl BD and Allis CD: The language of covalent histone modifications. *Nature* 403: 41-45, 2000.
97. Kouzarides T: Chromatin modifications and their function. *Cell* 128: 693-705, 2007.
98. Fang K, Huang W, Sun YM, Chen TQ, Zeng ZC, Yang QQ, Pan Q, Han C, Sun LY, Luo XQ, *et al*: Cis-acting lnc-eRNA SEELA directly binds histone H4 to promote histone recognition and leukemia progression. *Genome Biol* 21: 269, 2020.
99. Wang YQ, Jiang DM, Hu SS, Zhao L, Wang L, Yang MH, Ai ML, Jiang HJ, Han Y, Ding YQ and Wang S: SATB2-AS1 suppresses colorectal carcinoma aggressiveness by inhibiting SATB2-dependent snail transcription and epithelial-mesenchymal transition. *Cancer Res* 79: 3542-3556, 2019.
100. Chu C, Qu K, Zhong FL, Artandi SE and Chang HY: Genomic maps of long noncoding RNA occupancy reveal principles of RNA-chromatin interactions. *Mol Cell* 44: 667-678, 2011.
101. Li Q, Su Z, Xu X, Liu G, Song X, Wang R, Sui X, Liu T, Chang X and Huang D: AS1DHR54, a head-to-head natural antisense transcript, silences the DHR54 gene cluster in cis and trans. *Proc Natl Acad Sci USA* 109: 14110-14115, 2012.
102. Hui B, Ji H, Xu Y, Wang J, Ma Z, Zhang C, Wang K and Zhou Y: RREB1-induced upregulation of the lncRNA AGAP2-AS1 regulates the proliferation and migration of pancreatic cancer partly through suppressing ANKRD1 and ANGPTL4. *Cell Death Dis* 10: 207, 2019.
103. Luo W, Li X, Song Z, Zhu X and Zhao S: Long non-coding RNA AGAP2-AS1 exerts oncogenic properties in glioblastoma by epigenetically silencing TFPI2 through EZH2 and LSD1. *Aging (Albany NY)* 11: 3811-3823, 2019.
104. Li W, Sun M, Zang C, Ma P, He J, Zhang M, Huang Z, Ding Y and Shu Y: Upregulated long non-coding RNA AGAP2-AS1 represses LATS2 and KLF2 expression through interacting with EZH2 and LSD1 in non-small-cell lung cancer cells. *Cell Death Dis* 7: e2225, 2016.
105. Qi F, Liu X, Wu H, Yu X, Wei C, Huang X, Ji G, Nie F and Wang K: Long noncoding AGAP2-AS1 is activated by SPI and promotes cell proliferation and invasion in gastric cancer. *J Hematol Oncol* 10: 48, 2017.
106. Liu S, Zheng Y, Zhang Y, Zhang J, Xie F, Guo S, Gu J, Yang J, Zheng P, Lai J, *et al*: Methylation-mediated LINC00261 suppresses pancreatic cancer progression by epigenetically inhibiting c-Myc transcription. *Theranostics* 10: 10634-10651, 2020.
107. Salerno D, Chiodo L, Alfano V, Floriot O, Cottone G, Paturel A, Pallocca M, Plissonnier ML, Jeddari S, Belloni L, *et al*: Hepatitis B protein HBx binds the DLEU2 lncRNA to sustain cccDNA and host cancer-related gene transcription. *Gut* 69: 2016-2024, 2020.
108. Hota SK and Bruneau BG: ATP-dependent chromatin remodeling during mammalian development. *Development* 143: 2882-2897, 2016.
109. Wang Y, Zhu P, Luo J, Wang J, Liu Z, Wu W, Du Y, Ye B, Wang D, He L, *et al*: lncRNA HAND2-AS1 promotes liver cancer stem cell self-renewal via BMP signaling. *EMBO J* 38: e101110, 2019.
110. Tang Y, Wang J, Lian Y, Fan C, Zhang P, Wu Y, Li X, Xiong F, Li X, Li G, *et al*: Linking long non-coding RNAs and SWI/SNF complexes to chromatin remodeling in cancer. *Mol Cancer* 16: 42, 2017.
111. Ma X and Kang S: Functional implications of DNA methylation in adipose biology. *Diabetes* 68: 871-878, 2019.
112. Nishiyama A and Nakanishi M: Navigating the DNA methylation landscape of cancer. *Trends Genet* 37: 1012-1027, 2021.
113. Schübeler D: ESCI award lecture: Regulation, function and biomarker potential of DNA methylation. *Eur J Clin Invest* 45: 288-293, 2015.
114. Nan X, Ng HH, Johnson CA, Laherty CD, Turner BM, Eisenman RN and Bird A: Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393: 386-389, 1998.
115. Zhang Y, Yan H, Jiang Y, Chen T, Ma Z, Li F, Lin M, Xu Y, Zhang X, Zhang J and He H: Long non-coding RNA IGF2-AS represses breast cancer tumorigenesis by epigenetically regulating IGF2. *Exp Biol Med (Maywood)* 246: 371-379, 2021.
116. Ma F, Lei YY, Ding MG, Luo LH, Xie YC and Liu XL: lncRNA NEAT1 interacted with DNMT1 to regulate malignant phenotype of cancer cell and cytotoxic T cell infiltration via epigenetic inhibition of p53, cGAS, and STING in lung cancer. *Front Genet* 11: 250, 2020.
117. Tang J, Xie Y, Xu X, Yin Y, Jiang R, Deng L, Tan Z, Gangarapu V, Tang J and Sun B: Bidirectional transcription of linc00441 and RB1 via H3K27 modification-dependent way promotes hepatocellular carcinoma. *Cell Death Dis* 8: e2675, 2017.
118. Feng H and Liu X: Interaction between ACOT7 and lncRNA NMRAL2P via methylation regulates gastric cancer progression. *Yonsei Med J* 61: 471-481, 2020.
119. Li N, Zhao Z, Miao F, Cai S, Liu P, Yu Y and Wang B: Silencing of long non-coding RNA LINC01270 inhibits esophageal cancer progression and enhances chemosensitivity to 5-fluorouracil by mediating GSTP1 methylation. *Cancer Gene Ther* 28: 471-485, 2021.
120. Zhang C, Wang L, Jin C, Zhou J, Peng C, Wang Y, Xu Z, Zhang D, Huang Y, Zhang Y, *et al*: Long non-coding RNA lnc-LALC facilitates colorectal cancer liver metastasis via epigenetically silencing LZTS1. *Cell Death Dis* 12: 224, 2021.
121. O'Leary VB, Ovsepian SV, Smida J and Atkinson MJ: PARTICLE-the RNA podium for genomic silencers. *J Cell Physiol* 234: 19464-19470, 2019.
122. O'Leary VB, Hain S, Maugg D, Smida J, Azimzadeh O, Tapio S, Ovsepian SV and Atkinson MJ: Long non-coding RNA PARTICLE bridges histone and DNA methylation. *Sci Rep* 7: 1790, 2017.
123. Arab K, Park YJ, Lindroth AM, Schäfer A, Oakes C, Weichenhan D, Lukanova A, Lundin E, Risch A, Meister M, *et al*: Long noncoding RNA TARID directs demethylation and activation of the tumor suppressor TCF21 via GADD45A. *Mol Cell* 55: 604-614, 2014.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.